Research Article

Analgesic and Anti-Inflammatory Activities of Methanol Extract of *Ficus pumila* L. in Mice

Chi-Ren Liao,¹ Chun-Pin Kao,² Wen-Huang Peng,¹ Yuan-Shiun Chang,¹ Shang-Chih Lai,³ and Yu-Ling Ho⁴

¹School of Chinese Pharmaceutical Sciences and Chinese Medicine Resources, College of Pharmacy, China Medical University, Taichung 404, Taiwan

²Department of Nursing, Hsin Sheng College of Medical Care and Management, Taoyuan 325, Taiwan

³Department of Health and Nutrition Biotechnology, Asia University, Taichung 413, Taiwan

⁴Department of Nursing, Hungkuang University, Sha Lu, Taichung 433, Taiwan

Correspondence should be addressed to Yu-Ling Ho, elaine@sunrise.hk.edu.tw

Received 15 November 2011; Revised 29 February 2012; Accepted 29 February 2012

Academic Editor: William C. S. Cho

Copyright © 2012 Chi-Ren Liao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This study investigated possible analgesic and anti-inflammatory mechanisms of the methanol extract of *Ficus pumila* (FP_{MeOH}). Analgesic effects were evaluated in two models including acetic acid-induced writhing response and formalin-induced paw licking. The results showed FP_{MeOH} decreased writhing response in the acetic acid assay and licking time in the formalin test. The anti-inflammatory effect was evaluated by λ -carrageenan-induced mouse paw edema and histopathological analyses. FP_{MeOH} significantly decreased the volume of paw edema induced by λ -carrageenan. Histopathologically, FP_{MeOH} abated the level of tissue destruction and swelling of the edema paws. This study indicated anti-inflammatory mechanism of FP_{MeOH} may be due to declined levels of NO and MDA in the edema paw through increasing the activities of SOD, GPx, and GRd in the liver. Additionally, FP_{MeOH} also decreased the level of inflammatory mediators such as IL-1 β , TNF- α , and COX-2. HPLC fingerprint was established and the contents of three active ingredients, rutin, luteolin, and apigenin, were quantitatively determined. This study provided evidence for the classical treatment of *Ficus pumila* in inflammatory diseases.

1. Introduction

Inflammatory reaction, typically characterized by redness, swelling, heat, and pain, is one of the most important host defense mechanisms against invading pathogens. However, persistent or overinflammation leads to tissue damage and possibly failure of organs. Proinflammatory cytokines (e.g., TNF- α , IL-1 β , and IL-6) are produced in large quantities by activated macrophages/monocytes that stimulate cellular responses via increasing prostaglandins (PGs) and reactive oxygen species (ROS). Additionally, lipid peroxidation (malondialdehyde, MDA) is produced by free radicals attacking the cell membranes. Thus, inflammatory effect can lead to the accumulation of MDA [1].

Ficus pumila, a creeping vine like fig plant, is native to South China and Malaysia. Several studies have been performed on the composition of *Ficus pumila*, and a number of compounds have been identified such as apigenin, luteolin

[2], rutin, genistein, hesperidin, astragaloside, isoquercitrin, and chrysin [3]. Dried stems and leaves of *Ficus pumila* have been folklorically used in the treatment of rheumatoid arthritis, edema, tonic medicament, throat pain, and postpartum abdominal pain [4]. However, no research has been investigated on the analgesic and anti-inflammatory mechanisms of *Ficus pumila* yet.

In this study, we investigated the analgesic and anti-inflammatory activities of the methanol extract of *Ficus pumila* (FP_{MeOH}). The analgesic activity was evaluated by acetic acid-induced writhing response and formalin test. Anti-inflammatory activity was determined by using λ -carrageenan-induced mouse paw edema model and histopathological analysis. In order to evaluate the mechanism of anti-inflammatory effect, we also analyzed TNF- α , IL-1 β , COX-2, MDA, and NO levels in the edema tissue, as well as antioxidant enzyme activities of SOD (superoxidase dismutase),

GPx (glutathione peroxidase), and GRd (glutathione reductase) in the liver.

Many studies have indicated that flavonoids in herbs possess anti-inflammatory activities via scavenging ROS and reducing proinflammatory cytokines, such as rutin [5], luteolin [6], and apigenin [7]. These three ingredients have also been isolated from *Ficus pumila* in previous studies [2, 3]. In the phytochemical part of this study, not only did we reconfirm the presence of these three compounds in FP_{MeOH} by establishing its fingerprint chromatogram, but also the contents of these active ingredients were quantitatively determined.

2. Materials and Methods

2.1. Chemicals and Drugs. λ -carrageenan, indomethacin, and Griess reagent were purchased from Uni-Onward (distributor of Sigma-Aldrich Chemical Co. in Taipei, Taiwan). Formalin was purchased from Nihon Shiyaku Industry Ltd. (Taipei, Taiwan). SOD, GPx, GRd, and MDA assay kits were purchased from Eugene Chen Co. Ltd. (distributor of Randox Laboratory Ltd. in Taipei, Taiwan). IL-1 β , IL-6, COX-2, and TNF- α were obtained from Blossom Biotechnologies, Inc. (agency of Assay Designs Inc. in Taipei, Taiwan). LC grade acetonitrile was purchased from the branch company of Merck in Taipei, Taiwan. All other reagents used were of analytical grade.

2.2. Plant Material. *Ficus pumila* L. was collected from the herbal garden of China Medical University, Taichung, Taiwan, as described in Flora of Taiwan [8]. A plant specimen has been deposited in the School of Chinese Pharmaceutical Sciences and Chinese Medicine Resources. Dried stems and leaves (1.0 kg) of *Ficus pumila* were sliced into small pieces, soaked with 10 liters of methanol at room temperature for three days. After passing through filter paper, the filtrate was concentrated under reduced pressure to dryness. The above steps were repeated two additional times, and a total of 123.1 g of dry crude extract (yield ratio 12.31%) was produced. The dried crude extract was dissolved in 0.5% CMC (carboxymethyl cellulose) solution into three concentrations (100 mg/mL, 50 mg/mL, and 10 mg/mL) prior to pharmacological testing. Equal volume of 0.1 mL (extract of different concentrations)/10 g B.W. was given to each mouse in all subsequent animal experiments.

2.3. Chromatographic Analysis of FP_{MeOH} . The HPLC system consisted of a Waters 2695 Alliance LC with 996 PDA. Chromatographic separation was performed on X-Bridge RP18 (25 cm $\times$ 4.6 mm I.D., 5 μ m) with an injection volume of 10 μ L. The mobile phase consisted of a mixture of 0.1% formic acid (A) and acetonitrile (B) using a gradient elution. The gradient program was set as follows: 0–15 min, 20% B, 15–30 min, 40% B, 30–45 min, 60% B. The flow rate was set at 0.8 mL/min and the detection wavelength was set to 255 nm. The above conditions were used in both HPLC assay and HPLC fingerprint of FP_{MeOH} .

The contents of rutin, apigenin, and luteolin of FP_{MeOH} were qualified and quantified in the HPLC assay. In qualitative analyses, comparisons were made with the retention time

and maximum absorption of the standards. In quantitative analyses, comparisons were made with peak areas under the standard curves.

2.4. Experimental Animals. Male ICR mice aged between 5–6 weeks and weighing between 20–25 g were obtained from BioLASCO Taiwan Co. Ltd. and were kept in the animal center of China Medical University at a controlled temperature of $22 \pm 1^\circ\text{C}$, relative humidity $55 \pm 5\%$, and with 12 h light/12 h dark cycles for 1 week before the experiment. Animals were provided with rodent diet and clean water *ad libitum*. The experimental protocol was approved by the Committee on Animal Research, China Medical University. Animals were sacrificed by decapitation under ether anesthesia. All studies were conducted in accordance with the National Institutes of Health (NIH) Guide for the Care and Use of Laboratory Animals. All tests were conducted under the guidelines of the International Association for the Study of Pain [9].

2.5. Acute Toxicity Study. The acute toxicology test in mice was carried out according to the method of Liao et al. [10]. Male ICR mice were randomly divided into three groups (10 mice per group). Three groups of mice were administered orally with three concentrations of FP_{MeOH} (2.5 g, 5 g, and 10 g/kg), respectively. The experimental mice were kept under regular observation for 14 days for any mortality or behavioral changes.

2.6. Acetic Acid-Induced Writhing Response. The writhing test in mice was carried out according to the method of Koster et al. [11]. Five randomly-selected groups of mice were orally administered with solvent control (0.5% CMC), positive control (indomethacin at 10 mg/kg) or three different doses of FP_{MeOH} (0.1, 0.5, and 1.0 g/kg) 60 min prior to the chemical stimulus. The number of muscular contractions was counted 5 min after the injection of 1% acetic acid (v/v, 0.1 mL/10 g body weight, i.p.). The data collected represented the total number of writhes observed in a duration of 10 minutes (5–15 min after the injection).

2.7. Formalin Test. The formalin test was conducted based on the method of Tjølsen et al. [12]. Twenty microliters of 5% formalin in saline was injected subcutaneously into the right hind paw of each mouse. The time (in seconds) spent on licking and biting of the injected paw was recorded in both the early phase (0–5 min) and late phase (20–30 min) after the formalin injection. Solvent control (0.5% CMC), positive control (indomethacin at 10 mg/kg), or three different doses of FP_{MeOH} (0.1, 0.5 and 1.0 g/kg) was orally administered to the animals 60 min prior to the formalin injection.

2.8. λ -Carrageenan-Induced Mouse Paw Edema. The test was conducted according to the method of Vinegar et al. [13]. The basal volume of right hind paw was determined before the administration of any drug. Fifty microliters of 1% λ -carrageenan suspended in saline was injected into the plantar side of right hind paw, and the paw volume was measured at the 1st, 2nd, 3rd, 4th, and 5th h after the injection using a

plethysmometer. The degree of swelling was evaluated by the delta volume (a-b), where “a” is the volume of right hind paw after the chemical treatment and “b” is the volume before the treatment. CMC (0.5%), indomethacin (10 mg/kg), and FP_{MeOH} (0.1, 0.5 and 1.0 g/kg) were administered orally 60 min after the λ -carrageenan injection. In the secondary experiment, another set of mice were orally administered with 0.5% CMC, indomethacin, or FP_{MeOH} 1 h after λ -carrageenan had been injected into their right hindpaws. The right hindpaws of the animals were surgically removed under anesthesia 2 h following treatments. The paw tissue was rinsed in ice-cold normal saline and immediately placed in cold normal saline four times its volume before homogenization at 4°C. The homogenate was centrifuged at 12,000 rpm for 5 min. The supernatant was obtained and stored at -20°C for the determinations of MDA, NO, TNF- α , IL-1 β , and COX-2. Similarly, the whole liver tissue was rinsed in ice-cold normal saline and immediately placed in cold normal saline of equal volume before homogenization at 4°C. The homogenate was then centrifuged at 12,000 rpm for 5 min. The supernatant was obtained and stored at -20°C for later analyses of antioxidant enzyme (SOD, GPx, and GRd) activities.

2.9. Histological Analysis. The mice were orally administered with 0.5% CMC, indomethacin, or FP_{MeOH} 1 h after λ -carrageenan had been injected into their right hindpaws. The right hindpaws of the animals were surgically removed under anesthesia 2 h after treatments. Tissue slices were fixed in 10% formalin for 3 days, decalcified overnight and embedded in paraffin and sectioned into 4 μm tissue sections. Tissue sections were stained with hematoxylin and eosin (H&E stain) before being examined under a BX60 microscope (Olympus, Melville, NY) for pathological changes. Inflammatory reactions induced by λ -carrageenan, including paw swelling and enlarged cavities, were examined. The severity after FP_{MeOH} (0.1, 0.5 and 1.0 g/kg) and indomethacin (10 mg/kg) treatments was also examined. Images were captured with a Macrofire 599831 camera. The results were identified in the Animal Disease Diagnostic Center (ADDC), National Chung Hsing University, Taichung, Taiwan.

2.10. MDA Assay. The production of MDA was induced by λ -carrageenan injection, and evaluated by the thiobarbituric acid reacting substance (TBARS) method [14]. Briefly, MDA reacted with TBARS at high temperature and formed a red-complex TBARS. The absorbance of TBARS was recorded at 532 nm.

2.11. NO Assay. NO was measured according to the method of Moshage et al. [15]. NO_3^- was converted into NO_2^- by nitrate reductase, NO_2^- subsequently reacted with sulfanilic acid to produce diazonium ion and coupled with N-(1-naphthyl) ethylenediamine to form the chromophoric aeroderivative (purplish red) which could be recorded at 540 nm.

2.12. TNF- α and IL-1 β Assay. IL-1 β was measured by an enzyme-linked immunosorbent assay kit (Blossom Biotechnologies Inc.) [16]. The capture antibody of IL-1 β was added

to each well of a 96-well plate overnight. Next day, a second set of biotinylated antibody was incubated with sample tissues or standard antigens in the plate before streptavidin was added. The color of the reaction converted from purple to yellow and was recorded at 450 nm. TNF- α was detected using the same method as IL-1 β . Each sample was presented as pg/mg in TNF- α and IL-1 β concentrations.

2.13. COX-2 Assay. The content of COX-2 was determined by measuring the peroxidase activity of PGHS (prostaglandin endoperoxide H_2 synthase) [17]. Peroxidase activity of PGHS was determined by following the oxidation of N,N,N',N'-tetramethyl-p-phenylenediamine (TMPD) at 37°C using arachidonate as the substrate. The increase in color was recorded at 590 nm.

2.14. Measurement of Antioxidant Enzymes. SOD was measured according to the method of Vani et al. [18]. Xanthine and xanthine oxidase (XOD) generated superoxide radicals reacted with 2-(4-iodophenyl)-3-(4-nitrophenol)-5-phenyl-tetrazolium chloride (I.N.T.) to form a red formazan dye, and the color was recorded at 540 nm. GPx was measured according to the method of Ceballos-Picot et al. by detecting the contents of GR and NADPH [19]. Oxidation of NADPH into NADP^+ is accompanied by a decrease in absorbance recorded at 340 nm. GRd was measured according to the method of Ahmad and Holdsworth [20] which detects the decrease of glutathione (GSSG) in the presence of NADPH. NADPH oxidized into NADP^+ would result in a decrease in absorbance recorded at 340 nm.

2.15. Statistical Analysis. All data of each group were expressed as mean $\pm$ SD ($n = 8$). Statistical analyses were performed with SPSS software and were carried out using one-way ANOVA followed by Scheffe's multiple range test.

3. Results

3.1. Chromatographic Analysis of FP_{MeOH} . HPLC fingerprint profile was established for FP_{MeOH} (Figure 1). Three flavonoid components were identified as rutin, luteolin, and apigenin with retention times of 13.8 min, 35.2 min, and 40.4 min, respectively. The maximum absorbance was 255 nm, and the relative amounts for each gram of crude extract were in the order of rutin (24.41 mg), apigenin (14.11 mg), and luteolin (2.72 mg).

3.2. Acute Toxicity Study. Acute toxicity of FP_{MeOH} was evaluated in mice at the doses of 2.5, 5 and 10 g/kg. After 14 days of oral administration, FP_{MeOH} did not cause any behavioral changes, and no mortality was observed. Therefore, the LD_{50} value of FP_{MeOH} was concluded to be greater than 10 g/kg in mice, indicating it was practically not acutely toxic.

3.3. Acetic Acid-Induced Writhing Response. Figure 2 shows acetic acid-induced writhing responses in mice which serve as an indication of analgesic activities of FP_{MeOH} . Intraperitoneal injection of acetic acid produced 42.2 ± 5.6 writhes

TABLE 1: Effect of FP_{MeOH} and indomethacin on liver SOD, GPx, and GRd activities in mice injected with λ -carrageenan.

Groups	SOD (U/g protein)	GPx (U/mg protein)	GRd (U/mg protein)
Normal	121.45 $\pm$ 12.53	1.579 $\pm$ 0.136	0.106 $\pm$ 0.012
Car	67.69 $\pm$ 16.43	0.949 $\pm$ 0.144	0.063 $\pm$ 0.014
Car + Indomethacin	99.72 $\pm$ 10.36**	1.222 $\pm$ 0.109	0.085 $\pm$ 0.006**
Car + FP _{MeOH} (0.1 g/kg)	84.94 $\pm$ 13.82*	1.287 $\pm$ 0.133*	0.082 $\pm$ 0.008**
Car + FP _{MeOH} (0.5 g/kg)	93.89 $\pm$ 16.68*	1.332 $\pm$ 0.207**	0.085 $\pm$ 0.007**
Car + FP _{MeOH} (1.0 g/kg)	95.48 $\pm$ 17.87*	1.339 $\pm$ 0.224**	0.078 $\pm$ 0.006*

Each value represents the mean $\pm$ SD ($n = 8$). * $P < 0.05$ and ** $P < 0.01$ as compared with the λ -carrageenan (Car) group (one-way ANOVA followed by Scheffe's multiple range test).

TABLE 2: Effects of FP_{MeOH} and indomethacin on the severity of mouse paw edema.

Groups	Grade (mean $\pm$ SD)
Car	3.6 $\pm$ 0.55
Car + Indomethacin	2.0 $\pm$ 0.71*
Car + FP _{MeOH} (0.1 g/kg)	3.6 $\pm$ 0.55
Car + FP _{MeOH} (0.5 g/kg)	3.0 $\pm$ 0.00
Car + FP _{MeOH} (1.0 g/kg)	2.2 $\pm$ 0.45*

Each value represents the mean $\pm$ SD ($n = 5$). * $P < 0.05$ as compared with the λ -carrageenan (Car) group (one-way ANOVA followed by Scheffe's multiple range test).

FIGURE 1: HPLC fingerprint of FP_{MeOH}. The peaks represent 1. rutin (13.8 min), 2. luteolin (35.2 min), and 3. apigenin (40.4 min). The contents of rutin, apigenin, and luteolin in FP_{MeOH} were 24.41 mg/g, 14.11 mg/g, and 2.72 mg/g, respectively.

in the solvent control group. The writhing response was significantly reduced by pretreatments with FP_{MeOH} (0.1, 0.5 and 1 g/kg) and indomethacin (10 mg/kg).

3.4. Formalin Test. In the early phase, FP_{MeOH}- and indomethacin-treated groups did not show any significant changes as compared to the solvent control group (Figure 3(a)). In the second phase, subcutaneous injection of formalin induced licking and biting responses in the solvent control group lasted for a period of 155.7 ± 10.5 seconds. The time was significantly decreased by pretreatment with FP_{MeOH} (0.1, 0.5 and 1 g/kg) and indomethacin (10 mg/kg) (Figure 3(b)).

3.5. Effect of FP_{MeOH} on λ -Carrageenan-Induced Mouse Paw Edema. Following the λ -carrageenan injection, the volume of mouse paw increased as edema developed,

FIGURE 2: Analgesic effects of FP_{MeOH} and indomethacin on acetic acid-induced writhing response in mice. Each value represents mean $\pm$ SD ($n = 8$). The number of abdominal writhes was counted over the time period of 5–15 min after acetic acid injection. ** $P < 0.01$ and *** $P < 0.001$ as compared to the solvent control group (one-way ANOVA followed by Scheffe's multiple range test).

indicating inflammatory activities (Figure 4). However, indomethacin (10 mg/kg) and FP_{MeOH} (0.1–1.0 g/kg) significantly decreased paw edema at the 3rd, 4th, and 5th h after the injection. FP_{MeOH} at the concentrations of 0.5 and 1 g/kg had about equal amount of inhibition as indomethacin.

3.6. Histological Analysis. No inflammation, tissue destruction, and swelling phenomenon were observed in the paws of normal mice (Figure 5(a)). On the other hand, the λ -carrageenan control group displayed enlarged cavities in the paw tissue (Figure 5(b)). Edematous condition was obviously abated by treatment with 10 mg/kg of indomethacin and 1.0 g/kg of FP_{MeOH} (Figures 5(c) and 5(f)). The severity of mouse paw edema was also graded and summarized in Table 2 (mean $\pm$ SD).

3.7. Effect of FP_{MeOH} on MDA Level. MDA level obviously increased in the λ -carrageenan control group ($1.120 \pm 0.172 \mu\text{M/g}$); however, pretreatments with FP_{MeOH} (0.1, 0.5 and 1.0 g/kg) and indomethacin (10 mg/kg) significantly inhibited the increase of MDA levels (Figure 6).

FIGURE 3: Effects of FP_{MeOH} and indomethacin on (a) early and (b) late phases of formalin test in mice. Each value represents mean $\pm$ SD ($n = 8$). The time spent on licking and biting the injected paws was recorded in the time periods of 0–5 min (early phase) and 20–30 min (late phase) as indicators of pain. * $P < 0.05$ and *** $P < 0.001$ as compared to the solvent control group (one-way ANOVA followed by Scheffe's multiple range test).

3.8. Effect of FP_{MeOH} on NO Level. Pretreatments with FP_{MeOH} (0.5 and 1.0 g/kg) and indomethacin (10 mg/kg) significantly inhibited the increase of NO levels in the edema paws of mice, as compared to the λ -carrageenan control group ($10.08 \pm 1.45 \mu\text{M}$), as shown in Figure 7.

3.9. Effect of FP_{MeOH} on TNF- α and IL-1 β . TNF- α and IL-1 β levels in λ -carrageenan-induced edema paws were increased remarkably. FP_{MeOH} (0.5 and 1.0 g/kg) and indomethacin (10 mg/kg) significantly reduced the levels of TNF- α (Figure 8). Similarly, IL-1 β levels were significantly lowered by FP_{MeOH} (0.1, 0.5 and 1.0 g/kg) and indomethacin, as shown in Figure 9.

3.10. Effect of FP_{MeOH} on COX-2 Level. Figure 10 shows that COX-2 level was greatly raised ($37.04 \pm 3.04 \text{ U/mg}$) in λ -carrageenan induced edema paw. However, COX-2 levels

FIGURE 4: Effects of FP_{MeOH} and indomethacin on λ -carrageenan-induced mouse paw edema. The volumes of paw edema were detected at the 1st, 2nd, 3rd, 4th, and 5th h after λ -carrageenan injection. Each value represents mean $\pm$ SD ($n = 8$). ** $P < 0.01$ and *** $P < 0.001$ as compared to the λ -carrageenan control group (one-way ANOVA followed by Scheffe's multiple range test).

were decreased by treating with FP_{MeOH} (0.5 and 1.0 g/kg) as well as indomethacin (10 mg/kg).

3.11. Effect of FP_{MeOH} on the Activities of Antioxidant Enzymes. SOD, GPx, and GRd activities were increased by treating with FP_{MeOH} (0.5 and 1.0 g/kg) and indomethacin (10 mg/kg), as compared to the λ -carrageenan control group (Table 1).

4. Discussion

Ficus is a genus of about 800 species found in tropical and subtropical regions. Several *Ficus* species have been studied for their anti-inflammatory actions, for example, *F. aurantiacea* [20], *F. carica* [21], *F. glomerata* [22], *F. maxima* [23], and *F. obtusifolia* [24].

Traditional medicines have been popularly used in the treatment of various diseases in recent years. Many medicinal plants supply analgesic and anti-inflammatory activities to treat acute, chronic, or recurring illnesses. *Ficus pumila* is a Chinese herbal medicine, its dried stems and leaves have been commonly used in the treatment of rheumatoid arthritis, edema, throat pain, and postpartum abdominal pain [4]. Based on its therapeutic claims in traditional medicine, we investigated possible mechanisms of the analgesic and anti-inflammatory effects of FP_{MeOH}.

Analgesic effect of FP_{MeOH} was evaluated by two animal models, including acetic acid-induced writhing response and formalin test. The results of acetic acid induced writhing test showed that FP_{MeOH} (0.1–1 g/kg) and indomethacin (10 mg/kg) provided antinociceptive effects in mice. Acetic acid indirectly triggers the release of nociceptive endogenous mediators (such as bradykinin, serotonin, and

FIGURE 5: Histopathological examinations on λ -carrageenan-induced paw tissue swelling, edema and neutrophil infiltration: (a) normal, (b) λ -carrageenan, (c) indomethacin, (d) FP_{MeOH} (0.1 g/kg), (e) FP_{MeOH} (0.5 g/kg), and (f) FP_{MeOH} (1.0 g/kg).

FIGURE 6: Effects of FP_{MeOH} and indomethacin on MDA concentrations in the edema paws. Each value represents mean $\pm$ SD ($n = 8$). *** $P < 0.001$ as compared to the λ -carrageenan control group (one-way ANOVA followed by Scheffe's multiple range test).

FIGURE 7: Effects of FP_{MeOH} and indomethacin on NO concentrations in the mouse edema paws. Each value represents mean $\pm$ SD ($n = 8$). ** $P < 0.01$ and *** $P < 0.001$ as compared to the λ -carrageenan control group (one-way ANOVA followed by Scheffe's multiple range test).

prostaglandin) and proinflammatory cytokines (such as $\text{TNF-}\alpha$ and $\text{IL-1}\beta$) to cause painful sensation [25]. This nociceptive effect can be prevented by analgesic agents with central actions such as morphine as well as peripherally acting drugs like NSAID.

In order to evaluate whether FP_{MeOH} acted centrally or peripherally in the suppression of pain, we also conducted

the formalin test. The formalin test involves a biphasic response: the first phase (neurogenic nociceptive response) occurs in the first 5 min after the formalin injection, while the second phase (inflammatory nociceptive response) occurs between 15 to 30 min after formalin injection. Centrally

FIGURE 8: Effects of FP_{MeOH} and indomethacin on TNF- α concentrations in mouse paw edema. Each value represents mean $\pm$ SD ($n = 8$). ** $P < 0.01$ and *** $P < 0.001$ as compared to the λ -carrageenan control group (one-way ANOVA followed by Scheffe's multiple range test).

acting drugs can inhibit both phases; however, peripherally acting drugs, such as NSAID, only inhibit the second phase [26]. The treatments of FP_{MeOH} (0.1–1 g/kg) and indomethacin (10 mg/kg) were able to diminish the nociceptive response in the second phase induced by formalin. The results indicated that the antinociceptive effect of FP_{MeOH} was due to its peripheral anti-inflammatory effect.

In the inflammatory model involving λ -carrageenan injection, FP_{MeOH} significantly decreased the volume of paw edema. The anti-inflammatory effect was also evident when we compared histopathological examinations of FP_{MeOH}-treated groups with that of λ -carrageenan control group. Paw biopsy of the λ -carrageenan control group showed obvious edematous condition and enlarged cavities in the connective tissues; on the other hand, the groups that received 10 mg/kg of indomethacin or 1 g/kg of FP_{MeOH} had significant improvement in edematous condition and decreased intercellular spaces in connective tissues as shown in Figure 5 and Table 2.

In our follow-up experiments that explored the mechanisms underlying the anti-inflammatory effect of FP_{MeOH}, we suspected that suppression of inflammatory cytokines may play an important role. λ -Carrageenan-induced paw edema has also been characterized as a biphasic event [13]: histamine, bradykinin, and 5-hydroxytryptamine (5-HT) are released in the first phase of edema (0–1 h), while TNF- α , IL-1 β , COX-2, and PGs are produced more actively in the second phase (1–6 h). TNF- α and IL-1 β have been reported in several studies to be able to recruit leukocytes, such as neutrophils [27, 28]. Additionally, COX-2 is an enzyme responsible for increasing prostaglandin levels in inflammatory reactions, which could in turn worsen the degree of swelling [29]. In this study, FP_{MeOH} (dose dependent)

FIGURE 9: Effects of FP_{MeOH} and indomethacin on IL-1 β concentrations in mouse paw edema. Each value represents mean $\pm$ SD ($n = 8$). ** $P < 0.01$ and *** $P < 0.001$ as compared to the λ -carrageenan control group (one-way ANOVA followed by Scheffe's multiple range test).

FIGURE 10: Effects of FP_{MeOH} and indomethacin on COX-2 concentrations in mouse paw edema. Each value represented mean $\pm$ SD ($n = 8$). * $P < 0.05$ and *** $P < 0.001$ as compared to the λ -carrageenan control group (one-way ANOVA followed by Scheffe's multiple range test).

and indomethacin (10 mg/kg) showed significant anti-inflammatory effect in λ -carrageenan-induced mouse paw edema from the 3rd to 5th h. Moreover, the levels of TNF- α , IL-1 β , and COX-2 were also decreased by treating with FP_{MeOH} and indomethacin (10 mg/kg). Thus, a putative anti-inflammatory mechanism of FP_{MeOH} could be associated with the inhibition of inflammatory mediators, such as TNF- α , IL-1 β , and COX-2.

Another possible mechanism behind the anti-inflammatory effect of FP_{MeOH} could be due to its antioxidative activity. Increased production of $\text{TNF-}\alpha$, $\text{IL-1}\beta$, and other proinflammatory cytokines not only has the capacity to recruit leukocytes, but would also trigger their release of active substances such as ROS [30]. Nitric oxide, hydrogen peroxide, hydroxyl radicals, and superoxide anions play major roles in terms of producing cellular damage [31]. MDA is a reactive aldehyde caused by toxic stress in cells and form covalent protein adducts which are referred to as advanced lipid peroxidation end products (ALE). Inflammation would result in the accumulation of MDA; on the other hand, enhancing the level of GPx, GRd and SOD could reduce MDA production [32]. In this study, SOD, GPx, and GRd activities increased in the liver after treatment with FP_{MeOH} . Conversely, the level of MDA decreased significantly. Therefore, we believe that the suppression of MDA production was likely associated to the increase in SOD, GPx, and GRd activities.

The fingerprint chromatogram of FP_{MeOH} was established, and its three flavonoid contents (rutin, luteolin and apigenin) were quantitatively determined. Three active compounds of FP_{MeOH} , rutin, luteolin, and apigenin, have been previously demonstrated to possess anti-inflammatory and antinociceptive activities which may be responsible, at least in part, to the anti-inflammatory and antinociceptive effects of FP_{MeOH} [5–7].

This study demonstrated that *Ficus pumila* exhibited antinociceptive and anti-inflammatory activities. The anti-inflammatory mechanisms of FP_{MeOH} against λ -carrageenan-induced paw edema involved two possible pathways. The first pathway alleviated the levels of inflammatory factors, such as $\text{IL-1}\beta$, $\text{TNF-}\alpha$, and COX-2 in the edema paw induced by λ -carrageenan. The other pathway was likely associated to the decrease in MDA and NO levels in the edema paw via increasing the activities of SOD, GPx, and GRd in the liver. In conclusion, this study supported possible mechanisms of FP_{MeOH} used for mitigating inflammatory pain.

Acknowledgments

The authors would like to thank Dr. Jiunn-Wang Liao from National Chung Hsing University (Taichung, Taiwan) for the histological analysis of tissue biopsy. In addition, we would also like to express our appreciation to Dr. Chao-Lin Chen, former professor in the University of Florida, for his critical reading of the paper and editorial help.

References

- [1] D. R. Janero, "Malondialdehyde and thiobarbituric acid-reactivity as diagnostic indices of lipid peroxidation and peroxidative tissue injury," *Free Radical Biology and Medicine*, vol. 9, no. 6, pp. 515–540, 1990.
- [2] J. B. Harborne, *The Flavonoids: Advances in Research Since 1980*, Chapman and Hall, London, UK, 1986.
- [3] L. Pistelli, E. E. Chiellini, and I. Morelli, "Flavonoids from *Ficus pumila*," *Biochemical Systematics and Ecology*, vol. 28, no. 3, pp. 287–289, 2000.
- [4] Editorial Board of Zhong Hua Ben Cao, *Zhong Hua Ben Cao (China Herbal)*, Shanghai Scientific Technology Press, Shanghai, China, 1999.
- [5] L. Selloum, H. Bouriche, C. Tigrine, and C. Boudoukha, "Anti-inflammatory effect of rutin on rat paw oedema, and on neutrophils chemotaxis and degranulation," *Experimental and Toxicologic Pathology*, vol. 54, no. 4, pp. 313–318, 2003.
- [6] S. Hougee, A. Sanders, J. Faber et al., "Decreased pro-inflammatory cytokine production by LPS-stimulated PBMC upon in vitro incubation with the flavonoids apigenin, luteolin or chrysin, due to selective elimination of monocytes/macrophages," *Biochemical Pharmacology*, vol. 69, no. 2, pp. 241–248, 2005.
- [7] G. S. Jeong, S. H. Lee, S. N. Jeong, Y. C. Kim, and E. C. Kim, "Anti-inflammatory effects of apigenin on nicotine- and lipopolysaccharide-stimulated human periodontal ligament cells via heme oxygenase-1," *International Immunopharmacology*, vol. 9, no. 12, pp. 1374–1380, 2009.
- [8] T. S. Huang, *Flora of Taiwan*, Tah Jinn Printing Company Limited, Taipei, Taiwan, 1998.
- [9] M. Zimmermann, "Ethical guidelines for investigations of experimental pain in conscious animals," *Pain*, vol. 16, no. 2, pp. 109–110, 1983.
- [10] J. W. Liao, Y. C. Chung, J. Y. Yeh et al., "Safety evaluation of water extracts of *Toona sinensis* Roemor leaf," *Food and Chemical Toxicology*, vol. 45, no. 8, pp. 1393–1399, 2007.
- [11] R. Koster, M. Anderson, and E. J. Beer, "Acetic acid for analgesic screening," *Federation Proceeds*, vol. 18, pp. 412–416, 1959.
- [12] A. Tjølsen, O. G. Berge, S. Hunskaar, J. H. Rosland, and K. Hole, "The formalin test: an evaluation of the method," *Pain*, vol. 51, no. 1, pp. 5–17, 1992.
- [13] R. Vinegar, W. Schreiber, and R. Hugo, "Biphasic development of carrageenin edema in rats," *Journal of Pharmacology and Experimental Therapeutics*, vol. 166, no. 1, pp. 96–103, 1969.
- [14] V. L. Tatum, C. Changchit, and C. K. Chow, "Measurement of malondialdehyde by high performance liquid chromatography with fluorescence detection," *Lipids*, vol. 25, no. 4, pp. 226–229, 1990.
- [15] H. Moshage, B. Kok, J. R. Huizenga, and P. L. M. Jansen, "Nitrite and nitrate determinations in plasma: a critical evaluation," *Clinical Chemistry*, vol. 41, no. 6, pp. 892–896, 1995.
- [16] T. Ataoglu, M. Üngör, B. Serpek, S. Halilolu, H. Ataoglu, and H. Ari, "Interleukin- 1β and tumour necrosis factor- α levels in periapical exudates," *International Endodontic Journal*, vol. 35, no. 2, pp. 181–185, 2002.
- [17] N. Petrovic and M. Murray, "Using N,N,N',N'-tetramethyl-p-phenylenediamine (TMPD) to assay cyclooxygenase activity in vitro," *Methods in Molecular Biology*, vol. 594, pp. 129–140, 2010.
- [18] M. Vani, G. P. Reddy, G. R. Reddy, K. Thyagaraju, and P. Reddanna, "Glutathione-S-transferase, superoxide dismutase, xanthine oxidase, catalase, glutathione peroxidase and lipid peroxidation in the liver of exercised rats," *Biochemistry International*, vol. 21, no. 1, pp. 17–26, 1990.
- [19] I. Ceballos-Picot, J. M. Trivier, A. Nicole, P. M. Sinet, and M. Thevenin, "Age-correlated modifications of copper-zinc superoxide dismutase and glutathione-related enzyme activities in human erythrocytes," *Clinical Chemistry*, vol. 38, no. 1, pp. 66–70, 1992.
- [20] F. B. Ahmad and D. K. Holdsworth, "Medicinal plants of Sabah, Malaysia, Part II. The muruts," *International Journal of Pharmacognosy*, vol. 32, no. 4, pp. 378–383, 1994.

- [21] E. Sezik, E. Yesilada, M. Tabata et al., "Traditional medicine in Turkey VIII. Folk medicine in East Anatolia; Erzurum, Erzincan, Agri, Kars, Igdir provinces," *Economic Botany*, vol. 51, no. 3, pp. 195–211, 1997.
- [22] V. K. Singh, Z. A. Ali, S. T. H. Zaidi, and M. K. Siddiqui, "Ethnomedicinal uses of plants from Gonda district forests of Uttar Pradesh, India," *Fitoterapia*, vol. 67, no. 2, pp. 129–139, 1996.
- [23] D. L. Lentz, A. M. Clark, C. D. Hufford et al., "Antimicrobial properties of Honduran medicinal plants," *Journal of Ethnopharmacology*, vol. 63, no. 3, pp. 253–263, 1998.
- [24] X. A. Dominguez and J. B. Alcorn, "Screening of medicinal plants used by Huastec Mayans of North Eastern Mexico," *Journal of Ethnopharmacology*, vol. 13, no. 2, pp. 139–156, 1985.
- [25] M. N. Manjavachi, N. L. M. Quintao, M. M. Campos et al., "The effects of the selective and non-peptide CXCR2 receptor antagonist SB225002 on acute and long-lasting models of nociception in mice," *European Journal of Pain*, vol. 14, no. 1, pp. 23–31, 2010.
- [26] A. J. Reeve and A. H. Dickenson, "The roles of spinal adenosine receptors in the control of acute and more persistent nociceptive responses of dorsal horn neurones in the anaesthetized rat," *British Journal of Pharmacology*, vol. 116, no. 4, pp. 2221–2228, 1995.
- [27] D. Salvemini, Z. Q. Wang, P. S. Wyatt et al., "Nitric oxide: a key mediator in the early and late phase of carrageenan-induced rat paw inflammation," *British Journal of Pharmacology*, vol. 118, no. 4, pp. 829–838, 1996.
- [28] L. C. Loram, A. Fuller, L. G. Fick, T. Cartmell, S. Poole, and D. Mitchell, "Cytokine profiles during carrageenan-induced inflammatory hyperalgesia in rat muscle and hind paw," *Journal of Pain*, vol. 8, no. 2, pp. 127–136, 2007.
- [29] K. Seibert, Y. Zhang, K. Leahy et al., "Pharmacological and biochemical demonstration of the role of cyclooxygenase 2 in inflammation and pain," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 91, no. 25, pp. 12013–12017, 1994.
- [30] A. G. Semb, J. Vaage, and O. D. Mjos, "Oxygen free radical producing leukocytes cause functional depression of isolated rat hearts: role of leukotrienes," *Journal of Molecular and Cellular Cardiology*, vol. 22, no. 5, pp. 555–563, 1990.
- [31] A. Kumaran and R. J. Karunakaran, "Activity-guided isolation and identification of free radical-scavenging components from an aqueous extract of *Coleus aromaticus*," *Food Chemistry*, vol. 100, no. 1, pp. 356–361, 2007.
- [32] S. Cuzzocrea, G. Costantino, B. Zingarelli, E. Mazzon, A. Micali, and A. P. Caputi, "The protective role of endogenous glutathione in carrageenan induced pleurisy in the rat," *European Journal of Pharmacology*, vol. 372, no. 2, pp. 187–197, 1999.