Research Article

Green Tea Attenuates Oxidative Stress and Downregulates the Expression of Angiotensin II AT₁ Receptor in Renal and Hepatic Tissues of Streptozotocin-Induced Diabetic Rats

Martha Thomson, Khaled Al-Qattan, Mohamed H. Mansour, and Muslim Ali

Department of Biological Sciences, Faculty of Science, Kuwait University, P.O. Box 5969, Safat 13060, Kuwait

Correspondence should be addressed to Mohamed H. Mansour, mheshamshaker@yahoo.com

Received 13 August 2012; Accepted 13 October 2012

Academic Editor: Khalid Rahman

Copyright © 2012 Martha Thomson et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This study investigates the potential of green tea to modulate oxidative stress and angiotensin II AT₁ receptor expression in renal and hepatic tissues of diabetic rats. Three groups of rats were studied after 8 weeks following diabetes induction: normal, streptozotocin-induced diabetic (diabetic control), and green-tea-treated diabetic rats. Total antioxidant, catalase, and malondialdehyde levels were assayed by standard procedures. Levels of AT₁ receptor labeling, in renal and hepatic tissues of the three rat groups, were immunohistochemically investigated using an anti-AT₁ receptor antibody. Levels of total antioxidant and catalase were significantly reduced, whereas malondialdehyde levels and AT₁ receptor labeling were significantly increased in renal and hepatic tissues of diabetic control rats compared to normal rats. Compared to diabetic control rats, total antioxidant and catalase levels were significantly increased, whereas malondialdehyde levels and AT₁ receptor labeling in the green-tea-treated diabetic group were significantly reduced throughout hepatic lobules and renal cortical and medullary vascular and tubular segments to levels comparable to those observed in normal rats. The capacity of green tea to modulate diabetes-induced oxidative stress and AT₁ receptor upregulation may be beneficial in opposing the deleterious effects of excessive angiotensin II signaling, manifested by progressive renal and hepatic tissue damage.

1. Introduction

Brewing dried leaves and buds of the plant *Camellia sinensis* produces tea; one of the most commonly consumed beverages worldwide [1]. Green tea, about 20% of the tea produced annually, is predominantly consumed in Asian countries but has become increasingly popular in Western countries during the last decade [2]. All types of tea beverages contain polyphenols specifically flavonoids. Green tea contains mainly catechins, members of the flavan-3-ol class of flavonoids. The four major catechins in green tea are (–)-epicatechin (EC), (–)-epicatechin gallate (ECG), (–)-epigallocatechin (EGC), and (–)-epigallocatechin gallate (EGCG). A typical green tea serving contains about 90–100 mg of catechins of which EGCG accounts for 50–80% [3].

Many medicinal properties have been ascribed to green tea including anticancer, antidiabetic, and antihypertensive effects [4, 5]. Many of these beneficial effects have been attributed to the presence of catechins, which are potent antioxidants. Since oxidative stress has also been strongly correlated to the increased incidence of diseases such as cancer, cardiovascular diseases, and even in the aging process, green tea has been studied as a potential ameliorative agent in these conditions. Several studies have demonstrated that green tea components scavenge reactive oxygen species formed *in vitro* [6]. In addition, animal studies have suggested that green tea might protect against the development of coronary heart disease by reducing blood glucose levels and body weight [7]. Catechins also reduced plasma triglyceride levels in an oral glucose tolerance test in normal rats [5]. Several human and animal studies suggested

that green tea and its flavonoids have antidiabetic effects [8, 9].

Diabetes mellitus is the most common metabolic disorder that causes nearly 7% of all worldwide deaths annually [10]. Chronic hyperglycemia in diabetes is associated with long-term damage, dysfunction, and eventually the failure of organs, especially eyes, kidneys, nerves, and the cardiovascular system [11–13]. Dyslipidemia is also involved in the development of cardiovascular complications in diabetes, which are the major causes of morbidity and mortality [14, 15]. Experimental and clinical studies have shown that diabetic hyperglycemia is a major source of oxidative stress, and that increased generation of free radicals plays a major role in the pathogenesis and complications of diabetes [16–19]. Glucose autooxidation, formation of advanced glycation end products, and increased activity of the sorbitol pathway have been suggested as the basic mechanism by which chronic hyperglycemia promotes increased oxidative stress [18]. Oxidative stress in diabetes leads to tissue damage, with lipid peroxidation, inactivation of proteins, and protein glycation as intermediate mechanisms for its complications. Damage to small blood vessels (microvascular disease) can lead to the development of diabetic nephropathy which is characterized by progressive loss of kidney function, as a result of reduction in glomerular filtering capacity leading to proteinuria and albuminuria. Between 25 and 40% of diabetic patients have nephropathy, and 30–40% of newly diagnosed cases of end-stage renal disease requiring dialysis are characterized as having increased oxidative stress [20, 21] attributed to diabetes [22, 23]. As a prevalent and serious complication of diabetes, prevention or reversal of diabetic nephropathy would improve the prognosis of diabetic patients.

As manifested in diabetic patients and animal models, the renin-angiotensin system has been shown to progressively augment proteinuria and accelerate decline in renal function [24]. Through its multiple interactions with its specific AT₁ receptor, angiotensin II (Ang II), the major active product of this system synthesized both systemically and within the kidney, is a mediator of progressive injury in diabetic nephropathy. As previously documented, the hyperglycemia-induced AT₁ receptor upregulation in the adrenal and kidney mediates the increase of Ang II local effects in stimulating aldosterone production and over-activating renal sodium transporters causing sodium retention and hypertension [25]. In the diabetic liver, Ang II induces hepatic inflammation and stimulates a range of fibrogenic actions, including cell migration, cell proliferation, the production of intracellular reactive oxygen species, secretion of proinflammatory cytokines, and collagen synthesis predominantly through the AT₁ receptor [26]. Within the kidney, the upregulated AT₁ receptor signaling increases vascular resistance, glomerular capillary pressure, and mechanical stretch-induced glomerular injury and stimulates production of reactive oxygen species and extracellular matrix in the mesangium and tubulointerstitium [25, 27], which collectively perpetrates the development of diabetic nephropathy. Currently, there is accumulating evidence that pharmacological interference with the chronic activation of the renin-angiotensin system

represents an important therapeutic target, which may provide incremental end-organ protection. In particular, the use of AT₁ receptor blockers, solely or among other regimens, may prove beneficial in opposing the detrimental effects of AT₁ receptors in the progression of diabetic nephropathy as well as other diabetic complications [28].

The present study was designed to study the beneficial effects of green tea in the streptozotocin- (STZ-) induced diabetic rat model. In particular, the current study focused on the antioxidant potential and putative modulatory effect of green tea on AT₁ receptor expression in ameliorating diabetic complications in STZ-induced diabetic rats.

2. Materials and Methods

2.1. Animal Model and Experimental Design. Male Sprague-Dawley rats weighing 150–180 g were used for the study. The study conformed to the National Research Council (NRC) Guide for the Care and Use of Laboratory Animal [29]. All animals were maintained on a 12 h light/dark cycle and a temperature of $22 \pm 1^\circ\text{C}$. All animals received diet and water *ad libitum*. Thirty rats were injected with streptozotocin (STZ). The remaining rats were used as normal controls. For induction of diabetes, rats were weighed, fasted overnight, and injected intraperitoneally (IP) with a single dose of 60 mg STZ/kg body weight in a constant volume of 0.5 mL citrate buffer. The rats were divided into 3 groups ($n = 8$) and were treated for 8 weeks. Group 1 was normal control rats drinking only water (normal control), group 2 was diabetic rats drinking only water (diabetic control), and group 3 was diabetic rats drinking 0.1% green tea (diabetic + green tea). A drop of blood was obtained from each rat by tail-vein puncture before STZ injection and weekly after STZ injection to determine blood glucose levels. The blood glucose level was measured using a one touch UltraEasy glucometer (LifeScan, Johnson & Johnson, UK), and rats were determined to be diabetic if they had elevated plasma glucose concentrations ≥ 300 mg/dL five days postinjection, as described previously [30].

2.2. Antibodies and Preparation of Green Tea Extract. Polyclonal anti-AT₁ receptor antibody (sc-1173, rabbit IgG specific to an epitope mapping within the N-terminal extracellular domain of the human AT₁ polypeptide) and peroxidase-conjugated goat anti-rabbit IgG antibody were purchased from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA, USA). An eight-amino-acid peptide, corresponding to amino acids 14–21 (Ile-Gln-Asp-Asp-Cys-Pro-Lys-Ala) of the first extracellular domain, as deduced from the published AT₁ receptor cDNA sequence [31], was synthesized manually on the base-labile linker 4-(hydroxymethyl)-benzoyloxymethyl and supplied by The Protein/DNA Technology Center (The Rockefeller University, NY, USA). The octapeptide was conjugated to bovine serum albumin (BSA) and coupled to CNBr-activated Sepharose 4B as described previously [32].

A 0.1% green tea extract (Al-Wazah Tea, Ceylon) was prepared by weighing 1 gram of green tea leaves and suspending

them in 1000 mL of hot boiled distilled water. The extract was then stirred for 15 min and filtered using cheesecloth. The tea extract was prepared daily. The water content of the control group bottles was also changed daily. After 8 weeks, rats were sacrificed under sodium pentobarbitone anesthesia (10 mg/kg, Sagatal, May and Baker UK), blood was collected via cardiac puncture, and serum was obtained and stored in small aliquots at -80°C for further analysis. The liver and kidneys of each rat were individually collected, weighed, and either placed in vials containing 3 mL of Bouin's fixative for 24–48 h at room temperature or placed in separate small plastic bags and stored at -40°C for further analysis.

2.3. Preparation of Tissue Homogenates. One gram of the kidney or liver excised from normal, control diabetic, and/or green-tea-treated diabetic rats was cut into small pieces and mixed with 3 mL Tris HCl (0.05 M, pH = 7.6) buffer. The mixture was homogenized, allowed to stand on ice for few minutes, and then centrifuged for 15 min at $8000 \times g$ at 4°C . The supernatant stored in small aliquots at -40°C for further analysis.

2.4. Biochemical Assays. Protein concentration of serum and tissue homogenates was determined by the Coomassie blue dye-binding method using bovine serum albumin as standard [33]. Total serum or tissue antioxidant levels were determined as described in Drobiova et al. [30]. Tissue malonaldehyde (MDA) levels were determined according to the method of Ohkawa and coworkers [34]. Catalase activity was quantified by the method of Abei [35].

2.5. Preparation of Tissue Sections. Fixed kidneys and Livers of normal, control diabetic, and/or green-tea-treated diabetic rats were individually carried through a routine paraffin embedding technique, which included dehydration through a series of ethanol concentrations 50%, 70%, 90%, and 100%, clearing in toluene, embedding in paraffin wax, and finally sectioning of the paraffin blocks into 3–4 μm thick sections on a rotary microtome. The sections were picked up on clean slides after spreading them in a water bath at 40°C . The slides were air-dried to be used for subsequent staining.

2.6. Labeling of Tissue Sections with Anti-AT₁ Receptor Antibody. Tissue sections were examined for AT₁ receptor distribution by an indirect immunohistochemical labeling technique. Tissue sections were dewaxed in xylene, hydrated with a series of 90%, 75%, and 60% ethanol, and washed with PBS, pH 7.2. Sections were individually labeled for 45 min in a humidified chamber with 300 μL of rabbit anti-AT₁ receptor antibody (diluted to 1 : 100 with PBS, pH 7.2). After three washes with 200 μL PBS, pH 7.2, the sections were incubated for 45 min with 300 μL of peroxidase-conjugated goat anti-rabbit IgG antibody (diluted to 1 : 200 in PBS, pH 7.2) then washed three times with 200 μL PBS, pH 7.2, followed by a 5 min treatment with 400 μL of 3,3'-diaminobenzidine tablets (fast DAB) reconstituted in water. All sections were counterstained with 100 μL haematoxylin (Gill number1) for 1 min and examined by light microscopy

for positive labeling of cells expressing AT₁ receptors. Control sections were identically stained by replacing the specific antibody with anti-AT₁ antibody preabsorbed with an AT₁ octapeptide/BSA complex-coated CNBr-activated Sepharose 4B beads. To exclude nonspecific binding, all sections were treated with 300 μL of 0.3% hydrogen peroxide in PBS, pH 7.2 for 10 min to block endogenous peroxidase, followed by a 30-min block with 300 μL of 5% normal goat serum in PBS, pH 7.2, prior to the application of the anti-AT₁ antibody, to minimize off-target staining. Tissue sections prepared from normal control, diabetic control, and green-tea-treated diabetic rats were processed in parallel with strictly uniform incubation times and reagent volumes to ensure standardization of labeling between samples. Slides were quantitatively examined by using the Image-Pro Plus 5.1 software program (Media Cybernetics, Silver Spring, MD). Automatic object counting and measuring processes were used to quantify labeled areas, and values expressed as μm^2 . Each slide was analyzed for AT₁ receptor labeling in the liver and in the renal cortex and outer medulla, with three separate fields viewed in each region and four independent samples for each group. Data were expressed and analyzed as indicated below. Representative microphotographs were taken using Olympus AH-3 automated microscope (Tokyo, Japan), equipped with an Olympus Vanox camera.

2.7. Statistical Analysis. The data are expressed as mean $\pm$ standard error of the mean (SEM). The test of analysis of variance (ANOVA) with post hoc STD was employed to examine differences among readings between groups. The $P < 0.05$ level was selected as the level of significance. All statistical analyses were performed using SPSS (version 19).

3. Results

3.1. Effect of Drinking Green Tea on Blood Glucose Level and Body Weight. Figure 1 shows the effect of drinking green tea extract on blood glucose level. Blood glucose levels in diabetic control rats were significantly increased compared to normal control rats during the entire experimental period. In comparison, the green tea drinking rats showed significantly decreased blood glucose compared to diabetic control rats by 5 and 8-weeks. In comparison, Figure 2 depicts body weight changes during the 8 week experiment. Figures 1 and 2 show that as blood glucose levels increased in diabetic control rats, body weight decreased compared to normal control rats that nearly doubled their weight during the 8-week experiment (Figure 2). In comparison, green-tea-treated diabetic rats have lowered blood glucose levels and increased body weight compared to the diabetic controls.

3.2. Effect of Drinking Green Tea on Kidney and Liver Antioxidants, MDA, and Catalase. Figure 3 shows the effect of drinking green tea extract on the kidney total antioxidants and MDA levels. Kidney tissue exhibited a simultaneous significant decrease in total antioxidants and increase in MDA in diabetic control rats drinking water compared to normal control rats. The diabetic rats drinking green tea

FIGURE 1: Effect of green tea on blood glucose in diabetic rats. Blood glucose levels were measured at weeks 1, 5, and 8 of the experiment in normal control rats, STZ-induced diabetic rats (diabetic control), and 0.1% green-tea-treated STZ-induced diabetic rats (diabetic + green tea). Results are expressed as means $\pm$ SEM, $n = 8$. ^aSignificantly different from normal control. ^bSignificantly different from diabetic control. ^cNo significant difference from diabetic control.

showed a significant increase in total antioxidant levels and decrease in MDA when compared to diabetic control rats.

Liver MDA levels were significantly higher (Figure 4) as a result of STZ induction of diabetes (diabetic controls). In contrast, diabetic rats that drank green tea had liver MDA levels significantly lower than diabetic control rats. In contrast, liver total antioxidant levels were significantly decreased in diabetic control rats (Figure 4) and significantly elevated in diabetic rats drinking green tea.

Table 1 shows the effect of drinking green tea on kidney and liver catalase activity. Kidneys and livers of diabetic control rats exhibited a significant decrease in catalase activity compared to normal control rats. However, treating diabetic rats with green tea extract partially reversed this with a significant increase in both kidney and liver catalase observed in green tea drinking rats compared to diabetic control rats.

3.3. Effect of Drinking Green Tea on AT₁ Receptor Expression in the Kidney and Liver. In immunohistochemical labeling experiments of kidney sections of the three rat groups, light microscopy revealed that variable patterns of AT₁ receptor labeling were primarily associated with the cortex as well as the inner stripe of the outer medulla. As depicted in Figure 5(a), cortical nephron segments of normal control rats exhibited AT₁ receptor labeling of marked intensity in the outer rim of Bowman's capsule and the entire epithelial lining of the distal convoluted tubules, but of moderate intensity among glomeruli and uniformly diffused in the epithelial lining of the proximal convoluted tubules. In diabetic control rats, AT₁ receptor labeling was maintained

FIGURE 2: Effect of green tea on change in body weight in diabetic rats. Weights were taken for normal control rats, STZ-induced diabetic rats (diabetic control), and 0.1% green-tea-treated STZ-induced diabetic rats (diabetic + green tea). The animals were weighed before STZ injection (Before STZ), 1 week after STZ injection (after STZ), and at weeks 4 and 8 of the experiment. Weights are plotted as percentiles with the starting, weights all standardized to 100%. ^aSignificantly different from normal control. ^bSignificantly different from diabetic control. ^cNo significant difference from diabetic control.

at a high intensity in the epithelial lining of the distal convoluted tubules, but was markedly reduced in Bowman's capsule and glomeruli as opposed to its enhanced intensity in the epithelial lining of the proximal convoluted tubules, particularly at the apical side (Figure 5(b)). On the other hand, except for the persistent high intensity of the AT₁ receptor labeling confined to the epithelial lining of the distal convoluted tubules, the specific labeling was almost lacking in Bowman's capsule and glomeruli and appeared as faded patches at the basolateral side of some of the epithelial cells lining the proximal convoluted tubules in green-tea-treated diabetic rats (Figure 5(c)).

As in the cortical regions, distinct labeling patterns of the AT₁ receptor were also observed in the inner stripe of the outer medulla of the three rat groups. In normal control rats (Figure 6(a)), binding of the anti-AT₁ receptor antibody was evident in interstitial cells outlining the vasa recta bundles as well as the basolateral side of collecting tubules and, to a lesser extent, the apical side of collecting ducts, without any obvious involvement of the epithelial elements of ascending or descending Henle's loop segments. Compared to normal control rats, AT₁ receptor labeling was of relatively lower intensity in interstitial cells outlining the vasa recta bundles in diabetic control rats, but was accentuated in the cytoplasm of the entire epithelial lining of collecting tubules, collecting ducts, and Henle's loop segments (Figure 6(b)). As shown in Figure 6(c), AT₁ receptor labeling in interstitial cells outlining the vasa recta bundles was of significantly reduced intensity and was apparently faded and confined to the basolateral side of collecting tubules, collecting ducts, and Henle's loop segments in green-tea-treated diabetic rats.

FIGURE 3: Effect of green tea on kidney MDA and antioxidants in diabetic rats. Kidney MDA and antioxidant levels were measured at the end of week 8 of the experiment in normal control rats, STZ-induced diabetic rats (diabetic control), and 0.1% green-tea-treated STZ-induced diabetic rats (diabetic + green tea). Results are expressed as means $\pm$ SEM, $n = 8$. ^aSignificantly different from normal control. ^bSignificantly different from diabetic control.

None of these specific cortical and medullary labeling patterns were observed in renal sections treated with anti-AT₁ antibody preabsorbed with AT₁ octapeptide/BSA complex-coated CNBr-activated Sepharose 4B beads (Figures 5(d) and 6(d)). Quantification of AT₁ receptor staining by computer-based image analysis revealed a significant increase in expression of AT₁ receptors in the renal cortical and outer medullary regions of the control diabetic group compared with the normal control rats. Green-tea-treated diabetic rats were marked by a return of AT₁ receptor labeling to levels significantly lower than the control diabetic group and comparable to those observed in normal control rats (Table 2).

In liver sections, moderate AT₁ receptor labeling was primarily observed among hepatocytes arranged around the central vein within hepatic lobules, but was less evident in the bile duct and vessels within portal tracts in normal control as well as green-tea-treated diabetic rats (Figures 7(a) and 7(c)). In contrast, a more intense AT₁ receptor labeling, involving more hepatocytes in addition to bile ducts and vessels in portal tracts, was observed in control diabetic rats (Figure 7(b)). None of these specific labeling patterns were observed with liver sections treated with anti-AT₁ antibody preabsorbed with AT₁ octapeptide/BSA complex-coated CNBr-activated Sepharose 4B beads (Figure 7(d)). As depicted in Table 2, quantification of AT₁ receptor staining by computer-based image analysis revealed a significant increase in the expression of AT₁ receptors in the liver of control diabetic rats compared with either the normal control group or the green-tea-treated diabetic group.

FIGURE 4: Effect of green tea on liver MDA and antioxidants in diabetic rats. Liver MDA and antioxidant levels were measured at the end of week 8 of the experiment in normal control rats, STZ-induced diabetic rats (diabetic control), and 0.1% green-tea-treated STZ-induced diabetic rats (diabetic + green tea). Results are expressed as means $\pm$ SEM, $n = 8$. ^aSignificantly different from normal control. ^bSignificantly different from diabetic control.

4. Discussion

The present study was conducted to evaluate the role of green tea as a natural agent of an ameliorative potential to oxidative stress and progressive tissue damage in STZ-induced diabetic rats. Experimental diabetes induced by STZ is a well-established method used to evaluate the mechanisms involved in the alterations of physiopathology observed in diabetic patients. STZ destroys pancreatic β cells, resulting in a diabetic syndrome, similar to that seen in human type 1 diabetes and characterized by hyperglycemia, hypoinsulinemia, glucosuria, and loss in body weight [36–38]. In this study as in previous studies, the observed hyperglycemia in STZ-induced diabetic rats was paralleled by 40% reduction in body weight, whereas green tea extract treatments resulted in a significant reduction in blood glucose levels and prevention of weight loss in these animals. These observations are in accordance with the documented role of green tea in the regulation of body weight [38] and furthermore confirm the hypoglycemic effects of green tea reported in several other studies [7, 39, 40]. Green tea polyphenols have been shown to improve glucose tolerance in normal and obese rats and db/db mice [2, 41]. In addition, green tea has been reported to enhance basal and insulin-stimulated glucose uptake in rat adipocytes [42]. The major green tea catechin, EGCG, was shown to inhibit intestinal glucose uptake by sodium-dependent glucose transporter SGLT1 [43], whereas a catechin-rich green tea extract and EGCG have been reported to mimic insulin by decreasing the expression of genes that control gluconeogenesis [44]. Collectively, the antihyperglycemic effect of green tea observed in these

FIGURE 5: Immunohistochemical localization of AT₁ receptor expression in the renal cortex of normal control (a), diabetic control (b), and green-tea-treated diabetic (c) rats. Renal sections were treated with the polyclonal anti-AT₁ receptor antibody and counterstained with hematoxylin as detailed in Materials and Methods. In (a), positive labeling was detected with marked intensity in Bowman's capsule and distal convoluted tubules epithelia, but was of moderate intensity among glomeruli and uniformly diffused in proximal convoluted tubules epithelia. In (b), the labeling was maintained at a high intensity in the epithelial lining of the distal convoluted tubules, but was markedly reduced in Bowman's capsule and glomeruli as opposed to its enhanced intensity in the apical side of the proximal convoluted tubules epithelia. In (c), the labeling was almost lacking in Bowman's capsule and glomeruli and appeared as faded patches at the basolateral side of some of the proximal convoluted tubules epithelia, but was of persistent high intensity confined to the distal convoluted tubules epithelia. No specific labeling was observed in renal sections labeled with the anti-AT₁ antibody preabsorbed with an AT₁ octapeptide/BSA complex (d). Images are representative of four different samples for each rat group. DCT: distal convoluted tubule, G: glomerulus, and PCT: proximal convoluted tubule. Bar = 100 µm, $\times 100$.

TABLE 1: Effect of green tea on kidney and liver catalase in diabetic rats.

Group ($n = 8$)	Mean $\pm$ SEM (units/mg Protein)	
	Kidney	Liver
Normal control rats	0.317 ± 0.023	0.929 ± 0.09
Diabetic control rats	0.162 ± 0.038^a	0.493 ± 0.042^a
Diabetic + green tea rats	$0.252 \pm 0.038^{a,b}$	$0.575 \pm 0.070^{a,b}$

^aSignificantly different from normal control rats.

^bSignificantly different from diabetic control rats.

TABLE 2: AT₁ receptor quantification.

	Normal control rats	Diabetic control rats	Diabetic + green tea rats
Kidney			
Cortex	$6,421 \pm 541$	$10,825 \pm 447^*$	$5,101 \pm 375$
Outer medulla	$6,567 \pm 677$	$11,074 \pm 305^*$	$6,212 \pm 715$
Liver	$2,514 \pm 240$	$5,815 \pm 635^*$	$2,014 \pm 670$

Values are means $\pm$ SEM ($n = 4$). Each value represents the number of pixels that exceeded an arbitrary staining threshold per 300×300 pixel area.

* $P < 0.05$ versus green tea-treated diabetic rats and normal rats.

studies suggests an important clinical relevance to diabetes treatment.

Increased oxidative stress can be the result of several diabetes-induced abnormalities, including autooxidation of glucose, the formation of advanced glycation end products, and impairment of the scavenging system [18]. The impaired scavenging system is always associated with reduced antioxidant capacity, abnormal activity or reduced expression

of antioxidant enzymes in diabetes [19]. Among various antioxidant enzymes, catalase plays an important role in balancing the generation of reactive oxygen species and the overall tissue antioxidant capacity. In our study, the total antioxidant activity and catalase activity in the kidney and liver tissues were significantly compromised in diabetic rats compared to normal controls. This was paralleled by an

FIGURE 6: Immunohistochemical localization of AT₁ receptor expression in the outer medulla of the kidney of normal control (a), diabetic control (b), and green-tea-treated diabetic (c) rats. Renal sections were treated with the polyclonal anti-AT₁ receptor antibody and counterstained with hematoxylin as detailed in Materials and Methods. In (a), the labeling was evident in vasa recta bundles and, to a lesser extent, the basolateral side of collecting tubules and the apical side of collecting ducts, whereas Henle's loop segments were not labeled. In (b), the labeling was of relatively low intensity in the vasa recta bundles but was evident in the epithelial lining of collecting tubules and ducts as well as Henle's loop segments. In (c), labeling with faded intensity was observed in vasa recta bundles, and confined to the basolateral side of collecting tubules, collecting ducts, and Henle's loop epithelia. No specific labeling was observed in kidney tissue sections labeled with the anti-AT₁ antibody preabsorbed with an AT₁ octapeptide/BSA complex (d). Images are representative of four different samples for each rat group. A: ascending limb of Henle's loop, CD: collecting duct, CT: collecting tubule, D: descending limb of Henle's loop, and VR: vasa recta bundle. Bar = 100 μ m, $\times 100$.

increase in lipid peroxidation as evidenced by the elevated levels of MDA detected in kidney and liver tissues in diabetic rats compared to normal controls. Treatments of diabetic rats with green tea significantly increased both antioxidant levels and catalase activity, which was reflected by significantly lowering MDA levels in kidney and liver tissues compared to diabetic control rats. These observations are consistent with several studies showing the potential of green tea and/or catechins in reducing oxidative stress by either increasing total plasma antioxidant activity [45–47] or reducing MDA levels in tissues [48]. However, Hininger-Favier and coworkers [47] did not observe any change in liver MDA levels after treatment of insulin-resistant rats with green tea extract. Similarly, Yokazawa and coworkers reported no significant change in kidney MDA levels after green tea treatment of diabetic rats [49]. However, these conflicting findings might be due to differences in doses, concentrations of green tea, time of exposure, and type of green tea as well as the degree of hyperglycemia and the detection methods employed.

In diabetic patients and animal models, the altered expression of the Ang II AT₁ receptor has been implicated as a pivotal factor in the development of early changes associated with diabetic end-organ damages, particularly nephropathy [24–27]. In the present study, AT₁ receptor expression in the kidney and liver tissues was investigated immunohistochemically using a polyclonal anti-AT₁ antibody of proven specificity to an epitope mapping within the N-terminal extracellular domain of the AT₁ polypeptide [50]. In normal rats, the binding of this antibody was predominately observed among hepatocytes arranged around the central vein within hepatic lobules, in discrete kidney regions, including the glomeruli and proximal and distal convoluted tubules in the cortex, in interstitial cells outlining the vasa recta bundles, and to a lesser extent, collecting tubules and collecting ducts in the inner stripe of the outer medulla. These hepatic and renal sites corresponded well to those shown to express detectable levels of the AT₁ receptor by *in vitro* autoradiographic localization, *in situ* hybridization, and semiquantitative reverse transcription

FIGURE 7: Immunohistochemical localization of AT₁ receptor expression in the liver of normal control (a), diabetic control (b), and green-tea-treated diabetic (c) rats. Liver sections were treated with the polyclonal anti-AT₁ receptor antibody and counterstained with hematoxylin as detailed in Materials and Methods. In both (a) and (c), moderate AT₁ receptor labeling was selectively observed among hepatocytes arranged around the central vein (CV) within hepatic lobules. In (b), AT₁ receptor labeling with higher intensity was observed among hepatocytes as well as bile ducts and vessels in portal tracts (arrowheads). No specific labeling was observed in liver sections labeled with the anti-AT₁ antibody preabsorbed with an AT₁ octapeptide/BSA complex (d). Images are representative of four different samples for each rat group. Bar = 100 μ m, $\times 100$.

polymerase chain reaction [51]. Confirming the documented effect of hyperglycemia on AT₁ receptor expression [24, 27, 50], STZ treatment in the present study was paralleled by alterations in the pattern of AT₁ receptor expression in both renal and hepatic tissues. Compared to normal rats, a significant increase in AT₁ receptor expression was observed among the cytoplasmic and apical aspects of the proximal convoluted tubule epithelia in the renal cortex, the interstitial cells outlining the vasa recta bundles and tubular segments including Henle's loop and collecting tubules and ducts within the inner stripe of the outer renal medulla, and involved more frequent hepatocytes in addition to bile ducts and vessels in portal tracts in the liver of STZ-induced diabetic rats. As previously indicated, the upregulatory shift in AT₁ receptor expression in the renal cortex leads to higher stimulation of Na-K-ATPase and Na/H exchanger 3 in proximal segments leading to sodium retention and subsequent hypertension [52], as well as increased mechanical stretch-induced glomerular injury culminating in the development of glomerulosclerosis and proteinuria [27]. Overstimulated Ang II signaling through upregulated AT₁ receptors in the outer renal medulla and hepatic tissues has been directly implicated in excessive reactive oxygen species production and oxidative stress [26, 27] and in stimulating a range of fibrogenic action, including cell migration, cell hypertrophy

and proliferation, secretion of proinflammatory cytokines, and extracellular matrix production [24, 26, 27, 53]. In both organs, the direct consequences of the AT₁ receptor upregulation involves the over-stimulation of the profibrotic cytokine transforming growth factor- β , which seems to orchestrate the development and progression of diabetes-associated hepatic fibrosis and nephropathy [53].

A major observation of the present investigation was the significant downregulation of the AT₁ receptor expressed in renal and hepatic tissues of green-tea-treated diabetic rats. Compared to the control diabetic rats, the expressed levels of the AT₁ receptor in the green-tea-treated diabetic group were significantly reduced throughout renal cortical and medullary vascular and tubular segments, as well as hepatic lobules and portal tracts, to levels comparable to those observed in normal rats. Recently, the efficacy of green tea catechin treatments in attenuating diabetes-induced build-up of tissue-advanced glycation end products, and the subsequent generation of free radicals through autooxidation of glucose and glycated proteins has been reported [54]. As suggested by the present observations, the capacity of green tea in modulating diabetes-induced AT₁ receptor upregulation may also imply beneficial interference in the deleterious effects of excessive AT₁ receptor signaling, which is central in mediating excessive tissue remodeling

and alterations in renal hemodynamic and tubular functions, manifested by the development of nephropathy and hepatic fibrosis [4–27, 52, 53]. In this regard, our observations may be supported by the documented therapeutic potential of AT₁ receptor blockers not only in the management of progressive diabetic nephropathy [28] but also in alleviating diabetes-associated hepatic fibrosis [55]. Future studies on the significance of constituent green tea polyphenols may clarify the mechanisms that underlie their beneficial capacity in modulating the expression of AT₁ receptor molecules implicated in diabetes-associated disorders.

Acknowledgments

This study was partially supported by the College of Graduate Studies, Kuwait University, for which the authors are grateful. The authors also acknowledge the contribution of Mrs. Warda Assadzadeh and Mrs. Fatma AlTawari in data collection and biochemical analysis. Their thanks are due to Mrs. Divya J. S. for assistance with lab supervision and preparation of figures.

References

- [1] J. H. Weisburger, "Tea and health: a historical perspective," *Cancer Letters*, vol. 114, no. 1-2, pp. 315–317, 1997.
- [2] N. Chen, R. Bezzina, E. Hinch et al., "Green tea, black tea, and epigallocatechin modify body composition, improve glucose tolerance, and differentially alter metabolic gene expression in rats fed a high-fat diet," *Nutrition Research*, vol. 29, no. 11, pp. 784–793, 2009.
- [3] J. M. Hodgson and K. D. Croft, "Tea flavonoids and cardiovascular health," *Molecular Aspects of Medicine*, vol. 31, no. 6, pp. 495–502, 2010.
- [4] Y. Clement, "Can green tea do that? A literature review of the clinical evidence," *Preventive Medicine*, vol. 49, no. 2-3, pp. 83–87, 2009.
- [5] S. M. Chacko, P. T. Thambi, R. Kuttan, and I. Nishigaki, "Beneficial effects of green tea: a literature review," *Chinese Medicine*, vol. 5, article 13, 2010.
- [6] Q. Guo, B. Zhao, M. Li, S. Shen, and X. Wenjuan, "Studies on protective mechanisms of four components of green tea polyphenols against lipid peroxidation in synaptosomes," *Biochimica et Biophysica Acta*, vol. 1304, no. 3, pp. 210–222, 1996.
- [7] H. Tsuneki, M. Ishizuka, M. Terasawa, J. B. Wu, T. Sasaoka, and I. Kimura, "Effect of green tea on blood glucose levels and serum proteomic patterns in diabetic (db/db) mice and on glucose metabolism in healthy humans," *BMC Pharmacology*, vol. 4, article 18, 2004.
- [8] H. Iso, C. Date, K. Wakai et al., "The relationship between green tea and total caffeine intake and risk for self-reported type 2 diabetes among Japanese adults," *Annals of Internal Medicine*, vol. 144, no. 8, pp. 554–562, 2006.
- [9] S. Wolfram, D. Raederstorff, M. Preller et al., "Epigallocatechin gallate supplementation alleviates diabetes in rodents," *Journal of Nutrition*, vol. 136, no. 10, pp. 2512–2518, 2006.
- [10] International Diabetes Federation, *IDF Diabetes Atlas*, International Diabetes Federation, Brussels, Belgium, 5th edition, 2011, <http://www.idf.org/diabetesatlas>.
- [11] D. Aronson, "Hyperglycemia and the pathobiology of diabetic complications," in *Cardiovascular Diabetology: Clinical, Metabolic and Inflammatory Facets*, E. Z. Fisman and A. Tenenbaum, Eds., Advances in Cardiology, Basel, Switzerland, 2008.
- [12] American Diabetes Association, "Diagnosis and classification of diabetes mellitus," *Journal of Diabetes Care*, vol. 33, supplement 1, pp. S62–S69, 2010.
- [13] S. Tesfaye and G. Gill, "Chronic diabetic complications in Africa," *African Journal of Diabetes Medicine*, vol. 9, pp. 4–8, 2011.
- [14] Y. Xu, Z. He, and G. L. King, "Introduction of hyperglycemia and dyslipidemia in the pathogenesis of diabetic vascular complications," *Current Diabetes Reports*, vol. 5, no. 2, pp. 91–97, 2005.
- [15] A. P. Chintan, L. P. Nimish, B. Nayana, M. Bhavna, G. Mahendra, and T. Hardik, "Cardiovascular complication of diabetes mellitus," *Journal of Applied Pharmaceutical Science*, vol. 4, pp. 1–6, 2011.
- [16] A. C. Maritim, R. A. Sanders, and J. B. Watkins, "Diabetes, oxidative stress, and antioxidants: a review," *Journal of Biochemical and Molecular Toxicology*, vol. 17, no. 1, pp. 24–38, 2003.
- [17] R. Ahmed, "The physiological and biochemical effects of diabetes on the balance between oxidative stress and antioxidant defense system," *Medical Journal of Islamic World Academy of Sciences*, vol. 15, pp. 31–42, 2005.
- [18] P. Goycheva, V. Gadjeva, and B. Popov, "Oxidative stress and its complications in diabetes mellitus," *Trakia Journal of Sciences*, vol. 4, pp. 1–8, 2006.
- [19] V. Selvaraju, M. Joshi, S. Suresh, J. A. Sanchez, N. Maulik, and G. Maulik, "Diabetes, oxidative stress, molecular mechanism and cardiovascular disease—an overview," *Toxicology Mechanisms and Methods*, vol. 22, no. 5, pp. 330–335, 2012.
- [20] F. Locatelli, B. Canaud, K. U. Eckardt, P. Stenvinkel, C. Wanner, and C. Zoccali, "Oxidative stress in end-stage renal disease: an emerging treat to patient outcome," *Nephrology Dialysis Transplantation*, vol. 18, no. 7, pp. 1272–1280, 2003.
- [21] E. C. Samouilidou, E. J. Grapsa, I. Kakavas, A. Lagouranis, and B. Agrogiannis, "Oxidative stress markers and C-reactive protein in end-stage renal failure patients on dialysis," *International Urology and Nephrology*, vol. 35, no. 3, pp. 393–397, 2003.
- [22] E. Ritz, "Diabetic nephropathy," *Saudi Journal of Kidney Diseases and Transplantation*, vol. 17, no. 4, pp. 481–490, 2006.
- [23] S. C. W. Tang, "Diabetic nephropathy: a global and growing threat," *Hong Kong Medical Journal*, vol. 16, no. 4, pp. 244–245, 2010.
- [24] K. D. Burns, "Angiotensin II and its receptors in the diabetic kidney," *American Journal of Kidney Diseases*, vol. 36, no. 3, pp. 449–467, 2000.
- [25] C. Xue and H. M. Siragy, "Local renal aldosterone system and its regulation by salt, diabetes, and angiotensin II type 1 receptor," *Hypertension*, vol. 46, no. 3, pp. 584–590, 2005.
- [26] R. Bataller, P. Sancho-Bru, P. Ginès, and D. A. Brenner, "Liver fibrogenesis: a new role for the renin-angiotensin system," *Antioxidants and Redox Signaling*, vol. 7, no. 9-10, pp. 1346–1355, 2005.
- [27] H. M. Siragy, "AT₁ and AT₂ receptor in the kidney: role in health and disease," *Seminars in Nephrology*, vol. 24, no. 2, pp. 93–100, 2004.

- [28] L. Gnudi and D. Goldsmith, "Renin angiotensin aldosterone system (RAAS) inhibitors in the prevention of early renal disease in diabetes," *F1000 Medicine Reports*, vol. 2, no. 1, article 18, 2010.
- [29] National Research Council, Institute for Laboratory Animal Research, and Commission on Life Sciences, *Guide for the Care and Use of Animals*, The National Academies Press, Washington, DC, USA, 1996.
- [30] H. Drobiova, M. Thomson, K. Al-Qattan, R. Peltonen-Shalaby, Z. Al-Amin, and M. Ali, "Garlic increases antioxidant levels in diabetic and hypertensive rats determined by a modified peroxidase method," *Evidence-based Complementary and Alternative Medicine*, vol. 2011, Article ID 703049, 8 pages, 2011.
- [31] T. J. Murphy, R. W. Alexander, K. K. Griendling, M. S. Runge, and K. E. Bernstein, "Isolation of a cDNA encoding the vascular type-1 angiotensin II receptor," *Nature*, vol. 351, no. 6323, pp. 233–236, 1991.
- [32] K. K. Al-Qattan, S. J. Al-Akhawand, and M. H. Mansour, "Immunohistochemical localization of distinct angiotensin II AT₁ receptor isoforms in the kidneys of the Sprague-Dawley rat and the desert rodent Meriones Crassus," *Journal of Veterinary Medicine Series C*, vol. 35, no. 2, pp. 130–138, 2006.
- [33] M. M. Bradford, "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding," *Analytical Biochemistry*, vol. 72, no. 1–2, pp. 248–254, 1976.
- [34] H. Ohkawa, N. Ohishi, and K. Yagi, "Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction," *Analytical Biochemistry*, vol. 95, no. 2, pp. 351–358, 1979.
- [35] H. Abei, *Catalase in the Method of Enzymatic Analysis*, vol. 2, Academic Press, New York, NY, USA, 1974.
- [36] Z. M. Al-Amin, M. Thomson, K. K. Al-Qattan, and M. Ali, "Anti-diabetic and hypolipidaemic properties of ginger (*Zingiber officinale*) in streptozotocin-induced diabetic rats," *British Journal of Nutrition*, vol. 96, no. 4, pp. 660–666, 2006.
- [37] M. Thomson, Z. M. Al-Amin, K. K. Al-Qattan, L. H. Shaban, and M. Ali, "Anti-diabetic and hypolipidaemic properties of garlic (*Allium sativum*) in streptozotocin-induced diabetic rats," *International Journal of Diabetes and Metabolism*, vol. 15, no. 3, pp. 108–115, 2007.
- [38] P. V. Anandh Babu, K. E. Sabitha, and C. S. Shyamaladevi, "Green tea extract impedes dyslipidaemia and development of cardiac dysfunction in streptozotocin-diabetic rats," *Clinical and Experimental Pharmacology and Physiology*, vol. 33, no. 12, pp. 1184–1189, 2006.
- [39] R. A. Anderson and M. M. Polansky, "Tea enhances insulin activity," *Journal of Agricultural and Food Chemistry*, vol. 50, no. 24, pp. 7182–7186, 2002.
- [40] L. Y. Wu, C. C. Juan, L. T. Ho, Y. P. Hsu, and L. S. Hwang, "Effect of green tea supplementation on insulin sensitivity in sprague-dawley rats," *Journal of Agricultural and Food Chemistry*, vol. 52, no. 3, pp. 643–648, 2004.
- [41] H. Ortsater, N. Grankvist, S. Wolfram, N. Kuehn, and A. Sjöholm, "Diet supplementation with green tea extract epigallocatechin gallate prevents progression to glucose intolerance in db/db mice," *Nutrition and Metabolism*, vol. 9, pp. 11–17, 2012.
- [42] L. Y. Wu, C. C. Juan, L. S. Hwang, Y. P. Hsu, P. H. Ho, and L. T. Ho, "Green tea supplementation ameliorates insulin resistance and increases glucose transporter IV content in a fructose-fed rat model," *European Journal of Nutrition*, vol. 43, no. 2, pp. 116–124, 2004.
- [43] Y. Kobayashi, M. Suzuki, H. Satsu et al., "Green tea polyphenols inhibit the sodium-dependent glucose transporter of intestinal epithelial cells by a competitive mechanism," *Journal of Agricultural and Food Chemistry*, vol. 48, no. 11, pp. 5618–5623, 2000.
- [44] K. Yasui, H. Tanabe, N. Okada, R. Fukutomi, Y. Ishigami, and M. Isemura, "Effects of catechin-rich green tea on gene expression of gluconeogenic enzymes in rat hepatoma H4IIE cells," *Biomedical Research*, vol. 31, no. 3, pp. 183–189, 2010.
- [45] T. Yokozawa, T. Nakagawa, and K. Kitani, "Antioxidative activity of green tea polyphenol in cholesterol-fed rats," *Journal of Agricultural and Food Chemistry*, vol. 50, no. 12, pp. 3549–3552, 2002.
- [46] O. O. Ojo, O. Ladeji, and M. S. Nadro, "Studies of the antioxidative effects of green and black tea (*Camellia sinensis*) extracts in rats," *Journal of Medicinal Food*, vol. 10, no. 2, pp. 345–349, 2007.
- [47] I. Hiningier-Favier, R. Benaraba, S. Coves, R. A. Anderson, and A. M. Roussel, "Green tea extract decreases oxidative stress and improves insulin sensitivity in an animal model of insulin resistance, the fructose-fed rat," *Journal of the American College of Nutrition*, vol. 28, no. 4, pp. 355–361, 2009.
- [48] M. Roghani and T. Baluchnejadmojarad, "Hypoglycemic and hypolipidemic effect and antioxidant activity of chronic epigallocatechin-gallate in streptozotocin-diabetic rats," *Pathophysiology*, vol. 17, no. 1, pp. 55–59, 2010.
- [49] T. Yokozawa, T. Nakagawa, T. Oya, T. Okubo, and L. R. Juneja, "Green tea polyphenols and dietary fibre protect against kidney damage in rats with diabetic nephropathy," *Journal of Pharmacy and Pharmacology*, vol. 57, no. 6, pp. 773–780, 2005.
- [50] M. H. Mansour, K. Al-Qattan, M. Thomson, and M. Ali, "Garlic (*Allium sativum*) down-regulates the expression of angiotensin II AT₁ receptor in adrenal and renal tissues of streptozotocin-induced diabetic rats," *Inflammopharmacology*. In press.
- [51] J. Zhuo, C. Maric, P. J. Harris, D. Alcorn, and F. A. O. Mendelsohn, "Localization and functional properties of angiotensin II AT₁ receptors in the kidney: focus on renomedullary interstitial cells," *Hypertension Research*, vol. 20, no. 4, pp. 233–250, 1997.
- [52] A. A. Banday and M. F. Lokhandwala, "Oxidative stress-induced renal angiotensin AT₁ receptor upregulation causes increased stimulation of sodium transporters and hypertension," *American Journal of Physiology*, vol. 295, no. 3, pp. F698–F706, 2008.
- [53] K. Sharma, Y. Jin, J. Guo, and F. N. Ziyadeh, "Neutralization of TGF- β by anti-TGF- β antibody attenuates kidney hypertrophy and the enhanced extracellular matrix gene expression in STZ-induced diabetic mice," *Diabetes*, vol. 45, no. 4, pp. 522–530, 1996.
- [54] A. Elost, T. Ghous, and N. Ahmed, "Natural products as anti-glycation agents: possible therapeutic potential for diabetic complications," *Current Diabetes Reviews*, vol. 8, no. 2, pp. 92–108, 2012.
- [55] J. E. Toblli, M. C. Muñoz, G. Cao, J. Mella, L. Pereyra, and R. Mastai, "ACE inhibition and AT₁ receptor blockade prevent fatty liver and fibrosis in obese Zucker rats," *Obesity*, vol. 16, no. 4, pp. 770–776, 2008.