Research Article

Attenuation of CCl₄-Induced Oxidative Stress and Hepatonephrotoxicity by Saudi Sidr Honey in Rats

Mohammed Al-Yahya,¹ Ramzi Mothana,^{1,2} Mansour Al-Said,^{1,2}
Mohammed Al-Dosari,¹ Nawal Al-Musayeb,¹ Mohammed Al-Sohaibani,³
Mohammad Khalid Parvez,¹ and Syed Rafatullah²

¹ Department of Pharmacognosy, College of Pharmacy, King Saud University, P.O. Box 2457, Riyadh 11451, Saudi Arabia

² Medicinal, Aromatic and Poisonous Plants Research Center (MAPPRC), College of Pharmacy, King Saud University, P.O. Box 2457, Riyadh 11451, Saudi Arabia

³ Department of Pathology (32), King Khalid University Hospital, King Saud University, P.O. Box 2925, Riyadh 11461, Saudi Arabia

Correspondence should be addressed to Ramzi Mothana; r_mothana@yahoo.com

Received 19 December 2012; Accepted 16 January 2013

Academic Editor: Olumayokun A. Olajide

Copyright © 2013 Mohammed Al-Yahya et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The present study was undertaken to investigate the possible protective effect of Saudi Sidr honey (SSH) on carbon tetrachloride (CCl₄) induced oxidative stress and liver and kidney damage in rat. Moreover, the antioxidant activity and the phenolic and flavonoidal contents were determined. The hepatorenal protective activity of the SSH was determined by assessing biochemical, hematological, and histological parameters. Serum transaminases, ALP, GGT, creatinine, bilirubin, urea, uric acid, and MDA level in liver and kidney tissues were significantly elevated, and the antioxidant status of nonprotein sulfhydryls, albumin, and total protein levels in liver and kidney were declined significantly in CCl₄ alone treated animals. Pretreatment with SSH and silymarin prior to the administration of CCl₄ significantly prevented the increase of the serum levels of enzyme markers and reduced oxidative stress. SSH also exhibited a significant lipid-lowering effect and caused an HDL-C enhanced level in serum. The histopathological evaluation of the liver and kidney also revealed that honey protected incidence of both liver and kidney lesions. Moreover, SSH showed a strong antioxidant activity in DPPH and β -carotene-linoleic acid assays. SSH was found to contain phenolic compounds. Additionally, the SSH supplementation restored the hepatocytes viability against 2',7'-dichlorofluorescein (DCF) toxicity in *ex vivo* test.

1. Introduction

Aasal is the Arabic name for honey. It is a naturally sweet and flavorful product produced by honeybees, *Apis mellifera*, from the nectar of blossoms or from the exudates of trees and plants giving the nectar honey or honeydews [1]. Honey is a natural, unprocessed, and easily digested food in the diet [2]. Honey has been used in almost all cultures and traditions since ancient times as a food and medicinal product. It is considered to be used for the treatment of burns, faster wound healing [3], gastrointestinal disorders, asthma, skin, and eye diseases, besides being a natural food preservative and sweetening agent [4, 5]. Hippocrates, the

father of medicine, describes the nutritional and medicinal importance of honey [6].

Numerous nutritional [7] and biological effects [8] are attributed to honey, which include antibacterial [9], antioxidant [10], antiviral [11], antiparasitic [12], anti-inflammatory [13], anticancer [14], and immunosuppressive [15] activities. Apart from various therapeutic uses, honey is used in many folkloric traditions, for instance it is given to new born babies to clean their digestive system. Honey is considered to be the first in the line to treat jaundice in traditional medicine of different countries. In the present investigation, an attempt has been made to validate its use in both liver and kidney disorders.

2. Materials and Methods

Pure honey (Saudi Sidr variety) was purchased from an exclusive honey shop in Riyadh, Saudi Arabia.

2.1. Acute Toxicity Test. The acute toxicity test was performed on the mice using the oral route. SSH was dissolved in distilled water and administered at various doses, ranging from (500–5000 mg/kg), to different groups of mice. The animals were observed continuously for 1 h and then at half-hourly intervals for 4 h on the first day for clinical signs and symptoms of toxicity and further up to 72 h followed by 14 days for any mortality.

2.2. Animals and Study Design. Wistar albino rats (180–200 g) were obtained from the Experimental Animal Care Center of the College of Pharmacy, King Saud University, Riyadh. Animals were maintained on standard chow diet and housed in polycarbonate cages in a room free from any source of chemical contamination, artificially illuminated (12 h dark/light cycle) and thermally controlled ($25 \pm 2^\circ\text{C}$) at the animal facility. All animals received humane care in compliance with the guidelines of the Ethics Committee of the Experimental Animal Care Society, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia.

After a one-week acclimatization period, animals were randomly allocated into 5 groups and treated as follows: group (1), untreated control; group (2, 3, 4, and 5) received 0.25 mL of CCl_4 in liquid paraffin (1:1) 1.25 mL/kg body weight intraperitoneally (IP). Group 2 was administered only CCl_4 . Groups 3 and 4 received SSH 0.5 and 1.0 g/kg/day orally for 6 weeks. Rats in group 5 were treated with 10 mg/kg orally with silymarin for similar days. The blood was collected by cardiac puncture after 24 h following the administration of CCl_4 , allowed to clot and the serum was separated. After collecting the blood, the animals were sacrificed using ether anaesthesia. The liver and kidneys were dissected out and used for biochemical estimations and histological assessment.

2.2.1. Estimation of Marker Enzymes and Bilirubin. Serum glutamate oxaloacetate transaminase (SGOT), serum glutamate pyruvate transaminase (SGPT) [16], alkaline phosphatase (ALP) [17], gamma-glutamyl transferase (GGT) [18], hemoglobin, and bilirubin [19] were determined using Reflotron Plus Analyzer and Roche kits (Roche Diagnostics GmbH, Mannheim, Germany).

2.2.2. Estimation of Lipid Profile. Total cholesterol [20], triglycerides [21], high-density lipoproteins (HDL) [22], and glucose levels were estimated in serum using Roche diagnostic kits (Roche Diagnostics GmbH, Mannheim, Germany).

2.2.3. Determination of Malondialdehyde (MDA). The method reported by Utley et al. [23] was followed. In brief, the liver and kidney tissues were removed, and each tissue was homogenized in 0.15 M KCl (at 4°C ; Potter-Elvehjem type C homogenizer) to give a 10% w/v homogenate. The

absorbance of the solution was then read at 532 nm. The content of malondialdehyde (nmol/g wet tissue) was then calculated, by reference to a standard curve of malondialdehyde solution.

2.2.4. Estimation of Nonprotein Sulphydryls (NP-SH). Hepatic nonprotein sulphydryls were measured according to the method of Sedlak and Lindsay [24]. The liver and kidney were homogenized in ice-cold 0.02 mmol/L ethylenediaminetetraacetic acid (EDTA). The absorbance was measured within 5 min of addition of 5,5'-dithio-bis(2-nitrobenzoic acid) (DTNB) at 412 nm against a reagent blank.

2.2.5. Determination of Total Protein (TP). Total protein was estimated by the kit method, supplied by Crescent Diagnostics, Jeddah, Saudi Arabia. The absorbance of this complex at 546 nm is proportional to the protein concentration. The serum total protein was calculated using the equation:

$$\text{Serum total protein} = \frac{\text{ABS}_{\text{sample}}}{\text{ABS}_{\text{standard}}} \times \text{concentration of standard.} \quad (1)$$

2.2.6. Histopathological Evaluation. The liver and kidney tissue samples were fixed in neutral buffered formalin for 24 h. Sections of the liver tissue were histopathologically examined. These sections were stained with haematoxylin and eosin using routine procedures [25].

2.3. Ex Vivo Assay of SSH on Cultured Hepatocytes

2.3.1. Cells and Reagents. HepG2, a human hepatoma cell line was grown in RPMI medium (supplemented with 10% bovine serum, 1x penicillin-streptomycin, and 1x sodium pyruvate streptomycin (HyClone Laboratories)) at 37°C in a humidified chamber with 5% CO_2 .

2.3.2. Hepatotoxicity and Treatment. HepG2 cells were seeded (10^5 cells/well in triplicate) in a 96-well flat-bottom plate (Becton-Dickinson Labware) a day before treatment and grown. 2',7'-dichlorofluorescein (DCFH) (Sigma) commonly used to measure oxidative stress, *in vitro* [26], was used as a cytotoxic agent (IC_{50} : 100 μM , personal observation), prepared in DMSO (Sigma). An aqueous suspension of pure SSH (250 mg/mL stock) was prepared in RPMI medium followed by diluting to seven doses of SSH (0.1, 0.25, 0.5, 1.0, 2.5, 5, and 10 mg/mL). The cells (in triplicate) were replenished with RPMI containing 100 μM DCF plus a dose of SSH, including untreated as well as DCF-treated controls, and further incubated for 48 hours.

2.3.3. Microscopy. A direct visual observation was made under an inverted microscope (Optica, 40x and 100x) to see any morphological changes in the cells cultured with SSH and/or DCF on day 1 and 2.

2.3.4. Cell Proliferation and Viability Test. On day 2 of treatment, hepatocyte proliferation and viability test was

performed using TACS MTT Cell Proliferation and Viability Assay Kit (TACS).

2.4. Studies of the In Vitro Antioxidant Activity

2.4.1. DPPH Free Radical Scavenging Assay. The radical scavenging activity of the SSH against DPPH was evaluated as previously described [27]. The sample was redissolved in water, and various concentrations (31, 62, 125, 250, and 500 mg/mL) of the honey, 125 μ L prepared DPPH (1 mM in methanol), and 375 μ L solvent (methanol) were added. After 30 min incubation at 25°C, the decrease in absorbance was measured at $\lambda = 517$ nm. The radical scavenging activity was calculated from the equation:

$$\begin{aligned} &\% \text{ of radical scavenging activity} \\ &= \frac{\text{ABS}_{\text{control}} - \text{Abs}_{\text{sample}}}{\text{Abs}_{\text{control}}} \times 100. \end{aligned} \quad (2)$$

2.4.2. β -Carotene-Linoleic Acid Assay. The antioxidant activity of the SSH was evaluated, using the β -carotene bleaching method as described by Mothana et al. [28]. Rutin (1 mg/mL) was used as a standard. Absorbance was read at 470 nm at 15 min intervals, using a UV-visible spectrophotometer (UV mini-1240, Shimadzu, Japan). The antioxidant activity was calculated using the equation:

$$\% \text{ of antioxidant activity} = \frac{1 - (\text{Abs}_0 - \text{Abs}_t)}{(\text{Abs}_0 - \text{Abs}_t^{\circ})} \times 100, \quad (3)$$

where Abs_0 and Abs_0° are the absorbance values measured at zero time of incubation for sample extract and control, respectively.

Abs_t and Abs_t° are the absorbance values for sample extract and control, respectively, at $t = 120$ min.

2.5. Total Phenolic Content. The Folin-Ciocalteu method was used to determine the total phenolic content (TPC) of the honey according to Singleton et al. [29] and Liberato et al. [30]. Values of TPC were estimated by comparing the absorbance of each sample with a standard response curve generated using gallic acid (0, 12.5, 25, 50, 100, and 200 μ g/mL). The results were expressed as mg gallic acid equivalents (GAE)/100 g of honey. All the measurements were taken in triplicate, and the mean values were calculated.

2.6. Total Flavonoid Content. The total flavonoid content was determined by using a colorimetric assay as previously described [30]. Briefly, an aliquot of 5 mL of honey solution (0.02 mg/mL) or standard solution was mixed individually with the same volumes of solution of 2% aluminum chloride (AlCl_3) and allowed to stand at room temperature for 10 minutes. The absorbance was then read at 415 nm. A calibration curve was prepared with quercetin, and the results were expressed as mg quercetin equivalents (CE)/100 g of honey.

2.7. Statistical Analysis. Values are given as arithmetic means $\pm$ standard error of the mean (S.E.M.). Data was statistically analyzed by using one-way analysis of variance (ANOVA) followed by Student's t -test.

3. Results

3.1. Acute Toxicity Test. No toxicity symptoms were recorded. The LD_{50} value by oral route could not be determined as no lethality was observed up to 5.0 g/kg of the SSH in the animals.

3.2. In Vivo Effect of SSH on Liver and Kidney. The results indicated that animals treated with CCl_4 showed a significant increase in all biochemical parameters tested. However, animals treated with Saudi Sidr honey for 6 weeks before the intoxication with CCl_4 showed a significant decrease in serum GOT, GPT, ALP, GGT, creatinine, bilirubin, urea, and uric acid levels (Tables 1 and 3). CCl_4 -induced oxidative stress caused an elevation in lipid profile including cholesterol, triglycerides, LDL-C, and VLDL-C and reduction in the HDL-C levels in serum. The six-week pretreatment of rats with SSH in both doses, dose-dependently and significantly, reduced the cholesterol, triglycerides, LDL-C, and VLDL-C levels and significantly improved HDL-C level (Table 2). Silymarin, on the other hand, significantly prevent the CCl_4 -induced elevated levels of marker enzymes and lipid profile.

The results also indicated that treatment with CCl_4 resulted in a significant increase in MDA and a significant decrease in NP-SH and TP concentration in both liver and kidney tissues (Tables 4 and 5). Treatment of rats with SSH resulted in a significantly diminished level of MDA and significantly enhanced NP-SH and TP levels in both liver and kidney tissue.

Upon histopathological assessment of liver, the CCl_4 -induced rats showed an evidence of fatty changes with necrosis in liver cells, extensive fatty and inflammatory changes along with vascular congestion, and minimal fibrosis. The rats which received SSH (0.5 and 1.0 g/kg/day) and silymarin, as oral pretreatment showed a marked improvement in liver parenchyma with remnant degenerative changes of the cytoplasm and completely intact liver hepatocytes (Figure 1).

The histological examination of kidney sections of control rat showed normal glomeruli, interstitial tubules, and blood vessels. The kidney tissue slices in rats treated with CCl_4 exhibited glomerular congestion with vacuolization of the glomerular tuft and tubule with sloughing of the renal tubular lining. The kidney sections in rats pretreated with honey and silymarin showed vacuolization of glomerular tuft, evidence of tubular necrosis, and, regenerative and desquamation vacuolization (Figure 2).

3.3. Hepatoprotective Effect of SSH on Cultured Human Liver Cells

3.3.1. Microscopy. DCF exhibited severe cytotoxic effect on the human liver cells as reflected by altered morphology

TABLE 1: Effect of honey on CCl₄-induced hepatotoxicity-related parameters in rats.

Treatment group (n = 6)	SGOT (U/L)	SGPT (U/L)	GGT (U/L)	ALP (U/L)	Bilirubin (mg/dL)
Normal control	75.53 ± 2.86	26.93 ± 1.75	3.66 ± 0.37	282.00 ± 10.05	0.52 ± 0.02
CCl ₄ only (1.25 mL/kg)	300.00 ± 8.23 ^{***a}	209.50 ± 8.44 ^{*a}	15.45 ± 1.20 ^{***a}	671.33 ± 15.15 ^{***a}	3.45 ± 0.16 ^{***a}
Honey (0.5 g/kg) + CCl ₄	273.66 ± 10.28 ^b	182.83 ± 7.61 ^{*b}	12.81 ± 0.61 ^b	614.83 ± 13.41 ^{*b}	2.67 ± 0.14 ^{**b}
Honey (1.0 g/kg) + CCl ₄	200.00 ± 5.08 ^{***b}	140.83 ± 6.12 ^{***b}	10.38 ± 0.54 ^{**b}	605.16 ± 12.82 ^{**b}	2.01 ± 0.10 ^{***b}
Silymarin (10 mg/kg) + CCl ₄	133.58 ± 8.08 ^{***b}	106.78 ± 6.33 ^{***b}	5.08 ± 0.30 ^{***b}	416.16 ± 11.62 ^{***b}	1.17 ± 0.10 ^{***b}

All values represent mean ± SEM. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; ANOVA, followed by Dunnett's multiple comparison test.

^a As compared with normal group. ^b As compared with CCl₄ only group.

TABLE 2: Effect of honey on CCl₄-induced lipid profile changes in rats.

Treatment group (n = 6)	Cholesterol (mg/dL)	Triglycerides (mg/dL)	HDL (mg/dL)	LDL (mg/dL)	VLDL (mg/dL)
Normal control	13.79 ± 1.22	91.34 ± 4.57	52.56 ± 6.14	68.98 ± 6.14	105.33 ± 3.81
CCl ₄ only (1.25 mL/kg)	43.61 ± 1.58 ^{***a}	237.63 ± 10.61 ^{***a}	23.47 ± 1.09 ^{***a}	218.05 ± 7.94 ^{***a}	281.33 ± 11.01 ^{***a}
Honey (0.5 g/kg) + CCl ₄	34.35 ± 1.63 ^{**b}	172.29 ± 8.93 ^{***b}	31.28 ± 1.12 ^{**b}	171.29 ± 8.19 ^{**b}	206.66 ± 9.77 ^{***b}
Honey (1.0 g/kg) + CCl ₄	20.00 ± 1.17 ^{***b}	170.50 ± 6.46 ^{***b}	44.38 ± 1.97 ^{***b}	100.00 ± 5.87 ^{***b}	190.66 ± 6.97 ^{***b}
Silymarin (10 mg/kg) + CCl ₄	38.79 ± 0.80 ^{*b}	198.43 ± 9.82 ^{*b}	29.27 ± 1.55 ^{*b}	199.98 ± 4.02 ^{*b}	237.33 ± 9.16 ^{*b}

All values represent mean ± SEM. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; ANOVA, followed by Dunnett's multiple comparison test.

^a As compared with normal group, ^b As compared with CCl₄ only group.

TABLE 3: Effect of honey on CCl₄-induced kidney function test in serum.

Treatment group (n = 6)	Creatinine (mg/dL)	Uric Acid (mg/dL)	Urea (mmol/L)	Sodium (mEq/L)	Potassium (mEq/L)	Calcium (mg/dL)
Normal control	1.34 ± 0.09	1.28 ± 0.010	35.8 ± 1.90	54.96 ± 0.99	3.61 ± 0.18	4.42 ± 0.34
CCl ₄ only (1.25 mL/kg)	7.48 ± 0.31 ^{***a}	5.68 ± 0.36 ^{***a}	167.16 ± 4.65 ^{***a}	101.89 ± 2.26 ^{***a}	10.58 ± 0.48 ^{***a}	24.64 ± 0.81 ^{***a}
Honey (0.5 g/kg) + CCl ₄	6.48 ± 0.31 ^{*b}	4.25 ± 0.43 ^{*b}	157.5 ± 5.51 ^b	88.67 ± 2.35 ^{**b}	9.18 ± 0.27 ^{*b}	20.42 ± 0.98 ^{**b}
Honey (1.0 g/kg) + CCl ₄	5.11 ± 0.13 ^{***b}	2.89 ± 0.21 ^{***b}	131.83 ± 8.15 ^{**b}	78.14 ± 2.30 ^{***b}	7.31 ± 0.23 ^{***b}	13.76 ± 0.87 ^{***}
Silymarin (10 mg/kg) + CCl ₄	6.08 ± 0.27 ^{**b}	4.30 ± 0.25 ^{*b}	154.66 ± 5.85 ^b	93.44 ± 3.20 ^{*b}	10.05 ± 0.40 ^b	25.12 ± 0.84 ^b

All values represent mean ± SEM. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; ANOVA, followed by Dunnett's multiple comparison test.

^a As compared with normal group, ^b As compared with CCl₄ only group.

TABLE 4: Biochemical parameters (liver tissue) treated with honey.

Treatment group (n = 6)	Total protein (g/L)	MDA (nmol/g)	NP-SH (nmol/g)
Normal control	122.49 ± 4.61	0.97 ± 0.11	7.08 ± 0.34
CCl ₄ only (1.25 mL/kg)	41.52 ± 2.59 ^{***a}	11.88 ± 0.77 ^{***a}	3.81 ± 0.43 ^{***a}
Honey (0.5 g/kg) + CCl ₄	54.38 ± 4.29 ^{*b}	7.96 ± 0.77 ^{**b}	4.96 ± 0.47 ^b
Honey (1.0 g/kg) + CCl ₄	58.44 ± 3.81 ^{**b}	5.80 ± 0.58 ^{***b}	6.13 ± 0.37 ^{**b}
Silymarin (10 mg/kg) + CCl ₄	80.83 ± 3.28 ^{***b}	2.91 ± 0.23 ^{***b}	6.96 ± 0.29 ^{***b}

All values represent mean ± SEM. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; ANOVA, followed by Dunnett's multiple comparison test.

^a As compared with normal group, ^b As compared with CCl₄ only group.

compared to untreated cells. Interestingly, the DCF-treated cells supplemented with 10 mg/mL of SSH were morphologically different from the DCF-treated cells but comparable to untreated cells (data not shown).

3.3.2. Hepatocyte Protection and Viability Restoration by SSH. Our MTT test showed a protective effect of SSH (10 mg/mL) against DCF (100 μ M) induced hepatotoxicity at 48 hours posttreatment (Figure 3), and in line with our microscopic observation that confirmed the *ex vivo* hepatoprotection by pure SSH. Under these conditions DCF-toxicated cells were

recovered to about 70% with 10 mg/mL of SSH. The SSH supplementation restored the hepatocytes viability by 1.5-fold against DCF toxicity.

3.4. Antioxidant Activity and Phenolic and Flavonoidal Contents of the SSH. The potential antioxidant activity of the SSH was investigated on the basis of DPPH radical scavenging activity and of inhibition of linoleic acid oxidation. As demonstrated in Table 6, SSH was able to reduce the stable free radical DPPH to the yellow-colored DPPH at low concentrations (125 and 250 mg/mL), almost near to the

TABLE 5: Biochemical parameters (kidney tissue) treated with honey.

Treatment group (n = 6)	Total protein (g/L)	MDA (nmol/g)	NP-SH (nmol/g)
Normal control	89.75 ± 4.12	0.69 ± 0.13	7.23 ± 0.46
CCl ₄ only (1.25 mL/kg)	31.70 ± 3.45 ^{***a}	8.53 ± 0.91 ^{***a}	4.08 ± 0.35 ^{***a}
Honey (0.5 g/kg) + CCl ₄	41.89 ± 1.62 ^{*b}	5.77 ± 0.39 ^{*b}	4.76 ± 0.45 ^b
Honey (1.0 g/kg) + CCl ₄	46.17 ± 2.18 ^{**b}	3.95 ± 0.32 ^{***b}	5.80 ± 0.31 ^{**b}
Silymarin (10 mg/kg) + CCl ₄	64.72 ± 7.42 ^{***b}	1.89 ± 0.28 ^{***b}	6.29 ± 0.46 ^{**b}

All values represent mean ± SEM. **P* < 0.05; ***P* < 0.01; ****P* < 0.001; ANOVA, followed by Dunnett's multiple comparison test.

^aAs compared with normal group, ^bAs compared with CCl₄ only group.

TABLE 6: Free radical scavenging activity, antioxidant activity, total phenolic, and total flavonoidal contents of the honey sample.

Plant species	Radical scavenging activity (%)					Total antioxidant activity (%) 125 (mg/mL)	TPC (mg GAE /100 g)	TFC (mg QE/100 g)
	31.25	62.5	125	250	500			
Honey	28.0	45.0	72.7	89.3	91.8	90.5 ± 7.62	105.1 ± 4.04	48.3 ± 1.17
Ascorbic acid	18.5	74.1	88.9	93.0	94.0	—		
Rutin						91.9 ± 5.83		

TPC: Total phenolic content; TFC: Total flavonoidal content.

ascorbic acid. In addition to that, in the β -carotene/linoleic acid model system, the SSH was also able to inhibit the discoloration of β -carotene at a concentration of 125 mg/mL. The total antioxidant value was 90.8% (Table 6). The observed antioxidant activities were comparable to that of the positive control, rutin (Table 6). Moreover, the SSH showed high total phenolic and flavonoidal contents (105.1 ± 4.04 mg gallic acid equivalents/100 g and 48.3 ± 1.17 mg quercetin equivalents/100 g).

4. Discussion

The liver is an organ that not only performs physiological functions but also protect against the hazards of harmful drugs, chemicals and xenobiotics. The liver is one of the largest and highly complex organs with multifunction, including nutrient storage, maintenance of homeostasis, secretory and excretory function, and synthesis of proteins [31]. These functions also include the metabolism of certain hormones, lipid metabolism (cholesterol, triglycerides, and high-density lipoproteins (HDL)), protein metabolism, and detoxification of xenobiotics [32]. The model used in the present study has been subjected to thorough critical appraisal and validated animal models of hepatorenal protective activity of honey in rats against CCl₄ as a hepatorenal toxin to prove its claims in folklore practice against liver and kidney disorders.

Carbon tetrachloride, besides exerting its toxic effect on liver, also reportedly gets distributed at higher concentrations in the kidney than in the liver [33]. The mechanism of CCl₄ renal toxicity is almost the same as that of liver, but CCl₄ shows a high affinity to the kidney cortex which contains cytochrome P-450 predominantly [34, 35]. Due to CCl₄ hepatorenal injury, the transport function of hepatocytes and nephrotic cells gets disturbed resulting in the leakage

of plasma membrane, thereby causing an increased enzyme level in the serum [36].

In the present study CCl₄ administration to rats caused a significant increase in serum GOT, GPT, GGT, ALP, urea, uric acid, bilirubin, and creatinine as well as MDA in the liver and kidney levels accompanied with a significant decrease in total protein and NP-SH in liver and kidney, which indicates the extensive disruption of the structure and function of the liver and kidney. The tendency of these marker enzymes (SGPT, SGOT, ALPG, GGT) at a near normal level in the groups of rats treated with Saudi Sidr honey and silymarin is a clear manifestation of antihepatorenal toxic effect of the honey. Our findings are in agreement with an earlier study in which feeding of honey caused liver protective effect by depleting the elevated liver marker enzymes in CCl₄-treated rats [37]. On the other hand, treatment of rats with CCl₄ increases the levels of total lipids, triglycerides, and cholesterol in serum [38]. The significant diminution of lipids and total cholesterol in the SSH-treated rats and an increase in HDL-C level further indicates the hepatoprotective potential of the honey.

Furthermore, the efficacy of any hepatoprotective drug is essentially dependent on its capacity of either maintaining normal physiological function or diminishing the harmful effects which were affected by hepatotoxic agents [39]. Reduction in total protein (TP) content can be deemed as a useful index of severity of hepatocellular damage [40]. The lowered levels of total proteins in the serum of CCl₄-treated rats exhibited the severity of hepatopathy. Thus, this suggests that honey possesses the ability to promote protein synthesis leading to the higher concentration of protein in the liver [35].

The present study also revealed that the administration of CCl₄ caused marked impairment in renal function alongside with significant oxidative stress in the kidney [41]. Serum creatinine, urea, and uric acid concentrations were significantly higher in CCl₄-treated rats. Serum creatinine elevation was

FIGURE 1: Light micrographs showing the effect of SSH on CCl_4 -induced hepatotoxicity in rats. (a) Normal hepatocytes. (b) CCl_4 -induced severe necrosis and inflammation. (c) Pretreatment of rats with SSH 0.5 g/kg. (d) Pretreatment of rats with SSH 1.0 g/kg. (e) Pretreatment of rats with silymarin 10 mg/kg.

caused by CCl_4 due to altered kidney function. Urea is the main end product of protein catabolism. It is one of the waste products of the body which is passed into blood stream to be removed by kidney [42]. Honey significantly decreased the elevated levels of serum creatinine, urea, and uric acid, which indicates that the honey possibly protects kidney tissue against oxidative damages induced by CCl_4 [43] and indicates maintenance of renal function [44].

In vitro antioxidant activity of Saudi Sidr honey revealed a strong antioxidant activity in both the tests DPPH and β -carotene linoleic acid used. The observed hepatonephro

protective and antioxidant activity of honey might have been due to the presence of phenolic and flavonoidal contents. In the current study, the increase in MDA level in the liver and kidney by CCl_4 suggests enhanced lipid peroxidation (LPO) which is an important pathogenic event that damages biomembranes [45]. LPO is thought to be a consequence of oxidative stress which occurs when the dynamic balance between prooxidant and antioxidant mechanism is impaired [46]. Malondialdehyde is known to be a reliable marker of LPO and oxidative stress [47]. Treatment of rats with SSH significantly prevented CCl_4 -induced increase in MDA

FIGURE 2: Light micrographs showing the effect of SSH on CCl_4 -induced nephrotoxicity in rats. (a) Normal kidney tissues. (b) CCl_4 -induced severe glomerular congestion and vacuolization of the glomerular tuft and tubules with sloughing of the renal tubular lining. (c) Pretreatment of rats with SSH 0.5 g/kg. (d) Pretreatment of rats with SSH 1.0 g/kg. (e) Pretreatment of rats with silymarin 10 mg/kg.

concentration in both liver and kidney. Hence, the observed hepato-renal protective activity of honey may be due to its antioxidant property.

Nonprotein sulfhydryls are known to be involved in several defense processes against oxidative damage; protects cells against free radicals peroxides and various poisonous substances [48]. Thus, a deficiency of GSH within the living organisms can cause tissue injury and malfunction [49]. In the current study, the liver and kidney NP-SH level in CCl_4 -treated group was significantly diminished when compared with the control group. These findings are in accordance with earlier reports as sulfhydryl levels were significantly

depleted in different organs of rats, when exposed with CCl_4 [50]. Pretreatment of rats with SSH replenished NP-SH concentration in both liver and kidney homogenate as compared with CCl_4 only treated animals, suggesting the ameliorative effects of SSH on liver and kidney damage induced by CCl_4 , at least in part, due to its free radical scavenging activity. Furthermore, most of the parameters in the group which received CCl_4 plus Sidr honey having nearer value of the control group demonstrate Sidr honey to have antiradical effect [51]. This effect is considered to be related to the phenolic compounds and *in vitro* antioxidant activity of Saudi Sidr honey.

FIGURE 3: MTT test showing hepatoprotective effect of SSH (10 mg/mL) against DCF (100 µM)-induced toxicity of cultured human liver cells (HepG2).

On the other hand, DCFH is generally used to measure *in vitro* oxidative stress generated by free radicals through the principle of oxidation of DCFH to the fluorescent DCF. However, we used this agent because of its cytotoxic effect (unpublished). In this study, our *ex vivo* SSH protection against DCFH-induced human liver toxicity further supported the *in vivo* effects of SSH in CCl₄-induced rat liver injury. In this way, we have confirmed our results by using two different toxins in two different systems, at least for hepatoprotection.

5. Conclusions

In conclusion, the administration of Saudi Sidr honey (SSH) prevented biochemical and histomorphological alteration induced by CCl₄. This hepatorenal protective effect of SSH could be attributed to the presence of antioxidative factors for example, phenolic and flavonoidal compounds which cause significant lowering of the oxidative threat leading to normal physiological function. The findings support the use of SSH for the treatment of several liver and kidney ailments.

Conflict of Interests

The authors confirm that there is no conflict of interests.

Acknowledgments

The authors would like to express their gratitude to the Research Center, College of Pharmacy, and the Deanship of Scientific Research of King Saud University, Riyadh, for their assistance.

References

- [1] J. M. Alvarez-Suarez, S. Tulipani, S. Romandini, E. Bertoli, and M. Battino, "Contribution of honey in nutrition and human health: a review," *Mediterranean Journal of Nutrition and Metabolism*, vol. 3, no. 1, pp. 15–23, 2010.
- [2] X. Feás, J. Pires, A. Iglesias, and M. L. Estevinho, "Characterization of artisanal honey produced on the Northwest of Portugal by melissopalynological and physico-chemical data," *Food and Chemical Toxicology*, vol. 48, no. 12, pp. 3462–3470, 2010.
- [3] K. Lay-flurrie, "Honey in wound care: effects, clinical application and patient benefit," *British Journal of Nursing*, vol. 17, no. 11, pp. S30–S32, 2008.
- [4] M. L. Al, D. Daniel, A. Moise, O. Bobis, L. Laslo, and S. Bogdanov, "Physico-chemical and bioactive properties of different floral origin honeys from Romania," *Food Chemistry*, vol. 112, no. 4, pp. 863–867, 2009.
- [5] I. C. F. R. Ferreira, E. Aires, J. C. M. Barreira, and L. M. Estevinho, "Antioxidant activity of Portuguese honey samples: different contributions of the entire honey and phenolic extract," *Food Chemistry*, vol. 114, no. 4, pp. 1438–1443, 2009.
- [6] A. B. Jull, A. Rodgers, and N. Walker, "Honey as a topical treatment for wounds," *Cochrane Database of Systematic Reviews*, no. 4, p. CD005083, 2008.
- [7] S. Bogdanov, T. Jurendic, R. Sieber, and P. Gallmann, "Honey for nutrition and health: a review," *Journal of the American College of Nutrition*, vol. 27, no. 6, pp. 677–689, 2008.
- [8] A. E. Jeffrey and C. M. Echazarreta, "Medical uses of honey," *Reviews in Biomedical*, vol. 7, no. 1, pp. 43–49, 1996.
- [9] K. L. Allen, P. C. Molan, and G. M. Reid, "A survey of the antibacterial activity of some New Zealand honeys," *Journal of Pharmacy and Pharmacology*, vol. 43, no. 12, pp. 817–822, 1991.
- [10] L. Vela, C. De Lorenzo, and R. A. Pérez, "Antioxidant capacity of Spanish honeys and its correlation with polyphenol content and other physicochemical properties," *Journal of the Science of Food and Agriculture*, vol. 87, no. 6, pp. 1069–1075, 2007.

- [11] A. Jeddar, A. Kharsany, and U. G. Ramsaroop, "The antibacterial action of honey. An in vitro study," *South African Medical Journal*, vol. 67, no. 7, pp. 257–258, 1985.
- [12] B. Zeina, B. I. Zohra, and S. Al-assad, "The effects of honey on leishmania parasites: an in vitro study," *Tropical Doctor*, vol. 27, no. 1, pp. 36–38, 1997.
- [13] F. R. Greten, L. Eckmann, T. F. Greten et al., "IKK β links inflammation and tumorigenesis in a mouse model of colitis-associated cancer," *Cell*, vol. 118, no. 3, pp. 285–296, 2004.
- [14] M. López-Lázaro, "Dual role of hydrogen peroxide in cancer: possible relevance to cancer chemoprevention and therapy," *Cancer Letters*, vol. 252, no. 1, pp. 1–8, 2007.
- [15] P. Michaluart, J. L. Masferrer, A. M. Carothers et al., "Inhibitory effects of caffeic acid phenethyl ester on the activity and expression of cyclooxygenase-2 in human oral epithelial cells and in a rat model of inflammation," *Cancer Research*, vol. 59, no. 10, pp. 2347–2352, 1999.
- [16] S. A. Reitman and S. Frankel, "A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases," *American Journal of Clinical Pathology*, vol. 28, no. 1, pp. 56–63, 1957.
- [17] E. J. King and A. R. Armstrong, "Calcium, phosphorus and phosphate," in *Practical Clinical Biochemistry*, H. Varley, Ed., CBS publishers, New Delhi, India, 1988.
- [18] S. Fiala, A. E. Fiala, and B. Dixon, "Gamma-glutamyl transpeptidase in transplantable, chemically induced rat hepatomas and spontaneous mouse hepatomas," *Journal of the National Cancer Institute*, vol. 48, no. 5, pp. 1393–1401, 1972.
- [19] A. Stiehl, "Hyperbilirubinaemia in liver disease," *Fortschritte der Medizin*, vol. 100, no. 18, pp. 842–845, 1982.
- [20] P. N. M. Demacher and A. G. M. Hijamaus, "A study of the use of polyethylene glycol in estimating cholesterol," *Clinical Chemistry*, vol. 26, pp. 1775–1778, 1980.
- [21] L. B. Foster and R. T. Dunn, "Stable reagents for determination of serum triglycerides by a colorimetric Hantzsch condensation method," *Clinical Chemistry*, vol. 19, no. 3, pp. 338–340, 1973.
- [22] M. Burstein and H. R. Scholnick, "Turbidimetric estimation of chylomicrons and very low density lipoproteins in human sera after precipitation by sodium lauryl sulfate," *Biomedicine*, vol. 19, no. 1, pp. 16–19, 1973.
- [23] H. C. Utley, F. Bernheim, and P. Hochslein, "Effect of sulfhydryl reagent on peroxidation in microsome," *Archives of Biochemistry Biophysics*, vol. 260, pp. 521–531, 1967.
- [24] J. Sedlak and R. H. Lindsay, "Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent," *Analytical Biochemistry*, vol. 25, pp. 192–205, 1968.
- [25] C. F. Culling, *Handbook of Histopathological and Histochemical Techniques*, Butterworth, London, UK, 3rd edition, 1974.
- [26] C. Rota, C. F. Chignell, and R. P. Mason, "Evidence for free radical formation during the oxidation of 2'-7'-dichlorofluorescein to the fluorescent dye 2'-7'-dichlorofluorescein by horseradish peroxidase: possible implications for oxidative stress measurements," *Free Radical Biology and Medicine*, vol. 27, no. 7-8, pp. 873–881, 1999.
- [27] W. W. Brand-Williams, M. E. Cuvelier, and C. Berset, "Use of a free radical method to evaluate antioxidant activity," *Food Science and Technology*, vol. 28, no. 1, pp. 25–30, 1995.
- [28] R. A. A. Mothana, M. S. Al-Said, A. J. Al-Rehaily, T. M. Thabet, N. A. Awad, M. Lalk et al., "Anti-inflammatory, antinociceptive, antipyretic and antioxidant activities and phenolic constituents from *Loranthus regularis* Steud. Ex Sprague," *Food Chemistry*, vol. 130, no. 2, pp. 344–349, 2012.
- [29] V. L. Singleton, R. Orthofer, and R. M. Lamuela-Raventós, "Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent," *Methods in Enzymology*, vol. 299, pp. 152–178, 1998.
- [30] M. D. C. T. C. Liberato, S. M. De Moraes, S. M. C. Siqueira et al., "Phenolic content and antioxidant and antiacetylcholinesterase properties of honeys from different floral origins," *Journal of Medicinal Food*, vol. 14, no. 6, pp. 658–663, 2011.
- [31] E. M. Williamson, D. T. Okpako, and F. J. Evans, *The Liver and Biliary System: Selection, Preparation and Pharmacological Evaluation of Plant Material*, John Wiley and Sons, London, UK, 1996.
- [32] B. L. de Medeiros, K. S. Costa, J. Alves et al., "Liver protective activity of a hydroethanolic extract of *Arrabidaea chica* (Humb. and Bonpl.) B. Verl., (pariri)," *Pharmacognosy Research*, vol. 3, no. 2, pp. 79–84, 2011.
- [33] U. Y. Sanzgiri, V. Srivatsan, S. Muralidhara, C. E. Dallas, and J. V. Bruckner, "Uptake, distribution, and elimination of carbon tetrachloride in rat tissues following inhalation and ingestion exposures," *Toxicology and Applied Pharmacology*, vol. 143, no. 1, pp. 120–129, 1997.
- [34] F. Jaramillo-juárez, M. L. Rodríguez-Vázquez, A. R. Rincón-Sánchez, M. Consolación Martínez, G. G. Ortiz, J. Llamas et al., "Acute renal failure induced by carbon tetrachloride in rats with hepatic cirrhosis," *Annals of Hepatology*, vol. 7, no. 4, pp. 331–338, 2008.
- [35] P. Abraham, G. Wilfred, and S. P. Cathrine, "Oxidative damage to the lipids and proteins of the lungs, testis and kidney of rats during carbon tetrachloride intoxication," *Clinica Chimica Acta*, vol. 289, no. 1-2, pp. 177–179, 1999.
- [36] G. S. Achliya, S. G. Wadodkar, and A. K. Dorle, "Evaluation of hepatoprotective effect of Amalkadi Ghrita against carbon tetrachloride-induced hepatic damage in rats," *Journal of Ethnopharmacology*, vol. 90, no. 2-3, pp. 229–232, 2004.
- [37] B. Kilicoglu, C. Gencay, K. Kismet et al., "The ultrastructural research of liver in experimental obstructive jaundice and effect of honey," *American Journal of Surgery*, vol. 195, no. 2, pp. 249–256, 2008.
- [38] A. Seakins and D. S. Robinson, "The effect of the administration of carbon tetrachloride on the formation of plasma lipoproteins in the rat," *The Biochemical Journal*, vol. 86, pp. 401–407, 1963.
- [39] M. I. Kazeem, H. A. Bankole, and A. A. Fatai, "Protective effect of ginger in normal and carbon-tetrachloride induced hepatotoxic rats," *Annals of Biological Research*, vol. 2, no. 1, pp. 1–8, 2011.
- [40] Y. Aniya, T. Koyama, C. Miyagi et al., "Free radical scavenging and hepatoprotective actions of the medicinal herb, *Crossocephalum crepidioides* from the Okinawa Islands," *Biological and Pharmaceutical Bulletin*, vol. 28, no. 1, pp. 19–23, 2005.
- [41] S. O. Adewole, A. A. Salako, O. W. Doherty, and T. Naicker, "Effect of melatonin on carbon tetrachloride-induced kidney injury in Wistar rats," *African Journal of Biomedical Research*, vol. 10, no. 2, pp. 153–164, 2007.
- [42] S. R. Orth and E. Ritz, "The nephrotic syndrome," *Internist*, vol. 39, no. 12, pp. 1246–1252, 1998.
- [43] D. W. Cockcroft and M. H. Gault, "Prediction of creatinine clearance from serum creatinine," *Nephron*, vol. 16, no. 1, pp. 31–41, 1976.
- [44] A. M. Davison, J. S. Cameron, J.-P. Gruenfeld, D. N. S. Kerr, E. Ritz, and G. Winearls, "Oxford Textbook of Clinical Nephrology," pp. 36–39, Oxford University Press, Oxford, USA, 1998.

- [45] A. A. J. Nayagam, S. Manokaran, and N. Sudhakar, "Hepatoprotective efficacy of *Tricholepis radicans* DC. against CCl_4 induced liver toxicity in albino rats," *Journal of Pharmacy Research*, vol. 4, no. 4, pp. 1073–1075, 2011.
- [46] R. S. Kumar, K. Asokkumar, and N. V. Murthy, "Hepatoprotective effects and antioxidant role of *scutia myrtina* on paracetamol induced hepatotoxicity in rats," *Journal of Complementary and Integrative Medicine*, vol. 8, no. 1, p. 8, 2011.
- [47] D. D. Kitts, Y. V. Yuan, and D. V. Godin, "Plasma and lipoprotein lipid composition and hepatic antioxidant status in spontaneously hypertensive (SHR) and normotensive (WKY) rats," *Canadian Journal of Physiology and Pharmacology*, vol. 76, no. 2, pp. 202–209, 1998.
- [48] H. Sies, "Glutathione and its role in cellular functions," *Free Radical Biology and Medicine*, vol. 27, no. 9–10, pp. 916–921, 1999.
- [49] S. A. Ganie, E. Haq, A. Hamid et al., "Carbon tetrachloride induced kidney and lung tissue damages and antioxidant activities of the aqueous rhizome extract of *Podophyllum hexandrum*," *BMC Complementary and Alternative Medicine*, vol. 11, p. 17, 2011.
- [50] Y. Ohta, M. Kongo, E. Sasaki, K. Nishida, and I. Ishiguro, "Therapeutic effect of melatonin on carbon tetrachloride-induced acute liver injury in rats," *Journal of Pineal Research*, vol. 28, no. 2, pp. 119–126, 2000.
- [51] G. Eraslan, M. Kanbur, S. Silici, and M. Karabacak, "Beneficial effect of pine honey on trichlorfon induced some biochemical alterations in mice," *Ecotoxicology and Environmental Safety*, vol. 73, no. 5, pp. 1084–1091, 2010.