

## Research Article

# Evaluation on Anti-Inflammatory, Analgesic, Antitumor, and Antioxidant Potential of Total Saponins from *Nigella glandulifera* Seeds

Jun Zhao, Fang Xu, Hua Huang, Zhengyi Gu, Linlin Wang, Wei Tan, Jinhua He, Yan Chen, and Chenyang Li

Xinjiang Key Laboratory for Uighur Medicine, Institute of Materia Medica of Xinjiang, Urumqi 830004, China

Correspondence should be addressed to Zhengyi Gu; zhengyi087@126.com

Received 12 October 2012; Revised 15 December 2012; Accepted 16 January 2013

Academic Editor: Weena Jiratchariyakul

Copyright © 2013 Jun Zhao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Nigella glandulifera* seeds are used as a spice or remedy for the treatment of various inflammatory diseases. This study aimed to investigate analgesic (writhing test), anti-inflammatory (ear-induced edema, vascular permeability test), antioxidant, and antitumor activities of total saponins from this plant (TSN). TSN (6, 12, and 24 mg/kg) were exhibited analgesic and anti-inflammatory activities in a dose-dependent manner ( $P < 0.05$ ). In D-galactose-induced ageing model, TSN significantly increased the plasma superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px) activities ( $P < 0.05$ ) and decreased the malondialdehyde (MDA) level compared to control group ( $P < 0.05$ ). DPPH radical scavenging effect of TSN was also found. Moreover, TSN (20 mg/mL) showed 86.75% and 88.26% inhibition of the growth on Bel-7402 and Hela cells, respectively. Five compounds were further isolated and identified from TSN as Nigella A, B, C, D, and nige-glanoside, of which the content of Nigella A was  $60.36 \pm 1.25$  g/100 g TSN by HPLC-ELSD method. Altogether, these results suggest that TSN could be considered as a potential analgesic, anti-inflammatory, antitumor, and antioxidant agent.

## 1. Introduction

*Nigella* (*Nigella sativa* L., *Nigella glandulifera* Freyn et Sint, and *Nigella damascena*, L. etc.), belonging to the buttercup family Ranunculaceae, is commonly known as black cumin (black seeds) [1]. *Nigella* seeds are widely used for medicinal purposes as a natural remedy for a number of illnesses such as hypertension, diabetes, inflammation, bronchitis, headache, and gastrointestinal disturbances [2]. In recent years, voluminous research has been carried out on the medicinal properties of the seeds as antioxidant, antimicrobial, anti-inflammatory, and anticancer agents [3–5]. These properties have been attributed to a variety of active constituents in seeds and its fixed oil [6, 7], of which saponins are mainly characteristic compounds of water-soluble extracts from *Nigella* [8]. Pharmacological effects of saponins have been reported in many references, and these compounds are considered to be beneficial to the health of mankind [9, 10].

*N. glandulifera* Freyn is widely distributed in Xinjiang, Yunnan, and Tibet of China and is now mainly cultivated in the Taklimakan Desert edge region in Xinjiang [11]. The seeds of *N. glandulifera* are frequently added to “naan” (a kind of crusty pancake, favorite food of the Uygur and Kazak people) as a spice, and its water decoction is used in Uighur’s traditional medicine for the treatment of numerous disorders such as diuretic, analgesic, insomnia, dizziness, tinnitus, amnesia, and bronchial asthma [12, 13]. Several classes of compounds have been isolated from water-soluble extracts of *N. glandulifera*, such as alkaloids, flavonoids, and saponins [14, 15]. Saponins with the highest content (64.5%) are mainly characteristic compound of *N. glandulifera* (TSN). However, the biological activities of saponins from this plant were rarely reported so far. Therefore, the aim of this study was to investigate anti-inflammatory, analgesic, anti-tumor, and antioxidant potential of total saponins from *N. glandulifera* by experiments *in vivo* or *in vitro*.

## 2. Material and Methods

**2.1. Chemicals.** 2,2-Diphenyl-1-(2,4,6-trinitrophenyl)hydrazyl (DPPH) and dimethyl sulfoxide (DMSO) were purchased from Sigma-Aldrich (St. Louis, MO, USA). Superoxide dismutase (SOD), glutathione peroxidase (GSH-Pxs), and malondialdehyde (MDA) assay kits were supplied by Nanjing Jiancheng Bioengineering Institute (Nanjing, China). RPMI 1640 medium was obtained from Gibco Co. (USA). HPLC-grade acetonitrile was purchased from Fisher BioReagents (NJ, USA). Yemugua tablet (YMGP) was purchased from Hepin Pharmacy Co. (Guangdong, China). The other chemicals and solvents used in this experiment were of the highest quality available.

**2.2. Plant Material.** *N. glandulifera* were collected from Aksu, Xinjiang, in China, in July 2011. The plant material was identified by the associate researcher Jiang He, Xinjiang Institute of Material Medica. A voucher specimen was deposited at Xinjiang Institute of Material Medica in China.

**2.3. Preparation of TSN.** The powdered seeds (10.0 kg) were defatted at reflux condition with petroleum ether and extracted with 30% ethanol by exhaustive maceration to yield a dark brown residue (2.2 kg). After being dissolved in water, the extract was purified by AB-8 adsorption macroporous resin to obtain total saponin extracts from *N. glandulifera* (TSN, 210 g). TSN were applied to ODS RP-18 column and eluted with mixtures of MeOH:H<sub>2</sub>O (0:1 → 1:0) successively. Elutes were combined into five subfractions according to TLC behavior using two solvent systems CHCl<sub>3</sub>:MeOH:H<sub>2</sub>O (6:4:0.5) and BuOH:AcOH:H<sub>2</sub>O (4:1:1) (spots were visualized after spraying 10% H<sub>2</sub>SO<sub>4</sub>). Various fractions were repeatedly purified by Sephadex LH-20 column with methanol, and five saponins were isolated from TSN, and their structures were confirmed using MS, <sup>1</sup>H, and <sup>13</sup>C NMR (References). The purity of the saponins was determined to be more than 95% compared with the peak areas detected by HPLC-ELSD.

**2.4. HPLC-ELSD Analysis of TSN.** The high-performance liquid chromatography (HPLC) (LC-10A HPLC instrument, Shimadzu Co., Japan) was employed to analyze the percentage contents of Nigella A in TNS. A Cosmosil-C18 column (250 mm × 4.6 mm, 5 μm) and a Zorbax ODS C18 guard column (12.5 mm × 4.6 mm, 5 μm) were used at 40°C. A binary gradient elution system consisted of water (A) and acetonitrile (B) and separation was achieved using the following gradient program: 0–15 min, 20–40%B; 15–20 min, 40%B; 20–22 min, 40–20%B; and 22–25 min, 20%B. The flow rate was at 1.0 mL/min and the sample injection volume was 5 μL. The all tech ELSD impactor was set at ON mode, the drift tube temperature was 65°C, and the nebulizer nitrogen gas flow rate was at 2.0 L/min.

**2.5. Animals.** The study was conducted on Kunming mice weighting 18 ± 22 g (The Experimental Animal Center in Xinjiang, China; SCXK (Xin) 2003–2001). Animals were kept

under a 12 h/12 h light/dark cycle and allowed free access to food and water. The study protocols were approved by the Ethics Committee on Animal Experiment, Xinjiang Material Medica, China.

**2.6. Cells.** The human hepatoma carcinoma cell lines Bel 7402 and cervical carcinoma cell lines Hela were provided by the Institute of Materia Medica, Chinese Academy of Medical Sciences, and maintained with RPMI 1640 medium containing 10% fetal bovine serum and 100 ng/mL, each, of penicillin and streptomycin at 37°C in a humidified atmosphere with 5% CO<sub>2</sub>.

**2.7. Acute Oral Toxicity Study in Mice.** TSN were subjected to acute oral toxicity studies as per SFDA guidelines no. [Z] GPT2-1 [16]. Mice were randomly assigned to each of three groups of 30 mice (15 females and 15 males). They were fasted overnight (12 h) with free access to water prior to administration of doses (48 and 24 g/kg) of TSN dissolved in distilled water. The control group mice were p.o. administered with distilled water. The animals were observed continuously for first 72 hours and 14 days for any signs of mortality, body weight, toxicity, behavioral, and viscera changes. On day 15, mice were killed and all organs and tissues were observed macroscopically.

**2.8. Cytotoxicity Assays.** The human hepatoma carcinoma cell lines Bel-7402 and cervical carcinoma cell lines Hela were cultured in RPMI 1640 medium supplemented with 10% fetal bovine serum (FBS), 100 U/mL penicillin, and 100 μg/mL streptomycin, at 37°C in an incubator containing 5% CO<sub>2</sub>. Cells were passaged every 2 days using Trypsin (0.25%) solution. Exponentially growing cells were used for experimentation. Cell viability assay was performed with MTT photometric analysis, as first described by Mosmann [17], with slight modification. Briefly, 4.5 × 10<sup>5</sup>/mL was seeded in 96-well microtiter plates at 200 μL/well. Twenty-four hours after treatment, the supernatant was removed. TNS were made into 20, 10, 5, 2.5, 1.25, 0.625, 0.3125, and 0.156 mg/mL in cultured solution and added to 96-well microtiter plates. Forty-eight hours after treatment, 20 μL MTT reagent (5 mg/mL) was added. After 4 h incubation at 37°C, 200 μL of DMSO was added and plates were oscillated for 10 min in a balance oscillator. The extent of the MTT reduction was measured by a plate reader at a wavelength of 570 nm. The inhibitory rate of cell proliferation was calculated as follows: inhibitory rate (%) = (1 – experimental group A value/control group A value) × 100%.

**2.9. Determination of Antioxidant Activity by DPPH Radical Scavenging Ability.** The effects of TSN on DPPH radicals were estimated according to the method of Zhao et al. [18]. Aliquots of TSN at various concentrations were mixed with 2.0 vols of 6.5 × 10<sup>-5</sup> M solution of DPPH. The mixture was shaken vigorously and left to stand at room temperature for 30 min in the dark. The absorbance of the reaction solution was measured spectrophotometrically at 517 nm. The percentages of DPPH decolorization of the samples

were calculated according to the equation: radical scavenging activity (%) =  $[1 - (\text{ABS sample}/\text{ABS control})] \times 100$ . EC<sub>50</sub> value was defined as the concentration (in mg/mL) of the extract required to deplete the amount of DPPH radical by 50%. BHT was used as a positive control.

**2.10. Antioxidant Effect of TSN in Ageing Mice.** After 1 week of acclimatization, the mice were randomly divided into five groups (12 mice per group) and i.p. injected with 0.1 mL/10 g of 5% D-galactose once daily for 6 weeks except normal control group (i.p. injected with 0.3 mL of physiological saline) [19]. Simultaneously, normal control group mice and ageing model group mice were p.o. administered with 0.1 mL/10 g of 20% Arabic gum each, respectively; TSN treatment group mice were p.o. administered with a different dose of extract (8.8, 17.5, and 35 mg/kg). Following the 35-day treatment process and 24 h after the last administration, all 50 mice were weighed then sacrificed by the humane method of cervical dislocation. The liver was surgically excised from the animal, accurately weighed, and then homogenized immediately in ice-cold 0.9% NaCl solution (0.1 g tissue/mL solution). The suspension was centrifuged at 4000 rpm/min at 4°C for 10 min, and the supernatant was collected for further analysis. The activities of SOD, GSH-Px, and the level of MDA were measured using commercially available kits in accordance with instructions.

**2.11. Analgesic Activities.** The analgesic activity of TSN was investigated using acetic acid-induced writhing response in mice, and the test was carried out using the method of Muhammad et al. [20]. This method was used to preferentially evaluate possible peripheral effects of TSN as analgesic substance. Five groups of Kunming male mice ( $n = 10$ ) were fasted overnight prior the start of the experiment, and water ad libitum. The peripheral analgesic drug, YMGP, was used as a positive control. Group 1 received the vehicle-distilled water (10 mL/kg, p.o.), and group 2 was treated with YMGP (928 mg/kg, p.o.), whereas groups 3, 4, and 5 animals were orally administered with TSN at doses of 6, 12, and 24 mg/kg. Sixty minutes after treatment, the mice were injected (i.p.) with 0.1 mL/10 g body of 0.7% acetic acid solution to induce the characteristic writhings. After 5 min, the mice were placed in an observation box, and the number of writhes in a 15 min period was counted. Antinociception (analgesia) is expressed as the reduction of the number of writhing movements between control animals (acetic-acid-treated mice) and mice pretreated with these compounds and then acetic acid.

**2.12. Anti-Inflammatory Activity.** The anti-inflammatory activity of TSN was investigated using the following models.

**2.12.1. Xylene-Induced Ear Edema in Mice.** Antiacute inflammatory activity was determined by xylene-induced mice ear edema [21]. Fifty mice were equally divided into five groups randomly including control group (distilled water), YMGP-positive control group (928 mg/kg body wt), and TSN groups (6, 12, or 24 mg/kg body wt). The vehicle and drugs

were administered orally, respectively, once per day for 3 days. One and half hour after the last administration of drugs, inflammatory response was induced on the inner and external surface of the right ear (surface: about 1 cm<sup>2</sup>) by application of 20  $\mu$ L xylene. 30 min later, mice were sacrificed by cervical dislocation and a section ( $\varnothing$  6 mm) of ears was removed from both the treated (right) and the untreated (left) ears. Edema rate was measured as the percentage of the weight difference between the two ear discs compared to the untreated (left) ears. The anti-inflammatory activity was expressed as the percentage of inhibition in treated mice compared to the control mice.

**2.12.2. Vascular Permeability Test.** The method of Koo et al. [22] was used to evaluate the effect of the extract on vascular permeability in adult albino mice of both sexes. Fifty mice were equally divided into five groups randomly including control group (distilled water), YMGP-positive control group (928 mg/kg body wt), and TSN groups (6, 12, or 24 mg/kg body wt). One and half hour after oral administration of TSN and YMGP, 0.2 mL of Evans Blue dye (0.5% in Normal saline) was intravenously administered through the tail vein. Subsequently, animals received intraperitoneal injection of 1.0 mL/100 g of acetic acid (0.6%, v/v). Treated animals were sacrificed 20 min after acetic acid injection and the peritoneal cavity washed with normal saline (6.0 mL) into heparinized tubes and centrifuged 15 min at 3000 rpm. The dye content in the supernatant was measured at 590 nm using spectrophotometer.

**2.13. Statistical Analysis.** The data obtained were computed using SPSS 11.5 software and later analyzed using ANOVA of variance. The Duncan test with significance level of 0.05 between means was used.

### 3. Results and Discussion

**3.1. Phytochemical Study.** Five compounds were isolated from TSN and elucidated as 3-O-[D-xylopyranosyl-(1  $\rightarrow$  3)- $\alpha$ -L-rhamnopyranosyl-(1  $\rightarrow$  2)- $\alpha$ -L-arabinopyranosyl]-28-O-[ $\alpha$ -L-rhamnopyranosyl-(1  $\rightarrow$  4)- $\beta$ -D-glucopyranosyl-(1  $\rightarrow$  6)- $\beta$ -D-glucopyranosyl] hederagenin (Nigella A, 1), 3-O-[D-xylopyranosyl-(1  $\rightarrow$  3)- $\alpha$ -L-rhamnopyranosyl-(1  $\rightarrow$  2)- $\alpha$ -L-arabinopyranosyl]-28-O- $\beta$ -D-glucopyranosyl hederagenin (Nigella B, 2), 3-O-[ $\alpha$ -L-rhamnopyranosyl-(1  $\rightarrow$  2)- $\alpha$ -L-arabinopyranosyl]-28-O-[ $\beta$ -D-glucopyranosyl-(1  $\rightarrow$  6)- $\beta$ -D-glucopyranosyl] hederagenin (Nigella C, 3), 3-O-[D-xylopyranosyl-(1  $\rightarrow$  3)- $\alpha$ -L-rhamnopyranosyl-(1  $\rightarrow$  2)- $\alpha$ -L-arabinopyranosyl]-28-O-[ $\alpha$ -L-rhamnopyranosyl-(1  $\rightarrow$  4)- $\beta$ -D-glucopyranosyl-(1  $\rightarrow$  6)- $\beta$ -D-glucopyranosyl] hederagenin (Nigella D, 4), and nigelanoside (5) by extensive spectroscopic methods including 1D-(<sup>1</sup>H, <sup>13</sup>C) NMR experiments (Figure 1). All data were compared with those in previous literature [23–25]. Among these compounds, the content of Nigella A from TSN was far above others. Therefore, the content of Nigella A was determined by HPLC-ELSD, and its content is  $60.36 \pm 2.5$  g/100 g in TSN (Figure 2).




FIGURE 1: Chemical structure of compounds.

**3.2. Acute Toxicity.** Following administration of TSN, the mice were monitored daily for mortality, clinical signs, and gross changes in appearance and behavior. No deaths occurred in the test and control group. There were no significant differences in body weights between the TSN treated and control group in either sex (data not reported in this paper). The result indicated that treatment of TSN was safe under the maximum dose at 48 g TSN/kg body.

**3.3. Cytotoxicity Activity.** Two cell lines, that is, Bel-7402, and Hela cells, were used to test the effect of TSN on their cell proliferation. TSN showed significant inhibitory effects on these two cell lines, which were in a dose-dependant manner. TSN at the dose of 20 mg/mL showed 86.75% and 88.26% inhibition of the growth of Bel-7402 and Hela cells, respectively (Figure 3).

**3.4. DPPH Radical Scavenging Activity.** The scavenging capability of DPPH radical was determined by the decrease in




FIGURE 2: HPLC chromatogram of TSN solution.

its absorbance at 517 nm under effect of antioxidants. Due to rapid hydrogen-accepting ability, DPPH reacted with antioxidants and converted into 1,1-diphenyl-2-picrylhydrazin, shows decrease in absorbance simultaneously [26]. The degree of discoloration indicates the scavenging potential of the antioxidant extracts [27]. Figure 4 showed the DPPH free radical scavenging activities of TSN. DPPH was reduced

TABLE 1: Effect of TSN on MDA, SOD, and GSH-Px in liver in D-galactose-induced ageing mice.

Group	Dose (mg/kg-d)	MDA (nmol/mgprot)	SOD (U/mgprot)	GSH-Px (U/mgprot)
Control	—	1.181 ± 0.187	259.785 ± 34.007	577.273 ± 95.582
Model	—	1.789 ± 0.383 <sup>##</sup>	173.843 ± 21.939 <sup>##</sup>	270.775 ± 35.582 <sup>##</sup>
TSN	8.8	1.296 ± 0.206 <sup>**</sup>	181.220 ± 13.194	300.344 ± 26.338 <sup>**</sup>
	17.5	1.252 ± 0.398 <sup>**</sup>	194.609 ± 18.456 <sup>*</sup>	354.958 ± 69.549 <sup>**</sup>
	35	1.236 ± 0.315 <sup>**</sup>	203.477 ± 17.719 <sup>**</sup>	411.949 ± 57.553 <sup>**</sup>

Values are expressed as mean ± SD ( $n = 12$ ).

<sup>##</sup> $P < 0.01$ , compared with control group; <sup>\*</sup> $P < 0.05$ , <sup>\*\*</sup> $P < 0.01$ , compared with model group.




FIGURE 3: Inhibiting effect of TSN on human hepatoma carcinoma cell lines Bel 7402 and cervical carcinoma cell lines Hela growth. Results are mean ± SD ( $n = 4$ ).




FIGURE 4: DPPH radical scavenging activity of TSN. Results are mean ± SD ( $n = 3$ ).

by TSN in a concentration-dependent manner. Figure 4 shows the dose response curves of DPPH radical scavenging activities of TSN, at a concentration range of 0.2–1.6 mg/mL, and percent inhibition and the  $IC_{50}$  of TSN were 21.87–53.28% and 1.58 mg/mL, respectively. BHT as a reference drug scavenged the DPPH radical by inhibitions of 60.31–74.01% and the  $IC_{50}$  value of 58.07  $\mu$ g/mL.

**3.5. The Effect of TSN on the Activities of SOD, GSH-Px, and MDA in the Liver of Aged Mice.** Ageing is a natural process in all living organisms, and oxidative stress is one of root causes of the ageing process [28]. The importance of antioxidant enzymes is generally emphasized in the prevention of oxidative stresses by scavenging of ROS [29, 30]. The antioxidant system comprises several enzymes such as SOD and GSH-Px [31–33]. The effect of TSN on the activities of SOD, GSH-Px, and the levels of MDA in the liver of aged mice is shown in Table 1. A marked increase in MDA and significant decrease ( $P < 0.05$ ) of antioxidant enzymes activity (SOD, GSH-Px) were observed in the liver between the treatments of normal control group and model control group. TSN treatment significantly inhibited ( $P < 0.05$ ) the formation of MDA in the liver and raised the activity of antioxidant enzymes in a dose-dependent manner. The administration of TSN to the D-galactose-treated mice with 8.8, 17.5, and 35 mg/kg increased both SOD and GSH-Px enzymatic antioxidant liver activity ( $P < 0.01$  and  $P < 0.05$ ).

**3.6. Anti-Inflammatory and Analgesic Activities.** The peripheral analgesic effect may be mediated through inhibition of cyclo-oxygenases and/or lipoxygenases (and other inflammatory mediators), while the central analgesic action may be mediated through inhibition of central pain receptors [34]. The acetic-acid-induced writhes have often been used to evaluate the peripherally and centrally acting analgesic drugs [35]. YMGP is the most popular medication for rheumatism arthritis pain, sciatica, and prosopalgia in Chinese medicine. Therefore, we select YMGP as a reference drug.

Peripheral analgesic activity was assessed by acetic-acid-induced writhing test, which showed significant ( $P < 0.01$  and  $P < 0.05$ ) suppression of writhes (Table 2). Acetic acid is known to trigger the production of noxious substances within the peritoneum, which induces the writhing response. The effect of TSN against the noxious stimulus may be an indication that it depressed the production of irritants and thereby bringing a reduction in the number of writhes in animals. Injection of acetic acid into the control mice resulted in  $71 \pm 19$  writhes. Pretreatment with TSN at the doses of 6, 12, and 24 mg/kg reduced the number of writhes to  $38 \pm 8$  (46.5% inhibition),  $33 \pm 9$  (53.6% inhibition), and  $29 \pm 8$  (59.2% inhibition) at a dose-activity dependence relationship, respectively. Interestingly, the analgesic activity of high dose was higher than that of YMGP ( $33 \pm 8$  writhes, 54.5% inhibition). It was also observed that the onset of writhing

TABLE 2: Effects of TSN on acetic-acid-induced writhing response in mice.

Treatment group	Dose (mg/kg)	Number of writhes	Inhibition (%)
Control	—	71 ± 19	—
YMGP	928	33 ± 8**	54.5
	6	38 ± 8**	46.5
TSN	12	33 ± 9*	53.6
	24	29 ± 8**	59.2

Values are expressed as mean ± SD ( $n = 10$ ).

Control (vehicle): distilled water; YMGP: Yemugua tablet; TSN: total saponins from *Nigella glandulifera*. \*\* $P < 0.01$ , \* $P < 0.05$ , compared with corresponding control.

was delayed and the duration of writhing was shortened with TSN pretreatment.

Xylene-induced mouse ear edema reflects the oedematization during the early stages of acute inflammation, which was probably related to the release and inhibition of the inflammation factors [36]. In xylene-induced ear edema, the oral administration of TSN suppressed significantly xylene-induced ear oedema in mice. The oedema inhibitory rates of casticin were 16.6%, 25.9%, and 35.5% at doses of 6, 12 and 24 mg/kg, respectively, whereas YMGP (928 mg/kg) produced 30.53% inhibitory rate compared with control (Figure 5).

The inflammatory response is a physiological characteristic of vascularized tissues also [37]. Exudation, which is a consequence of increased vascular permeability, is considered a major feature of acute inflammation. In the inflammatory reaction, increased vascular permeability leads to exudation of fluid rich in plasma proteins including immunoglobulins, coagulation factors, and cells into the injured tissues [38] (with subsequent edema at the site). Therefore, inhibition of increased vascular permeability can modulate the extent and magnitude of the inflammatory reaction [39, 40]. Chemical-induced vascular permeability causes an immediate sustained reaction that is prolonged over 24 h, and its inhibition suggests that the extract may effectively suppress the exudative phase of acute inflammation [41], and acetic-acid-induced increased vascular permeability in mouse model is a typical capillary permeability assay [42, 43]. In this study, TSN significantly reduced the increased peritoneal vascular permeability, indicating the suppression of the vascular response in the process of acute inflammation (Figure 6). Oral administration of 6, 12, and 24 mg/kg of TSN evoked a significant ( $P < 0.05$ ) dose-related inhibition of vascular permeability induced by acetic acid in mice. At the doses of 6, 12, and 24 mg/kg, TSN produced 17.9%, 23.9%, and 29.4% inhibition of dye leakage, respectively, with dose-dependent manner, while YMGP produced 26.9% inhibition.

#### 4. Conclusion

In summary, our results suggest that TSN possesses anti-inflammatory, analgesic, antitumor, and free radical scavenging activities and provide some scientific evidence to support




FIGURE 5: Effects of TSN and YMGP on xylene-induced ear oedema in mice. Values are mean ± SD of differences in weight between right and left ears of animals.  $n = 10$ . Control (vehicle): distilled water; YMGP: Yemugua tablet; TSN: total saponins from *Nigella glandulifera*. \* $P < 0.05$  and \*\* $P < 0.01$ , compared with corresponding control.




FIGURE 6: Effects of TSN and YMGP on acetic-acid-induced increased vascular permeability in mice. Values are mean ± SD,  $n = 10$ . Control (vehicle): distilled water; YMGP: Yemugua tablet; TSN: total saponins from *Nigella glandulifera*. \* $P < 0.05$  and \*\* $P < 0.01$ , compared with corresponding control.

the folk medicinal utilization of *N. glandulifera*. Furthermore, TSN is a good candidate for the development of new anti-inflammatory and analgesic medicine compared with efficacy of YMGP. Therefore, TSN may be worth further investigating and elucidating.

#### Conflict of Interests

All authors declare that they have no competing interests.

#### Acknowledgment

This work was supported by a grant from the project science and technology support programme of the Xinjiang Uygur Autonomous Region in China (no. 201033121).

#### References

- [1] L. F. D'Antuono, A. Moretti, and A. F. S. Lovato, "Seed yield, yield components, oil content and essential oil content and composition of *Nigella sativa* L. and *Nigella damascena* L.," *Industrial Crops and Products*, vol. 15, no. 1, pp. 59–69, 2002.

- [2] M. F. Ramadan, "Nutritional value, functional properties and nutraceutical applications of black cumin (*Nigella sativa* L.): an overview," *International Journal of Food Science and Technology*, vol. 42, no. 10, pp. 1208–1218, 2007.
- [3] B. H. Ali and G. Blunden, "Pharmacological and toxicological properties of *Nigella sativa*," *Phytotherapy Research*, vol. 17, no. 4, pp. 299–305, 2003.
- [4] L. S. Mohamed, "Immunomodulatory and therapeutic properties of the *Nigella sativa* L. seed," *International Immunopharmacology*, vol. 5, pp. 1749–1770, 2005.
- [5] G. Fico, L. Panizzi, G. Fiamin et al., "Biological screening of *Nigella damascena* for antimicrobial and molluscicidal activities," *Phytotherapy Research*, vol. 18, no. 6, pp. 468–470, 2004.
- [6] I. Merfort, V. Wray, H. H. Barakat, S. A. M. Hussein, M. A. M. Nawwar, and G. Willuhn, "Flavonol triglycosides from seeds of *Nigella sativa*," *Phytochemistry*, vol. 46, no. 2, pp. 359–363, 1997.
- [7] L. Kokoska, J. Havlik, I. Valterova, A. Nepovim, V. Rada, and T. Vanek, "Chemical composition of the essential oil of *Nigella orientalis* L. seeds," *Flavour and Fragrance Journal*, vol. 20, no. 4, pp. 419–420, 2005.
- [8] Z. Tian, Y. M. Liu, S. B. Chen et al., "Cytotoxicity of two triterpenoids from *Nigella glandulifera*," *Molecules*, vol. 11, no. 9, pp. 693–699, 2006.
- [9] B. Dinda, S. Debnath, B. C. Mohanta, and Y. Harigaya, "Naturally occurring triterpenoid saponins," *Chemistry and Biodiversity*, vol. 7, no. 10, pp. 2327–2580, 2010.
- [10] J. M. Augustin, V. Kuzina, S. B. Andersen, and S. Bak, "Molecular activities, biosynthesis and evolution of triterpenoid saponins," *Phytochemistry*, vol. 72, no. 6, pp. 435–457, 2011.
- [11] Editorial Committee of Flora of China, Chinese Academy of Science, *Flora of China*, vol. 27, Science Press, Urumqi, China, 1979.
- [12] Chinese Pharmacopoeia Commission of Sanitary Ministry of People's Republic of China, *Chinese Pharmacopoeia, Part 1*, Chemical Industry Publishing House, Beijing, China, 2005.
- [13] Y. M. Liu, *Pharmacography of Uighur, Part 1*, Xinjiang Science & Technology & Hygiene Publishing House, Urumuqi, China, 1999.
- [14] H. F. Hao, L. J. Ren, and Y. W. Chen, "Studies on the chemical constituents of seed from *Nigella glandulifera*," *Acta Pharmaceutica Sinica*, vol. 31, pp. 689–694, 1996.
- [15] X. L. Xin, H. A. Aisa, and H. Q. Wang, "Flavonoids and phenolic compounds from seeds of the Chinese plant *Nigella glandulifera*," *Chemistry of Natural Compounds*, vol. 44, no. 3, pp. 368–369, 2008.
- [16] Center for Drug Evaluation of State Food and Drug Administration of People's Republic of China, "Technical guidelines of acute toxicity for traditional Chinese medicine and natural drugs," SFDA guidelines No. [Z] GPT2-1.
- [17] T. Mosmann, "Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays," *Journal of Immunological Methods*, vol. 65, no. 1-2, pp. 55–63, 1983.
- [18] J. Zhao, T. Liu, L. Ma et al., "Antioxidant and preventive effects of extract from *Nymphaea candida* flower on *in vitro* immunological liver Injury of rat primary hepatocyte cultures," *Evidence-Based Complementary and Alternative Medicine*, vol. 2011, Article ID 497673, 8 pages, 2011.
- [19] J. H. Liu, S. C. Ho, T. H. Lai, T. H. Liu, P. Y. Chi, and R. Y. Wu, "Protective effects of Chinese herbs on D-galactose-induced oxidative damage," *Methods and Findings in Experimental and Clinical Pharmacology*, vol. 25, no. 6, pp. 447–452, 2003.
- [20] A. Muhammad, G. Gaurav, K. Imran et al., "Anti-inflammatory and analgesic potential of a novel steroidal derivative from *Bryophyllum pinnatum*," *Fitoterapia*, vol. 83, pp. 853–858, 2012.
- [21] H. Hosseinzadeh, M. Ramezani, and G. A. Salmani, "Antinociceptive, anti-inflammatory and acute toxicity effects of *Zataria multiflora* Boiss extracts in mice and rats," *Journal of Ethnopharmacology*, vol. 73, no. 3, pp. 379–385, 2000.
- [22] H. J. Koo, K. H. Lim, H. J. Jung, and E. H. Park, "Anti-inflammatory evaluation of gardenia extract, geniposide and genipin," *Journal of Ethnopharmacology*, vol. 103, no. 3, pp. 496–500, 2006.
- [23] Y. M. Liu, J. S. Yang, and Q. H. Liu, "Studies on chemical constituents in the seeds of *Nigella glandulifera*," *China Journal of Chinese Materia Medica*, vol. 30, no. 13, pp. 893–896.
- [24] J. J. Ni, H. Y. Gao, B. H. Sun, Z. X. Wang, and L. J. Wu, "Chemical constituents of active extracts from the seeds of *Nigella glandulifera* Freyn," *Asian Journal of Traditional Medicines*, vol. 1, pp. 3–4, 2006.
- [25] I. Merfort, V. Wray, H. H. Barakat, S. A. M. Hussein, M. A. M. Nawwar, and G. Willuhn, "Flavonol triglycosides from seeds of *Nigella sativa*," *Phytochemistry*, vol. 46, no. 2, pp. 359–363, 1997.
- [26] K. Marxen, K. H. Vanselow, S. Lippemeier, R. Hintze, A. Ruser, and U. P. Hansen, "Determination of DPPH radical oxidation caused by methanolic extracts of some microalgal species by linear regression analysis of spectrophotometric measurements," *Sensors*, vol. 7, no. 10, pp. 2080–2095, 2007.
- [27] M. Philip, "The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity," *Songklanakarinn Journal of Science and Technology*, vol. 26, pp. 211–219, 2004.
- [28] K. U. Ansari, "Free radical induced diseases," *Journal of the Indian Medical Association*, vol. 94, no. 6, pp. 238–239, 1996.
- [29] P. Uma Devi and P. Chinnaswamy, "Age and gender related changes in free radical pathology and antioxidant defense in Schizophrenia," *Asian Journal of Biochemistry*, vol. 3, no. 3, pp. 143–152, 2008.
- [30] G. Vendemiale, I. Grattagliano, and E. Altomare, "An update on the role of free radicals and antioxidant defense in human disease," *International Journal of Clinical and Laboratory Research*, vol. 29, no. 2, pp. 49–55, 1999.
- [31] M. Kasapoglu and T. Özben, "Alterations of antioxidant enzymes and oxidative stress markers in aging," *Experimental Gerontology*, vol. 36, no. 2, pp. 209–220, 2001.
- [32] B. B. Warner, L. Stuart, S. Gebb, and J. R. Wispe, "Redox regulation of manganese superoxide dismutase," *American Journal Physiology*, vol. 271, pp. 150–158, 1996.
- [33] C. Ke, D. Qiao, D. Gan, Y. Sun, H. Ye, and X. Zeng, "Antioxidant activity *in vitro* and *in vivo* of the capsule polysaccharides from *Streptococcus equi* subsp. *zooepidemicus*," *Carbohydrate Polymers*, vol. 75, no. 4, pp. 677–682, 2009.
- [34] H. Matsumoto, H. Naraba, A. Ueno et al., "Induction of cyclooxygenase-2 causes an enhancement of writhing response in mice," *European Journal of Pharmacology*, vol. 352, no. 1, pp. 47–52, 1998.
- [35] R. M. Gené, L. Segura, T. Adzet, E. Marin, and J. Iglesias, "Heterotheca inuloides: anti-inflammatory and analgesic effect," *Journal of Ethnopharmacology*, vol. 60, no. 2, pp. 157–162, 1998.
- [36] C. O. Okoli and P. A. Akah, "Mechanisms of the anti-inflammatory activity of the leaf extracts of *Culcasia scandens* P. Beauv (Araceae)," *Pharmacology Biochemistry and Behavior*, vol. 79, no. 3, pp. 473–481, 2004.

- [37] C. R. Wu, Y. C. Hseu, J. C. Lien, L. W. Lin, Y. T. Lin, and H. Ching, "Triterpenoid contents and anti-inflammatory properties of the methanol extracts of *Ligustrum* species leaves," *Molecules*, vol. 16, no. 1, pp. 1–15, 2011.
- [38] C. O. Okoli, P. A. Akah, S. V. Nwafor, A. I. Anisiobi, I. N. Ibegbunam, and O. Erojikwe, "Anti-inflammatory activity of hexane leaf extract of *Aspilia africana* C.D. Adams," *Journal of Ethnopharmacology*, vol. 109, no. 2, pp. 219–225, 2007.
- [39] K. B. Glaser, M. L. A. Sung, D. A. Hartman et al., "Cellular and topical *in vivo* inflammatory murine models in the evaluation of inhibitors of phospholipase A2," *Skin Pharmacology*, vol. 8, no. 6, pp. 300–308, 1995.
- [40] A. P. Rogerio, C. Fontanari, M. C. C. Melo et al., "Anti-inflammatory, analgesic and anti-oedematous effects of *Lafoensia pacari* extract and ellagic acid," *Journal of Pharmacy and Pharmacology*, vol. 58, no. 9, pp. 1265–1273, 2006.
- [41] P. D. Collins, D. T. Connolly, and T. J. Williams, "Characterization of the increase in vascular permeability factor *in vivo*," *British Journal of Pharmacology*, vol. 109, no. 1, pp. 195–199, 1993.
- [42] X. H. Liu, A. Kirschenbaum, S. Yao et al., "Upregulation of vascular endothelial growth factor by cobalt chloride-simulated hypoxia is mediated by persistent induction of cyclooxygenase-2 in a metastatic human prostate cancer cell line," *Clinical and Experimental Metastasis*, vol. 17, no. 8, pp. 687–694, 1999.
- [43] S. Karim, E. Berrou, S. Lévy-Toledano, M. Bryckaert, and J. Maclouf, "Regulatory role of prostaglandin E2 in induction of cyclo-oxygenase-2 by a thromboxane A2 analogue (U46619) and basic fibroblast growth factor in porcine aortic smooth-muscle cells," *Biochemical Journal*, vol. 326, no. 2, pp. 593–599, 1997.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

