Research Article

Pleurotus eryngii Ameliorates Lipopolysaccharide-Induced Lung Inflammation in Mice

Junya Kawai,¹ Tsugunobu Andoh,² Kenji Ouchi,¹ and Satoshi Inatomi¹

¹ Mushroom Research Laboratory, Hokuto Corporation, 800-8 Shimokomazawa, Nagano 381-0008, Japan

² Department of Applied Pharmacology, Graduate School of Medicine and Pharmaceutical Sciences, University of Toyama, Toyama 930-0194, Japan

Correspondence should be addressed to Tsugunobu Andoh; andoht@pha.u-toyama.ac.jp

Received 2 December 2013; Revised 17 February 2014; Accepted 4 March 2014; Published 1 April 2014

Academic Editor: Evelin Tiralongo

Copyright © 2014 Junya Kawai et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Pleurotus eryngii (*P. eryngii*) is consumed as a fresh cultivated mushroom worldwide and demonstrated to have multiple beneficial effects. We investigated the anti-inflammatory effect of *P. eryngii* in mice with acute lung injury (ALI). Intranasal instillation of lipopolysaccharide (LPS) (10 µg/site/mouse) induced marked lung inflammation (increase in the number of inflammatory cells, protein leakage, and production of nitric oxide in bronchoalveolar lavage fluid) as well as histopathological damage in the lung, 6 h after treatment. Mice administered heat-treated *P. eryngii* (0.3–1 g/kg, p.o. (HTPE)) 1 h before LPS challenge showed decreased pulmonary inflammation and ameliorated histopathological damage. These results suggest that HTPE has anti-inflammatory effects against ALI. Thus, *P. eryngii* itself may also have anti-inflammatory effects and could be a beneficial food for the prevention of ALI induced by bacterial infection.

1. Introduction

Pleurotus eryngii is an edible mushroom native to Europe. Its natural habitat is the dead roots of the weed *Eryngium campestre*. *P. eryngii* is cultivated widely, and its production has been increasing in Asia, including Japan [1]. *P. eryngii* is considered to be a health food because it is low in fat and calories but rich in amino acids, vitamins, and dietary fiber. *P. eryngii* is also bioactive, with hypolipidemic [2], antitumor [3], antioxidant [4–6], and antiallergic activities [1]. In particular, *in vitro* studies have demonstrated that the antiallergic activity of *P. eryngii* is caused by the down-regulation of allergy-related signaling proteins (including inflammation-related proteins) by inhibition of the nuclear factor of activated T cells, nuclear factor-kappa B (NF-κB), and high-affinity immunoglobulin E receptor (FcεRI) mediated signaling in antigen-stimulated mast cells. However, whether *P. eryngii* has clinical effectiveness remains to be determined.

Acute respiratory distress syndrome is a result of acute inflammation of the lung and noncardiogenic pulmonary

edema that often leads to multiorgan system failure and death [7, 8]. Lipopolysaccharide (LPS) is present in the outer membrane of Gram-negative bacteria. LPS can cause acute inflammation of the lung because of neutrophil recruitment and pulmonary edema [9, 10]. Intranasal instillation of LPS in mice (as an animal model of acute lung injury (ALI)) has been shown to result in the release of proinflammatory cytokines, which cause aggregation of inflammatory cells and, consequently, injury to lung tissue [11, 12]. There have been reports with regard to these mechanisms and the features of this model. Such reports have shown that LPS activates alveolar macrophages directly and stimulates neutrophils to migrate into the lung and that the proinflammatory mediators released from these inflammatory cells recruit lymphocytes to the lung [7, 13]. The critical feature of LPS-induced ALI is the destruction of vascular integrity and the subsequent upregulated permeability results in protein leakage and pulmonary edema [9, 10, 14]. It has also been reported that nitric oxide (NO) has important roles in the pathogenesis of ALI because inhibitors of NO synthase inhibit LPS-induced damage [9, 15–17].

FIGURE 1: The fresh fruiting body of *Pleurotus eryngii* and freeze-dried powder of heat-treated *P. eryngii* (HTPE).

The aim of this study was to examine the anti-inflammatory effect of *P. eryngii* *in vivo* using an LPS-induced ALI model in mice. We used the whole *P. eryngii* to illustrate the functional utility of this mushroom as food.

2. Materials and Methods

2.1. Animals. Male BALB/c mice (6 weeks of age; Japan SLC, Ltd., Shizuoka, Japan) were used. They were kept under controlled temperature (21–23°C) and humidity (45–65%). The room was lit from 7:00 am to 7:00 pm and during the behavioral test. Food and water were available *ad libitum*. The study was approved by the Committee for Animal Experiments at the University of Toyama (Toyama, Japan).

2.2. Agents. LPS (*Escherichia coli* 0111:B4), dexamethasone, and pentobarbital sodium were purchased from Sigma-Aldrich (St. Louis, MO, USA). Other agents used in this study were purchased from Wako Pure Chemical Industries Ltd. (Osaka, Japan). LPS was dissolved in phosphate-buffered saline (PBS) and instilled intranasally. Dexamethasone, which was dissolved in saline with 10% ethanol, was given to mice in the dexamethasone group by intraperitoneal injection 1 h before LPS administration. Pentobarbital sodium was dissolved in saline containing 0.4% propylene glycol and 1.05% ethanol, and given by intraperitoneal injection.

2.3. Preparation of *P. eryngii* Intakes. The fruiting body of *P. eryngii* (Figure 1) was obtained from Hokuto Co. (Nagano, Japan). It was then cut into small pieces and boiled in an equal amount of distilled water for 10 min. Heat-treated *P. eryngii* (HTPE) was freeze-dried and powdered. HTPE was resuspended in tap water and administered orally 1 h before intranasal administration of LPS.

2.4. LPS-Induced ALI Model in Mice. BALB/c mice were challenged with intranasal instillation (i.n.) of LPS (10 µg in 50 µL PBS per mouse) to induce lung inflammation. Control mice were given PBS (i.n.) without LPS. After 6 h, collection of bronchoalveolar lavage fluid (BALF) was carried out following the method of Chu et al. under anesthesia (sodium

pentobarbital, 80 mg/kg, i.p.) [14]. After centrifugation (25×g, 4°C, 5 min), cell pellets were resuspended in PBS for total cell counts using a hemacytometer. The supernatant was used for NO analyses and protein analyses.

2.5. Measurement for Protein Concentration. The protein concentration in BALF was determined using a protein assay kit (Bio-Rad Laboratories, Hercules, CA, USA).

2.6. Measurement for NO Production. A metabolite of NO, nitrite (NO²⁻), in BALF was measured using Griess reagent (1% sulfanilamide, 0.1% N-(1-naphthyl)ethylenediamine dihydrochloride, 2.5% phosphoric acid). Briefly, 50 µL of BALF and 50 µL of Griess reagent were mixed in a 96-well plate. The azo dye formed was determined with a spectrophotometer (Multiskan FC; Thermo Fisher Scientific K.K., Yokohama, Japan) at 540 nm using sodium nitrite as the standard.

2.7. Hematoxylin and Eosin (H&E) Staining. Anesthesia (sodium pentobarbital, 80 mg/kg, i.p.) was induced in mice 6 h after LPS treatment. Mice were then decapitated. Lungs were removed and, after washing with PBS, placed in 10% formalin solution. Preparation of paraffin-embedded sections and staining with H&E were undertaken using standard procedures. Staining was observed under a light microscope (AX80; Olympus, Osaka, Japan) with a charge-coupled device camera (Axio Cam; Carl Zeiss, Jena, Germany).

2.8. Immunohistochemical Staining. Anesthesia (sodium pentobarbital, 80 mg/kg, i.p.) was induced in mice 6 h after LPS treatment. Mice were then decapitated. Lungs were removed and, after washing with PBS, placed in 10% formalin solution. Preparation of paraffin-embedded sections and deparaffinization were undertaken using standard procedures. Deparaffinized sections were treated with methanol containing 0.3% hydrogen peroxide and then with 0.2% Triton X-100 in PBS. After treatment with 0.3% fetal bovine serum in PBS to block immunoglobulin binding, sections were incubated with rabbit anti-myeloperoxidase (MPO) antibody (DAKO, Glostrup, Denmark) or rat anti-Mac-2 antibody (Cedarlane, Ontario, Canada) at 4°C overnight, followed by horseradish peroxidase-conjugated anti-rabbit IgG antibody (DAKO) or horseradish peroxidase-conjugated anti-rat IgG antibody (DAKO). Color was developed using DAKO liquid with a 3,3-diaminobenzidine tetrahydrochloride (DAB) substrate chromogen system (DAKO) and counterstained with hematoxylin. These stained sections were observed under a light microscope (AX-80; Olympus) coupled to a CCD camera (Axio Cam; Carl Zeiss).

2.9. Statistical Analyses. Data are presented as the mean ± standard error of the mean (SEM). Statistical significance between groups was assessed using one-way analysis of variance and *post hoc* Holm-Sidak multiple comparisons. *P* < 0.05 was considered significant. Statistical analyses were done using Sigmaplot v11 (Systat Software, Inc., Chicago, IL, USA).

FIGURE 2: Effect of HTPE on the total number of cells in the BALF of LPS-induced ALI mice. Mice were administered, via the oral and intraperitoneal routes, HTPE (0.1–1 g/kg) and dexamethasone (DEX: 5 mg/kg, positive control group), respectively, 1 h before intranasal administration of LPS (10 μ g/site). Vehicle (VH: tap water, vehicle for HTPE) was also administered orally 1 h before intranasal administration of LPS (LPS group) or saline (vehicle for LPS, negative control group). BALF was collected 6 h after LPS challenge to measure the total number of cells. Values are presented as the mean and SEM ($n = 6$ for VH or HTPE; $n = 4$ for DEX). [#] $P < 0.05$ versus negative control group (VH + saline), ^{*} $P < 0.05$ versus LPS group (VH + LPS) (Holm-Sidak multiple comparisons).

3. Results

3.1. Effects of HTPE on the Aggregation of Inflammatory Cells Induced by LPS in the Lung. LPS-induced ALI resulted in an increase in the number of inflammatory cells, such as neutrophils and macrophages, in the lung [14]. Six hours after intranasal instillation of LPS, the number of inflammatory cells in BALF increased significantly compared with intranasal instillation of saline (vehicle for LPS) (Figure 2). Oral pretreatment with HTPE (0.1–1.0 g/kg) significantly inhibited the LPS-induced increase in the number of cells in BALF compared with that of the vehicle (for HTPE) pretreated group (Figure 2). Intraperitoneal pretreatment with dexamethasone (5 mg/kg) also significantly decreased the number of cells in the BALF of LPS-treated mice compared with that of the vehicle (for HTPE) pretreated group (Figure 2).

3.2. Effect of HTPE on LPS-Induced Vascular Permeability in the Lung. One of the major pathological changes observed in LPS-induced ALI is increased vascular permeability, which results in increased protein leakage in BALF [14]. Six hours after intranasal instillation of LPS, the concentration of protein in BALF increased significantly compared with the intranasal instillation of saline (vehicle for LPS) (Figure 3). Oral pretreatment with HTPE (0.1–1.0 g/kg) significantly inhibited LPS-induced protein leakage in BALF compared with that of the vehicle (for HTPE) pretreated group (Figure 3). Intraperitoneal pretreatment with dexamethasone (5 mg/kg) also significantly attenuated the concentration of protein in the BALF of LPS-treated mice compared with that of the vehicle (for HTPE) pretreated group (Figure 3).

3.3. Effect of HTPE on LPS-Induced NO Production in the Lung. It has been reported that NO plays an important part in the pathogenesis of ALI [8, 15–17]. Six hours after

intranasal instillation of LPS, the concentration of nitrite and metabolites of NO in BALF increased significantly with intranasal instillation of saline (vehicle for LPS) (Figure 4). Oral pretreatment with HTPE (0.1–1.0 g/kg) significantly inhibited LPS-induced NO production in the lung compared with that of the vehicle (for HTPE) pretreated group (Figure 4). Intraperitoneal pretreatment with dexamethasone (5 mg/kg) showed a tendency toward inhibition of LPS-induced NO production, but not in a significant manner ($P = 0.051$) (Figure 4).

3.4. Effect of HTPE on LPS-Induced Pulmonary Histopathological Changes. To evaluate the histopathological changes in LPS-treated mice, lung sections 6 h after LPS treatment were stained with H&E. Normal pulmonary histology was observed in mice that underwent intranasal instillation of saline (vehicle for LPS) (Figure 5(a)). LPS-treated lungs exhibited an apparent increase in infiltration of inflammatory cells, interstitial edema, and hyperemic thickening of the alveolar wall (Figure 5(b)). In particular, the infiltrated inflammatory cells in LPS-treated lung were neutrophils (MPO-immunoreactive cells) (Figure 5(c)) and macrophages (Mac-2-immunoreactive cells) (Figure 5(d)). However, these histopathological changes in lungs treated with LPS were ameliorated by pretreatment with HTPE (1 g/kg) (Figure 5(e)) or dexamethasone (5 mg/kg) (Figure 5(f)).

4. Discussion

We evaluated the anti-inflammatory activities of HTPE using a LPS-induced model of ALI in mice. Pretreatment with HTPE as well as dexamethasone decreased the number of total cells, protein leakage, and NO production in the BALF of LPS-induced ALI mice. Histopathological studies revealed infiltration of inflammatory cells (such as neutrophils and macrophages), interstitial edema, and thickening

FIGURE 3: Effect of HTPE on total protein concentration in the BALF of LPS-induced ALI mice. Mice were administered, via the oral and intraperitoneal routes, HTPE (0.1–1 g/kg) and dexamethasone (DEX: 5 mg/kg, positive control group), respectively, 1 h before intranasal administration of LPS (10 μ g/site). Vehicle (VH: tap water, vehicle for HTPE) was also administered orally 1 h before intranasal administration of LPS (LPS group) or saline (vehicle for LPS, negative control group). BALF was collected 6 h after LPS challenge to measure total protein concentration. Values are presented as the mean and SEM ($n = 6$ for VH or HTPE; $n = 4$ for DEX). [#] $P < 0.05$ versus negative control group (VH + saline), ^{*} $P < 0.05$ versus LPS group (VH + LPS) (Holm-Sidak multiple comparisons).

FIGURE 4: Effect of HTPE on production of nitric oxide in the BALF of LPS-induced ALI mice. Mice were administered, via the oral and intraperitoneal routes, HTPE (0.1–1 g/kg) and dexamethasone (DEX: 5 mg/kg, positive control group), respectively, 1 h before intranasal administration of LPS (10 μ g/site). Vehicle (VH: tap water, vehicle for HTPE) was also administered orally 1 h before intranasal administration of LPS (LPS group) or saline (vehicle for LPS, negative control group). BALF was collected 6 h after LPS challenge to measure the nitrite (a metabolite of nitric oxide) concentration. Values are presented as the mean and SEM ($n = 6$ for VH or HTPE; $n = 4$ for DEX). [#] $P < 0.05$ versus negative control group (VH + saline), ^{*} $P < 0.05$ versus LPS group (VH + LPS) (Holm-Sidak multiple comparisons).

of the alveolar walls in the lungs of LPS-induced ALI mice. These histopathological changes were also prevented in mice given HTPE and dexamethasone. These results suggest that HTPE improves the lung injury induced by LPS in mice through inhibition of the recruitment of inflammatory cells and overproduction of NO.

Mushrooms are low in calories, abundant in amino acids, vitamins, and dietary fiber, and are popular foods worldwide. It has been reported that some edible mushrooms have a wide range of pharmaceutical properties, including anti-inflammatory and antioxidant activities [4–6, 18–20].

It is well known that *P. eryngii* contains mainly β -glucans, including β -(1,3)-(1,6)-glucans [21, 22]. The *P. eryngii* (per 100 g of fresh fruiting body) used in the present study contained β -glucan (1.9 g). Polysaccharides such as β -glucans are known to be biologically active substances [23–25]. β -Glucans have been reported to possess immunomodulatory/immunostimulatory activities, and β -glucans usually have a β -(1,3)-linked main chain and β -(1,6)-linked branches [23, 24, 26, 27]. The frequency of branching varies, and immunomodulatory/immunostimulatory activities are dependent upon the structure of β -glucans [28, 29]. It has been reported that β -(1,6) branches could contribute to the stimulatory activity of β -glucans [28]. In the present study, we did not have information on the function and structure of β -glucans. However, *P. eryngii* contains β -(1,3)-(1,6)-glucans [21, 22]. Lentinan is a β -glucan from the fruiting bodies of *Lentinus edodes* and a β -(1,3)-glucan

with β -(1,6) branching. Lentinan has been shown to suppress LPS-induced secretion of NO and tumor necrosis factor- α from RAW264.7 macrophages [25]. Thus, β -(1,3)-(1,6)-glucans in HTPE may have important roles in anti-inflammatory actions and NO production in lungs treated with LPS. In the present study, nitrite concentration was correlated with total cell numbers in BALF ($r = 0.716$, $P \leq 0.001$). However, the decrease in nitrite concentration by HTPE was not completely correlated with the decrease in total cell numbers. β -Glucans inhibit LPS-induced NO production [25]. Thus, it is suggested that the decrease in nitrite concentration by HTPE is associated with the decrease in total cell numbers and inhibition of NO production by HTPE in inflammatory cells. The *P. eryngii* (per 100 g of fresh fruiting body) used in the present study also contained ergosterol (45.5 mg) and vitamin D2 (1.9 μ g). The fungal sterol ergosterol (known as provitamin D2) is abundant in mushrooms, as is its peroxide [30, 31]. Sterols suppress LPS-induced inflammatory responses in RAW264.7 macrophages through inhibition of the transcriptional activity of NF- κ B and CCAAT-enhancer-binding protein (C/EBP) β as well as phosphorylation of mitogen-activated protein kinases (MAPKs) [31]. Supplementation with vitamin D results in reductions in the levels of proinflammatory cytokines such as interleukin (IL)-4, IL-5, and IL-13 in the BALF of mice challenged with ovalbumin [32, 33]. Moreover, it has been reported that *P. eryngii* has a significantly higher amount of total phenolic compounds and has a close relationship

FIGURE 5: Effect of HTPE on histological changes in the lungs of LPS-induced ALI mice. Mice were administered, via the oral and intraperitoneal routes, HTPE (1 g/kg) and dexamethasone (DEX: 5 mg/kg, positive control group), respectively, 1 h before intranasal administration of LPS (10 μ g/site). Vehicle (tap water, vehicle for HTPE) was also administered orally 1 h before intranasal administration of LPS (LPS group) or saline (vehicle for LPS, saline group (negative control group)). Lungs ($n = 3-4$) from each experimental group were processed for histological evaluation 6 h after LPS challenge. The result in each group was almost identical. Lung sections were stained with hematoxylin and eosin ((a), (b), (e), and (f)). Lung sections in (c) and (d) show the immunoreactivity of myeloperoxidase (a marker of neutrophils) and Mac-2 (a marker of macrophages). Scale bar: 100 μ m. (a) Saline group (negative control group); ((b), (c), and (d)) LPS group; (e) HTPE-treated LPS group; (f) DEX-treated LPS group (positive control group).

with antioxidant activity and 2,2-diphenylpicrylhydrazyl-scavenging activity [6]. Taken together, these findings suggest that the components of *P. eryngii* may contribute to the anti-inflammatory actions of HTPE *in vivo*.

We used HTPE obtained from the whole fresh fruiting bodies of *P. eryngii*. HTPE (0.3–1.0 g/kg) was effective against ALI in mice. HTPE at 0.3–1.0 g/kg is equivalent to a fresh fruiting body at 3–10 g/kg. The effective dose in mice cannot be applied directly for humans, but a fresh fruiting body of *P. eryngii* (180–600 g per person (60 kg body weight)) should

be eaten to protect against ALI. We consider that intake of *P. eryngii* (which has anti-inflammatory effects and which many people can obtain) on a daily basis could be a useful prevention strategy against ALI.

5. Conclusion

HTPE obtained from the heat-treated fresh fruiting body of *P. eryngii* prevented LPS-induced ALI through inhibition

of infiltration of inflammatory cells, destruction of vascular integrity, and overproduction of NO and led to alleviation of histopathological damage. Our study supports the hypothesis that *P. eryngii* could be a beneficial food for the prevention of ALI associated with bacterial infection.

Conflict of Interests

The authors state that there is no conflict of interests.

Authors' Contribution

Junya Kawai and Tsugunobu Andoh contributed equally to this work.

Acknowledgment

The authors thank Mr. K. Matsunaga of the University of Toyama for technical assistance with the preparation of lung sections.

References

- [1] E. H. Han, Y. P. Hwang, H. G. Kim et al., "Inhibitory effect of *Pleurotus eryngii* extracts on the activities of allergic mediators in antigen-stimulated mast cells," *Food and Chemical Toxicology*, vol. 49, no. 6, pp. 1416–1425, 2011.
- [2] T. Mizutani, S. Inatomi, A. Inazu, and E. Kawahara, "Hypolipidemic effect of *Pleurotus eryngii* extract in fat-loaded mice," *Journal of Nutritional Science and Vitaminology*, vol. 56, no. 1, pp. 48–53, 2010.
- [3] Z. Yang, J. Xu, Q. Fu et al., "Antitumor activity of a polysaccharide from *Pleurotus eryngii* on mice bearing renal cancer," *Carbohydrate Polymers*, vol. 95, no. 2, pp. 615–620, 2013.
- [4] F. S. Reis, A. Martins, L. Barros, and I. C. Ferreira, "Antioxidant properties and phenolic profile of the most widely appreciated cultivated mushrooms: a comparative study between *in vivo* and *in vitro* samples," *Food and Chemical Toxicology*, vol. 50, no. 5, pp. 1201–1207, 2012.
- [5] S. Li and N. P. Shah, "Effects of various heat treatments on phenolic profiles and antioxidant activities of *Pleurotus eryngii* extracts," *Journal of Food Sciences*, vol. 70, no. 8, pp. C1122–C1129, 2013.
- [6] K. K. Mishra, R. S. Pal, R. Arunkumar, C. Chandrashekar, S. K. Jain, and J. C. Bhatt, "Antioxidant properties of different edible mushroom species and increased bioconversion efficiency of *Pleurotus eryngii* using locally available casing materials," *Food Chemistry*, vol. 138, no. 2–3, pp. 1557–1563, 2013.
- [7] K. Tsushima, L. S. King, N. R. Aggarwal, A. de Gorordo, F. R. D'Alessio, and K. Kubo, "Acute lung injury review," *Internal Medicine*, vol. 48, no. 9, pp. 621–630, 2009.
- [8] L. B. Ware and M. Herridge, "Acute lung injury," *Seminars in Respiratory Critical Care Medicine*, vol. 34, no. 4, pp. 439–440, 2013.
- [9] K. Sato, M. B. Kadiiska, A. J. Ghio et al., "In vivo lipid-derived free radical formation by NADPH oxidase in acute lung injury induced by lipopolysaccharide: a model for ARDS," *FASEB Journal*, vol. 16, no. 13, pp. 1713–1720, 2002.
- [10] J. Reutershan, A. Basit, E. V. Galkina, and K. Ley, "Sequential recruitment of neutrophils into lung and bronchoalveolar lavage fluid in LPS-induced acute lung injury," *The American Journal of Physiology—Lung Cellular and Molecular Physiology*, vol. 289, no. 5, pp. L807–L815, 2005.
- [11] L. G. Rochelle, B. M. Fischer, and K. B. Adler, "Concurrent production of reactive oxygen and nitrogen species by airway epithelial cells *in vitro*," *Free Radical Biology and Medicine*, vol. 24, no. 5, pp. 863–868, 1998.
- [12] G. Matute-Bello, C. W. Frevert, and T. R. Martin, "Animal models of acute lung injury," *The American Journal of Physiology—Lung Cellular and Molecular Physiology*, vol. 295, no. 3, pp. L379–L399, 2008.
- [13] Z. Ilidromiti, D. Zygouris, S. Sifakis et al., "Acute lung injury in preterm fetuses and neonates: mechanisms and molecular pathways," *Journal of Maternal-Fetal and Neonatal Medicine*, vol. 26, no. 17, pp. 1696–1704, 2013.
- [14] X. Chu, K. Song, K. Xu et al., "Ceftiofur attenuates lipopolysaccharide-induced acute lung injury," *International Immunopharmacology*, vol. 10, no. 5, pp. 600–604, 2010.
- [15] Y. Fujii, P. Goldberg, and S. N. A. Hussain, "Contribution of macrophages to pulmonary nitric oxide production in septic shock," *The American Journal of Respiratory and Critical Care Medicine*, vol. 157, no. 5, pp. 1645–1651, 1998.
- [16] S. Cheng, W. M. Yan, B. Yang, J. D. Shi, M. M. Song, and Y. Zhao, "A crucial role of nitric oxide in acute lung injury secondary to the acute necrotizing pancreatitis," *Human and Experimental Toxicology*, vol. 29, no. 4, pp. 329–337, 2010.
- [17] K. Mikawa, K. Nishina, Y. Takao, and H. Obara, "ONO-1714, a nitric oxide synthase inhibitor, attenuates endotoxin-induced acute lung injury in rabbits," *Anesthesia and Analgesia*, vol. 97, no. 6, pp. 1751–1755, 2003.
- [18] J. S. Lee, S. Y. Park, D. Thapa et al., "*Grifola frondosa* water extract alleviates intestinal inflammation by suppressing TNF- α production and its signaling," *Experimental and Molecular Medicine*, vol. 42, no. 2, pp. 143–154, 2010.
- [19] A. Jedinak, S. Dudhgaonkar, Q. Wu, J. Simon, and D. Sliva, "Anti-inflammatory activity of edible oyster mushroom is mediated through the inhibition of NF- κ B and AP-1 signaling," *Nutrition Journal*, vol. 10, no. 1, article 52, 2011.
- [20] Y. O. Kim, S. W. Lee, C. H. Oh, and Y. H. Rhee, "Hericium erinaceus suppresses LPS-induced pro-inflammation gene activation in RAW264.7 macrophages," *Immunopharmacology and Immunotoxicology*, vol. 34, no. 3, pp. 504–512, 2012.
- [21] E. R. Carbonero, A. H. P. Gracher, F. R. Smiderle et al., "A β -glucan from the fruit bodies of edible mushrooms *Pleurotus eryngii* and *Pleurotus ostreatus*," *Carbohydrate Polymers*, vol. 66, no. 2, pp. 252–257, 2006.
- [22] J. Nitschke, H. Modick, E. Busch, R. W. von Rekowski, H. Altenbach, and H. Mölleken, "A new colorimetric method to quantify β -1,3-1,6-glucans in comparison with total β -1,3-glucans in edible mushrooms," *Food Chemistry*, vol. 127, no. 2, pp. 791–796, 2011.
- [23] F. R. Smiderle, L. M. Olsen, E. R. Carbonero et al., "Anti-inflammatory and analgesic properties in a rodent model of a (1 $\rightarrow$ 3),(1 $\rightarrow$ 6)-linked β -glucan isolated from *Pleurotus pulmonarius*," *European Journal of Pharmacology*, vol. 597, no. 1–3, pp. 86–91, 2008.
- [24] L. S. Queiroz, M. S. Nascimento, A. K. M. Cruz et al., "Glucans from the *Caripia montagnei* mushroom present anti-inflammatory activity," *International Immunopharmacology*, vol. 10, no. 1, pp. 34–42, 2010.
- [25] X. Xu, M. Yasuda, S. Nakamura-Tsuruta, M. Mizuno, and H. Ashida, " β -glucan from *Lentinus edodes* inhibits nitric oxide

- and tumor necrosis factor- α production and phosphorylation of mitogen-activated protein kinases in lipopolysaccharide-stimulated murine RAW 264.7 macrophages,” *The Journal of Biological Chemistry*, vol. 287, no. 2, pp. 871–878, 2012.
- [26] J. Chen and R. Seviour, “Medicinal importance of fungal β -(1 $\rightarrow$ 3), (1 $\rightarrow$ 6)-glucans,” *Mycological Research*, vol. 111, no. 6, pp. 635–652, 2007.
- [27] R. Tada, A. Tanioka, H. Iwasawa et al., “Structural characterisation and biological activities of a unique type β -D-glucan obtained from *Aureobasidium pullulans*,” *Glycoconjugate Journal*, vol. 25, no. 9, pp. 851–861, 2008.
- [28] E. Sonck, B. Devriendt, B. Goddeeris, and E. Cox, “Varying effects of different β -glucans on the maturation of porcine monocyte-derived dendritic cells,” *Clinical and Vaccine Immunology*, vol. 18, no. 9, pp. 1441–1446, 2011.
- [29] W. Chanput, M. Reitsma, L. Kleinjans, J. J. Mes, H. F. Savelkoul, and H. J. Wichers, “ β -glucans are involved in immune-modulation of THP-1 macrophages,” *Molecular Nutrition and Food Research*, vol. 56, no. 5, pp. 822–833, 2012.
- [30] Y. Yaoita, Y. Yoshihara, R. Kakuda, K. Machida, and M. Kikuchi, “New sterols from two edible mushrooms, *Pleurotus eryngii* and *Panellus serotinus*,” *Chemical and Pharmaceutical Bulletin*, vol. 50, no. 4, pp. 551–553, 2002.
- [31] M. Kobori, M. Yoshida, M. Ohnishi-Kameyama, and H. Shinmoto, “Ergosterol peroxide from an edible mushroom suppresses inflammatory responses in RAW264.7 macrophages and growth of HT29 colon adenocarcinoma cells,” *The British Journal of Pharmacology*, vol. 150, no. 2, pp. 209–219, 2007.
- [32] T. Agrawal, G. K. Gupta, and D. K. Agrawal, “Vitamin D supplementation reduces airway hyperresponsiveness and allergic airway inflammation in a murine model,” *Clinical and Experimental Allergy*, vol. 43, pp. 672–683, 2013.
- [33] D. Parekh, D. R. Thickett, and A. M. Turner, “Vitamin D deficiency and acute lung injury,” *Inflammation and Allergy—Drug Targets*, vol. 12, no. 4, pp. 253–261, 2013.