Research Article

Endangered Uyghur Medicinal Plant *Ferula* Identification through the Second Internal Transcribed Spacer

Congzhao Fan,¹ Xiaojin Li,¹ Jun Zhu,¹ Jingyuan Song,^{2,3} and Hui Yao²

¹ Xinjiang Institute of Chinese Materia Medica and Ethnical Materia, 9 Xinming Road, Urumqi, Xinjiang 830002, China

² The National Engineering Laboratory for Breeding of Endangered Medicinal Materials, Institute of Medicinal Plant Development, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100193, China

³ Chongqing Institute of Medicinal Plant Cultivation, Chongqing 408335, China

Correspondence should be addressed to Xiaojin Li; xjlxj@126.com

Received 10 June 2014; Revised 6 October 2014; Accepted 6 October 2014

Academic Editor: Robert Henry

Copyright © 2015 Congzhao Fan et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The medicinal plant *Ferula* has been widely used in Asian medicine, especially in Uyghur medicine in Xinjiang, China. Given that various substitutes and closely related species have similar morphological characteristics, *Ferula* is difficult to distinguish based on morphology alone, thereby causing confusion and threatening the safe use of *Ferula*. In this study, internal transcribed spacer 2 (ITS2) sequences were analyzed and assessed for the accurate identification of two salable *Ferula* species (*Ferula sinkiangensis* and *Ferula fukangensis*) and eight substitutes or closely related species. Results showed that the sequence length of ITS2 ranged from 451 bp to 45 bp, whereas guanine and cytosine contents (GC) were from 53.6% to 56.2%. A total of 77 variation sites were detected, including 63 base mutations and 14 insertion/deletion mutations. The ITS2 sequence correctly identified 100% of the samples at the species level using the basic local alignment search tool 1 and nearest-distance method. Furthermore, neighbor-joining tree successfully identified the genuine plants *F. sinkiangensis* and *F. fukangensis* from their succedaneum and closely related species. These results indicated that ITS2 sequence could be used as a valuable barcode to distinguish Uyghur medicine *Ferula* from counterfeits and closely related species. This study may broaden DNA barcoding application in the Uyghur medicinal plant field.

1. Introduction

The endangered Uyghur medicinal herb *Ferula* has a long history of application in China. It was first utilized in the *Tang Materia Medica* during the Tang Dynasty (AD 659) because of its effectiveness in removing food residues, as well as its dispersed stuffiness and chlordime form [1]. To date, *Ferula* is traditionally employed for treating different diseases such as asthma, epilepsy, stomach ache, flatulence, intestinal parasites, weak digestion, and influenza in many countries [2, 3]. Pharmacological and biological studies have shown that *Ferula* has antioxidant [4], antiviral [5], antifungal [6], cancer chemopreventive [7], antidiabetic [8], antispasmodic [6], hypotensive [9], and molluscicidal effects [10]. *Ferula* is now included in the Uyghur Medicine Criteria [11] and calendar version of the Chinese Pharmacopoeia [12]. The original plants for this medicine are *Ferula sinkiangensis* and *Ferula fukangensis*. In addition to a close relationship with

traditional Chinese medicine, Uyghur medicine has absorbed the essences of Indian, Arabian, Iran, and ancient Greek medicine, thereby forming a complete system of medicine with distinctive ethnic characteristics. However, its medicinal parts and functions are not precisely the same. For instance, traditional Chinese medicine utilizes *Ferula* resin, whereas Uyghur medicine utilizes *Ferula* resin, roots, and seeds [13] as clinical parts.

In China, the *Ferula* genus is mostly distributed in Xinjiang, but has been destroyed in this area as a result of excessive collection. In addition to seed plant breeding, change in reclamation, irrigation, road building, and deterioration of the original habitat contribute to the annual shrinking of *Ferula* resources. Currently, *F. fukangensis* is on the verge of extinction, and *F. sinkiangensis* is listed as a class 3 endangered plant and class 2 protected wild medicinal species. Given its high cost and scarce resources, various substitutes and adulterants have emerged. *F. sinkiangensis* and *F. fukangensis*

have been misused and substituted by other species such as *F. ferulaeoides* Korov [14]. Furthermore, numerous species from the *Ferula* genus have been utilized as traditional Chinese medicines and ethical materials. For example, *F. soongarica* has been utilized by Kazakhs in treating headaches, colds, and stomach aches; *F. caspica* in treating nervous breakdown; and *F. lehmannii* in treating parasitic malnutrition and cold pain in the heart and abdomen [15]. When mixed with substitutes, adulterants, and closely related species, the original medicinal plant could not be traced to the medicinal herb market. A famous line that is widely circulated in the herbal medicine field is “there is no false *Scutellaria*, no true *Ferula*.” Thus, investigating the authenticity of the Uyghur medicinal herb *Ferula* is urgently required.

The *Ferula* genus is usually recognized as monophyletic; the delimitation of species requires examination of complete specimens with roots, stem bases, basal leaves, inflorescence, flowers, and ripe fruits based on the observations of living plants [16]. Given that members are similar in habit and morphology, flowers, inflorescences, and fruit anatomy are often hardly discernible taxa [17, 18]. Thus, the infrageneric classification method of the Uyghur medicinal herb *Ferula* urgently requires a solution. To address this problem, microscopic identification and powder identification techniques have been utilized for *F. sinkiangensis* and its closely related species [19–21]. However, these techniques could not solve the problem of authenticating the Uyghur medicinal herbs *F. sinkiangensis* and *F. fukanensis* from other traditional medicinal herbs and closely related *Ferula* species. By comparison, DNA barcoding employs a short DNA sequence from a standard locus as a species identification tool [22], which has been widely applied in forage identification [23] and animal genetic relationship identification [24]. The internal transcribed spacer (ITS) region of this barcode, which comprises ITS1, 5.8S, and ITS2, has been widely utilized as a core DNA barcode to identify different herbal medicinal materials [25]. Researchers have proposed ITS2 as a standard DNA barcode for medicinal plant authentication [26, 27]. However, molecular identification of *F. sinkiangensis* and *F. fukanensis* from their closely related species utilizing the ITS2 sequence has not been reported. In the present study, DNA barcoding technology was applied for the first time to distinguish *F. sinkiangensis* and *F. fukanensis* from their succedaneum and other closely related species. The application of DNA barcoding in trade and market management could ensure the safe use of the pharmacopoeia herb *Ferula*.

2. Samples and Methods

2.1. Materials and Methods for Sampling of Plant Materials. Seventy-three leaves and root samples, which represented 10 *Ferula* species, were collected from different locations. Nine *Ferula* species were acquired from Xinjiang and one species of *Ferula litwinowiana* was obtained from Kazakhstan. The 10 species were given specimen collection numbers (Table S1) (see Supplementary Material available online at <http://dx.doi.org/10.1155/2015/479879>) and authenticated by Researcher Guanmian Shen (Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences) and Associate

Fellow Guoping Wang (Xinjiang Institute of Chinese Traditional Medical and Ethical Materia Medica). All corresponding voucher samples were deposited in the herbarium of the Xinjiang Institute of Chinese Traditional Medical and Ethical Materia Medica. In addition, one published sequence was downloaded from GenBank for analysis (Table S1).

2.2. DNA Extraction, Amplification, and Sequencing. Genomic DNA was isolated from 30 mg of leaves or roots according to the protocol of the Plant Genomic DNA Kit (Tiangen Biotech Co., China). Specific regions of the ITS2 sequences were amplified from dried leaves and root extracts; universal primer and PCR conditions for ITS2 were obtained from previous studies [28]. The purified PCR products were sequenced using a 3730XL sequencer (Applied Biosystems, Foster City, CA, USA).

2.3. Sequence Alignment and Analysis. The CodonCode Aligner v 4.0.4 was utilized to edit the sequences and assemble the contigs and shear sequence using universal primers (5′-3′: ATTACACCAAGTATCGCAT and 3′-5′: ATTGTAGTCTGGAGAAGCGTC). The ITS2 core region could be obtained using the HMMer annotation method based on the hidden Markov model (HMM) to remove the 5.8S and 28S sections at both ends of the sequences [29]. The interspecific/intraspecific variation of the samples was calculated according to Kress et al. [22]. A phylogenetic tree was constructed utilizing MEGA 5.0. Neighbor-joining algorithm (NJ tree) was used to evaluate the capability of the ITS2 sequence to authenticate the studied species. Bootstrap tests were conducted utilizing 1,000 resampling to assess the confidence of the phylogenetic relationships using employed MEGA 5.0 [30, 31]. BLAST1 and nearest-distance methods were performed to identify species, as previously described [32, 33].

3. Results

3.1. Universality and Sequence Characteristics. DNA was successfully extracted from 73 samples. Gel electrophoretic analysis revealed that all sample extractions emitted bright bands. ITS2 sequences were successfully amplified and sequenced; high-quality bidirectional sequences were obtained. The ITS2 sequence, including partial the 5.8S, ITS2 region, and partial 28S, ranged from 451 bp to 455 bp, and GC contents were 53.6% to 56.2%. The partial 5.8S length was 84 bp; the GC contents ranged from 57.1% to 58.3%. The length of the ITS2 region was 226 bp to 230 bp; the GC contents ranged from 53.0% to 56.8%. The partial 28S was 141 bp and the GC contents ranged from 51.8% to 53.9%.

3.2. Intraspecific/Interspecific Variations. Sequence length, variable sites, and average interspecific K2P distance of ITS2 regions were analyzed and summarized (Table 1). Most of the 10 *Ferula* species in this study were represented by several samples because they were collected from different localities. *F. sinkiangensis* was represented by 13 samples (Table S1), *F. fukanensis* by 6 samples, *F. ferulaeoides* by 14

FIGURE 1: Variable sites in the haplotypes of the 10 *Ferula* species.

samples, *F. soongarica* by 6 samples, *F. caspica* by 11 samples, and so on (Table 1). The results showed that each *Ferula* species only possesses a haplotype, and their intraspecific distances were all zero.

Among the 74 ITS2 sequences, a total of 77 variation sites were detected in the region, including 63 base mutations and 14 insertion/deletion mutations. The following contents were detected among the 63 base mutation sites: C/T (26), C/G (5), A/C (12), G/A (10), T/A (5), T/G (4), C/G/A (1), and C/T/A (4) substitutions. A total of 76 distinctive variable sites were discovered among the 77 variation sites, which could precisely authenticate different *Ferula* species (Figure 1). For instance, the C–T transition at site 133 and the A–G transition at site 206 could distinguish *F. sinkiangensis* from the other nine *Ferula* species. The C–T transition at the 397 bp site could distinguish *F. soongarica* and the C–T transition at site 176 could individually classify *F. caspica*. The interspecific divergence calculated utilizing the K2P model indicated that the ITS2 sequences of *F. sinkiangensis* possessed an average interspecific distance of 0.030; *F. fukanensis* possessed an average intraspecific distance of 0.027 (Table 1).

3.3. Species Identification Capability of ITS2 Barcode and NJ Tree Analysis. BLAST1, nearest distance, and tree-based methods were utilized to determine whether the certified medicinal herb *Ferula* could be identified by ITS2 sequences. The results indicated that ITS2 correctly identified 100% of the samples at the species level through the BLAST1 method. The nearest genetic distance method also resulted in 100% success rate of identification.

Up to 74 ITS2 sequences of *Ferula* species were chosen to build the trees. NJ trees (Figure 2) were constructed based on the ITS2 sequences. The results showed that the Uyghur medicinal herbs *F. fukangensis* and *F. sinkiangensis* presented good monophyly; they could be correctly distinguished from their succedaneum *F. ferulaeoides* and other closely related species. Furthermore, the remaining *Ferula* species could

be distinguished from one another. Thus, the ITS2 barcode could correctly identify two Uyghur medicinal herb plants from their adulterants and closely related species.

4. Discussion

4.1. *ITS2 Region Is a Useful DNA Barcode to Differentiate 10 Ferula Species.* The genus *Ferula* is an important ephemeral plant that possesses highly ecological and medicinal values. Our results emphasized the advantages of using the ITS2 region as a DNA barcode, including high interspecific divergence but small intraspecific variation, favorable universality, and short length, which lead to easy amplification and sequencing efficiency. Given that pharmacologists often traded *Ferula* genus plants internationally for their medicinal values, our research provides a convenient tool for validating the quality products of *Ferula*. In a previous study [18], 83 accessions (74 species) of *Ferula* were used to ascertain the phylogenetic position of the genus within the family. Most species had one sample; hence, the intraspecific divergence could not be determined. The ITS1 and ITS2 regions are separately amplified, and then assembled to successfully redress several misplaced *Ferula* genus plants. Phylogenetic analyses of tree-based method successfully corrected the three accessions of misplaced *Ferula* species. In our study, 74 ITS2 sequences of 10 *Ferula* species were used to authenticate *F. sinkiangensis* and *F. ferulaeoides* from their succedaneum and closely related species, most of which were newly reported sequences and each species was represented by two or more samples. Therefore, intraspecific variation was clearly displayed.

Morphological characteristics are an important basis for identifying the original *Ferula* plants. However, morphological identification often relies on abundant experience, which could be easily affected by biocoenosis and geographical environment [17]. As a result, identification of traditional Chinese medicine *Ferula* has been limited. Genomic sequence is not

FIGURE 2: NJ tree of *F. sinkiangensis* and *F. fukanensis* and their closely related species constructed with the ITS2 sequence. The bootstrap scores (1,000 replicates) are shown ($\geq 50\%$) for each branch.

TABLE 1: ITS2 sequence characteristics and K2P distances of *Ferula* used in this research.

Species	Sample number	GC content (%)	Sequence length (bp)	Distinctive variable site	Interspecific K2P distance (mean)
<i>F. sinkiangensis</i>	13	55.0	451	7	0.009–0.163 (0.030)
<i>F. fukanensis</i>	6	54.7	455	0	0.002–0.161 (0.027)
<i>F. ferulaeoides</i>	14	54.6	454	0	0.002–0.150 (0.023)
<i>F. soongarica</i>	6	54.4	454	1	0.002–0.154 (0.025)
<i>F. caspica</i>	11	54.3	455	3	0.009–0.161 (0.030)
<i>F. lehmannii</i>	5	54.9	455	0	0.002–0.157 (0.028)
<i>F. syreitschikowi</i>	10	54.3	455	3	0.007–0.145 (0.028)
<i>F. akitschkensis</i>	4	56.2	452	53	0.145–0.170 (0.156)
<i>F. dubjanskyi</i>	3	53.6	455	5	0.016–0.151 (0.035)
<i>F. litwinowiana</i>	2	54.9	455	4	0.007–0.170 (0.033)

influenced by individual characteristics and developmental stages; therefore, DNA barcode technology has been widely used in recent years [34]. DNA barcoding is an effective supplement to traditional morphological methods, and, in our study, the phylogenetic tree (Figure 2) demonstrated that the ITS2 region has an effective role in authenticating the Uyghur medicinal herb *Ferula* and differentiating 10 *Ferula* species from each other.

4.2. Identification of Uyghur Medicine Utilizing ITS2 Sequence. According to the records, Uyghur medicine includes 1,000 types of herbal medicines and up to 450 types are frequently utilized. Nevertheless, limited by rare resources and confusing plant origins, only about 200 species of Uyghur medicinal herbs have been formulated based on national and local criteria. Aside from the Uyghur medicinal herb *Ferula*, numerous other Uyghur medicinal herbs, such as *Cichorium intybus* L. and *Cichorium glandulosum* Boiss. et Huet, are facing the same dilemma [14]. These situations are significant obstacles to the standardization and industrialization of Uyghur medicine. Although morphological examination and chemical analysis are routine practices in identifying Uyghur medicinal herbs, these methods are influenced by biological and physical factors [35]. DNA barcoding and forensically informative nucleotide sequencing are less affected by these elements. The ITS2 barcode has been proposed as a universal DNA barcode for medicinal herb and animal identification [26, 27, 36, 37]. Thus, based on this study, DNA barcoding would have a broad prospect in authenticating more Uyghur medicinal herbs, which provides a new method for the Food and Drug Administration to control Uyghur medicine and ensure its safe and efficient use and has a considerable role in developing the Uyghur medicine industry.

4.3. Significance of DNA Barcode in Food Safety Control. Although several *Ferula* genus plant roots are used traditional medicine or folk medicine in Xinjiang, few *Ferula* species are edible. Starch extracted from *F. ferulaeoides* and *F. dubjanskyi* could be made into food [2]. The tender leaves of *F. lehmannii* could be used as wild herbs to make cold dishes and dumplings [38]. DNA barcoding has been used for food safety control [39–41]. The NJ tree demonstrated that the ITS2 barcode could accurately perform food safety control. Sequencing the ITS2 DNA barcode region represents a new technique that guarantees food and drug safety in *Ferula* species. This technology is easy to learn and utilized by food managers [42].

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The authors thank Professor S. L. Chen from the Institute of Medicinal Plant Development, Chinese Academy of Medicinal Sciences, for his valuable comments. This study was

supported by the National Natural Science Foundation of China (Grant nos. 81060265 and 81373922) and the Natural Science Foundation of the Xinjiang Uyghur Autonomous Region (Grant no. 2011211B44).

References

- [1] E. P. Korovin, “*Ferula* L.,” in *Flora Uzbekistana*, vol. 4, pp. 399–438, 1959.
- [2] M. Iranshahy and M. Iranshahi, “Traditional uses, phytochemistry and pharmacology of asafoetida (*Ferula assa-foetida* oleogum-resin)—a review,” *Journal of Ethnopharmacology*, vol. 134, no. 1, pp. 1–10, 2011.
- [3] P. Mahendra and S. Bisht, “*Ferula asafoetida*: traditional uses and pharmacological activity,” *Pharmacognosy Reviews*, vol. 6, no. 12, pp. 141–146, 2012.
- [4] S. Ben Salem, A. Jabrane, F. Harzallah-Skhiri, and H. Ben Jannet, “New bioactive dihydrofuranocoumarins from the roots of the Tunisian *Ferula lutea* (Poir.) Maire,” *Bioorganic and Medicinal Chemistry Letters*, vol. 23, no. 14, pp. 4248–4252, 2013.
- [5] Z. E. Nazari and M. Iranshahi, “Biologically active sesquiterpene coumarins from *Ferula* species,” *Phytotherapy Research*, vol. 25, no. 3, pp. 315–323, 2011.
- [6] G. Kavooosi, A. Tafsiry, A. A. Ebdam, and V. Rowshan, “Evaluation of antioxidant and antimicrobial activities of essential oils from *Carum copticum* Seed and *Ferula assafoetida* latex,” *Journal of Food Science*, vol. 78, no. 2, pp. T356–T361, 2013.
- [7] M. Iranshahi, A. Sahebkar, S. T. Hosseini, M. Takasaki, T. Konoshima, and H. Tokuda, “Cancer chemopreventive activity of diversin from *Ferula diversivittata* in vitro and in vivo,” *Phytomedicine*, vol. 17, no. 3–4, pp. 269–273, 2010.
- [8] A. S. Abu-Zaiton, “Anti-diabetic activity of *Ferula assafoetida* extract in normal and alloxan-induced diabetic rats,” *Pakistan Journal of Biological Sciences*, vol. 13, no. 2, pp. 97–100, 2010.
- [9] M. Fatehi, F. Farifteh, and Z. Fatehi-Hassanabad, “Antispasmodic and hypotensive effects of *Ferula assafoetida* gum extract,” *Journal of Ethnopharmacology*, vol. 91, no. 2–3, pp. 321–324, 2004.
- [10] S. Kumar, D. K. Singh, and V. K. Singh, “Tertiary combination of freeze-dried urine of Indian breeds of cow with plant products against snail *Lymnaea acuminata*,” *Pakistan Journal of Biological Sciences*, vol. 15, no. 20, pp. 992–996, 2012.
- [11] Xinjiang Food and Drug Administration, *The Xinjiang Uyghur Autonomous Region Traditional Chinese Medicine and Uyghur Medicine Pieces Concoct Standard*, Xinjiang People’s Saitary Press, 2010.
- [12] Chinese Pharmacopoeia Commission, *Chinese Pharmacopoeia*, vol. 1, Chinese Medicine Science and Technology Press, 2010.
- [13] Chinese Materia Medica Editorial Committee, *The Chinese Materia Medica (Uyghur Volume)*, Shanghai Science and Technology Press, 2005.
- [14] C. Liu, W. J. Yang, Z. Y. Gu et al., “Research progress of Uyghur medicine easy mixed species arrangement and the application of molecular biology,” *Northwest Pharmaceutical Journal*, vol. 28, no. 4, pp. 440–442, 2013.
- [15] D. Bandyopadhyay, B. Basak, A. Chatterjee et al., “Saradaferin, a new sesquiterpenoid coumarin from *Ferula assafoetida*,” *Natural Product Research*, vol. 20, no. 10, pp. 961–965, 2006.
- [16] E. P. Korovin, *Generis Ferula (Tourn.) L. Monographia illustrata*, Academiae Scientiarum UzRSS, Taschkent, Uzbekistan, 1947.

- [17] Y. Ajani, A. Ajani, J. M. Cordes, M. F. Watson, and S. R. Downie, "Phylogenetic analysis of nrDNA ITS sequences reveals relationships within five groups of Iranian *Apiaceae* subfamily Apioideae," *Taxon*, vol. 57, no. 2, pp. 383–401, 2008.
- [18] R. Kurzyńska-Młynik, A. A. Oskolski, S. R. Downie, R. Kopacz, A. Wojewódzka, and K. Spalik, "Phylogenetic position of the genus *Ferula* (Apiaceae) and its placement in tribe Scandiceae as inferred from nrDNA ITS sequence variation," *Plant Systematics and Evolution*, vol. 274, no. 1–2, pp. 47–66, 2008.
- [19] Y. Tan, T.-T. Gao, Y. Ma, X. Ding, and Y.-H. Cheng, "Pharmacognostic identification of *Ferula syreitschikowii*," *Journal of Chinese Medicinal Materials*, vol. 34, no. 11, pp. 1694–1696, 2011.
- [20] H. Y. Han, J. M. Xu, G. Y. Li et al., "Microscopic identification of *Ferula dissecta* (Ledeb.) Ledeb," *Modern Chinese Medicine*, vol. 12, no. 12, pp. 17–19, 2010.
- [21] X. Ding, Y. Tan, Y. Zhang et al., "Pharmacognostic identification of *Ferula sinkiangensis*," *China Pharmacy*, vol. 35, pp. 3307–3309, 2011.
- [22] W. J. Kress, K. J. Wurdack, E. A. Zimmer, L. A. Weigt, and D. H. Janzen, "Use of DNA barcodes to identify flowering plants," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 102, no. 23, pp. 8369–8374, 2005.
- [23] I. Ganopoulos, P. Madesis, and A. Tsaftaris, "Universal ITS2 barcoding DNA region coupled with High-Resolution Melting (HRM) analysis for seed authentication and adulteration testing in Leguminous forage and pasture species," *Plant Molecular Biology Reporter*, vol. 30, no. 6, pp. 1322–1328, 2012.
- [24] R. Zomuanpuui, L. Ringnaghi, S. Brindha, G. Gurusubramanian, and N. Senthil Kumar, "ITS2 characterization and *Anopheles* species identification of the subgenus *Cellia*," *Acta Tropica*, vol. 125, no. 3, pp. 309–319, 2013.
- [25] M. Li, H. Cao, P. P. H. But, and P. C. Shaw, "Identification of herbal medicinal materials using DNA barcodes," *Journal of Systematics and Evolution*, vol. 49, no. 3, pp. 271–283, 2011.
- [26] S. Chen, H. Yao, J. Han et al., "Validation of the ITS2 region as a novel DNA barcode for identifying medicinal plant species," *PLoS ONE*, vol. 5, no. 1, Article ID e8613, 2010.
- [27] H. Yao, J. Song, C. Liu et al., "Use of ITS2 region as the universal DNA barcode for plants and animals," *PLoS ONE*, vol. 5, no. 10, Article ID e13102, 2010.
- [28] Z. Sun and S. Chen, "Identification of cortex herbs using the DNA barcode nrITS2," *Journal of Natural Medicines*, vol. 67, no. 2, pp. 296–302, 2013.
- [29] A. Keller, T. Schleicher, J. Schultz, T. Müller, T. Dandekar, and M. Wolf, "5.8S-28S rRNA interaction and HMM-based ITS2 annotation," *Gene*, vol. 430, no. 1–2, pp. 50–57, 2009.
- [30] C. Koetschan, T. Hackl, T. Müller, M. Wolf, F. Förster, and J. Schultz, "ITS2 Database IV: interactive taxon sampling for internal transcribed spacer 2 based phylogenies," *Molecular Phylogenetics and Evolution*, vol. 63, no. 3, pp. 585–588, 2012.
- [31] K. Tamura, D. Peterson, N. Peterson, G. Stecher, M. Nei, and S. Kumar, "MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods," *Molecular Biology and Evolution*, vol. 28, no. 10, pp. 2731–2739, 2011.
- [32] L. Xiang, J. Song, T. Xin et al., "DNA barcoding the commercial Chinese caterpillar fungus," *FEMS Microbiology Letters*, vol. 347, no. 2, pp. 156–162, 2013.
- [33] H. A. Ross, S. Murugan, and W. L. S. Li, "Testing the reliability of genetic methods of species identification via simulation," *Systematic Biology*, vol. 57, no. 2, pp. 216–230, 2008.
- [34] C. Liu, D. Liang, T. Gao et al., "PTIGS-IdIt, a system for species identification by DNA sequences of the psbA-trnH intergenic spacer region," *BMC Bioinformatics*, vol. 12, no. 13, article S4, 2011.
- [35] P. C. Shaw, F. N. Ngan, and P. P. H. But, *Authentication of Chinese Medicinal Materials by DNA Technology*, World Scientific Publishing, Singapore, 2002.
- [36] P. D. N. Hebert, S. Ratnasingham, and J. R. DeWaard, "Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species," *Proceedings of the Royal Society B: Biological Sciences*, vol. 270, supplement 1, pp. S96–S99, 2003.
- [37] M. Li, P. P. H. But, and P. C. Shaw, "Molecular identification of traditional medicinal materials," in *Molecular Pharmacognosy*, pp. 45–66, Springer, Dordrecht, The Netherlands, 2013.
- [38] China Flora Editorial Board of Cas, *Flora China*, vol. 14, Sciences Press, Beijing, China, 1992.
- [39] L. Jaakola, M. Suokas, and H. Häggman, "Novel approaches based on DNA barcoding and high-resolution melting of amplicons for authenticity analyses of berry species," *Food Chemistry*, vol. 123, no. 2, pp. 494–500, 2010.
- [40] M. Á. Pavón, I. González, R. Martín, and T. G. Lacarra, "ITS-based detection and quantification of *Alternaria* spp. in raw and processed vegetables by real-time quantitative PCR," *Food Microbiology*, vol. 32, no. 1, pp. 165–171, 2012.
- [41] T. Xin, H. Yao, H. Gao et al., "Super food *Lycium barbarum* (Solanaceae) traceability via an internal transcribed spacer 2 barcode," *Food Research International*, vol. 54, no. 2, pp. 1699–1704, 2013.
- [42] M. Y. Stoeckle, C. C. Gamble, R. Kirpekar, G. Young, S. Ahmed, and D. P. Little, "Commercial teas highlight plant DNA barcode identification successes and obstacles," *Scientific Reports*, vol. 1, article 42, 2011.