Review Article

Evaluation of Chemical Constituents and Important Mechanism of Pharmacological Biology in *Dendrobium* Plants

Yau Lam,¹ Tzi Bun Ng,² Ren Ming Yao,¹ Jun Shi,¹ Kai Xu,¹
Stephen Cho Wing Sze,¹ and Kalin Yanbo Zhang¹

¹School of Chinese Medicine, LKS Faculty of Medicine, The University of Hong Kong, 10 Sassoon Road, Hong Kong

²School of Biomedical Sciences, Faculty of Medicine, The Chinese University of Hong Kong, Shatin, NT, Hong Kong

Correspondence should be addressed to Tzi Bun Ng; b021770@mailserv.cuhk.edu.hk, Stephen Cho Wing Sze; stephens@hku.hk, and Kalin Yanbo Zhang; ybzhang@hkucc.hku.hk

Received 24 November 2014; Accepted 9 February 2015

Academic Editor: Wagner Vilegas

Copyright © 2015 Yau Lam et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dendrobium species, commonly known as “Shihu” or “Huangcao,” represents the second largest genus of Orchidaceae, which are used commonly as tonic herbs and healthy food in many Asian countries. The aim of this paper is to review the history, chemistry, and pharmacology of different *Dendrobium* species on the basis of the latest academic literatures found in Google Scholar, PubMed, Sciencedirect, Scopus, and SID.

1. Introduction

The origin of orchids (Orchidaceae) probably has to be backdated to 120 million years ago. It is the largest family group among angiosperms, as well as the most highly evolved family of the flowering plants, with approximately 25,000 to 35,000 species under 750 to 900 genera [1–4]. *Dendrobium* species, commonly known as “Shihu” or “Huangcao,” is the second largest genus in Orchidaceae. Most *Dendrobium* species grow best in relatively high and mountainous areas, at 1400–1600 m above sea level, at a mild temperature, and in a humid and foggy environment. Characterized by a broad geographical distribution, which allows *Dendrobium* species to grow into tremendous diversities producing a large number of interspecific hybrids with different morphological features, they are widely distributed in Asia, Australia, and Europe, for instance, in India, Sri Lanka, China, Japan, Korea, New Guinea, and New Caledonia [5]. In the early age, using molecular approaches would delimit the subtribe Dendrobiinae, affecting approximately 1100 species (900 in *Dendrobium* Sw.) from Indo-Asian and Pacific regions [6, 7], and more than 1200 species in Australasia are from various *Dendrobium* species. Today, Indo-Asian and Pacific regions have one of the largest and most diverse and taxonomically problematic orchid groups [8]. Interestingly, there has been

a long history of the usage of first-rate herbs and folk traditional herbs in India and China [9].

Since the ancient times (600 B.C.) in India, the oldest references regarding the use of medicinal herbs are found in the Sanskrit literature, namely, “Charaka Samhita.” The earliest treatise on “Ayurveda” describes the property of plant drugs and their uses. In the Ayurvedic system of medicine, *Flickingeria macraei* is used in “Ayurveda.” It is commonly named as “jeevanti” and is used as an astringent to the bowels, as an aphrodisiac, and in asthma and bronchitis [10]. Other commonly used orchid drugs in the Ayurvedic system are salem, including jewanti (*Dendrobium alpestre*). Similarly, it is the first time for China to regard orchids as herbal medicines [11]. The Emperor “Shen-Nung” advised on the medicinal properties of *Dendrobium* species in “materia medica” in the 28th century B.C. [12]. As early as 200 B.C., the Chinese pharmacopoeia “The Sang Nung Pen Tsao Ching” mentioned *Dendrobium* as a source of tonic, astringent, analgesic, and anti-inflammatory substances [13]. In Song Dynasty (960–1279 A.D.). It mentioned the medicinal uses of orchids, namely *Dendrobium* species according A Diagnosis of Medical Herbs from the “Zheng Lei Ben Cao”. In Ming Dynasty (1368–1644) many references on the use of orchids as medicinal herbs were available [14].

2. The Chemical Compounds of *Dendrobium* Species

The stem of *Dendrobium* species has been used in traditional Chinese medicine as a tonic and antipyretic since ancient days for treating human disorders. However, misidentification and adulteration led to a loss of therapeutic potency and potential intoxication. For decades, fast developing molecular techniques using DNA fingerprinting, DNA sequencing, and DNA microarray have been applied extensively to authenticate Chinese medicinal materials, including various *Dendrobium* species [15], namely, *D. aphyllum*, *D. candidum*, *D. chrysanthum*, *D. densiflorum*, *D. huoshanense*, *D. gratiosissimum*, *D. longicornu*, *D. nobile*, *D. secundum*, *D. chrysotoxum*, *D. crystallinum*, *D. fimbriatum*, and others [16]. In the early years, Williams and Harborne conducted a major survey of leaf flavonoids at the Plant Science Laboratories of the University of Reading in UK. They conducted research on 142 species belonging to 75 genera and found that the most common constituents were flavone C-glycoside and flavonols [17]. Since then, about 100 compounds from 42 *Dendrobium* species including 32 alkaloids, 6 coumarins, 15 bibenzyls, 4 fluorenones, 22 phenanthrenes, and 7 sesquiterpenoids constituents have been discussed and reviewed [18–31] (Table 1). To date, various *Dendrobium* species are known to produce a variety of secondary metabolites. The biological activities and pharmacological actions of all of these compounds were investigated in detail [32–39].

3. The Pharmacological Effects of *Dendrobium* Species

3.1. Antioxidant Activities. Oxidative stress is induced by free radicals that participate in a variety of chemical reactions with biomolecules, leading to a pathological condition. Reactive oxygen species (ROS) is one of the major free radicals. It mainly comprises superoxide (O_2^-) and nitric oxide (NO) radicals, including (1) catalase (CAT), (2) peroxidase (POD), (3) ascorbate peroxidase (APX), (4) 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) or ABTS, (5) hydroxyl, and (6) 1,1-diphenyl-2-picrylhydrazyl (DPPH) radicals. During these processes, it was found that the content of malondialdehyde (MDA) was increased.

However, cells possess two distinctive antioxidant defense systems to counteract the damage, including enzymatic antioxidants and nonenzymatic antioxidants. Enzymatic antioxidants comprise catalase, superoxide dismutase (SOD), glutathione peroxidase (GPx), and others associated with enzymes/molecules. Nonenzymatic antioxidants include ascorbic acid (vitamin C), α -tocopherol (vitamin E), and β -carotene, which play a key role in removing reactive oxygen species.

Oxidative stresses have been classified as exogenous factors and endogenous factors. Factors related to environmental stress include water/soil drought stress, chilling injury stress, and sound wave stress. There were studies on the effects of cold storage on cut *Dendrobium* inflorescences, showing that chilling injury symptoms for floral buds and open flowers

of *Dendrobium* could lead to oxidative stress, stemming from the production of reactive oxygen species. In this process, the decrease of cellular functions by peroxidation of membrane lipids when lipoxygenase enzymes are degraded in cells. However, this physiological process in floral buds and open flowers is part of antioxidant defense systems, which can decrease the content of oxygen free radicals and other oxygen compounds. It indicates the protective cellular functions and antioxidant capacity. This is an adaptive nature to ensure some plants survive in the freezing winter [62].

In addition, sound wave stress is one of the environmental stresses, similar to the low temperature stress. Studies of the effects of sound wave stress on *D. candidum* yielded data indicated it as a lipid peroxidation parameter which can regulate the levels of MDA. Firstly, the MDA content in different organs is increased and then declines afterwards, which will then be followed by an increase again. That gives a net increase of the level of MDA. It was noted that the MDA level appears to be the lowest when the activities of antioxidant enzymes are the highest. However, the levels of MDA are yet to be fully understood. We believe the antioxidant enzymes could protect plant cells from oxidative damage in sound wave stress [63]. However, the correlation between the mechanisms of antioxidant action and the sound wave stress and chilling injury stress has yet to be identified.

Water deficit caused by soil drought is one of the most frequent environmental stresses. The effects of exogenous drought menace, prompting further increases in the activity of ROS and decrease in the malondialdehyde (MDA) content, to prevent the breakage of DNA and protein degradation from doing damage to plant life [64]. It is well known that nitric oxide (NO) is a ubiquitous signal molecule involved in many life processes of plants, seed germination, hypocotyl elongation, leaf, expansion, root growth, lateral roots initiation, and apoptosis, and so forth. It is also involved in multiple plant responses to environmental stress. Data demonstrated that, at lower concentrations of exogenous NO with 50 mmol L⁻¹ SNP, the activation of POD, SOD, and CAT was significantly increased, and the MDA content decreased with 50mM SNP. NO could protect *D. huoshanense* against the oxidative insult caused by a drought stress; meanwhile, a high level of RWC can be maintained. Furthermore, it suggests that the molecular messenger NO can trigger epigenetic variation and increase the demethylation ratio of methylated sites for *D. huoshanense* by using DNA analysis. These results may imply that the expression of some genes is involved in the response to drought stress triggered by NO [65].

Numerous *Dendrobium* species such as *D. nobile*, *D. denneanum*, *D. huoshanense*, *D. chrysotoxum*, *D. moniliforme*, *D. tosaense*, *D. linawianum*, *D. candidum*, *D. loddigesii*, and *D. fimbriatum* have polysaccharides as the active compounds. Polysaccharides play important roles in many biological processes and are used to treat various diseases [66, 67]. Polysaccharides isolated from *D. nobile*, *D. huoshanense*, *D. chrysotoxum*, and *D. fimbriatum* species manifest antioxidant and free radical scavenging activities. *D. nobile* polysaccharide displayed the highest scavenging activity toward hydroxyl, ABTS, and DPPH free radicals [40, 68, 69]. However, *D. fimbriatum* polysaccharide has significant scavenging

TABLE 1: The compounds of various *Dendrobium* species.

Species	Constituent	Chemical formula	Structures of compound	Reference
<i>D. amoenum</i>	Bibenzyls derivatives			
	Amoenylin	$R_1 = R_3 = \text{Me}, R_2 = \text{OH}, R_4 = \text{H}, R_5 = \text{OMe}$		
	Isoamoenylin	$R_1 = R_3 = \text{Me}, R_2 = \text{OMe}, R_4 = \text{OH}, R_5 = \text{H}$		
	Moscatilin (analogues)	—		
	3,4'-Dihydroxy-5-methoxybibenzyl	—		
	—	$R_1 = R_2 = R_4 = \text{H}, R_3 = \text{Me}, R_5 = \text{OH}$		[19]
	—	$R_1 = R_2 = \text{Me}, R_2 = \text{OH}, R_4 = \text{H}, R_5 = \text{OMe}$		
	—	$R_1 = R_3 = \text{Me}, R_2 = \text{OAc}, R_4 = \text{H}, R_5 = \text{OMe}$		
	—	$R_1 = R_3 = \text{Me}, R_2 = \text{OMe}, R_4 = \text{OAc}, R_5 = \text{H}$		
	—	$R_1 = \text{Ac}, R_2 = R_4 = \text{H}, R_3 = \text{Me}, R_5 = \text{OAc}$		
	—	$R_1 = R_3 = \text{Me}, R_2 = R_5 = \text{OH}, R_4 = \text{OMe}$		
	—	$R_1 = R_3 = \text{Me}, R_2 = R_4 = \text{OMe}, R_5 = \text{OH}$		
	—	$R_1 = R_3 = \text{Me}, R_2 = R_5 = \text{OMe}, R_4 = \text{OAc}$		
	—	$R_1 = R_2 = R_5 = \text{H}, R_3 = \text{Me}, R_4 = \text{OH}$		
	—	$R_1 = \text{Ac}, R_2 = R_5 = \text{H}, R_3 = \text{Me}, R_4 = \text{OAc}$		
	—	$R_1 = R_2 = R_3 = R_4 = \text{H}, R_5 = \text{OH}$		
<i>D. candidum</i>	Bibenzyls derivatives			
	Dendrocandin C	$R_1 = R_2 = R_5 = \text{OH}, R_3 = R_6 = \text{OCH}_3,$ $R_4 = \text{H}$		[20]
	((S)-3,4,4-trihydroxy-5,α-dimethoxybibenzyl)	$R_1 = R_2 = R_5 = \text{OH}, R_3 = \text{OCH}_3, R_6 = \text{OCH}_2,$ $\text{CH}_3, R_4 = \text{H}$		
	Dendrocandin D	$\text{CH}_3, R_4 = \text{H}$		
	((S)-3,4,4-trihydroxy-5-methoxy-α-ethoxybibenzyl)	$R_1 = R_2 = R_4 = R_5 = \text{OH}, R_3 = \text{OCH}_3,$ $R_6 = \text{H}$		
	Dendrocandin E			
	(3,3,4,4-tetrahydroxy-5-methoxybibenzyl)			
	Amotin (type 1, sesquiterpenoids)	—		[19]

TABLE 1: Continued.

Species	Constituent	Chemical formula	Structures of compound	Reference
	Amoenin (type 2 sesquiterpenoids)	—		[19]
	Flaccidin (9,10-dihydrophenanthropyran)	—		[19]
	Bibenzyls derivatives			
	4,4-Dihydroxy-3,5-dimethoxybibenzyl	$R_2 = R_5 = OH, R_1 = R_3 = OCH_3, R_4 = R_6 = OH$		
	3,4-Dihydroxy-5,4 dimethoxybibenzyl	$R_1 = R_2 = OH, R_3 = R_5 = OCH_3, R_4 = R_6 = OH$		
	3-O-Methylgigantol	$R_1 = OH, R_3 = R_4 = R_5 = OCH_3, R_2 = R_6 = H$		
	Dendrophanol	$R_2 = R_4 = R_5 = OH, R_1 = R_3 = OCH_3, R_6 = H$		
	Gigantol	$R_1 = R_5 = OH, R_3 = R_4 = OCH_3, R_2 = R_6 = H$		[21]
	New compounds			
	Dendrocandin A	$R_1 = R_2 = OH, R_3 = R_5 = R_6 = OCH_3,$ $R_4 = H$		
	((R)-3,4-dihydroxy-5,4,α-trimethoxybibenzyl)			
	Dendrocandin B			[21]
	(4-[2-((2S,3S)-3-(4-hydroxy-3,5-dimethoxyphenyl))-2-hydroxymethyl]-8-methoxy-2,3-dihydrobenzo[1,4]dioxin-6-yl]ethyl)-1-methoxybenzene)	—		

TABLE 1: Continued.

Species	Constituent	Chemical formula	Structures of compound	Reference
<i>D. apiphyllum</i>	Aromatic compounds			
	Flavanthrin	—		[22]
	Coelonin Iusanthridin	$R_1 = OH, R_2 = OMe$ $R_1 = OMe, R_2 = OH$		[22]
	Moscatin	—		[22]
	Gigantol Batatasin III	$R_1 = R_3 = OMe, R_2 = OH$ $R_1 = OH, R_2 = H, R_3 = OMe$		[22]

TABLE 1: Continued.

Species	Constituent	Chemical formula	Structures of compound	Reference
<i>D. cariniferum</i>	Dibutyl phthalate	$R = CH_2CH_2CH_2CH_3$	 	[22]
	Diisobutyl phthalate	$R = CH_2CH(CH_3)_2$		
<i>D. cariniferum</i>	<i>P</i> -hydroxyphenylpropionic methyl ester	—		[22]
<i>D. chrysotoxum</i>	Phenanthrenequinone	—		[23]
	Dendronone (5-hydroxy-7-methoxy-9,10-dihydro-1,4-phenanthrenequinone)	—		
<i>D. chrysotoxum</i>	Aromatic organic compounds (Fluorenones)	—		[24]
	Dendroflorin Denchrysan A	$R_1 = H, R_2 = OH$ $R_1 = OH, R_2 = H$		
<i>D. chrysotoxum</i>	1,4,5-trihydroxy-7-methoxy-9H-fluoren-9-one	—		[24]

TABLE 1: Continued.

Species	Constituent (Fluorenol)	Chemical formula	Structures of compound	Reference
<i>D. densiflorum</i>	(9R)-4methoxy-9H-fluorene-2,5,9-triol	R = H		[25]
	Phenanthrene			
	2,7-dihydroxy-8methoxyphenanthro[4,5-bcd]pyran-5(5H)-one	—		[25]
	Bibenzyl			
<i>D. densiflorum</i>	Densiflorol A	—		[26]
	Phenanthrenedione			
	Densiflorol B	R = H		[26]

TABLE I: Continued.

Species	Constituent	Chemical formula	Structures of compound	Reference
	Dendroflorin	R = OH		[26]
Known as oleanolic acid and b-sitosterol compounds				
	(A) Dengibsin	R = H		[26]
	(B) Cypripedin	R = OMe		[26]
	(C) Gigantol (D) Moscatilin (E) Tristin	R = OH, R = H, R = OMe R = R = OMe, R = OH R = R = OH, R = H		[26]
	(F) Naringenin (G) Homoeriodictyol	R = H R = OMe		[26]

TABLE 1: Continued.

Species	Constituent	Chemical formula	Structures of compound	Reference
	(H) Moscatin	—		[26]
	(I) 2,6-dihydroxy-1,5,7-trimethoxyphenanthrene	—		[26]
	(J) 4,7-dihydroxy-2-methoxy-9,10-dihydrophenanthrene	—		[26]
	(K) Scopoletin (L) Scopoletin (M) Ayapin	$R = R = \text{OMe}$ $R = \text{OMe}, R = \text{OH}$ $R + R = \text{OCH}_2\text{O}$		[26]
<i>D. longicornu</i>				
	Monoaromatic compounds			
	Bis(2-ethylhexyl)phthalate Dibutyl phthalate	$\text{CH}_2\text{CH}(\text{C}_2\text{H}_5)(\text{CH}_2)_3\text{CH}_3$ $(\text{CH}_2)_3\text{CH}_3$		[27]
	Ethyl haematommate	$\text{C}_{11}\text{H}_{12}\text{O}_5$		[27]

TABLE I: Continued.

Species	Constituent	Chemical formula	Structures of compound	Reference
	Methyl <i>B</i> -orcinol carboxylate	$C_{10}H_{12}O_4$		[27]
	<i>N</i> -Docosyl <i>trans</i> -ferulate Ferulaldehyde	$R = COOCH_2(CH_2)_{20}CH_3$ $R = CHO$		[27]
<i>D. nobile</i>	Sesquiterpenes (Dendronobilins A)			[28]
	Copacamphane-type	(1 <i>R</i> ,2 <i>R</i> ,4 <i>S</i> ,5 <i>S</i> ,6 <i>S</i> ,8 <i>S</i> ,9 <i>R</i>)-2,8-Dihydroxycopacamphan-15-one		[28]
	Picrotoxane-type	(2 <i>b</i> ,3 <i>b</i> ,4 <i>b</i> ,5 <i>b</i>)-2,4,11-Trihydroxy-picrotoxano-3(15)-lactone		[28]

TABLE 1: Continued.

Species	Constituent	Chemical formula	Structures of compound	Reference
	Picrotoxane-type	(2b,3b,5b,9a,11b)-2,11-Epoxy-9,11,13-trihydroxypicrotoxano-3(15)-lactone		[28]
	Picrotoxane-type	(2b,3b,5b,12R*)-2,11,13-Trihydroxypicrotoxano-3(15)-lactone		[28]
	Picrotoxane-type	(2b,3b,5b,12S*)-2,11,13-Trihydroxypicrotoxano-3(15)-lactone		[28]

TABLE 1: Continued.

Species	Constituent	Chemical formula	Structures of compound	Reference
Picrotoxane-type		(2b,3b,5b,9a)-,10-Cyclo-2,11,13-trihydroxy-picrotoxano-3(15)-lactone		[28]
Murolene-type		(9b,10a)-Murol-4-ene-9,10,11-triol		[28]
Alloaromadendrene-type		(10a)-Alloaromadendrene-10,12,14-triol		[28]
Cyclocopacamphane-type		(5b)-Cyclocopacamphane-5,12,15-triol		[28]

TABLE 1: Continued.

Species	Constituent	Chemical formula	Structures of compound	Reference
Bibenzyl derivatives				
Dendronophenol A		2,2',9,9'-Tetramethoxy13',14'-peroxy-1,1'-bis(bibenzyl)-6,6'-diol		[29]
			[29]
Dendronophenol B		1-(2',6'-Dimethoxy7',8'-peroxyphenyl-propyl)-2,9-dimethoxybibenzyl-6,9'-diol		
<i>D. thyrsiflorum</i>				
Bi-bicyclic and bi-tricyclic compounds				
Denthyrsin		[3-(5',6'-Dimethoxybenzofuran-2'-yl)-6,7-dimethoxy-2H-chromen-2-one		[30]
			[30]

TABLE 1: Continued.

Species	Constituent	Chemical formula	Structures of compound	Reference
	Denthyrsinone	(7,4',7'-Trihydroxy-2,2',8'-trimethoxy- [5,1' biphenanthrenyl]-1,4-dione)		[30]
	Denthyrsinin	1,5,7-Trimethoxyphenanthrene-2,6-diol		[30]
	Scoparone	—		[30]
	b-Sitosterol	—		[30]
	Daucosterol	—		[30]

TABLE 2: Polysaccharides DDP1-1, DDP2-1, and DDP3-1 isolated from *D. denneanum* and DNP1-1 DNP2-1, DNP3-1, and DNP4-2 isolated from *D. nobile*.

	ABTS radicals	Hydroxyl radicals	DPPH radicals
<i>D. denneanum</i>			
DDP1-1	(LW)	(LW)	(LW)
DDP2-1	(LW)	(HH)	(HH)
DDP3-1	(LW)	(LW)	(LW)
<i>D. nobile</i>			
DNP1-1	(LW)	(LW)	(LW)
DNP2-1	(LW)	(LW)	(LW)
DNP3-1	(LW)	(LW)	(LW)
DNP4-2	(HH)	(HH)	(HH)

*Notes: Low (LW); High (HH).
Up the table: the antioxidant effect of DNP4-2 on ABTS, hydroxyl, and DPPH free radicals, suggesting the levels of DNP4-2 higher than DNP1-1, DNP2-1, and DNP3-1; however, the antioxidant effects of DDP2-1 on hydroxyl and DPPH free radicals were higher than ABTS free radicals. Moreover, the DDP1-1 and DDP 3-1 on inhibitory effect were low than other compounds. We believe that the focus on antioxidant potential of DNP4-2 and DDP2-1 structures in near future.

action toward ABTS free radicals, and it also shows a weak DPPH free radical scavenging action. On the contrary, *D. denneanum* polysaccharide exerts a powerful DPPH free radical scavenging action, but its ABTS scavenging effect is not obvious [54]. In addition, both *D. huoshanense* and *D. chrysotoxum* polysaccharides reveal an insufficient ABTS free radical scavenging effect. However, it manifests potential hydroxyl radical scavenging activity. Overall, the free radical scavenging activities of polysaccharides on hydroxyl and DPPH free radicals remain at a high level, but the inhibitory effect on ABTS free radicals is weak. The above results, compared with other types of polysaccharides from *D. denneanum* and *D. nobile*, are shown in Tables 2 and 3, and others are shown in Table 4 [41, 53]. The polysaccharide of *D. huoshanense* has a backbone of (1 → 4)-linked α-D-Glcp, (1 → 6)-linked α-D-Glcp, and (1 → 4)-linked β-D-Manp from mannose (Man), glucose (Glc), and a trace of galactose (Gal). Its antioxidant effect in the livers of CCl4-treated mice was evidenced by a decrease of malondialdehyde (MDA) and increase of superoxide dismutase (SOD), catalase (CAT), and glutathione peroxidase (GPx) [70].

3.1.1. The Antioxidant Activities of Bibenzyl Derivatives, Phenanthrenes, and Stilbenes from *D. candidum* and *D. loddigesii*. Four new bibenzyl derivatives, dendrocandin F, dendrocandin G, dendrocandin H, and dendrocandin I, are extracted from *D. candidum*. They act as antioxidant agents to clear up the free radicals. Accordingly, the IC50 values of dendrocandin F and dendrocandin G are 55.8 mM and 32.4 mM, respectively [49]. A series of structurally related compounds from *D. loddigesii*, known as loddigesiinol A–D, suppress the production of nitric oxide (NO) with IC50 values of 2.6 mM for loddigesiinol A, 10.9 mM for loddigesiinol B, and 69.7 mM for loddigesiinol D. Further study

TABLE 3: Assessment of the ABTS, hydroxyl, and DPPH scavenging abilities of DNP and compounds DNP4-2, DNP2-1, DNP3-1, and DNP1-1 derived from DNP.

DNP, DNP4-2, DNP2-1, DNP3-1, and DNP1-1			
0.5 mg for DPPH scavenging (%)		1 mg for DPPH scavenging (%)	
Type	Scavenging (%)	Type	Scavenging (%)
DNP	20	DNP	20–30
DNP4-2	20–30	DNP4-2	30–40
DNP2-1	5–10	DNP2-1	10–20
DNP3-1	About 5	DNP3-1	<5
DNP1-1	<5	DNP1-1	About 5
0.5 mg for ABTS scavenging (%)		1 mg for ABTS scavenging (%)	
Type	Scavenging (%)	Type	Scavenging (%)
DNP	30–40	DNP	<60
DNP4-2	About 40	DNP4-2	>60
DNP2-1	<20	DNP2-1	About 40
DNP3-1	≥20	DNP3-1	30–40
DNP1-1	10	DNP1-1	30
0.5 mg for hydroxyl scavenging (%)		1 mg for hydroxyl scavenging (%)	
Type	Scavenging (%)	Type	Scavenging (%)
DNP	30–40	DNP	35–40
DNP4-2	20–30	DNP4-2	30–35
DNP2-1	10	DNP2-1	10–15
DNP3-1	5–10	DNP3-1	10
DNP1-1	0	DNP1-1	5
Based on data for DNP, DNP4-2, DNP2-1, DNP3-1, and DNP1-1 scavenging (%)			
DPPH scavenging % (0.5 mg content): DNP4-2 > DNP > DNP2-1 > DNP3-1 > DNP1-1			
DPPH scavenging % (1 mg content): DNP4-2 > DNP > DNP2-1 > DNP1-1 > DNP3-1			
ABTS scavenging % (0.5 mg content): DNP4-2 > DNP > DNP3-1 > DNP2-1 > DNP1-1			
ABTS scavenging % (1 mg content): DNP4-2 > DNP > DNP2-1 > DNP3-1 > DNP1-1			
Hydroxyl scavenging % (0.5 mg content): DNP > DNP4-2 > DNP2-1 > DNP3-1 > DNP1-1			
Hydroxyl scavenging % (1 mg content): DNP > DNP4-2 > DNP2-1 > DNP3-1 > DNP1-1			

Up the table: it is shown that DPPH and ABTS free radical scavenging (%) of DNP4-2 are better than other compounds; however, the hydroxyl free radical scavenging (%) of DNP is better than other compounds.

indicated that phenanthrenes and dihydrophenanthrenes in *D. loddigesii* have significant effects on inhibiting the production of NO and DPPH free radicals. These compounds are known as (numbered as 1a–7a) phenanthrenes (1a), phenanthrenes (2a), phenanthrenes (3a), phenanthrenes (4a), dihydrophenanthrenes (5a), dihydrophenanthrenes (6a), and dihydrophenanthrenes (7a). Based on the above results, the scavenging activity of loddigesiinols toward NO free radicals

TABLE 4: Comparison of the DPPH, hydroxyl, and ABTS free radical scavenging abilities of polysaccharides from *D. huoshanense*, *D. chrysotoxum*, and *D. fimbriatum* species.

<i>Dendrobium</i> species	Chemical compounds	DPPH concentration (mg/mL)			
		0.5%	1%	2%	3%
<i>D. huoshanense</i> and <i>D. chrysotoxum</i>	Polysaccharides (scavenging effects %)	45–50	AB 80	90–95	—
<i>D. fimbriatum</i>	Polysaccharides (scavenging effects %)	—	10	—	40–50
<i>Dendrobium</i> species	Chemical compounds	Hydroxyl concentration (mg/mL)			
		0.5%	1%	2%	3%
<i>D. huoshanense</i> and <i>D. chrysotoxum</i>	Polysaccharides (scavenging effects %)	50	60–65	80	—
<i>D. fimbriatum</i>	Polysaccharides (scavenging effects %)	—	55–60	—	70–75
<i>Dendrobium</i> species	Chemical compounds	ABTS concentration (mg/mL)			
		0.5%	1%	2%	3%
<i>D. huoshanense</i> and <i>D. chrysotoxum</i>	Polysaccharides (scavenging effects %)	AB 10	AB 20	AB 25	—
<i>D. fimbriatum</i>	Polysaccharides (scavenging effects %)	—	AB 70	—	AB 90

* Notes: —: no tests and about: AB.

The scavenging hydroxyl free radicals abilities of polysaccharides from *D. huoshanense* and *D. chrysotoxum* are stronger than *D. fimbriatum*, and the scavenging ABTS free radicals abilities of polysaccharides from *D. fimbriatum* are stronger than *D. huoshanense* and *D. chrysotoxum* polysaccharides. Results showed that polysaccharides (scavenging DPPH effects %) from *D. huoshanense* and *D. chrysotoxum* are higher than hydroxyl and ABTS. And polysaccharides (scavenging ABTS effects %) from *D. fimbriatum* are higher than DPPH and hydroxyl free radicals.

is pronounced. The inhibitory actions of phenanthrenes and dihydrophenanthrenes from *D. loddigesii* on the production of NO and DPPH free radicals are recognised. The relevant data and information are shown in Table 5 [71].

3.2. Anti-Inflammatory Activity

3.2.1. Inhibition of Nitric Oxide (NO) by *D. nobile* and *D. chrysanthum* Constituents. It is known that, upon lipopolysaccharide (LPS) stimulation, macrophages produce a large amount of inflammatory factors, such as tumor necrosis factor α (TNF- α), interleukin-1 beta (IL-1 β), interleukin 6 (IL-6), interferon (IFN), and other cytokines. LPS induces endogenous nitric oxide (NO) biosynthesis through induction of inducible nitric oxide synthase (iNOS) in macrophages, which is involved in inflammatory responses [42]. *D. nobile* and *D. chrysanthum* constituents strongly inhibit TNF- α , IL-1 β , IL-6, and the NO production.

3.2.2. *D. nobile* Constituents. From *D. nobile* two new bibenzyl derivatives, namely, nobilin D (given number 1) and nobilin E (given number 2), and a new fluorenone, namely, nobilone (given number 3), together with seven known compounds (given numbers 4–10), including $R_1 = R_2 = OCH_3$, $R_3 = OH$, $R_4 = H$ (4), $R_1 = R_2 = R_3 = OCH_3$, $R_4 = H$ (5), $R_1 = R_2 = OH$, $R_3 = R_4 = H$ (6), $R_1 = R_3 = OCH_3$, $R_2 = OH$, $R_4 = H$ (7), $R_1 = OCH_3$, $R_2 = R_3 = OH$, $R_4 = H$ (8), $R_1 = R_3 = OH$, $R_2 = R_4 = H$ (9), and $R_1 = R_3 = OH$, $R_2 = HR_4 = OCH_3$ (10), have been identified. Compounds 1–3, 5, and 8–10 can inhibit NO production. On the other hand, compounds 2 and 10 can exhibit a strong to resveratrol, while enhanced cytotoxic potential has been found in compounds 4 and 7. Compounds 1–10 act as NO inhibitors, as evidenced by oxygen radical absorbency capacity (ORAC) assays. These findings are focused on all of these compounds, except

compound 6, and their inhibitory effects on NO production, in murine macrophages (RAW 264.7), activated by LPS and interferon (IFN- γ) are shown in Table 5 [38]. Furthermore, it was found that a new phenanthrene, together with nine known phenanthrenes and three known bibenzyls, manifests an inhibitory effect on NO production in RAW 264.7 cells. The structures of all of the compounds (given numbers A–M), including $R_1 = R_3 = OH$, $R_2 = R_5 = OCH_3$, $R_4 = H$ (A), $R_1 = R_4 = R_5 = OH$, $R_2 = H$, $R_3 = OCH_3$ (B), $R_1 = R_5 = OH$, $R_2 = R_3 = OCH_3$, $R_4 = H$ (C), $R_1 = R_2 = OCH_3$, $R_3 = R_5 = OH$, $R_4 = H$ (D), $R_1 = H$, $R_2 = R_3 = OH$ (E), $R_1 = H$, $R_2 = OH$, $R_3 = OCH_3$ (F), $R_1 = OCH_3$, $R_2 = R_4 = OH$, $R_3 = R_5 = H$ (G), $R_1 = R_3 = OH$, $R_2 = OCH_3$ (H), $R_1 = R_5 = OH$, $R_2 = R_4 = H$, $R_3 = OCH_3$ (I), $R_1 = R_3 = OCH_3$, $R_2 = R_5 = OH$, $R_4 = H$ (J), $R_1 = R_2 = R_4 = OH$, $R_3 = OCH_3$, $R_5 = H$ (K), $R_1 = OCH_3$, $R_2 = R_4 = H$, $R_3 = R_5 = OH$ (L), and $R_1 = R_2 = R_3 = OH$, $R_4 = R_5 = H$ (M), were elucidated by analysis of the spectroscopic data, including extensive two-dimensional nuclear magnetic resonance spectroscopy (2D-NMR) and mass spectrometry (MS). All of compounds C, D, I, K, and L are relatively strong inhibitors of NO production, and their potencies are better than those of compounds A and E–H [58]. Compounds C, D, I, K, and L could inhibit NO synthesis which might contribute to the suppression of LPS-induced TNF- α and IL-1 β production.

3.2.3. *D. chrysanthum* Constituents. Anti-inflammatory compounds from ethyl acetate extracts of *D. chrysanthum* species were evaluated. These structures were elucidated on the basis of the high-resolution mass spectrometry, NMR spectroscopy, and X-ray crystal diffraction analysis. A novel phenanthrene phenol derivative with a spironolactone ring (dendrochrysanene), namely, 2-hydro-7,70,100-trihydroxy-4-40-dimethoxylspiro[(1H)-cyclopenta[a]naphthalene-3,30-(20H)-phenanthro[2,1-b]furan]-1,20-dione, was identified. Results showed anti-inflammatory activity of dendrochry-

TABLE 5: Inhibition of compounds from *D. loddigesii* and *D. nobile* species on NO and DPPH formation.

Species	Chemical constituent	NO	DPPH
<i>D. loddigesii</i>	Phenanthrenes (1a)	2.6	26.1
	Phenanthrenes (2a)	6.4	59.8
	Phenanthrenes (3a)	5.3	12
	Phenanthrenes (4a)	10.9	—
	Dihydrophenanthrenes (5a)	4.6	62.2
	Dihydrophenanthrenes (6a)	29.1	—
	Dihydrophenanthrenes (7a)	29.2	—
	Loddigesiinols A	2.6	—
	Loddigesiinols B	10.9	—
	Loddigesiinols C	—	23.7
<i>D. nobile</i>	Loddigesiinols D	69.7	—
	Nobilin D (1)	15.3	—
	Nobilin E (2)	19.2	—
	Nobilone F (3)	38.1	—
	Phenanthrenes and bibenzyls related-composition (4–10)	—	—
	RO $R_1 = R_2 = \text{OCH}_3$ $R_3 = \text{OH}$ $R_4 = \text{H}$ (4)	—	—
	RO $R_1 = R_2 = R_3 = \text{OCH}_3$ $R_4 = \text{H}$ (5)	48.2	—
	RO $R_1 = R_2 = \text{OH}$ $R_3 = R_4 = \text{H}$ (6)	—	—
	RO $R_1 = R_3 = \text{OCH}_3$ $R_2 = \text{OH}$ $R_4 = \text{H}$ (7)	—	—
	RO $R_1 = \text{OCH}_3$ $R_2 = R_3 = \text{OH}$ $R_4 = \text{H}$ (8)	36.8	—
	RO $R_1 = R_3 = \text{OH}$ $R_2 = R_4 = \text{H}$ (9)	32.9	—
	RO $R_1 = R_3 = \text{OH}$ $R_2 = \text{H}$ $R_4 = \text{OCH}_3$ (10)	13.4	—

*Notes: —: no tests; RO: relation composition.

Inhibitory effects of different compounds from *D. loddigesii* and *D. nobile* species on NO production. Result shown in Table 5, ranked from high to low as follows: loddigesiinol D > $R_1 = R_2 = R_3 = \text{OCH}_3$ $R_4 = \text{H}$ (5) > nobilone F (3) > $R_1 = \text{OCH}_3$ $R_2 = R_3 = \text{OH}$ $R_4 = \text{H}$ (8) > $R_1 = R_3 = \text{OH}$ $R_2 = R_4 = \text{H}$ (9) > dihydrophenanthrenes (7a) > dihydrophenanthrenes (6a) > nobilin E (2) > nobilin D (1) > $R_1 = R_3 = \text{OH}$ $R_2 = \text{H}$ $R_4 = \text{OCH}_3$ (10) > phenanthrenes (4a) = loddigesiinols B > phenanthrenes (2a) > phenanthrenes (3a) > dihydrophenanthrenes (5a) > phenanthrenes (1a) > loddigesiinols A, and all of the compounds inhibit DPPH production (from high to low): dihydrophenanthrenes (5a) > phenanthrenes (2a) > phenanthrenes (1a) > loddigesiinol C > phenanthrenes (3a).

sanene on iNOS mRNA induced by LPS in mouse peritoneal macrophages. Meanwhile, there are several inflammatory cytokines, such as TNF- α , IL-8, and IL-10, which were inhibited. Thus, the beneficial effects of dendrochrysane compounds as anti-inflammatory agents were identified [72].

3.3. Sjögren's Syndrome (SS). Sjögren's syndrome (SS) is a chronic autoimmune disease with disorder of the exocrine glands. The symptoms are dry eyes, dry mouth, dry throat and thirst with blurred vision, and so forth. The abnormal distribution of salivary glands in SS leads to an inflammatory effect and pathological processes triggering lymphocyte infiltration and apoptosis. Lymphocyte infiltration and apoptotic pathways, shown by regulation of Bax, Bcl-2, and caspase-3, have been reported in submandibular glands (SG). Furthermore, in pathological processes were also found

that the subsequent signal of cell expression to mRNA of proinflammatory cytokines, such as tumor necrosis (TNF- α), interleukin (IL-1 β), and IL-6. As well as, a high level of expression of aquaporin-5 (AQP-5) is identified. AQP-5 is one of a water channel protein, and plays an important role in 31 the salivary secretions [73, 74].

Treatment with *D. officinale* polysaccharides (DP) could suppress the progression of lymphocyte infiltration and apoptosis in SS and rectify the chaos of proinflammatory cytokines including TNF- α , IL-1 beta, and IL-6 in SG [71, 75] (Figure 1). mRNA expression of TNF- α was inhibited. The process of regulation resulted in a series of marked responses, such as translocation of NF- κ B, prolonged MAPK, cytochrome C release, and caspase-3 activation, which have been identified [55, 76]. In addition to the findings on DP treatment, there are reports on the findings on the extracts of *D. candidum* and *D. nobile*.

A clinical study found that, after 1 week of oral administration of the extract of *D. candidum*, salivary secretion was increased by approximately 65%. The extract promoted the expression of aquaporin-5 useful for treatment of Sjögren's syndrome (SS).

Chrysotoxine isolated from *D. nobile* can increase the expression of AQP-5 in dry eyes, one of the symptoms of SS, and restore the distribution of AQP-5 in lacrimal glands and corneal epithelia, by inhibiting the release of cytokines, such as IL-1, IL-6, and TNF- α . In the meantime, it can activate the mitogen-activated protein kinase (MAPK) signaling pathways. Furthermore, it elicits an increase in the production of matrix metalloproteinase-9 (MMP-9). Ultimately, it leads to an increase of saliva and tear secretion. The medical efficacy of the extracts of *D. officinale*, *D. nobile*, and *D. candidum* species has been demonstrated [43] (Figure 1).

3.4. Neuroprotective Effect (Parkinsonian Syndrome). Five bioactive derivatives isolated from *Dendrobium* species, namely, chrysotoxine (CTX), moscatilin, crepidatin, nobilin B, and chrysotobibenzyl, are potential neuroprotective compounds with antioxidant activity which can be used in the treatment of Parkinson's disease (PD). The IC_{50} values of DPPH free radical scavenging activities of chrysotoxine (CTX), moscatilin, crepidatin, and nobilin B were 20.8 ± 0.9 , 28.1 ± 7.1 , 38.2 ± 3.5 , and 22.2 ± 1.4 , respectively. The above data show that crepidatin is better than other compounds in DPPH free radical scavenging capacity. CTX could selectively antagonize MPP⁺ in dopaminergic pathways in the brain; also 6-hydroxydopamine (6-OHDA) has been inhibited by mitochondrial protection and NF- κ B modulation in SH-SY5Y cells. Thus, it could explain how it may be beneficial in preventing PD [44, 77].

In addition, results of CTX compound in *Dendrobium nobile* Lindl. used in treatment of PD have also been clearly reported. We evaluated the pharmacokinetics of oral (100 mg/kg) and intravenous (25 mg/kg) administration of CTX preparation using high performance liquid chromatography-tandem mass spectrometric (HPLC-MS/MS) method in animal tests [78]. Results indicate efficacy

TABLE 6: The chemical compounds from *D. nobile* and *D. moniliforme* that inhibited/activated T cells and B cells.

Dendrobium species	Compound/structure	Lymphocytes	
		T cell	B cell
<i>D. nobile</i>	Dendroside D	(AD)	(AD)
	Dendroside E	(AD)	(AD)
	Dendroside F	(AD)	(AD)
	Dendroside G	(AD)	(AD)
<i>D. moniliforme</i>	Dendroside A	(IY)	(AD)
	Dendroside C	(IY)	(AD)
	Vanilloloside	(IY)	(AD)
	Denbinobin	(IY)	(AD)
	2,6-Dimethoxy 1,4,5,8-phenanthradiquinone	(IY)	(AD)

* Notes: activation: AD; inhibition: IY.

Up the table, showed T cell and B cell activated by chemical compounds from *D. nobile*, and inhibit T cell and activated on B cell by chemical compounds from *D. moniliforme*.

evidence on regulation of hematopoietic growth factors and cytokines factors in the immune system by different polysaccharides from *D. huoshanense* [81, 82]. Interestingly, the total polysaccharides of *D. huoshanense* induced the expression of interleukin-1 receptor antagonist (IL-1ra) in human monocytes. Which the IL-1ra was regulated by 37 a series of signaling pathways, like ERK/ELK, p38 MAPK, PI3K, and NF- κ B [83].

In addition, the water-soluble polysaccharides (DTP) derived from *D. tosaense* were isolated by using HPAEC-PAD, HP-SEC, GC-MS, and NMR spectroscopy. They significantly increased the number of natural killer (NK) cells, NK cytotoxicity, macrophage phagocytosis, and cytokine induction in splenocytes, when administered orally to BALB/c mice for 3 weeks [59].

3.6. Treatment of Diabetes. The compounds isolated from *Dendrobium huoshanense* (DHP) have been evaluated for treatment of diabetes in recent years. The hypoglycemic and antioxidative activities of three polysaccharides, namely, *D. officinale* (DOP), *D. nobile* (DNP), and *D. chrysotoxum* (DCP), have been compared in diabetic mice. Result showed that the hypoglycemic effect of DHP was better than those of DNP and DOP. In addition, the antioxidant effect of DHP was better than those of DOP and DNP by regulating the superoxide dismutase, catalase, malonaldehyde, and l-glutathione levels in the liver and kidney. Thus, the hypoglycemic and antioxidant activities of DHP need to be studied more extensively in the future [60].

3.7. Antitumor Activity

3.7.1. The Anticancer Effect of *D. chrysotoxum* and *D. nobile*. Fluorenone derivatives, namely, dendroflorin and denchrysan A, isolated from stems of *D. chrysotoxum* species, show significant inhibitory effects on the growth of human

hepatoma BEL-7402 cells. The IC₅₀ values of 1,4,5-trihydroxy-7-methoxy-9H-fluoren-9-one, denchrysan A 1, and dendroflorin were 1.49 μ g/mL, 1.38 μ g/mL, and 0.97 μ g/mL, respectively.

On the other hand, erianin, a phenanthrene from *D. chrysanthum*, is recognized as an antitumor agent. Reports showed that it has potent inhibitory activity in hepatoma Bel7402, melanoma A375, and HL-60 cells, as evidenced by inhibition of angiogenesis and induction of endothelial cytoskeletal disorganization *in vivo* and *in vitro*, but did not show antitumor activity [56, 61]. Thus, further investigation is needed to identify if the antitumor mechanisms of erianin and denbinobin are similar.

The antitumor effects of polysaccharides isolated from stems of *D. nobile*, namely, DNP-W, DNP-OH, and DNP-H, were studied. DNP-W was further fractionated to give six subfractions including DNP-W1, DNP-W2, DNP-W3, DNP-W4, DNP-W5, and DNP-W6 by using anion exchange chromatography. The results show that DNP-W, DNP-OH and DNP-H, DNP-W1, DNP-W2, and DNP-W3, especially DNP-W1 and DNP-W3, have a strong antitumor action in inhibiting sarcoma 180 *in vivo* and HL-60 cells *in vitro* [48]. In addition, denbinobin is a major phenanthrene isolated from stems of *D. nobile*. The inhibitory mechanisms of denbinobin in SNU-484 cells have been observed. It significantly decreased the expressions of matrix metalloproteinase (MMP-2) and (MMP-9) and induced apoptosis through downregulation of Bcl-2 and upregulation of Bax in cancer cells. These findings may be used to provide reference for further studies on the pharmacological mechanisms from denbinobin and polysaccharides in *D. nobile* [50, 51, 84].

3.7.2. Antilymphoma Activity of *Dendrobium formosum*. Lymphoma is a cancer of the immune system. The ethanolic extract of *Dendrobium formosum* showed antilymphoma activity *in vitro* as evaluated by the MTT assay. It also significantly prolonged survival time in Dalton's lymphoma bearing mice [85].

3.7.3. Anti-Lung-Cancer Activity of *Dendrobium draconis*. The bibenzyl compound gigantol has been isolated from *Dendrobium draconis*. It inhibits several cancer cell lines and inhibits filopodia formation of the non-small-cell lung cancer cells. The molecular mechanism of gigantol includes (1) downregulating caveolin-1 (Cav-1), (2) activating ATP-dependent tyrosine kinase, and (3) regulating cell division cycle 42 (Cdc42) [86].

3.8. Antimicrobial Activity and Antifungal Activity. *Dendrobium* is an important economic plant; however, there have been certain cultivation issues yet to be resolved, particularly fungal infection [87]. *Pythium vexans* has been reported to cause stem rot, crown rot, root rot, damping-off, and patch canker in *Dendrobium* plants [88]. The diseases could lead to damage to the seedlings of *D. chrysotoxum*, *D. chrysanthum*, *D. thyrsiflorum*, and *D. aurantiacum*, characterized by water-soaked, brown or yellowish lesions with a brown margin, resulting in dieback of the plants within a few days

TABLE 7: The pharmacological effects of *Dendrobium* species.

Species	Chemical compounds/structures	Pharmacological activities	References
<i>D. nobile</i>	Polysaccharides	Scavenging effect of hydroxyl, ABTS, and DPPH	[40, 41]
(SR)	Nobilin D, nobilin E, nobilone	Inhibitory on NO production	[42]
(SR)	Phenanthrenes	Inhibition of LPS-induced of nitric oxide production	[38]
(SR)	Dendroside A, Dendronobilosides A	Immunomodulatory activity	[43, 44]
(SR)	Glycosides (dendrosides D–G)	Significant Stimulation of the proliferation of mouse T and/or B lymphocytes	[43, 44]
(SR)	Polysaccharides (DNP-W1, DNP-W2, DNP-W3, DNP-W4)	Anticancer activity	[45]
(SR)	4,7-Dihydroxy-2-methoxy-9,10-dihydrophenanthrene, denbinobin	Anticancer activity:	[46]
(SR)	(SR)	(SNU-484 human gastric cancer cells)	[46]
(SR)	(SR)	(human lung carcinoma), SK-OV-3	[47]
(SR)	(SR)	(human ovary adenocarcinoma)	[47]
(SR)	(SR)	HL-60 (human promyelocytic leukemia) cell lines	[47]
(SR)	Bibenzyl	Inhibit on furylfuramide	[48]
(SR)	(SR)	4-nitroquinoline-1-oxide (4NQO)	[48]
(SR)	(SR)	N-Methyl-N'-nitro-N-nitrosoguanidine	[48]
(SR)	(SR)	UV irradiation	[48]
(SR)	(SR)	3-Amino-1,4-dimethyl-5H-pyrido[4,3b]indole (Trp-P-1)	[48]
(SR)	(SR)	Benzo[a]pyrene (B[a]P)	[48]
(SR)	(SR)	Aflatoxin B(1)	[48]
(SR)	Nobilin D	Anti-inflammatory activity	[42]
(SR)	Nobilin E	(SR)	[42]
(SR)	Nobilone	(SR)	[42]
(SR)	$R_1 = R_2 = \text{OCH}_3$ $R_3 = \text{OH}$ $R_4 = \text{H}$	(SR)	[42]
(SR)	$R_1 = R_2 = R_3 = \text{OCH}_3$ $R_4 = \text{H}$	(SR)	[42]
(SR)	$R_1 = R_3 = \text{OCH}_3$ $R_2 = \text{OH}$ $R_4 = \text{H}$	(SR)	[42]
(SR)	$R_1 = \text{OCH}_3$ $R_2 = R_3 = \text{OH}$ $R_4 = \text{H}$	(SR)	[42]
(SR)	$R_1 = R_3 = \text{OH}$ $R_2 = R_4 = \text{H}$	(SR)	[42]
(SR)	$R_1 = R_3 = \text{OH}$ $R_2 = \text{H}$ $R_4 = \text{OCH}_3$	(SR)	[42]
<i>D. loddigesii</i>	Phenanthrenes	Inhibition of nitric oxide (NO)	[49]
(SR)	Loddigesiinols A–D	(SR)	[49]
(SR)	Moscatilin, moscatin, moscatilin diacetate	Inhibited both AA and collagen-induced platelet aggregations	[50, 51]
(SR)	—	Inhibition of aggregation of rabbit platelets induced by arachidonic acid and collagen	[50, 51]
<i>D. huoshanense</i>	Polysaccharides	Inducing several cytokines, including IFN- γ , IL-10, IL-6, and IL-1	[52]
(SR)	(SR)	Immunostimulating activity	[52]
(SR)	(SR)	Increase of TNF- α production	[52]
(SR)	(SR)	Inducing several cytokines, including hematopoietic growth factors GM-CSF and G-CSF	[52]
<i>D. denneanum</i>	Polysaccharides	Scavenging effect of hydroxyl, ABTS, and DPPH	[53]
<i>D. fimbriatum</i>	Polysaccharides	Scavenging effect of hydroxyl, ABTS, DPPH	[54]
<i>D. candidum</i>	Bibenzyl	Antioxidant activity	[20]
<i>D. officinalis</i>	—	Immunomodulatory activity	[55]
<i>D. findlayanum</i>	—	Inhibition of <i>Alternaria alternata</i> and other	[56]

TABLE 7: Continued.

Species	Chemical compounds/structures	Pharmacological activities	References
<i>D. densiflorum</i>	Moscatilin, homoeriodictyol, scoparone, scopoletin, gigantol	Antiplatelet aggregation activity	[50, 51]
<i>D. nobile</i> , <i>D. chrysanthum</i>	Erianin	Anticancer activity: hepatoma Bel7402 melanoma A375 and HL-60 cells	[57]
<i>D. chrysanthum</i> , <i>D. huoshanense</i>	Dendrochrysanene	TNF- α , IL-8, IL-10, and iNOS mRNAs were induced readily from mouse peritoneal macrophages by LPS	[58]
<i>D. devonianum</i> , <i>D. thyrsoflorum</i>	<i>Epicoccum</i> sp., epicorazine A Epicorazine B	Inhibition of <i>S. aureus</i> , <i>E. coli</i> , and <i>B. subtilis</i>	[59, 60]
<i>D. devonianum</i> , <i>D. thyrsoflorum</i>	Pyrenophorol derivatives	Inhibition of <i>E.coli</i> , <i>Bacillus megaterium</i> , and <i>Microbotryum violaceum</i>	[61]

* Notes: —: no tests; SR: similar.

[89]. Interestingly, it also resulted in antimicrobial activity from endophytic bacteria isolated from *Dendrobium* species. Screening of various endophytic fungi from roots and stems of different *Dendrobium* species, including 53 endophytic fungi from *D. devonianum*; 23 endophytic fungi from *D. thyrsoflorum* by way of morphological and/or molecular biological methods, showed that 10 endophytic fungi from *D. devonianum* and 11 endophytic fungi from *D. thyrsoflorum* exhibited strong antimicrobial activity. *Phoma*, *Epicoccum*, and *Fusarium* isolated from two abovementioned *Dendrobium* species demonstrated antibacterial and antifungal activities. The pyrenophorol derivatives of *Phoma* possessed antagonistic activity against *E. coli*, *Bacillus megaterium*, and *Microbotryum violaceum*, and *Epicoccum* sp. slightly inhibited *S. aureus*, *E. coli*, and *B. subtilis*. *Fusarium* demonstrated inhibitory effect on different pathogens. Moreover, epicorazine A and epicorazine B derivatives of *Epicoccum nigrum* isolated from *D. thyrsoflorum* possessed antibacterial activity against *S. aureus* and *B. subtilis* and also inhibited the growth of Gram-positive and Gram-negative bacteria, with a potency stronger than that of ampicillin sodium [90–93].

3.9. Antimalarial and Antiherpetic Activities

3.9.1. Antimutagenic Activity. Moscatilin obtained from *D. nobile* is a naturally occurring antimutagenic bibenzyl compound. This compound can exert a suppressive effect on the mutagenic activity of furylfuramide, 4-nitroquinoline-1-oxide (4NQO), N-methyl-N'-nitro-N-nitrosoguanidine, UV irradiation, 3-amino-1,4-dimethyl-5H-pyrido[4,3b]indole (Trp-P-1), benzo[a]pyrene (B[a]P), and aflatoxin B(1) [94].

3.10. Diabetic Retinopathy (DR). Diabetes mellitus (DM) is one of the metabolic diseases. It can cause many complications, such as kidney failure, stroke, and foot ulcers. The ethanol extract of *Dendrobium chrysotoxum* (DC) used in treatment of diabetic retinopathy reduced the expression of a series of growing factors in DC-treated diabetic rats. It decreased the retinal mRNA expression level of vascular endothelial growth factor (VEGF), vascular endothelial growth factor receptor 2 (VEGFR2), serum growth factor

(SVEGF), matrix metalloproteinase (MMP) 2/9, basic fibroblast growth factor (bFGF), platelet-derived growth factor (PDGF) A/B, insulin-like growth factor 1 (IGF-1), interleukin 1 β (IL-1 β), interleukin 6 (IL-6), and phosphorylation of p65. Moreover, a decrease in the expression of intercellular adhesion molecule-1 (ICAM-1) has been reported [95].

3.11. Antiplatelet Aggregating Activity. Studies on the aggregation of platelets induced through thrombin, arachidonic acid (AA), thrombin, collagen, and platelet-activating factor (PAF) were conducted by using 4 different aggregation inducers. Animal tests disclosed that different *Dendrobium* species including *Dendrobium loddigesii* and *Dendrobium densiflorum* possess antiplatelet aggregating activity. The results indicated some compounds isolated from two *Dendrobium* species more effectively inhibited AA-induced aggregation than they inhibited aggregation induced by PAF and collagen [96]. In animal tests, the antiplatelet aggregation of *D. loddigesii* species, which those compounds in part from moscatilin, moscatin and diacetate. They inhibit platelet aggregation was induced by AA was completely abolished by fraction 4 (50 μ g/mL) in rabbit platelets. Meanwhile, further demonstrated there are strongly inhibited both AA and collagen-induced platelet aggregations about 100 μ g/mL. However, in PAF there are no significant effects. In addition, comparison of different compounds from *D. densiflorum* species, such as moscatilin, homoeriodictyol, scoparone, scopoletin, and gigantol, which the antiplatelet aggregation activity of their were identified *in vitro*. Especially for scoparone has been reported to possess potent antiplatelet aggregation who more than other compounds. Thus, it possible interacted by compounds-compounds for antiplatelet aggregation [26, 97, 98] (Table 7).

4. Conclusion

In this paper, the history, chemistry, and pharmacology of different *Dendrobium* species are reviewed. Especially for biological pharmacology (Table 7). The pharmacological activities examined include antioxidant, anti-inflammatory, immunomodulatory, antitumor, antimicrobial/antifungal,

antimutagenic activities, and antiplatelet aggregation activities. The ABTS, DPPH, and hydroxyl radical scavenging effects of different polysaccharides such as DNP, DNP2-1, DNP1-1, DNP3-1, and DNP4-2 have been investigated. Loddigesiinol, nobileone, dihydrophenanthrenes, and phenanthrenes have been evaluated for nitric oxide (NO) scavenging effect. Five types of polysaccharides from *D. huoshanense*, *D. chrysotoxum*, and *D. fimbriatum* species were compared. *D. huoshanense* and *D. chrysotoxum* polysaccharides exhibited stronger DPPH and hydroxyl scavenging activity while *D. fimbriatum* polysaccharides have stronger ABTS scavenging activity. Dihydrophenanthrenes and loddigesiinol D were isolated from *D. loddigesii* and *D. nobile* species. Dihydrophenanthrenes significantly inhibited DPPH production. Loddigesiinol D significantly inhibited NO production.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

This study was supported by Seed Funding Programme for Basic Research from The University of Hong Kong Project no. 201311159022 and Seed Funding Programme for Basic Research from the University of Hong Kong Project no. 201111159043.

References

- [1] R. L. Dressler, *Phylogeny and Classification of the Orchid Family*, Cambridge University Press, New York, NY, USA, 1993.
- [2] D. K. Singh, "Morphological diversity of the orchids of Orissa/Sarat Misra," in *Orchids: Science and Commerce*, P. Pathak, R. N. Sehgal, N. Shekhar, M. Sharma, and A. Sood, Eds., p. 35, BSMPs, New Delhi, India, 2001.
- [3] S. P. Vaughan, A. M. Grisoni, A. M. H. Kumagai, and A. A. R. Kuehnle, "Characterization of Hawaiian isolates of Cymbidium mosaic virus (CymMV) co-infecting *Dendrobium orchid*," *Archives of Virology*, vol. 153, no. 6, pp. 1185–1189, 2008.
- [4] M. M. Hossain, "Traditional therapeutic uses of some indigenous orchids of Bangladesh," *Medicinal and Aromatic Plant Science and Biotechnology*, vol. 42, no. 1, pp. 101–106, 2009.
- [5] H. P. Wood, *The Dendrobiums*, Timber Press, Portland, Ore, USA, 2006.
- [6] K. M. Cameron, M. W. Chase, W. M. Whitten et al., "A phylogenetic analysis of the Orchidaceae: evidence from rbcL nucleotide sequences," *American Journal of Botany*, vol. 86, no. 2, pp. 208–224, 1999.
- [7] R. L. Dressler, *The Orchids: Natural History and Classification*, Harvard University Press, Cambridge, Mass, USA, 1981.
- [8] M. A. Clements, "Molecular phylogenetic systematics in the Dendrobieinae (Orchidaceae), with emphasis on *Dendrobium* section *Pedilonum*," *Telopea*, vol. 10, pp. 247–298, 2003.
- [9] J. Xu, J. Guan, X. J. Chen, J. Zhao, and S. P. Li, "Comparison of polysaccharides from different *Dendrobium* using saccharide mapping," *Journal of Pharmaceutical and Biomedical Analysis*, vol. 55, no. 5, pp. 977–983, 2011.
- [10] K. R. Kirtiker and B. D. Basu, *Indian Medicinal Plants*, IV: Bishen Singh Mohendra Pal Singh, Dehradun, India, 2nd edition, 1975.
- [11] C. J. Bulpitt, "The uses and misuses of orchids in medicine," *QJM*, vol. 98, no. 9, pp. 625–631, 2005.
- [12] C. L. Withner, *The Orchid: A Scientific Survey*, Ronald Press Company, New York, NY, USA, 1959.
- [13] S. P. Das and S. K. Bhattacharjee, "Orchids," in *Herbaceous Perennials and Shade Loving Foliage Plants*, S. K. Bhattacharjee, Ed., Pointer Publishers, Jaipur, India, 2006.
- [14] J.-M. Kong, N.-K. Goh, L.-S. Chia, and T.-F. Chia, "Recent advances in traditional plant drugs and orchids," *Acta Pharmacologica Sinica*, vol. 24, no. 1, pp. 7–21, 2003.
- [15] M. Li, K. Y.-B. Zhang, P. P.-H. But, and P.-C. Shaw, "Forensically informative nucleotide sequencing (FINS) for the authentication of Chinese medicinal materials," *Chinese Medicine*, vol. 6, article 42, 2011.
- [16] T. B. Ng, J. Liu, J. H. Wong et al., "Review of research on *Dendrobium*, a prized folk medicine," *Applied Microbiology and Biotechnology*, vol. 93, no. 5, pp. 1795–1803, 2012.
- [17] C. A. Williams, "The leaf flavonoids of the Orchidaceae," *Phytochemistry*, vol. 18, no. 5, pp. 803–813, 1979.
- [18] H. J. Zhang, J. J. Zhou, and X. S. Li, "Study advance of gastrodia elata b1," *Amino Acids and Biotic Resources*, vol. 25, no. 1, pp. 17–20, 2003.
- [19] P. L. Majumder, S. Guha, and S. Sen, "Bibenzyl derivatives from the orchid *Dendrobium amoenum*," *Phytochemistry*, vol. 52, no. 7, pp. 1365–1369, 1999.
- [20] Y. Li, C.-L. Wang, Y.-J. Wang et al., "Three new bibenzyl derivatives from *Dendrobium candidum*," *Chemical and Pharmaceutical Bulletin*, vol. 57, no. 2, pp. 218–219, 2009.
- [21] Y. Li, C. L. Wang, S. X. Guo, J. S. Yang, and P. G. Xiao, "Two new compounds from *Dendrobium candidum*," *Chemical and Pharmaceutical Bulletin*, vol. 56, pp. 1477–1499, 2008.
- [22] Y. Chen, J. Li, L. Wang, and Y. Liu, "Aromatic compounds from *Dendrobium aphyllum*," *Biochemical Systematics and Ecology*, vol. 36, no. 5–6, pp. 458–460, 2008.
- [23] Y. Chen, Y. Liu, J. Jiang, Y. Zhang, and B. Yin, "Dendronone, a new phenanthrenequinone from *Dendrobium cariniferum*," *Food Chemistry*, vol. 111, no. 1, pp. 11–12, 2008.
- [24] Y. Chen, Y. Li, C. Qing, Y. Zhang, L. Wang, and Y. Liu, "1,4,5-Trihydroxy-7-methoxy-9H-fluoren-9-one, a new cytotoxic compound from *Dendrobium chrysotoxum*," *Food Chemistry*, vol. 108, no. 3, pp. 973–976, 2008.
- [25] H. Yang, G.-X. Chou, Z.-T. Wang, Y.-W. Guo, Z.-B. Hua, and L.-S. Xu, "Two new compounds from *Dendrobium chrysotoxum*," *Helvetica Chimica Acta*, vol. 87, no. 2, pp. 394–399, 2004.
- [26] C. Fan, W. Wang, Y. Wang, G. Qin, and W. Zhao, "Chemical constituents from *Dendrobium densiflorum*," *Phytochemistry*, vol. 57, no. 8, pp. 1255–1258, 2001.
- [27] J.-T. Li, B.-L. Yin, Y. Liu, L.-Q. Wang, and Y.-G. Chen, "Monoaromatic constituents of *Dendrobium longicornu*," *Chemistry of Natural Compounds*, vol. 45, no. 2, pp. 234–236, 2009.
- [28] X. Zhang, H.-W. Liu, H. Gao et al., "Nine new sesquiterpenes from *Dendrobium nobile*," *Helvetica Chimica Acta*, vol. 90, no. 12, pp. 2386–2394, 2007.
- [29] Q.-F. Liu, W.-L. Chen, J. Tang, and W.-M. Zhao, "Novel bis(bibenzyl) and (propylphenyl)bibenzyl derivatives from *Dendrobium nobile*," *Helvetica Chimica Acta*, vol. 90, no. 9, pp. 1745–1750, 2007.

- [30] G.-N. Zhang, L.-Y. Zhong, S. W. A. Bligh et al., "Bi-bicyclic and bi-tricyclic compounds from *Dendrobium thyrsiflorum*," *Phytochemistry*, vol. 66, no. 10, pp. 1113–1120, 2005.
- [31] Y. Liu, J.-H. Jiang, B.-L. Yin, and Y.-G. Chen, "Chemical constituents of *Dendrobium cariniferum*," *Chemistry of Natural Compounds*, vol. 45, no. 2, pp. 237–238, 2009.
- [32] S.-F. Lo, V. Mulabagal, C.-L. Chen, C.-L. Kuo, and H.-S. Tsay, "Bioguided fractionation and isolation of free radical scavenging components from *in vitro* propagated chinese medicinal plants *Dendrobium tosaense* Makino and *Dendrobium moniliforme* SW," *Journal of Agricultural and Food Chemistry*, vol. 52, no. 23, pp. 6916–6919, 2004.
- [33] P. L. Majumder and R. C. Sen, "Moscaticin, a bibenzyl derivative from the orchid *Dendrobium moscatum*," *Phytochemistry*, vol. 26, no. 7, pp. 2121–2124, 1987.
- [34] P. L. Majumder and S. Chatterjee, "Crepidatin, a bibenzyl derivative from the orchid *Dendrobium crepidatum*," *Phytochemistry*, vol. 28, no. 7, pp. 1986–1988, 1989.
- [35] P. L. Majumder and S. Pal (Née Ray), "Rotundatin, a new 9,10-dihydrophenanthrene derivative from *Dendrobium rotundatum*," *Phytochemistry*, vol. 31, no. 9, pp. 3225–3228, 1992.
- [36] C. Honda and M. Yamaki, "Phenanthrenes from *Dendrobium plicatile*," *Phytochemistry*, vol. 53, no. 8, pp. 987–990, 2000.
- [37] K. Krohn, U. Looock, K. Paavilainen, B. M. Hausen, H. W. Schmalle, and H. Kiesele, "Synthesis and electrochemistry of annoquinone-A, cypripedin methyl ether, denbinobin and related 1,4-phenanthrenequinones," *Arkivoc*, vol. 2001, no. 1, pp. 88–130, 2001.
- [38] X. Zhang, J.-K. Xu, J. Wang et al., "Bioactive bibenzyl derivatives and fluorenones from *Dendrobium nobile*," *Journal of Natural Products*, vol. 70, no. 1, pp. 24–28, 2007.
- [39] Y. Li, C.-L. Wang, S.-X. Guo, J.-S. Yang, and P.-G. Xiao, "Two new compounds from *Dendrobium candidum*," *Chemical and Pharmaceutical Bulletin*, vol. 56, no. 10, pp. 1477–1479, 2008.
- [40] A. X. Luo, X. J. He, S. D. Zhou, Y. J. Fan, T. He, and Z. Chun, "In vitro antioxidant activities of a water-soluble polysaccharide derived from *Dendrobium nobile* Lindl. extracts," *International Journal of Biological Macromolecules*, vol. 45, no. 4, pp. 359–363, 2009.
- [41] A. Luo and Y. Fan, "In vitro antioxidant of a water-soluble polysaccharide from *Dendrobium fimbriatum* Hook. var. *oculatum* Hook," *International Journal of Molecular Sciences*, vol. 12, no. 6, pp. 4068–4079, 2011.
- [42] R. M. Chen, T. G. Chen, T. L. Chen et al., "Anti-inflammatory and antioxidative effects of propofol on lipopolysaccharide-activated macrophages," *Annals of the New York Academy of Sciences*, vol. 1042, pp. 262–271, 2005.
- [43] L. Xiao, T. B. Ng, Y.-B. Feng et al., "*Dendrobium candidum* extract increases the expression of aquaporin-5 in labial glands from patients with Sjögren's syndrome," *Phytomedicine*, vol. 18, no. 2-3, pp. 194–198, 2011.
- [44] J.-X. Song, P.-C. Shaw, C.-W. Sze et al., "Chrysotoxine, a novel bibenzyl compound, inhibits 6-hydroxydopamine induced apoptosis in SH-SY5Y cells via mitochondria protection and NF- κ B modulation," *Neurochemistry International*, vol. 57, no. 6, pp. 676–689, 2010.
- [45] Q. Ye, G. Qin, and W. Zhao, "Immunomodulatory sesquiterpene glycosides from *Dendrobium nobile*," *Phytochemistry*, vol. 61, no. 8, pp. 885–890, 2002.
- [46] C. Zhao, Q. Liu, F. Halaweish, B. Shao, Y. Ye, and W. Zhao, "Copacamphane, picrotoxane, and alloaromadendrane sesquiterpene glycosides and phenolic glycosides from *Dendrobium moniliforme*," *Journal of Natural Products*, vol. 66, no. 8, pp. 1140–1143, 2003.
- [47] X. Q. Zha, J. P. Luo, S. Z. Luo, and S. T. Jiang, "Structure identification of a new immunostimulating polysaccharide from the stems of *Dendrobium huoshanense*," *Carbohydrate Polymers*, vol. 69, no. 1, pp. 86–93, 2007.
- [48] J.-H. Wang, J.-P. Luo, X.-Q. Zha, and B.-J. Feng, "Comparison of antitumor activities of different polysaccharide fractions from the stems of *Dendrobium nobile* Lindl," *Carbohydrate Polymers*, vol. 79, no. 1, pp. 114–118, 2010.
- [49] Y. Li, C.-L. Wang, Y.-J. Wang et al., "Four new bibenzyl derivatives from *Dendrobium candidum*," *Chemical and Pharmaceutical Bulletin*, vol. 57, no. 9, pp. 997–999, 2009.
- [50] J. I. Song, Y. J. Kang, H. Y. Yong, Y. C. Kim, and A. Moon, "Denbinobin, a phenanthrene from *Dendrobium nobile*, inhibits invasion and induces apoptosis in SNU-484 human gastric cancer cells," *Oncology Reports*, vol. 27, no. 3, pp. 813–818, 2012.
- [51] J. I. Song, Y. J. Kang, H.-Y. Yong, Y. C. Kim, and A. Moon, "Denbinobin, a phenanthrene from *Dendrobium nobile*, inhibits invasion and induces apoptosis in SNU-484 human gastric cancer cells," *Oncology Reports*, vol. 27, no. 3, pp. 813–818, 2012.
- [52] J.-X. Song, S. C.-W. Sze, T.-B. Ng et al., "Anti-Parkinsonian drug discovery from herbal medicines: what have we got from neurotoxic models?" *Journal of Ethnopharmacology*, vol. 139, no. 3, pp. 698–711, 2012.
- [53] A. Luo, Z. Ge, Y. Fan, Z. Chun, and X. Jin He, "In vitro and in vivo antioxidant activity of a water-soluble polysaccharide from *Dendrobium denneanum*," *Molecules*, vol. 16, no. 2, pp. 1579–1592, 2011.
- [54] Y. Fan, X. He, S. Zhou, A. Luo, T. He, and Z. Chun, "Composition analysis and antioxidant activity of polysaccharide from *Dendrobium denneanum*," *International Journal of Biological Macromolecules*, vol. 45, no. 2, pp. 169–173, 2009.
- [55] X. Lin, P.-C. Shaw, S. C.-W. Sze, Y. Tong, and Y. B. Zhang, "*Dendrobium* officinale polysaccharides ameliorate the abnormality of aquaporin 5, pro-inflammatory cytokines and inhibit apoptosis in the experimental Sjögren's syndrome mice," *International Immunopharmacology*, vol. 11, no. 12, pp. 2025–2032, 2011.
- [56] T.-H. Lin, S.-J. Chang, C.-C. Chen, J.-P. Wang, and L.-T. Tsao, "Two phenanthraquinones from *Dendrobium moniliforme*," *Journal of Natural Products*, vol. 64, no. 8, pp. 1084–1086, 2001.
- [57] D. T. W. Lau, M. Poon, H. Leung, and K. Ko, "Immunopotentiating activity of *Dendrobium* species in mouse splenocytes," *Chinese Medicine*, vol. 2, no. 3, pp. 103–108, 2011.
- [58] J. S. Hwang, S. A. Lee, S. S. Hong et al., "Phenanthrenes from *Dendrobium nobile* and their inhibition of the LPS-induced production of nitric oxide in macrophage RAW 264.7 cells," *Bioorganic and Medicinal Chemistry Letters*, vol. 20, no. 12, pp. 3785–3787, 2010.
- [59] L. C. Yang, T. J. Lu, C. C. Hsieh, and W. C. Lin, "Characterization and immunomodulatory activity of polysaccharides derived from *Dendrobium tosaense*," *Carbohydrate Polymers*, vol. 111, pp. 856–863, 2014.
- [60] L.-H. Pan, X.-F. Li, M.-N. Wang et al., "Comparison of hypoglycemic and antioxidative effects of polysaccharides from four different *Dendrobium* species," *International Journal of Biological Macromolecules*, vol. 64, pp. 420–427, 2014.
- [61] Y. Chen, Y. Li, C. Qing, Y. Zhang, L. Wang, and Y. Liu, "1,4,5-Trihydroxy-7-methoxy-9H-fluoren-9-one, a new cytotoxic compound from *Dendrobium chrysotoxum*," *Food Chemistry*, vol. 108, no. 3, pp. 973–976, 2008.

- [62] S. Phetsirikoon, S. Ketsa, and W. G. van Doorn, "Chilling injury in *Dendrobium inflorescences* is alleviated by 1-MCP treatment," *Postharvest Biology and Technology*, vol. 67, pp. 144–153, 2012.
- [63] B. Li, J. Wei, X. Wei et al., "Effect of sound wave stress on antioxidant enzyme activities and lipid peroxidation of *Dendrobium candidum*," *Colloids and Surfaces B: Biointerfaces*, vol. 63, no. 2, pp. 269–275, 2008.
- [64] X. Tian and Y. Lei, "Nitric oxide treatment alleviates drought stress in wheat seedlings," *Biologia Plantarum*, vol. 50, no. 4, pp. 775–778, 2006.
- [65] H. Fan, T. Li, L. Guan et al., "Effects of exogenous nitric oxide on antioxidation and DNA methylation of *Dendrobium huoshanense* grown under drought stress," *Plant Cell, Tissue and Organ Culture*, vol. 109, no. 2, pp. 307–314, 2012.
- [66] S. F. Lo, S. M. Nalawade, V. Mulabagal et al., "In vitro propagation by asymbiotic seed germination and 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging activity studies of tissue culture raised plants of three medicinally important species of *Dendrobium*," *Biological and Pharmaceutical Bulletin*, vol. 27, no. 5, pp. 731–735, 2004.
- [67] S.-F. Lo, V. Mulabagal, C.-L. Chen, C.-L. Kuo, and H.-S. Tsay, "Bioguided fractionation and isolation of free radical scavenging components from *in vitro* propagated chinese medicinal plants *Dendrobium tosaense* Makino and *Dendrobium moniliforme* SW," *Journal of Agricultural and Food Chemistry*, vol. 52, no. 23, pp. 6916–6919, 2004.
- [68] J.-H. Wang, X.-Q. Zha, J.-P. Luo, and X.-F. Yang, "An acetylated galactomannoglucan from the stems of *Dendrobium nobile* Lindl," *Carbohydrate Research*, vol. 345, no. 8, pp. 1023–1027, 2010.
- [69] W.-G. Zhang, R. Zhao, J. Ren et al., "Synthesis and antiproliferative in-vitro activity of two natural dihydrostilbenes and their analogues," *Archiv der Pharmazie*, vol. 340, no. 5, pp. 244–250, 2007.
- [70] C. C. Tian, X. Q. Zha, L. H. Pan, and J. P. Luo, "Structural characterization and antioxidant activity of a low-molecular polysaccharide from *Dendrobium huoshanense*," *Fitoterapia*, vol. 91, pp. 247–255, 2013.
- [71] M. Ito, K. Matsuzaki, J. Wang et al., "New phenanthrenes and stilbenes from *Dendrobium loddigesii*," *Chemical and Pharmaceutical Bulletin*, vol. 58, no. 5, pp. 628–633, 2010.
- [72] L. Yang, L. H. Qin, S. W. A. Bligh et al., "A new phenanthrene with a spiro lactone from *Dendrobium chrysanthum* and its anti-inflammatory activities," *Bioorganic and Medicinal Chemistry*, vol. 14, no. 10, pp. 3496–3501, 2006.
- [73] R. I. Fox, "Sjögren's syndrome," *The Lancet*, vol. 366, no. 9482, pp. 321–331, 2005.
- [74] Y. P. Ding, L. S. Wu, and L. W. Yu, "Optimized harvesting period for *Dendrobium candidum*," *China journal Chinese Materia Medica*, vol. 23, pp. 458–460, 1998.
- [75] N. Roescher, P. P. Tak, and G. G. Illei, "Cytokines in Sjögren's syndrome: potential therapeutic targets," *Annals of the Rheumatic Diseases*, vol. 69, no. 6, pp. 945–948, 2010.
- [76] L. Xiang, C. W. Stephen Sze, T. B. Ng et al., "Polysaccharides of *Dendrobium officinale* inhibit TNF- α -induced apoptosis in A-253 cell line," *Inflammation Research*, vol. 62, no. 3, pp. 313–324, 2013.
- [77] J.-X. Song, P.-C. Shaw, N.-S. Wong et al., "Chrysotoxine, a novel bibenzyl compound selectively antagonizes MPP⁺, but not rotenone, neurotoxicity in dopaminergic SH-SY5Y cells," *Neuroscience Letters*, vol. 521, no. 1, pp. 76–81, 2012.
- [78] J. Fan, L. Guan, Z. Kou, F. Feng, Y. B. Zhang, and W. Liu, "Determination of chrysotoxine in rat plasma by liquid chromatography-tandem mass spectrometry and its application to a rat pharmacokinetic study," *Journal of Chromatography B*, vol. 967, pp. 57–62, 2014.
- [79] M. Liu, J. Li, F. Kong, J. Lin, and Y. Gao, "Induction of immunomodulating cytokines by a new polysaccharide-peptide complex from culture mycelia of *Lentinus edodes*," *Immunopharmacology*, vol. 40, no. 3, pp. 187–198, 1998.
- [80] W. Zhao, Q. Ye, X. Tan et al., "Three new sesquiterpene glycosides from *Dendrobium nobile* with immunomodulatory activity," *Journal of Natural Products*, vol. 64, no. 9, pp. 1196–1200, 2001.
- [81] Y. S.-Y. Hsieh, C. Chien, S. K.-S. Liao et al., "Structure and bioactivity of the polysaccharides in medicinal plant *Dendrobium huoshanense*," *Bioorganic and Medicinal Chemistry*, vol. 16, no. 11, pp. 6054–6068, 2008.
- [82] F. Li, S. H. Cui, X. Q. Zha et al., "Structure and bioactivity of a polysaccharide extracted from protocorm-like bodies of *Dendrobium huoshanense*," *International Journal of Biological Macromolecules*, vol. 72, pp. 664–672, 2015.
- [83] J. Lin, Y.-J. Chang, W.-B. Yang, A. L. Yu, and C.-H. Wong, "The multifaceted effects of polysaccharides isolated from *Dendrobium huoshanense* on immune functions with the induction of interleukin-1 receptor antagonist (IL-1ra) in monocytes," *PLoS ONE*, vol. 9, no. 4, Article ID e94040, 2014.
- [84] H. L. You, J. D. Park, N. I. Baek, S. I. Kim, and B. Z. Ahn, "In vitro and in vivo antitumoral phenanthrenes from the aerial parts of *Dendrobium nobile*," *Planta Medica*, vol. 61, no. 2, pp. 178–180, 1995.
- [85] R. Prasad and B. Koch, "Antitumor activity of ethanolic extract of *Dendrobium formosum* in T-cell lymphoma: an in vitro and in vivo study," *BioMed Research International*, vol. 2014, Article ID 753451, 11 pages, 2014.
- [86] S. Charoenrungruang, P. Chanvorachote, B. Sritularak, and V. Pongrakhananon, "Gigantol, a bibenzyl from *Dendrobium draconis*, inhibits the migratory behavior of non-small cell lung cancer cells," *Journal of Natural Products*, vol. 77, no. 6, pp. 1359–1366, 2014.
- [87] P. Benjaphorn, T. Lalita, N. Ratchaya, and P. Cholakan, "Efficacy of crude extract of antifungal compounds produced from *Bacillus subtilis* on prevention of anthracnose disease in *Dendrobium orchid*," *EnvironmentAsia*, vol. 5, no. 1, pp. 32–38, 2012.
- [88] Y. Tao, F. Zeng, H. Ho et al., "Pythium vexans causing stem rot of *Dendrobium* in Yunnan Province, China," *Journal of Phytopathology*, vol. 159, no. 4, pp. 255–259, 2011.
- [89] M. E. Barrow Lucero Jr., I. Reyes-Vera, and K. M. Havstad, "Do symbiotic microbes have a role in plant evolution, performance and response to stress?" *Communicative & Integrative Biology*, vol. 1, pp. 1–5, 2008.
- [90] M. A. Baute, G. Deffieux, R. Baute, and A. Neveu, "New antibiotics from the fungus *Epicoccum nigrum*. I. Fermentation, isolation and antibacterial properties," *The Journal of Antibiotics*, vol. 31, no. 11, pp. 1099–1101, 1978.
- [91] Y. Zhang, S. Liu, Y. Che, and X. Liu, "Epicoccins A-D, epipolythiodioxopiperazines from a *Cordyceps*-colonizing isolate of *Epicoccum nigrum*," *Journal of Natural Products*, vol. 70, no. 9, pp. 1522–1525, 2007.
- [92] W. Zhang, K. Krohn, H. Egold, S. Draeger, and B. Schulz, "Diversity of antimicrobial pyrenophorol derivatives from an endophytic fungus, *Phoma* sp.," *European Journal of Organic Chemistry*, no. 25, pp. 4320–4328, 2008.

- [93] N. Sattayasai, R. Sudmoon, S. Nuchadomrong et al., "Dendrobium findleyanum agglutinin: production, localization, antifungal activity and gene characterization," *Plant Cell Reports*, vol. 28, no. 8, pp. 1243–1252, 2009.
- [94] M. Miyazawa, H. Shimamura, S.-I. Nakamura, W. Sugiura, H. Kosaka, and H. Kameoka, "Moscatilin from *Dendrobium nobile*, a naturally occurring bibenzyl compound with potential antimutagenic activity," *Journal of Agricultural and Food Chemistry*, vol. 47, no. 5, pp. 2163–2167, 1999.
- [95] C. Y. Gong, Z. Y. Yu, B. Lu et al., "Ethanol extract of *Dendrobium chrysotoxum* Lindl ameliorates diabetic retinopathy and its mechanism," *Vascular Pharmacology*, vol. 62, no. 3, pp. 134–142, 2014.
- [96] C.-C. Chen, Y.-L. Huang, and C.-M. Teng, "Antiplatelet aggregation principles from *Ephemerantha lonchophylla*," *Planta Medica*, vol. 66, no. 4, pp. 372–374, 2000.
- [97] Y. Okada, N. Miyauchi, K. Suzuki et al., "Search for naturally occurring substances to prevent the complications of diabetes. II. Inhibitory effect of coumarin and flavonoid derivatives on bovine lens aldose reductase and rabbit platelet aggregation," *Chemical and Pharmaceutical Bulletin*, vol. 43, no. 8, pp. 1385–1387, 1995.
- [98] J.-M. Hu, J.-J. Chen, H. Yu, Y.-X. Zhao, and J. Zhou, "Two novel bibenzyls from *Dendrobium trigonopus*," *Journal of Asian Natural Products Research*, vol. 10, no. 7-8, pp. 653–657, 2008.