Corrigendum

Corrigendum to “Caffeic Acid Phenethyl Ester Inhibits Oral Cancer Cell Metastasis by Regulating Matrix Metalloproteinase-2 and the Mitogen-Activated Protein Kinase Pathway”

Chih-Yu Peng,^{1,2} Hui-Wen Yang,^{1,2} Yin-Hung Chu,³ Yu-Chao Chang,^{1,2} Ming-Ju Hsieh,⁴ Ming-Yung Chou,^{1,2} Kun-Tu Yeh,⁵ Yueh-Min Lin,⁵ Shun-Fa Yang,^{3,6} and Chiao-Wen Lin^{2,7}

¹School of Dentistry, Chung Shan Medical University, Taichung 40201, Taiwan

²Department of Dentistry, Chung Shan Medical University Hospital, Taichung 40201, Taiwan

³Institute of Medicine, Chung Shan Medical University, Taichung 402, Taiwan

⁴School of Medical Laboratory and Biotechnology, Chung Shan Medical University, Taichung 40201, Taiwan

⁵Department of Pathology, Changhua Christian Hospital, Changhua 50006, Taiwan

⁶Department of Medical Research, Chung Shan Medical University Hospital, Taichung 40201, Taiwan

⁷Institute of Oral Sciences, Chung Shan Medical University, Taichung 40201, Taiwan

Correspondence should be addressed to Shun-Fa Yang; ysf@csmu.edu.tw

Received 11 November 2015; Accepted 6 January 2016

Copyright © 2016 Chih-Yu Peng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We have noticed an inadvertent error in our paper “Caffeic Acid Phenethyl Ester Inhibits Oral Cancer Cell Metastasis by Regulating Matrix Metalloproteinase-2 and the Mitogen-Activated Protein Kinase Pathway” [1].

We found a misplaced figure in Figure 3(a). We have attached a corrected version of Figure 3(a). The correction does not affect the findings or conclusion of the study.

References

- [1] C.-Y. Peng, H.-W. Yang, Y.-H. Chu et al., “Caffeic Acid phenethyl ester inhibits oral cancer cell metastasis by regulating matrix metalloproteinase-2 and the mitogen-activated protein kinase pathway,” *Evidence-Based Complementary and Alternative Medicine*, vol. 2012, Article ID 732578, 10 pages, 2012.

FIGURE 3: Effect of CAPE on in vitro wound closure in oral cancer cells. (a) SCC-9 cells were wounded and then treated with vehicle (DMSO) or CAPE (0, 5, 10, 20, and 40 μM) for 0 h, 24 h, 48 h, and 72 h in 0.5% FBS-containing medium. At 0 h, 24 h, 48, and 72 h, phase-contrast pictures of the wounds at three different locations were taken. (b) Cells migrating into the wound area were counted using the dashed line as time zero. A quantitative assessment of the mean number of cells in the denuded zone is the mean $\pm$ SD ($n = 3$).