Research Article

¹H-NMR Based Serum Metabolomics Study to Investigate Hepatoprotective Effect of Qin-Jiao on Carbon Tetrachloride-Induced Acute Hepatotoxicity in Rats

Zeyun Li, Ying Li, Lingpan Lu, Zhiheng Yang, Wenhua Xue, Xin Tian, and Xiaojian Zhang

Department of Pharmacy, The First Affiliated Hospital of Zhengzhou University, Zhengzhou 450052, China

Correspondence should be addressed to Xin Tian; tianx@zzu.edu.cn and Xiaojian Zhang; zhangxj9758@126.com

Received 4 January 2017; Revised 29 May 2017; Accepted 7 June 2017; Published 1 November 2017

Academic Editor: Yohandra R. Torres

Copyright © 2017 Zeyun Li et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Gentiana macrophylla Radix, commonly known as Qin-Jiao (QJ), was recorded alone to treat jaundice in Compendium of Materia Medica and has been frequently prescribed for treatment of liver disease in China. However, the underlying mechanism remains unknown. In the present work, QJ of 1,2 g/kg or silybin of 40 mg/kg (positive control) was orally given to rats for 7 days to verify the protective effect on acute liver damage induced by tetrachloride (CCl_4). Together with serum biochemistry and histopathological examination, ¹H-NMR based metabolomics work was carried out to investigate the efficacy. It turned out that QJ of 2 g/kg exerted comparable protective effect with positive control and partially recovered disturbed metabolism by CCl_4 . Multivariate analysis was conducted and metabolites altered significantly among groups were assigned and discussed, including betaine, glucose, lactate, creatine, and LDL/VLDL. Metabolic regulations involved in QJ or silybin treatment were as follows: tricarboxylic acid (TCA) cycle, synthesis of LDL/VLDL, and gluconeogenesis were enhanced, while betaine metabolism, glycolysis, creatine metabolism, synthesis of ketone bodies, amino acids metabolism, and β -oxidation of fatty acids were suppressed. For the first time hepatoprotective effect of QJ on acute liver damage was revealed by ¹H-NMR based metabolomics, prompting understanding of the underlying mechanism.

1. Introduction

Liver diseases constitute a global concern and considerable interest has been rising in research of traditional Chinese medicine (TCM) with liver protective efficacy [1, 2]. *Gentiana Macrophyllae* Radix, dried root of *Gentiana dahurica* Fisch. (Fam. Gentianeae) commonly known as Qin-Jiao (QJ), has been an important herb medicine since ancient time in China and was frequently prescribed for treatment of rheumatism, arthralgia, and jaundice [3, 4]. Modern researches have revealed that QJ processed the efficiency of anti-inflammation [5], sedation [6], liver protection [7], and so on. Clinic use of QJ for liver disease has a long history, recorded to be used alone to treat jaundice in Compendium of Materia Medica. However, hepatoprotective efficacy of QJ was not well investigated and the mechanism underlying remains unknown.

Metabolomics, profiling various types of biofluids and reflecting the function and metabolic changes of complete

organisms in an holistic view, has attracted a great deal of interest in toxicological [8] or pharmaceutical [9] research. Its holistic view coincides with the integrative thinking of TCM and exhibits great potential in revealing activity, toxicity, and underlying mechanism of TCM [10, 11]. With the advantages of unbiased analysis, simple sample handling, and high reproducibility [12], NMR based metabolomics emerges as one of the most prominent and convenient platforms employed in TCM study [13].

Carbon tetrachloride (CCl_4) has been extensively used in animal models to explore hepatic injury and to investigate hypoprotective effect of candidates [14]. Many herb medicines or natural products have been proven to exhibit hypoprotective effect on CCl_4 induced acute liver injury models [15, 16]. In the present work, ¹H-NMR based metabolomics combined with conventional serum chemistry analysis as well as tissue section examination were carried out to investigate the protective effect of QJ (1,2 g/kg) on

acute hepatic damaged rats induced by CCl_4 . Silybin, a well-known natural product with hepatoprotective effect, was given at dosage of 40 mg/kg as positive control. Multivariate data analyses, such as principal components analysis (PCA) and orthogonal partial least-squares discriminant analysis (OPLS-DA), were utilized to reveal pharmacodynamics and to find out related biomarkers responding to liver damage or QJ treatment, which may facilitate the understanding of the pathological changes in acute liver damage and the hepatoprotective mechanisms of QJ.

2. Experimental

2.1. Chemicals, Reagents, and Herbal Materials Procession. Analytical grade ethanol and carboxymethylcellulose (CMC-Na) were purchased from Sinopharm Chemical Reagent Co., Ltd. (Shanghai, China). D_2O (99.9% D) was bought from Cambridge Isotopes, Inc. (Andover, MA). Silybin (purity > 98%) was commercially obtained from Chengdu Biopurify Phytochemicals Ltd. (Chengdu, China). CCl_4 and coil oil were purchased from Fluka Chimica (Milan, Italy). Kits for serum alanine aminotransferase (ALT) and aspartate aminotransferase (AST) test were purchased from Nanjing Jiancheng Biotech Inc. (Nanjing, China). QJ (*Gentiana dahurica* Fisch.) was purchased from Tong Ren Tang drug store in Zhengzhou (Zhengzhou, China) and identified by chief pharmacist Juan Wang (Department of Pharmacy, The First Affiliated Hospital of Zhengzhou University). A voucher specimen was deposited in the author's laboratory. The roots of QJ (2,000 g) were grinded into powder (<80 mesh) and were extracted with 8 volume of 70% ethanol under reflux (ca. 80°C) for 3 times, each time for 1 hour. After filtering, the filtrates were merged and concentrated to a volume equivalent to 5 g/mL of QJ in 0.3% CMC-Na solution and stored at 4°C.

2.2. Animal Experiment and Sample Collection. A total of 35 male Sprague-Dawley (SD) rats (220 ± 20 g, 7 weeks old) were obtained from Hunan Slac Jingda Laboratory Animal Co. Ltd (Changsha, China) and housed at a certified animal experimental laboratory, with a 12 h light/dark cycle and a constant temperature of 25 ± 1°C. Animals were allowed free access to food and water. After acclimatization for 1 week, the rats were randomly assigned to five groups ($n = 7$): a control group (Con), a model group (Mod), a positive control group (SYL), and 2 QJ treated groups (QJ1, QJ2). Con and Mod groups were subjected to oral gavage with 0.3% CMC-Na solution. SYL and QJ treated groups were orally given silybin of 40 mg/kg or QJ of 1,2 g/kg in 0.3% CMC-Na, respectively. Drugs or solvent were administered once a day for 7 consecutive days. 2 h after the last administration, Mod, SYL, and QJ treated groups were intraperitoneally injected with CCl_4 (2 mg/kg) in corn oil (1:1 v/v) to induce acute liver damage, while Con group was let alone. 24 h later, all rats were sacrificed to collect liver tissues and serum samples for histopathological, biochemical assays or metabolomics analysis. Liver tissues were fixed in 10% formalin solution and serum samples were stored at -80°C before analysis.

2.3. Histopathology and Serum Biochemistry Assessments. The formalin-fixed liver biopsies were embedded in paraffin wax, sectioned (3–4 μm), and stained with hematoxylin and eosin (H&E) for assessment by the Department of Pathology, The First Affiliated Hospital of Zhengzhou University. Generally, two or five slices were examined for each sample. The serum levels of ALT and AST were tested according to the manufacturer's instructions. Each sample was assayed in duplicate. Statistical analysis was performed using one-way analysis of variance (ANOVA) followed by least-significant difference (LSD) post hoc test (SPSS, Chicago, IL, USA). A probability of $P < 0.05$ was considered to be a statistically significant difference between two groups.

2.4. Sample Preparation and ^1H -NMR Spectroscopy Acquisition. Serum samples were thawed at room temperature, 200 μL of which was mixed with 200 μL D_2O and 100 μL distilled water. The resulting solution was well mixed and centrifuged at 4°C at 2,000g (4,300 rpm) for 5 min to remove any sediment. Then 450 μL of final solution was transferred in a 5-mm NMR tube (Norell, Landisville, NJ, USA) for spectra acquisition.

^1H -NMR spectra of serum were collected at 298 K on a Bruker 600-MHz AVANCE III NMR spectrometer (Bruker, Germany), equipped with a 5.0-mm BBO probe, operating at 600.13 MHz for ^1H . The NMR spectrum was recorded using the water-presaturated standard one-dimensional Carr-Purcell-Meiboom-Gill (CPMG) pulse sequence (recycle delay-90°-(τ -180°- τ) $_n$ -acquisition) to eliminate interference by macromolecules, with a total spin-spin relaxation delay (2τ) of 350 ms. A total of 128 free induction decays (FID) were collected into 64 k data points over a spectral width of 10 kHz for each spectrum, with a relaxation delay (D1) of 2 s and power level for presaturation (PLW9) of 33 dB. A line-broadening factor of 0.3 Hz was applied to FID before Fourier transformation. Additional 2-dimensional NMR experiments were performed for the purpose of confirming chemical shift assignments, including homo-nuclear total correlation spectroscopy (2D ^1H - ^1H TOCSY) and hetero-nuclear single quantum coherence spectroscopy (2D ^1H - ^{13}C HSQC) acquired by Bruker's standard pulse sequences mlevphpr and hsqcphpr.

2.5. Data Processing and Statistical Analysis. All the NMR spectra were phased and baseline-corrected manually using TOPSPIN 3.0 (Bruker, Germany). The spectra were referenced internally to the chemical shift of creatinine at 3.03 ppm. Each ^1H -NMR spectrum over the ranged 0.5–9.5 ppm was reduced to 225 regions of equal width (0.04 ppm) and the signal intensity in each region integrated using AMIX (Bruker, Germany). The region of 4.84–5.28 ppm was removed prior to any statistical analysis in order to eliminate any spurious effects of water suppression. Following removal of water regions, data was normalized in AMIX by dividing each integrated segment by the total area of the spectrum to reduce any significant concentration difference. Output data in ASCII data format was imported to Microsoft Excel (Microsoft Office 2007), pareto-scaled, and

FIGURE 1: Photomicrographs of the rat liver sections (hematoxylin and eosin staining, $\times 200$). *Note.* Con, control group; Mod, acute liver damage model; SYL, silybin treated group (40 mg/kg); QJ1-2, QJ treated groups (1,2 g/kg); blue arrow marked vacuolation of live cells, and yellow arrow marked inflammatory cell infiltration.

then imported into SIMCA-P version 13.0.3 (Umetrics, Umeå, Sweden) for multivariate statistical analysis [17].

PCA, a classical unsupervised multivariate pattern recognition method, was employed to examine the intrinsic variation within a group and to assess the clustering behavior among groups. Subsequently, OPLS-DA, a supervised pattern recognition method, was further performed to maximize the variation between each two groups and to determine the variables that contribute to this variation. The quality of models was validated by determining R^2 (goodness of fit parameter) and Q^2 (goodness of prediction parameter) values. The results were visualized in the form of score plots, where each point represents an individual sample (to show the group clusters), or S-plots, where each coordinate represents one $^1\text{H-NMR}$ spectral region (to identify the variables contributing to the classification). The corresponding variables with variable importance in the projection value (VIP) >1.0 were chosen as major metabolites, whose intensities were compared to indicate the metabolic alteration between groups. Statistical analysis was also performed using ANOVA followed by LSD post hoc test (SPSS, Chicago, IL, USA). A probability of $P < 0.05$ was considered to be a statistically significant difference between two groups.

2.6. Identifications of Major Metabolites. According to the $^1\text{H-NMR}$ spectra of serum samples, major metabolites were assigned by comparison with databases of HMDB (<http://www.hmdb.ca>; <http://www.bml-nmr.org>) and the library of the Chenomx software (Vers. 7.6, Chenomx, Edmonton, Canada). Biochemical reactions involving the identified

metabolites were found through the Kyoto Encyclopedia of Genes and Genomes (KEGG) and the Human Metabolome Database (HMDB).

3. Results and Discussion

3.1. Histopathology. As shown in Figure 1, the histological tissue sections of the Con group showed normal liver histological morphology after H&E staining, while the CCl_4 induced Mod group exhibit abnormal histopathological changes, with significant vacuolation and degenerative changes in liver, ballooning, and fatty changes of hepatocytes and inflammatory cell infiltration. On the other hand, silybin treated group presents normalization of fatty changes, cellular infiltration, and necrosis, confirming the liver protective effect. The QJ1 (1 g/kg) group showed eliminated vacuolation and inflammatory infiltration, suggesting mild improvement in contrast to Mod group, while the tissue sections of QJ2 group (2 mg/kg) showed significant symptomatic relief compared with Mod group, with vacuolation induced by CCl_4 remarkably reversed, showing a marked protective effect against CCl_4 -induced liver toxicity. In conclusion, moderate hepatic protection against CCl_4 was achieved by QJ treatment.

3.2. Serum Biochemistry Assessments. Serum aminotransferases, including ALT and AST levels, were used as biochemical indicators of liver damage. As shown in Table 1, serum ALT and AST activities significantly increased in Mod group compared to the Con group ($P < 0.01$), indicating occurrence of liver damage. Not surprisingly, silybin treatment significantly reduced serum ALT and AST by ca. 30%

TABLE 1: Serum biochemistry test results ($n = 7$).

	Con	Mod	SYL	QJ1	QJ2
AST	17.83 $\pm$ 3.82	515.68 $\pm$ 132.64**	170.11 $\pm$ 101.19**	299.66 $\pm$ 97.98*	169.44 $\pm$ 29.26**
ALT	19.11 $\pm$ 10.56	139.17 $\pm$ 58.34**	108.54 $\pm$ 20.33**	130.79 $\pm$ 28.32	121.88 $\pm$ 43.21*

Note. Con, control group; Mod, acute liver damage model; SYL, silybin treated group (40 mg/kg); QJ1-2, QJ treated groups (1,2 g/kg); * means $P < 0.05$, ** means $P < 0.01$.

FIGURE 2: Typical 600 MHz ^1H -NMR spectra of rat serum from Con (a), Mod (b), and QJ2 (c) groups. Note. Con, control group; Mod, acute liver damage model; SYL, silybin treated group (40 mg/kg); QJ2, QJ treated groups (2 g/kg). Metabolites: (1) LDL/VLDL, (2) isoleucine, (3) leucine, (4) valine, (5) 3-hydroxybutyrate, (6) lactate, (7) alanine, (8) lysine, (9) acetate, (10) acetone, (11) acetoacetate, (12) glutamate, (13) pyruvate, (14) succinate, (15) glutamine, (16) citrate, (17) N,N-dimethylglycine, (18) creatine, (19) choline, (20) O-phosphocholine, (21) betaine, (22) glycine, (23) glucose, (24) serine, (25) fumarate, (26) tyrosine, (27) τ -methylhistidine, (28) phenylalanine, and (29) format.

and 60%, in contrast to the Mod group. The reduction was consistent with previous report, explained by the antioxidant capacity of silybin [18]. In parallel, QJ treatment significantly reduced the serum activity level of AST or ALT by 20–60%, which was in line with previous report and can be further supported by similar report of gentiopicoside [19]. Although no significant difference in AST or ALT activities among QJ treated groups was noticed, it seemed that QJ2 group exhibit better reduction of AST or ALT than QJ1 group, inferring better liver protection of 2 g/kg, corresponding to ca. twice the clinical dosage of QJ recorded in Chinese Pharmacopoeia (2015).

3.3. ^1H -NMR Based Metabolomics Analysis. Representative serum ^1H -NMR spectra of Con, Mod, and QJ2 groups were shown in Figure 2, with major metabolites identified. The identified metabolites, chemical shifts, and related metabolic pathways were shown in Table 2. Detailed metabolomics information and differences among groups were revealed by PCA or OPLS-DA model in a more holistic way.

PCA score plot of samples from Con, Mod, SYL, QJ1, and QJ2 groups was shown in Figure 3(a), with R^2 0.846 and Q^2 0.660. As can be noticed, samples of Mod group clustered far away from those of Con group, indicating successfully established acute liver damage model and confirming histopathology and serum biochemistry result. On the other hand, QJ or SYL treated groups deviated from Mod group and located close to Con group, suggesting

that the metabolism disorder of Mod group was partially recovered. In addition, QJ2 group clustered closer to Con group compared with QJ1 group, inferring better protection effect of 2 g/kg. For SYL treatment, the metabolism restoration effect was consistent with previous report [20], and the metabolism change explained histopathology and serum biochemistry result. For QJ treatment, this was the first metabolic revealing of the liver protective effect and may provide further understanding of the protective efficacy. The loading plot (Figure 3(b)) revealed the correlations between groups and variables. Accordingly, glucose and LDL/VLDL were considered to be in high abundance in Con group, while betaine, creatine, and lactate were considered to be characteristics of Mod group.

Subsequently, to explore the metabolic alterations associated with acute liver damage, OPLS-DA model was carried out between Con and Mod groups. The model was well fitted with R^2Y 0.979 and Q^2 0.951 (permutations test result are shown in Figure S1 in Supplementary Material, available online at <https://doi.org/10.1155/2017/6091589>). The score plot (Figure 4(a)) shows good separation between the two groups, indicating significant metabolism abnormalities. Corresponding S-plot was shown in Figure 4(b), where coordinates in the lower-left quadrant were metabolites significantly increased in Con group compared with Mod group, while those in the upper-right quadrant represent the decreased ones. Meanwhile, VIP values of variables which denote the influence of metabolite on the classification, were

TABLE 2: Assignment results of the identified metabolites, chemical shifts, and related metabolic pathways.

Number	Metabolites	Moieties	$\delta^1\text{H}$ (ppm) and multiplicity	Related pathway
(1)	LDL/VLDL	CH_2	0.82(m), 0.86(m), 1.26(m)	Synthesis of LDL/VLDL
(2)	Isoleucine	$\gamma\text{CH}_3, \alpha\text{CH}_3$	0.93(t), 1.00(d)	Gluconeogenesis of glucogenic amino acid/proteolysis
(3)	Leucine	$\delta\text{CH}_3, \delta\text{CH}_3$	0.94(d), 0.96(d)	Ketogenesis of ketogenic amino acids/proteolysis
(4)	Valine	$\gamma\text{CH}_3, \gamma\text{CH}_3$	1.03(d); 0.97(d)	Gluconeogenesis of glucogenic amino acid/proteolysis
(5)	3-Hydroxybutyrate	γCH_3	1.19(d)	Synthesis and degradation of ketone bodies
(6)	Lactate	$\beta\text{CH}_3, \text{CH}$	1.33(d), 4.06(q)	Glycolysis and gluconeogenesis
(7)	Alanine	βCH_3	1.46(d)	Gluconeogenesis of glucogenic amino acid/proteolysis
(8)	Lysine	$\alpha\text{CH}_2, \gamma\text{CH}_2$	1.88(m), 1.72(m)	Ketogenesis of ketogenic amino acids
(9)	Acetate	COCH_3	1.90(d)	β -Oxidation of fatty acids
(10)	Acetone	$\text{CO}(\text{CH}_3)_2$	2.22(s)	Synthesis and degradation of ketone bodies
(11)	Acetoacetate	COCH_3	2.28(s)	Synthesis and degradation of ketone bodies
(12)	Glutamate	βCH_2	2.34(m)	Glutamate metabolism/nitrogen metabolism
(13)	Pyruvate	COCH_3	2.37(s)	Glycolysis/gluconeogenesis/TCA cycle
(14)	Succinate	$(\text{COCH}_2)_2$	2.40(s)	TCA cycle
(15)	Glutamine	βCH_2	2.40–2.48(m)	Glutamate metabolism/nitrogen metabolism
(16)	Citrate	$1/2\text{CH}_3, 1/2\text{CH}_2$	2.65–2.70(d); 2.20–2.55(d)	TCA cycle
(17)	N,N-Dimethylglycine	$\text{N}(\text{CH}_3)_2$	2.90(s)	Glycine, serine and threonine metabolism
(18)	Creatine	$\text{N-CH}_3, \text{N-CH}_2$	4.05(s); 3.04(s)	Creatine metabolism
(19)	Choline	$\text{N}(\text{CH}_3)_3$	3.18–3.19(s)	Choline metabolism
(20)	O-Phosphocholine	$\text{N}(\text{CH}_3)_3$	3.22(s)	Choline metabolism
(21)	Betaine	$\text{N}(\text{CH}_3)_3$	3.26(s)	Betaine metabolism
(22)	Glycine	CH_2	3.54(s),	Glycine, serine, and threonine metabolism
(23)	Glucose	CH	5.20(d), 4.66(d), 3.70–3.90(m), 3.30–3.50(m), 3.84(m)	Glycolysis and gluconeogenesis
(24)	Serine	CH_2	3.94(d)	Glycine and serine metabolism/ketogenesis of ketogenic amino acids
(25)	Fumarate	CH=CH	6.50(s)	TCA cycle
(26)	Tyrosine	ph-H	7.20(d), 6.86(d)	Tyrosine metabolism/ketogenesis of ketogenic amino acids
(27)	τ -Methylhistidine	CH-N-CH=	7.65(s), 7.02(s)	Histidine metabolism
(28)	Phenylalanine	ph-H	7.30(m)	Phenylalanine and tyrosine metabolism
(29)	Formate	HCOO	8.46(s)	Formate metabolism

FIGURE 3: PCA score plot (a), loading plot (b) of serum ^1H -NMR spectra obtained from Con, Mod, SYL, and QJ1-2 groups. Note. Con, control group; Mod, acute liver damage model; SYL, silybin treated group (40 mg/kg); QJ2, QJ treated group (2 g/kg).

calculated by SIMCA-P software. Variables with VIP value >1.0 between two groups were selected and discussed (Table S1). Likewise, OPLS-DA models were carried out between QJ2 and Mod or SYL and Mod groups to reveal the liver protective mechanism of QJ or SYL. The models were well fitted with R^2Y and Q^2 value 0.845, 0.572, or 0.918, 0.796 (permutation test is shown in Figure S1). The score plot (Figures 4(c) and 4(e)) shows good separation and confirms the liver protective effect. From the S-plot (Figures 4(d) and 4(f)), metabolites with VIP >1.0 between two groups were selected and listed (Table S1).

On the basis of aforementioned OPLS-DA analysis, statistical analysis of potential biomarkers among groups was performed and shown in Figure 5, further detailing alteration trends of metabolites. Alterations and related metabolic pathways were discussed, and it turned out that following metabolic pathways were involved in CCl_4 induced liver damage and liver protective efficacy: betaine metabolism, synthesis of LDL/VLDL, gluconeogenesis and glycolysis, tri-carboxylic acid (TCA) cycle, creatine metabolism, synthesis of ketone bodies, amino acids metabolism, and β -oxidation of fatty acids.

Betaine, with the chemical shift of 3.26 ppm and the highest VIP value among each OPLS-DA model, was considered as the most predominant biomarker associated with liver damage or restoration regulations. Betaine comes either from the diet or by the oxidation of choline, with major fate of being phosphorylated or as a donor of methyl-groups. Oxidation of choline to betaine was catalyzed mainly in the mitochondria of liver cells via a series of enzymes, including choline dehydrogenase (CHDH) [21]. CHDH was reported to be related to abnormal mitochondrial function and upregulated in the mitochondrial proteome of rats with fatty liver [22, 23]. The upregulation could be a compensatory response, since betaine is believed to have a hepatoprotective effect [24]. Meanwhile, elevated plasma betaine may promote

upregulation of multiple macrophage scavenger receptors or deteriorate liver functions, and thus lead to toxic responses of liver or kidney [25]. In addition, elevated betaine level was also revealed for the toxic responses of processed *Aconitum carmichaelii* Debx. [26]. Consistent with previous reports, in our study, serum betaine level was noticed to increase by 3.18-fold in Mod group, which may be derived from excessive choline released due to CCl_4 caused membranolysis or elevated levels of related oxidation enzymes. On the other hand, after SYL or QJ treatment, betaine levels were significantly decreased by a half compared with Mod group, which indicated that betaine metabolism disturbed by CCl_4 inducement was partially restored. The restoration was probably due to the fact that related oxidation enzymes were downregulated.

Another metabolism form of choline was to be phosphorylated, especially as phosphatidylcholine (PtdC), which is necessary for the packaging and export of triglycerides in very low density lipoprotein (VLDL) [27] and for the solubilization of bile salts for secretion [28]. In the case of acute liver damage, betaine may increase at the expense of reduced synthesis of PtdC, and it led to invalid exportation of triglycerides to the blood and accumulation of lipid in the liver. Besides, administration of CCl_4 causes an increase of triglycerides synthesis in the liver [29]. Enhanced synthesis and unbalanced import and export of triglycerides led to increased lipid and triglycerides levels as characteristic caused by the CCl_4 administration [30]. As have been verified by histopathological examination, fat accumulation was observed by the formation of foam cells in acute liver damage models. Coincidentally, ^1H -NMR metabolomics analysis has also revealed that serum LDL/VLDL levels were significantly decreased by 70% in Mod group with VIP value of 3.09 in OPLS-DA models of Mod and Con groups. Such alteration was explainable and supported by previous report where VLDL level was found significantly decreased in blood during

FIGURE 4: Score plots and S-plots of OPLS-DA model between Con and Mod (a, b), SYL and Mod (c, d), and QJ2 and Mod (e, f) groups.

FIGURE 5: Fold changes of potential biomarkers detected between different groups ($n = 7$). * $P < 0.05$, ** $P < 0.01$. Note. Mod, Con, SYL, and QJ1-2 represent liver damage model, control, silybin (40 mg/kg), and QJ (1.2 g/kg) separately. DMA is short for *N,N*-dimethylamine. XXX/YYY means integral of metabolite in XXX group was divided by that of YYY group. The ratio over 1.00 indicated an increase, while ratio less than 1.00 indicated a decrease. Corresponding cell was colored according to the fold change using the color bar labeled at the right side. Statistical analysis was performed by one-way analysis of variance followed by LSD test. * $P < 0.05$; ** $P < 0.01$.

toxic phase [30]. On the other hand, by SYL or QJ treatment, serum levels of LDL/VLDL were significantly elevated 2-3-fold, revealing that abnormal formation or excretion of triglycerides was restored. The restoration can be attributed to suppressed triglycerides formation or improved excretion of triglycerides from liver. Besides, the increase of LDL/VLDL level may reflect relieve of oxidative stress caused by CCl_4 administration.

Glucose, with VIP value, was ranked the second primary metabolite altered in acute liver damage models and was decreased by a half in Mod group comparing with Con group. The decrease was consistent with previous report [30], which may be associated with enhanced energy demand and glycolysis, depressed gluconeogenesis. However, by QJ or SYL treatment, serum glucose levels were partially recovered to different degrees, inferring downregulated glycolysis or enhanced gluconeogenesis.

In TCA cycle intermediate, citrate was noticed to decrease in Mod group, indicating suppressed TCA cycle. The suppression was consistent with previous report and the TCA cycle was disturbed by CCl_4 treatment [31]. On the other hand, suppression of TCA cycle was relieved by QJ or SYL treatment, with citrate level elevated. As end product of glycolysis, lactate

was noticed to increase by 30% in Mod group, with VIP value ranked top 4 of all metabolites, reflecting the enhanced glycolysis, which may be explained by increasing energy consumption or reduced plasma oxygen due to inflammatory response, while, after QJ treatment, serum lactate levels were decreased to different degree, especially for QJ2 group with significant improvement, confirming the best liver protective effect of dosage of 2 g/kg. However, SYL treatment did not alter lactate levels compared with Mod group, which may infer that glycolysis was not altered by SYL treatment. This may be difference of regulations by QJ and SYL, and the underlying mechanism remains to be revealed.

Pyruvate, an important intermediate of glycolysis and gluconeogenesis, can enter into TCA cycle or be converted to alanine via alanine aminotransferase (ALT) and to lactate via lactate dehydrogenase (LDH) [32]. In case of acute liver damage, TCA cycle was suppressed, yet glycolysis was enhanced [33]. As previously reported, excess acetyl-CoA accumulated by β -oxidization of fatty acid inhibits the utilization of pyruvate, finally leading to elevated levels of pyruvate [32], while after treatment of QJ or SYL serum pyruvate levels were decreased by 20%, suggesting improvement of pyruvate utilization, which may benefit form upregulated TCA cycle or enhanced synthesis of alanine.

Another energy metabolism related metabolite, creatine, was detected to be elevated 3-fold in Mod group and with VIP value ranked 5th metabolites altered between Mod and Con group. As is well known, creatinine is a nonenzymatic breakdown product of creatine and phosphocreatine, and the creatine-phosphocreatine system is crucial for cellular energy transportation [34]. Together with glucose and lactate, alterations of creatine indicated increasing energy demands. More importantly, it is generally known that creatinine is used as a routine detection index for renal dysfunction [35]. Thus, it was indicated that CCl_4 treatment may further cause kidney function disruption, not just liver disease alone. Interestingly, though not turning back to normal level, QJ or SYL treatment significantly reduced the creatine levels by ca. 20%, inferring mild improvement of energy consumption.

Amino acid metabolisms were also involved in acute liver damage model and protective effect of QJ or SYL treatment. In our study, alanine, valine, leucine, isoleucine, lysine, serine, and glutamine were detected to be altered significantly and with VIP value >1 . Comparing Mod group with Con group, serum levels of abovementioned amino acids were elevated, which was in accordance with previous report [36]. Such alterations could be attributed to muscle proteolysis as well as to liver parenchyma necrosis [36]. Besides, alanine, isoleucine, and valine are glucogenic amino acids. The increase in these amino acids indicated that gluconeogenesis was suppressed in case of acute liver damage, partly explaining the decreased glucose level. Leucine, lysine, serine, and tyrosine are ketogenic amino acids, which can be transformed to generate ketone bodies. We have noticed that ketogenic amino acids levels were significantly increased in Mod group, which could indirectly explain the accumulation of ketone bodies. Glutamine can transport ammonia, as it is the mechanism of ammonia transportation and storage [37]. The increase of glutamine level may be associated with

FIGURE 6: Potential metabolic pathways disturbed in acute liver damage induced by CCl_4 (red arrow) and restoration regulation by silybin (40 mg/kg, blue arrow) or QJ (2 g/kg, black arrow). * $P < 0.05$, ** $P < 0.01$.

an outflow of AST from hepatocellular mitochondrion, proteolysis, or negative nitrogen balance [38]. On the other hand, treatment of QJ or SYL significantly recovered the elevated amino acid levels, with reduction of different degrees. The restoration indicated that abnormal amino acid metabolism was recovered and impaired hepatic regulating function was improved.

Ketone bodies, including acetone and acetoacetate, were produced by the liver from fatty acids and then converted into acetyl-CoA which then enters the TCA cycle. In case of fasting, starvation, or enhanced fatty acids beta-oxidation, synthesis of ketone bodies was upregulated [39]. In current research, serum acetone and acetoacetate levels were detected to increase by 1- to 2-fold in Mod group, which was consistent with previous report where appearance of ketone bodies was considered to be one of the biomarkers for liver injury induced in rats by CCl₄ exposure [40]. Such increase may be attributed to excessive fatty acid oxidation and serves to eliminate excess amount of acetyl moieties produced. However, after QJ or SYL treatment, serum ketone body levels were detected to decrease, indicating that upregulated synthesis of ketone bodies was suppressed.

Acetate is an end product of fatty acid oxidation, and the doubled serum level of acetate in Mod group further indicate enhanced fatty acids β -oxidation [41]. While, by QJ or SYL treatment, serum acetate levels were decreased by ca. 20%, suggesting that acids β -oxidation was suppressed, which may also explain the decreased ketone body levels.

In summary, QJ treatment partially recovered the abnormal metabolism induced by CCl_4 administration. QJ of 2 g/kg especially exerts comparable liver protective effect with SYL of 40 mg/kg. According to aforementioned analysis, metabolic pathways regulated by QJ treatment were visualized in Figure 6, with the following metabolic pathways involved: betaine metabolism, excretion of LDL/VLDL, gluconeogenesis and glycolysis, tricarboxylic acid (TCA) cycle,

creatinine metabolism, synthesis of ketone bodies, amino acids metabolism, and β -oxidation of fatty acids.

4. Conclusions

In the present work, liver protective effect of QJ was confirmed by serum biochemistry, histopathological examination, and ^1H -NMR based metabolomics analysis. Metabolites significantly altered among groups were identified and discussed. It turned out that QJ treatment can partially recover abnormal metabolism of liver damage through the following metabolic pathways: TCA cycle, synthesis of LDL/VLDL, and gluconeogenesis were enhanced; while betaine metabolism, glycolysis, creatine metabolism, synthesis of ketone bodies, amino acids metabolism, and β -oxidation of fatty acids were suppressed. For the first time ^1H -NMR based metabolism regulation of QJ on acute liver damage was revealed, which provided further understanding for underlying mechanism and demonstrated ^1H -NMR based metabolomics as a useful platform for modern research of TCM.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

This work was supported by the National Nature Science Foundation of China (Grant no. 81603287).

References

- [1] C. Q. Zhao, Y. Zhou, J. Ping, and L. M. Xu, "Traditional Chinese medicine for treatment of liver diseases: progress, challenges and opportunities," *Journal of Integrative Medicine*, vol. 12, no. 5, pp. 401–408, 2014.

- [2] M.-L. Liu, L.-Y. Chien, C.-J. Tai, K.-C. Lin, and C.-J. Tai, "Effectiveness of traditional Chinese medicine for liver protection and chemotherapy completion among cancer patients," *Evidence-Based Complementary and Alternative Medicine*, vol. 2011, Article ID 291843, pp. 1–8, 2011.
- [3] F. Yu, F. Yu, R. Li, and R. Wang, "Inhibitory effects of the *Gentiana macrophylla* (Gentianaceae) extract on rheumatoid arthritis of rats," *Journal of Ethnopharmacology*, vol. 95, no. 1, pp. 77–81, 2004.
- [4] C.-Y. Huang, T.-C. Hsu, W.-W. Kuo et al., "The root extract of *Gentiana macrophylla* Pall. alleviates cardiac apoptosis in lupus prone mice," *PLoS ONE*, vol. 10, no. 5, Article ID e0127440, 2015.
- [5] K. S. Kyoung, H. I. Rhee, E. K. Park et al., "Anti-inflammatory effects of *Radix Gentianae Macrophyllae* (Qinjiao), *Rhizoma Coptidis* (Huanglian) and *Citri Unshiu Pericarpium* (Wenzhou migan) in animal models," *Chinese Medicine*, vol. 3, article 10, 2008.
- [6] X. J. N. Zhang, C. Sun, W. Zhao, and W. Sun, "Anti-inflammatory and analgesic activity of *G. macrophylla* Pall flower and *G. straminea* maxim flower," *Northwest Pharmaceutical Journal*, vol. 27, no. 4, pp. 341–343, 2012.
- [7] X. M. K. Zhang, P. Zhang, and L.-s. Xu, "Screening of active part of *Gentiana macrophylla* Pall. in the liver protection," *China Journal of Traditional Chinese Medicine and Pharmacy*, vol. 29, no. 12, 2014.
- [8] M. Bouhifd, T. Hartung, H. T. Hogberg, A. Kleensang, and L. Zhao, "Review: toxicometabolomics," *Journal of Applied Toxicology*, vol. 33, no. 12, pp. 1365–1383, 2013.
- [9] L. Puchades-Carrasco and A. Pineda-Lucena, "Metabolomics in pharmaceutical research and development," *Current Opinion in Biotechnology*, vol. 35, pp. 73–77, 2015.
- [10] M. Wang, R.-J. A. N. Lamers, H. A. A. J. Korthout et al., "Metabolomics in the context of systems biology: bridging traditional Chinese medicine and molecular pharmacology," *Phytotherapy Research*, vol. 19, no. 3, pp. 173–182, 2005.
- [11] J. Shi, B. Cao, X.-W. Wang et al., "Metabolomics and its application to the evaluation of the efficacy and toxicity of traditional Chinese herb medicines," *Journal of Chromatography B*, vol. 1026, pp. 204–216, 2015.
- [12] J. C. Lindon and J. K. Nicholson, "Spectroscopic and statistical techniques for information recovery in metabolomics and metabolomics," *Annual Review of Analytical Chemistry*, vol. 1, no. 1, pp. 45–69, 2008.
- [13] Z. Pan and D. Raftery, "Comparing and combining NMR spectroscopy and mass spectrometry in metabolomics," *Analytical and Bioanalytical Chemistry*, vol. 387, no. 2, pp. 525–527, 2007.
- [14] B. Jalali Ghassam, H. Ghaffari, H. S. Prakash, and K. R. Kini, "Antioxidant and hepatoprotective effects of *Solanum xanthocarpum* leaf extracts against CCl₄-induced liver injury in rats," *Pharmaceutical Biology*, vol. 52, no. 8, pp. 1060–1068, 2014.
- [15] P. Zhao, C. Qi, G. Wang, X. Dai, and X. Hou, "Enrichment and purification of total flavonoids from *Cortex Juglandis Mandshuricae* extracts and their suppressive effect on carbon tetrachloride-induced hepatic injury in Mice," *Journal of Chromatography B*, vol. 1007, pp. 8–17, 2015.
- [16] H. Rivera, M. Shibayama, V. Tsutsumi, V. Perez-Alvarez, and P. Muriel, "Resveratrol and trimethylated resveratrol protect from acute liver damage induced by CCl₄ in the rat," *Journal of Applied Toxicology*, vol. 28, no. 2, pp. 147–155, 2008.
- [17] Z.-Y. Li, L.-L. Ding, J.-M. Li et al., "¹H-NMR and MS based metabolomics study of the intervention effect of curcumin on hyperlipidemia mice induced by high-fat diet," *PLoS ONE*, vol. 10, no. 3, Article ID e0120950, 2015.
- [18] P. Lettérón, G. Labbe, C. Degott et al., "Mechanism for the protective effects of silymarin against carbon tetrachloride-induced lipid peroxidation and hepatotoxicity in mice. Evidence that silymarin acts both as an inhibitor of metabolic activation and as a chain-breaking antioxidant," *Biochemical Pharmacology*, vol. 39, no. 12, pp. 2027–2034, 1990.
- [19] C. C. L. Zhanwen, J. Ruomin, S. Guoqing, S. Chunqing, and H. Zhibi, "Studies on liver-protection and promoting bile secretion of gentiopicroside," *Chinese Traditional and Herbal Drugs*, vol. 33, no. 1, pp. 47–50, 2004.
- [20] Q. Liang, C. Wang, B. Li, and A.-H. Zhang, "Metabolic fingerprinting to understand therapeutic effects and mechanisms of silybin on acute liver damage in rat," *Pharmacognosy Magazine*, vol. 11, no. 43, pp. 586–593, 2015.
- [21] T. W. Chew, X. Jiang, J. Yan et al., "Folate intake, Mthfr genotype, and sex modulate choline metabolism in mice," *Journal of Nutrition*, vol. 141, no. 8, pp. 1475–1481, 2011.
- [22] A. R. Johnson, C. N. Craciunescu, Z. Guo et al., "Deletion of murine choline dehydrogenase results in diminished sperm motility," *The FASEB Journal*, vol. 24, no. 8, pp. 2752–2761, 2010.
- [23] K. K. Andringa, A. L. King, H. B. Eccleston et al., "Analysis of the liver mitochondrial proteome in response to ethanol and S-adenosylmethionine treatments: Novel molecular targets of disease and hepatoprotection," *American Journal of Physiology-Gastrointestinal and Liver Physiology*, vol. 298, no. 5, pp. G732–G745, 2010.
- [24] T. Murakami, Y. Nagamura, and K. Hirano, "The recovering effect of betaine on carbon tetrachloride-induced liver injury," *Journal of Nutritional Science and Vitaminology*, vol. 44, no. 2, pp. 249–255, 1998.
- [25] M. Lever, P. M. George, J. L. Elmslie et al., "Betaine and secondary events in an acute coronary syndrome cohort," *PLoS ONE*, vol. 7, no. 5, Article ID e37883, 2012.
- [26] Y. Tan, J. Ko, X. R. Liu et al., "Serum metabolomics reveals betaine and phosphatidylcholine as potential biomarkers for the toxic responses of processed *Aconitum carmichaelii* Debx," *Molecular BioSystems*, vol. 10, no. 9, pp. 2305–2316, 2014.
- [27] A. A. Noga and D. E. Vance, "A gender-specific role for phosphatidylethanolamine N-methyltransferase-derived phosphatidylcholine in the regulation of plasma high density and very low density lipoproteins in mice," *The Journal of Biological Chemistry*, vol. 278, no. 24, pp. 21851–21859, 2003.
- [28] Z. Li, L. B. Agellon, and D. E. Vance, "Phosphatidylcholine homeostasis and liver failure," *The Journal of Biological Chemistry*, vol. 280, no. 45, pp. 37798–37802, 2005.
- [29] H. Maling, A. Frank, and M. Horning, "Effect of carbon tetrachloride on hepatic synthesis and release of triglycerides," *Biochimica et Biophysica Acta*, vol. 64, no. 3, pp. 540–545, 1962.
- [30] A. Zira, S. Kostidis, S. Theocharis et al., "¹H NMR-based metabolomics approach in a rat model of acute liver injury and regeneration induced by CCl₄ administration," *Toxicology*, vol. 303, pp. 115–124, 2013.
- [31] P. Guo, D. Wei, J. Wang et al., "Chronic toxicity of crude ricinine in rats assessed by ¹H NMR metabolomics analysis," *RSC Advances*, vol. 5, no. 34, pp. 27018–27028, 2015.
- [32] X. Song, J. Wang, P. Wang, N. Tian, M. Yang, and L. Kong, "¹H NMR-based metabolomics approach to evaluate the effect of Xue-Fu-Zhu-Yu decoction on hyperlipidemia rats induced by high-fat diet," *Journal of Pharmaceutical and Biomedical Analysis*, vol. 78–79, pp. 202–210, 2013.

- [33] C.-Y. Jiang, K.-M. Yang, L. Yang, Z.-X. Miao, Y.-H. Wang, and H.-B. Zhu, "A ^1H NMR-Based Metabonomic Investigation of Time-Related Metabolic Trajectories of the Plasma, Urine and Liver Extracts of Hyperlipidemic Hamsters," *PLoS ONE*, vol. 8, no. 6, Article ID e66786, 2013.
- [34] Y.-Y. Zhao, Y.-L. Feng, X. Bai, X.-J. Tan, R.-C. Lin, and Q. Mei, "Ultra performance liquid chromatography-based metabonomic study of therapeutic effect of the surface layer of *poria cocos* on adenine-induced chronic kidney disease provides new insight into anti-fibrosis mechanism," *PLoS ONE*, vol. 8, no. 3, Article ID e59617, 2013.
- [35] N. Monteiro, M. Branco, S. Peres, F. Borges, and K. Mansinho, "The impact of tenofovir disoproxil fumarate on kidney function: four-year data from the HIV-infected outpatient cohort," *Journal of the International AIDS Society*, vol. 17, supplement 3, no. 4, 2014.
- [36] N. J. Waters, C. J. Waterfield, R. D. Farrant, E. Holmes, and J. K. Nicholson, "Metabonomic deconvolution of embedded toxicity: application to thioacetamide hepato- and nephrotoxicity," *Chemical Research in Toxicology*, vol. 18, no. 4, pp. 639–654, 2005.
- [37] J. T. Brosnan, "Interorgan amino acid transport and its regulation," *Journal of Nutrition*, vol. 133, supplement 1, no. 6, pp. 2068S–2072S, 2003.
- [38] X. Wang, J. Lin, T. Chen, M. Zhou, M. Su, and W. Jia, "Metabolic profiling reveals the protective effect of diammonium glycyrrhizinate on acute hepatic injury induced by carbon tetrachloride," *Metabolomics*, vol. 7, no. 2, pp. 226–236, 2011.
- [39] L. Laffel, "Ketone bodies: a review of physiology, pathophysiology and application of monitoring to diabetes," *Diabetes/Metabolism Research and Reviews*, vol. 15, no. 6, pp. 412–426, 1999.
- [40] L. L. De Zwart, J. Venhorst, M. Groot et al., "Simultaneous determination of eight lipid peroxidation degradation products in urine of rats treated with carbon tetrachloride using gas chromatography with electron-capture detection," *Journal of Chromatography B: Biomedical Sciences and Applications*, vol. 694, no. 2, pp. 277–287, 1997.
- [41] V. S. M. Bernson and D. G. Nicholls, "Acetate, a Major End Product of Fatty-Acid Oxidation in Hamster Brown-Adipose-Tissue Mitochondria," *European Journal of Biochemistry*, vol. 47, no. 3, pp. 517–525, 1974.