

Research Article

CBS-Induced H₂S Generation in Hippocampus Inhibits EA-Induced Analgesia

Wen-Jing Ren,^{1,2} Jia Fu,³ Hai-Yan Yin ^{1,2}, Neng-Gui Xu ⁴, Chun-Zhi Tang,⁴ Li-Zhou Liu,⁵ Shu-Guang Yu ¹ and Yong Tang ^{1,2,3}

¹School of Acupuncture and Tuina, Chengdu University of Traditional Chinese Medicine, Chengdu, Sichuan, China

²Acupuncture & Chronobiology Key Laboratory of Sichuan Province, Chengdu, Sichuan, China

³College of Medicine, Chengdu University, Chengdu, Sichuan, China

⁴School of Acupuncture and Tuina, Guangzhou University of Chinese Medicine, Guangzhou, China

⁵Centre for Health, Activity, and Rehabilitation Research, School of Physiotherapy, University of Otago, Dunedin, New Zealand

Correspondence should be addressed to Yong Tang; tangyong@cdutcm.edu.cn

Received 11 November 2019; Revised 13 March 2020; Accepted 3 April 2020; Published 30 April 2020

Academic Editor: Manel Santafe

Copyright © 2020 Wen-jing Ren et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Hydrogen sulfide (H₂S) is an important mediator participating in both physiological and pathological systems and related to the inflammatory process. Acupuncture has a therapeutic effect on inflammatory pain. However, whether H₂S generated in the central nervous system (CNS) is a mediator of electroacupuncture (EA) treatment for inflammatory pain is unknown. We injected complete Freund's adjuvant (CFA) to induce inflammatory pain and applied EA treatment as an interventional strategy for pain relief. The results presented here show that S-adenosyl-L-methionine (SAM), an allosteric activator of cystathionine-β-synthetase (CBS), may reverse the therapeutic effect of EA. CBS-induced H₂S generation might get involved in the mechanism of EA-induced analgesia in the hippocampus on chronic inflammatory pain.

1. Introduction

Hydrogen sulfide (H₂S) is the most recently accepted endogenously produced gasotransmitter that plays an important role in human health and physiology. Emerging evidence indicates that physiological concentrations of H₂S might have an anti-inflammatory effect, while higher concentrations can exert proinflammatory effects [1]. H₂S is synthesized by three enzymes: cystathionine γ-lyase (CSE), cystathionine β-synthetase (CBS), and 3-mercaptopyruvate sulfurtransferase (3-MST) [2]. The expression of CBS, CSE, and 3-MST shows tissue-specific dominance. Both CBS and 3-MST, as the primary physiological source of H₂S in the central nervous system, predominantly are localized in the brain; CSE abundantly exists at the vascular or nonvascular smooth muscle in the mammalian cardiovascular and respiratory system; therefore, H₂S-producing enzymes are generally ubiquitously expressed in mammalian tissues and impact a wide range of cellular processes [3, 4]. Homocysteine and

cysteine are substrates for H₂S production, and H₂S-synthesizing enzymes are responsible for converting these substrates to H₂S; in the presence of cysteine, especially homocysteine, CBS catalyzes the production of H₂S [5].

H₂S and its synthesizing enzymes got involved in a variety of conditions, including pain and inflammation [6]. It has been indicated that H₂S has an effect on pro- and anti-inflammation, and it is a mediator of peripheral and neurogenic inflammation [1]. The same as inflammation, H₂S participating in modulating visceral pain is also controversial [6]. H₂S-generating enzyme CBS is also involved in several pain conditions. CBS has been found with an increased expression in the dorsal root ganglion (DRGs) of the IBS model induced by neonatal colonic inflammation (NCI) [7]. Inhibition of the CBS-H₂S signaling pathway markedly attenuated heterotypical intermittent stress-induced visceral hyperalgesia; particularly, aminoxyacetic acid (AOAA), one of the CBS inhibitors, suppressed voltage-gated sodium channel currents of the colon-specific DRG neurons and

reversed the enhanced expression of NaV1.7 and NaV1.8 subtypes [8]. These findings indicated that CBS plays a crucial role in visceral hyperalgesia. Furthermore, CBS expression can be also upregulated in neuropathic pain and inflammatory pain. Chronic constriction injury (CCI) surgery caused significantly increased expression of CBS in the ipsilateral ventral and dorsal horn of L4-6 spinal cord segments [9]. Inhibiting CBS by intrathecal injection of AOAA attenuated mechanical and thermal hypersensitivity, and AOAA injecting suppressed the expression of NaV1.7 and NaV1.8 in DRG neurons [10]. Additionally, application of AOAA was observed to alleviate CFA-induced hyperalgesia on the temporomandibular joint and reduce CBS-induced H₂S production in trigeminal ganglion by reversing the enhancement of neural hyperexcitability and increasing the voltage-gated potassium currents [11]. These findings suggest that H₂S and CBS could play a pivotal role in a variety of pain conditions.

Acupuncture has been applied in various pain conditions, and the neurobiological mechanism has been the important focus to be investigated [12, 13]. When the needle is connected with an electrode, it is called as electroacupuncture (EA). EA is also very popular in the basic research and clinic practice because the stimulus is under control [14, 15]. EA stimulation has been found to induce time-dependent cumulative analgesia in rats with neuropathic pain, and the mechanism was demonstrated to be related to the activation of hippocampal MEK1 [16]. Recent studies have also shown that repeated EA intervention leads to synaptic remodeling of hippocampal neurons, and in consequence, the expression of calcium/calmodulin kinase II in the hippocampal CA3 region is upregulated, which contributed to acupuncture analgesia [17]. However, whether H₂S induced by CBS in the hippocampus would be able to get involved in acupuncture analgesia still remains unclear. In this study, we hypothesized that EA might modulate the H₂S generation induced by CBS in the hippocampus to alleviate chronic inflammatory pain.

2. Materials and Methods

2.1. Animals. Male BALB/c mice (8–9 weeks, 20–23 g) were purchased from Chengdu Da Shuo Experimental Animals Co. Ltd. All mice were adapted to the standard laboratory conditions (24 ± 2°C room temperature and 65 ± 5% humidity on 12/12 h light-dark cycles) with drinking water and food available *ad libitum*. The experimental procedures were conducted in accordance with the National Institutes of Health (NIH) Guidelines for the Care and Use of Laboratory Animals and approved by the Animal Ethics Committee of Chengdu University of Traditional Chinese Medicine. After adaptive domestication for one week, mice were randomly divided into different groups based on random numbers generated by SPSS software.

2.2. Chronic Inflammatory Pain Model. Mice were anesthetized with 1% isoflurane, and complete Freund's adjuvant (CFA, 20 µl) or saline (20 µl) was subcutaneously injected

into the right hind paw [18]. Thermal hyperalgesia was detected at the first day after injection [19].

2.3. EA Treatment. An electroacupuncture apparatus (HANS-200A acupoint nerve stimulator, Nanjing Jisheng Medical Co., Ltd) was used for EA treatment. EA stimulation was provided at the first day after CFA injection and subsequently lasted for one week. Mice were shaved to expose ST36, located at the posterolateral part of the knee about 2 mm below the fibular head [20], and two stainless steel needles (0.18 mm in diameter, 13 mm in length) were inserted directly at ST36 of both hind limbs about 3 mm deep. The stimulated parameters of EA included 0 Hz, 2 Hz, 100 Hz, and 2/100 Hz (0.2 mA intensity), lasting for 30 min during one session. During treatment, all mice were restrained on a specific device [21].

2.4. Intraperitoneal Injection. Mice were randomly divided into seven groups: CFA + Saline (*n* = 6), CFA + NaHS (*n* = 6), CFA + HA (*n* = 7), CFA + AOAA (*n* = 6), CFA + SAM (*n* = 7), CFA + IC2A (*n* = 7), and CFA + L-aspartate (*n* = 6). Saline, NaHS (H₂S donor, 5.6 mg/kg), hydroxylamine (HA, one of CBS inhibitors, 50 mg/kg), AOAA (one of CBS inhibitors, 45 mg/kg), SAM (CBS agonist, 100 mg/kg), and IC2A (7.9 mg/kg) or L-aspartate (1.25 mg/kg) were injected intraperitoneally one day after CFA injection and lasted for one week. No abnormalities were observed during the period of injection.

2.5. Surgical Procedures and Microinjection. Stereotaxic surgery was performed 7 days prior to experiments. Mice were anesthetized with isoflurane and fixed on a stereotaxic platform. Coordinates of A-P: 2.5 mm relative to the bregma, lateral: ±2.5 mm relative to the midline, and depth: 2 mm from the duramater were used for implanting guide cannulas (RWD, China) into the bilateral hippocampi [22]. After recovery from surgery, HA (9 nmol/1 µl [23]), AOAA (9 nmol/1 µl [24, 25]), or SAM (10 pmol/1 µl [26]) was bilaterally microinjected into the hippocampi. DAO and CAT inhibitors followed the same procedure. A 5 µl microsyringe was held by a micromanipulator on the stereotaxic apparatus (RWD, China), and injection was performed at a rate of 0.5 µl/min. The volume of each microinjection was 1 µl. Afterwards, the needle was kept at the injection place for another 5 minutes after each injection to allow for sufficient diffusion. No adverse effect was observed during each injection.

2.6. Thermal Withdrawal Latency (TWL). Mice were placed in behavior boxes on a glass platform before the behavior test [27]. Thermal hyperalgesia was assessed by measuring the TWL with a Plantar Test Apparatus (10% light intensity, 10 sec baseline latency, and 20 sec cutoff time. Hargreaves method, PL-200, Tai Meng, China). After 30-min of acclimatization, a mobile radiant heat source was focused on the plantar surface of the right hind paw. Each mouse was tested three times with intervals of 5 min. Pain threshold was tested




FIGURE 1: EA at a low frequency of 2 Hz had analgesic effects on thermal hyperalgesia in mice with inflammatory pain. (a) Schematic drawing for CFA injection and the location of acupoint ST36 in a mouse. (b) EA (0 Hz) had no significant therapeutic effect on chronic inflammatory pain ($P > 0.05$); $F(4, 40) 0.3040$, $P=0.8736$ (interaction), $F(4, 40) 123.7$, $P < 0.0001$ (time), and $F(1, 10) 0.6222$, $P=0.4485$ (column factor). (c) Chronic inflammatory pain could be alleviated by EA at low frequency (2 Hz) ($P < 0.01$); $F(4, 44) 8.287$, $P < 0.0001$ (interaction), $F(4, 44) 53.98$, $P < 0.0001$ (time), and $F(1, 11) 13.39$, $P=0.0038$ (column factor). (d) EA at 100 Hz failed to relieve hyperalgesia ($P > 0.05$); $F(4, 44) 2.366$, $P=0.0674$ (interaction), $F(4, 44) 116.6$, $P < 0.0001$ (time), and $F(1, 11) 0.008546$, $P=0.9280$ (column factor). (e) EA at 2/100 Hz failed to relieve hyperalgesia ($P > 0.05$); $F(4, 44) 3.416$, $P=0.0162$ (interaction), $F(4, 44) 98.86$, $P < 0.0001$ (time), and $F(1, 11) 2.898$, $P=0.1167$ (column factor). Data MEANS \pm SEM

as the baseline at day 1 before CFA injection and day 1, 3, 5, and 7 after CFA injection. In order to eliminate observational bias, all behavioral data were collected and recorded by the same investigator who was not responsible for drug injection or EA treatment.

2.7. *Drugs.* Complete Freund's adjuvant (CFA, F5881), sodium hydrogen sulfide (NaHS, 161527), hydroxylamine (HA, 438227), aminoxyacetic acid (AOAA, C13408), and L-aspartate (A9256) were purchased from Sigma (Sigma-Aldrich St. Louis, MO, USA). IC2A (I129126) was purchased




FIGURE 2: Inhibiting CBS activity alleviated pain sensation on the CFA mouse model. (a) H₂S donor NaHS failed to reduce thermal pain behavior ($P > 0.05$); $F(4, 40) 0.3923$, $P = 0.8129$ (interaction), $F(4, 40) 62.42$, $P < 0.0001$ (time), and $F(1, 10) 0.2892$, $P = 0.6025$ (column factor). (b) CBS agonist SAM failed to attenuate thermal pain behavior ($P > 0.05$); $F(4, 44) 0.08392$, $P = 0.9870$ (interaction), $F(4, 44) 42.73$, $P < 0.0001$ (time), and $F(1, 11) 0.3571$, $P = 0.5622$ (column factor). (c) HA notably attenuated thermal hypersensitivity ($P < 0.01$); $F(4, 44) 6.669$, $P = 0.0003$ (interaction), $F(4, 44) 24.06$, $P < 0.0001$ (time), and $F(1, 11) 18.68$, $P = 0.0012$ (column factor). (d) AAOA also remarkably relieved pain behavior ($P < 0.05$); $F(4, 40) 7.503$, $P = 0.0001$ (interaction), $F(4, 40) 27.84$, $P < 0.0001$ (time), and $F(1, 10) 12.95$, $P = 0.0049$ (column factor). (e) The injection of IC2A and L-aspartate had no effect on relieving thermal pain ($P > 0.05$); $F(8, 64) 0.8894$, $P = 0.5306$ (interaction), $F(4, 64) 81.15$, $P < 0.0001$ (time), and $F(2, 16) 2.716$, $P = 0.0965$ (column factor). Data MEANS \pm SEM

from Aladin (Shanghai Aladin Biochemical Technology Co. LTD). S-adenosine-L- methionine (SAM, S9990) was purchased from Solarbio (Beijing Solarbio Technology Co. LTD). Saline was purchased from Kelun (Sichuan Kelun Pharmaceutical Co. Ltd.).

2.8. Statistical Analysis. All data were analyzed and graphed using Graphpad Prism 6 (GraphPad Software, Inc., La Jolla, CA, USA). Data were expressed as data = MEANS \pm SEM (standard error of means). Two-way repeated-measure ANOVA followed by the Tukey-Kramer test was performed to compare the differences between groups. A value of $P < 0.05$ was considered statistically significant.

3. Results

3.1. Low-Frequency EA Stimulation Alleviated Chronic Inflammatory Pain. To investigate the efficacy of EA in chronic inflammatory pain, we injected CFA in the right plantar surface of the hind paw and measured responses to the thermal stimuli in separate groups at a different time point. We chose

the ST36 acupuncture point, which is one of the most frequently used acupoints for pain relief [28, 29] (Figure 1(a)). After one day, different groups received acupuncture treatments with different stimulated parameters at ST36, and then we measured changes of TWL on mice (Figures 1(b)–1(e)). On the first day after CFA injection, we found that the baseline value of TWL was significantly decreased (Figures 1(b)–1(e)). On the 3rd day after acupuncture treatment, EA at low frequency (2 Hz) had an analgesic effect on thermal hyperalgesia compared with EA at high-frequency stimulation (100 Hz) and alternating frequency (2/100 Hz) on mice with inflammatory pain ($P < 0.05$) (Figures 1(c)–1(e)). In addition, acupuncture without electrostimulation also failed to alleviate pain sensation ($P > 0.05$) (Figure 1(b)). The result suggests that low-frequency EA has a therapeutic effect on CFA-induced inflammatory pain, but acupuncture and high or alternating frequency EA stimulation failed to alleviate pain.

3.2. Intraperitoneal Injection of CBS Inhibitors Attenuated Thermal Hypersensitivity. To study the relationship between H₂S and inflammation, NaHS (5.6 mg/kg [30, 31]) was

intraperitoneally injected one day after CFA injection, and results indicated that NaHS failed to alleviate pain behavior (Figure 2(a)). Similar to NaHS, SAM (100 mg/kg) [32, 33] had no therapeutic effect on inflammatory pain (Figure 2(b)). However, HA (50 mg/kg) [34] and AOAA (45 mg/kg) [34] attenuated thermal hypersensitivity (Figures 2(c)–2(d)). A recent finding indicated that H₂S could also be produced by 3-MST from 3-mercaptopyruvate in the brain [35, 36]. Therefore, we applied a specific cysteine aminotransferase (CAT) inhibitor L-aspartate and D-amino acid oxidase (DAO) inhibitor IC2A [37]. However, they also failed to alleviate inflammatory pain (Figure 2(e)). These results indicated that CBS-induced H₂S induced the pain behavior and CBS inhibitors could alleviate pain sensation.

3.3. Hippocampal Injection of CBS Inhibitors Attenuated Thermal Hypersensitivity. To study the relationship between CBS-induced H₂S in the brain and thermal hypersensitivity, the CBS inhibitor HA (9 nmol) [23] or AOAA (9 nmol) [24, 25] was bilaterally microinjected into the hippocampus. Moreover, the CBS-specific agonist SAM (10 pmol) [26] or saline was also bilaterally injected into the hippocampus one day after CFA injection, and CAT and DAO inhibitors [37] followed the same procedure (Figure 3(a)). Results showed that HA and AOAA microinjection significantly increased the TWL and attenuated thermal hypersensitivity (Figures 3(b) and 1(c)). However, saline, SAM, IC2A, or L-aspartate injection failed to alleviate thermal pain behavior (Figures 3(d) and 3(e)). Therefore, suppressing H₂S induced by CBS in the hippocampus might relieve CFA-induced chronic inflammatory pain.

3.4. SAM Reversed the Therapeutic Effect of Low-Frequency EA. To determine whether CBS influences EA therapeutic effect on chronic inflammatory pain, EA intervention (2 Hz, 0.2 mA) was performed immediately after SAM injection or microinjection. However, EA stimulation failed to alleviate thermal pain behavior after SAM injection (Figures 4(a) and 4(b)). These results indicated that activating CBS reversed EA therapeutic effect on inflammatory pain, and CBS-induced H₂S release might inhibit EA therapeutic effect.

4. Discussion

Our data demonstrated that low-frequency EA stimulation notably alleviated the thermal pain behavior, and injection of CBS inhibitors might have a therapeutic effect on thermal pain. As an essential synthetase of H₂S, CBS is mostly responsible for the production of H₂S in brain tissue and is highly expressed in the hippocampus [38, 39]. Therefore, H₂S generation in the hippocampus might participate in the CFA-induced chronic inflammatory pain. Our data further showed that SAM, a CBS agonist, could reverse EA therapeutic effect on chronic inflammatory pain. Taken together, the downregulation of CBS-induced H₂S generation might be one of the mechanisms of EA treatment for chronic inflammatory pain.

H₂S is the third gasotransmitter discovered after NO, CO. It is considered as an important signaling molecule in the regulation of pathophysiological processes, including pain and inflammation [34, 40, 41]. However, the precise role of H₂S in inflammation is not clear. It was supposed to have pro- or anti-inflammatory effects under different conditions. In general, H₂S at physiological concentrations might have an anti-inflammatory effect, while increased concentrations of H₂S can exert proinflammatory effects [1]. Intraplantar administration of NaHS, the donor of H₂S, evoked mechanical hyperalgesia [42]. It was found that H₂S might activate Cav3.2 T-type Ca²⁺ channels that led to sensitization of nociceptive processing and hyperalgesia [43]. H₂S synthetase CSE and CBS also play a pivotal role in pain and inflammation. Pretreatment with the CSE inhibitor propargylglycine significantly reduced hind paw edema and decreased granulocyte infiltration into the tissue in response to an injection of carrageenan [44]. In the visceral pain rat model, the expression of CBS in the spinal cord was notably increased [45]. Some investigations also indicated that CBS was upregulated by TLR4 in NCI rats and mediated by the NF- κ B signaling pathway, thus contributing to visceral hypersensitivity [46]. In addition, the expression of CBS also has an effect on inflammatory pain. It was found that the expression of CBS in the rat dorsal root ganglia (DRG) was significantly upregulated after intraplantar administration of CFA, and the mechanical hyperalgesia was expectedly attenuated by intraperitoneal injection of CBS inhibitors in a dose-dependent manner [34]. These investigations suggest that H₂S and its enzymes in the peripheral nervous system are regulated by nociception. However, analgesic effect of H₂S and its enzymes on the brain are yet to be investigated in future studies.

CBS is responsible for most of the H₂S production in the brain, and it is highly expressed in the hippocampus. Multiple evidences indicated that the hippocampal formation responded to external nociceptive stimuli [47–49]. Previously, investigations showed that SAM resulted in an up to eight or ten-fold increase of the enzymatic activity of CBS [50–52], and CBS inhibitors HA and AOAA suppressed H₂S production in brain homogenates which can be enhanced by the CBS activator [39]. In addition, several investigations found that intraperitoneal injection of CBS inhibitors could decrease CBS expression and activity in brain tissue. For instance, Liu and his colleagues found that intraperitoneal injection of HA inhibited CBS protein expression as examined by western blot analysis and immunohistochemical analysis [53]. AOAA (i.p) also reduced CBS activity in brain tissue [54]. Thus, in order to study the relationship between CBS-induced H₂S and inflammatory pain, HA or AOAA was intraperitoneally injected into mice. Results has shown that injecting HA or AOAA led to relieve pain by inhibiting CBS. It is in consistent with the previous research [34]. However, analgesic effect of CBS-induced H₂S on the brain is yet to be investigated in future studies. To further investigate whether CBS located in the hippocampus would affect pain sensation, HA or AOAA was bilaterally microinjected into the hippocampus. We found that inhibiting H₂S generation in the hippocampus alleviated




FIGURE 3: Inhibiting CBS activity in the hippocampus alleviated pain sensation on the CFA mice model. (a) Schematic drawing of microinjection in the hippocampus. (b) The injection of HA notably attenuated thermal hypersensitivity ($P < 0.05$); $F(5, 50) 3.790$, $P = 0.0055$ (interaction), $F(5, 50) 50.14$, $P < 0.0001$ (time), and $F(1, 10) 10.78$, $P = 0.0082$ (column factor). (c) AOOA remarkably relieved pain behavior ($P < 0.05$); $F(5, 50) 3.760$, $P = 0.0057$ (interaction), $F(5, 50) 40.77$, $P < 0.0001$ (time), and $F(1, 10) 13.44$, $P = 0.0043$ (column factor). (d) SAM failed to alleviate thermal pain behavior ($P > 0.05$); $F(5, 50) 0.3387$, $P = 0.8870$ (interaction), $F(5, 50) 78.47$, $P < 0.0001$ (time), and $F(1, 10) 0.004093$, $P = 0.9502$ (column factor). (e) IC2A and L-aspartate injection failed to alleviate thermal pain behavior ($P > 0.05$); $F(10, 75) 1.313$, $P = 0.2394$ (interaction), $F(5, 75) 125.1$, $P < 0.0001$ (time), and $F(2, 15) 1.092$, $P = 0.3608$ (column factor). Data MEANS \pm SEM

inflammatory pain and activated H_2S generation by microinjected SAM had no effect on inflammatory pain. However, how CBS in the hippocampus participates in pain sensation is yet to be investigated. According to previous studies, CBS converted homocysteine to cysteine [55]. CBS inhibitors could promote accumulation of L-homocysteine and L-cysteine. It was reported that reducing L-cysteine and L-homocysteine induced time and dose-dependent peripheral hyperalgesia which could be modulated by 3bOH, a potent T-type Ca^{2+} channel blocker [56]. Therefore, homocysteine and cysteine might participate in inflammatory pain process, but it also needs to be further investigated.

Acupuncture has a therapeutic effect on inflammatory pain, and EA therapy has a cumulative effect [16, 57, 58]. The frequency of EA is one of the crucial factors to significantly influence its therapeutic effects. EA with different frequencies is regarded as distinct therapeutic methods and have been used in different diseases. High-frequency (100 Hz) stimulation was more effective as compared with the low frequency (2 Hz) in amelioration of muscle spasticity, and it might be mediated by dynorphin generating from the central nervous system [59]. Low-frequency EA

released β -endorphin and enkephalins, while high-frequency EA released the dynorphins to suppress pain sensation [60]. However, low-frequency EA produced the significant rapid decrease in ankle edema after ankle sprain which is an important feature of inflammation [61]. Furthermore, low-frequency EA (10 Hz) significantly reduced CFA-induced hind paw edema to suppress inflammation by activating the hypothalamus-pituitary-adrenal axis (HPA); however, 100 Hz EA had no effect on it [62]. Some investigations demonstrated that low-frequency EA could regulate the expression level of genes in the nucleus arcuate region, and low-frequency EA at ST36 in rats markedly relieved pain [63, 64]. Low-frequency EA could also suppress spinal long-term potential (LTP) to alleviate pain sensation in colitis rats [65]. Additionally, low-frequency stimulation, rather than high frequency, invoked endogenous mechanisms for the release of 5-hydroxytryptamine (5-HT) in the spinal cord, which activated 5-HT receptors to reduce hyperalgesia in rats with joint inflammation [66]. In accordance with these results, we found that low-frequency EA delivered to ST36 could alleviate pain. It has been reported that EA applied to ST36 could influence hemodynamic




FIGURE 4: Treatment of CBS agonist SAM reverses the therapeutic effect of EA. (a) Compared with EA(2 Hz), EA failed to alleviate pain sensation after SAM intraperitoneal injection ($P < 0.01$); $F(8, 64) = 3.965$, $P = 0.0007$ (interaction), $F(4, 64) = 77.51$, $P < 0.0001$ (time), and $F(2, 16) = 7.916$, $P = 0.0041$ (column factor). (b) EA also failed to alleviate pain sensation after SAM microinjection ($P > 0.05$); $F(5, 50) = 3.571$, $P = 0.0077$ (interaction), $F(5, 50) = 202.1$, $P < 0.0001$ (time), and $F(1, 10) = 0.2729$, $P = 0.6128$ (column factor). Data = MEANS \pm SEM.

signaling in the hippocampus [42]. Additionally, the previous report indicated that EA protected against hypoxic ischemic brain damage in immature rats via decreasing H_2S generation in brain tissue [53]. Also, EA treatment decreased CBS expression level and increased HO-1 and HIF-1 α expression levels in perinatal rat cortex cells [67]. Thus, EA protected against hypoxic damage via the hydrogen sulfide/CBS-CO/HO-1-HIF-1 α system.

To further determine the role of H_2S in EA stimulation, EA treatment was performed after injection of the CBS agonist SAM on a CFA-induced inflammatory mouse model, and we found that the EA therapeutic effect was suppressed by SAM. H_2S was expected to play an important role in modulating EA effect on inflammatory pain. Although the H_2S /CBS system participates in the inflammatory pain and it is related to EA treatment, the mechanisms of how EA modulates H_2S release need to be confirmed in our future work. Previous research has shown that intracellular Ca^{2+} can be regulated by the cAMP/PKA pathway, and H_2S can regulate $[Ca^{2+}]_i$ through the cAMP/PKA system [68]. EA pretreatment could produce an antiarrhythmic effect by regulating the L-type Ca^{2+} channel [69]. Additionally, AOAA, an inhibitor of CBS, could also suppress the potentiation of ATP-induced intracellular calcium signals in DRG neurons to markedly attenuate pain hypersensitivity [70]. Some investigators also showed that EA pretreatment could inhibit Ca^{2+} influx to alleviate the LPS-induced inflammation in rats [71]. Therefore, we speculated that acupuncture regulation of H_2S production might be related to calcium channel opening. This needs to be proved in our future work.

5. Conclusions

Current data indicated that the CBS-specific agonist SAM may reverse the therapeutic effect of EA. Downregulation of CBS might be a mediator of EA-induced analgesia on chronic inflammatory pain.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Authors' Contributions

Wen-Jing Ren, Jia Fu, and Hai-Yan Yin contributed equally to this work.

Acknowledgments

The authors appreciate Prof. Peter Illes (Universität Leipzig, Germany) for English language corrections. This work was supported by the grants from the National Natural Science Foundation of China (nos 81774437 and 81373735), the Project First-Class Disciplines Development of the Chengdu University of TCM (CZYHW1901), the Science & Technology Department of Sichuan Province (nos 2018HH0123 and 2019YFH0108), and the 2017-18 NZ-China NCD Emerging Researcher Travel Fellowship.

References

- [1] L. Li, M. Bhatia, and P. Moore, "Hydrogen sulphide—a novel mediator of inflammation?" *Current Opinion in Pharmacology*, vol. 6, no. 2, pp. 125–129, 2006.
- [2] M. D. Hartle and M. D. Pluth, "A practical guide to working with H_2S at the interface of chemistry and biology," *Chemical Society Reviews*, vol. 45, no. 22, pp. 6108–6117, 2016.

- [3] R. Wang, "Physiological implications of hydrogen sulfide: a whiff exploration that blossomed," *Physiological Reviews*, vol. 92, no. 2, pp. 791–896, 2012.
- [4] Q. Yang and G. He, "Imbalance of homocysteine and H₂S: significance, mechanisms, and therapeutic promise in vascular injury," *Oxidative Medicine and Cellular Longevity*, vol. 2019, Article ID 7629673, 2019.
- [5] H. Zhu, S. Blake, K. T. Chan, R. B. Pearson, and J. Kang, "Cystathionine β -synthase in physiology and cancer," *BioMed Research International*, vol. 2018, Article ID 3205125, 11 pages, 2018.
- [6] J. L. Wallace and R. Wang, "Hydrogen sulfide-based therapeutics: exploiting a unique but ubiquitous gasotransmitter," *Nature Reviews Drug Discovery*, vol. 14, no. 5, pp. 329–345, 2015.
- [7] L. Zhu, L. Zhao, R. Qu et al., "Adrenergic stimulation sensitizes TRPV1 through upregulation of cystathionine β -synthetase in a rat model of visceral hypersensitivity," *Scientific Reports*, vol. 5, Article ID 16109, 2015.
- [8] Y. Wang, R. Qu, S. Hu, Y. Xiao, X. Jiang, and G.-Y. Xu, "Upregulation of cystathionine β -synthetase expression contributes to visceral hyperalgesia induced by heterotypic intermittent stress in rats," *PLoS One*, vol. 7, no. 12, Article ID e53165, 2012.
- [9] Y. Gui, A. Li, B. Qiu et al., "Endogenous CBS-H₂S pathway contributes to the development of CCI-induced neuropathic pain," *Neurochemical Research*, vol. 41, no. 6, pp. 1381–1389, 2016.
- [10] J. Yan, S. Hu, K. Zou et al., "Inhibition of cystathionine beta-synthetase suppresses sodium channel activities of dorsal root ganglion neurons of rats with lumbar disc herniation," *Scientific Reports*, vol. 6, Article ID 38188, 2016.
- [11] X. Miao, X. Meng, G. Wu et al., "Upregulation of cystathionine- β -synthetase expression contributes to inflammatory pain in rat temporomandibular joint," *Molecular Pain*, vol. 10, p. 9, 2014.
- [12] B. Y. K. Loyeung and D. M. Cobbin, "Investigating the effects of three needling parameters (manipulation, retention time, and insertion site) on needling sensation and pain profiles: a study of eight deep needling interventions," *Evidence-Based Complementary and Alternative Medicine*, vol. 2013, Article ID 136763, 12 pages, 2013.
- [13] M. Huang, X. Wang, B. Xing et al., "Critical roles of TRPV2 channels, histamine H₁ and adenosine A₁ receptors in the initiation of acupoint signals for acupuncture analgesia," *Scientific Reports*, vol. 8, no. 1, p. 6523, 2018.
- [14] Y. Tang, H.-Y. Yin, P. Rubini, and P. Illes, "Acupuncture-induced analgesia," *The Neuroscientist*, vol. 22, no. 6, pp. 563–578, 2016.
- [15] T. Hou, H. Xiang, L. Yu et al., "Electroacupuncture inhibits visceral pain via adenosine receptors in mice with inflammatory bowel disease," *Purinergic Signalling*, vol. 15, no. 2, pp. 193–204, 2019.
- [16] Y.-H. Gao, C.-W. Li, J.-Y. Wang et al., "Activation of hippocampal MEK1 contributes to the cumulative antinociceptive effect of electroacupuncture in neuropathic pain rats," *BMC Complementary Medicine and Therapies*, vol. 16, no. 1, p. 517, 2016.
- [17] Q. Xu, T. Liu, S. Chen et al., "The cumulative analgesic effect of repeated electroacupuncture involves synaptic remodeling in the hippocampal CA3 region," *Neural Regeneration Research*, vol. 7, no. 18, pp. 1378–1385, 2012.
- [18] D. F. Martins, R. N. Brito, J. Stramosk et al., "Peripheral neurobiologic mechanisms of antiallodynic effect of warm water immersion therapy on persistent inflammatory pain," *Journal of Neuroscience Research*, vol. 93, no. 1, pp. 157–166, 2015.
- [19] P. B. Jacobson, S. J. Morgan, D. M. Wilcox et al., "A new spin on an old model: in vivo evaluation of disease progression by magnetic resonance imaging with respect to standard inflammatory parameters and histopathology in the adjuvant arthritic rat," *Arthritis & Rheumatism*, vol. 42, no. 10, pp. 2060–2073, 1999.
- [20] W. Kim, M. Kim, D. Go, B. Min, H. Na, and S. Kim, "Combined effects of bee venom acupuncture and morphine on oxaliplatin-induced neuropathic pain in mice," *Toxins*, vol. 8, no. 2, p. 33, 2016.
- [21] C.-Y. Zuo, P. Lv, C.-S. Zhang et al., "Ipsi- and contralateral moxibustion generate similar analgesic effect on inflammatory pain," *Evidence-based Complementary and Alternative Medicine*, vol. 2019, Article ID 1807287, 7 pages, 2019.
- [22] G. Paxinos and K. B. J. Franklin, *The Mouse Brain in Stereotaxic Coordinates*, Elsevier Academic Press, San Diego, CA, USA, 2nd edition, 1982.
- [23] Y.-F. Liang, D.-D. Zhang, X.-J. Yu et al., "Hydrogen sulfide in paraventricular nucleus attenuates blood pressure by regulating oxidative stress and inflammatory cytokines in high salt-induced hypertension," *Toxicology Letters*, vol. 270, pp. 62–71, 2017.
- [24] J. P. J. Sabino, R. N. Soriano, A. F. Donatti et al., "Involvement of endogenous central hydrogen sulfide (H₂S) in hypoxia-induced hypothermia in spontaneously hypertensive rats," *Canadian Journal of Physiology and Pharmacology*, vol. 95, no. 2, pp. 157–162, 2017.
- [25] J. P. J. Sabino, R. N. Soriano, B. M. Santos et al., "Central administration of aminoxyacetate, an inhibitor of H₂S production, affects thermoregulatory but not cardiovascular and ventilatory responses to hypercapnia in spontaneously hypertensive rats," *Respiratory Physiology & Neurobiology*, vol. 263, pp. 38–46, 2019.
- [26] H. Yu, H. Xu, X. Liu et al., "Superoxide mediates depressive effects induced by hydrogen sulfide in rostral ventrolateral medulla of spontaneously hypertensive rats," *Oxidative Medicine and Cellular Longevity*, vol. 2015, Article ID 927686, 8 pages, 2015.
- [27] E. S. Fernandes, F. A. Russell, K. M. Alawi et al., "Environmental cold exposure increases blood flow and affects pain sensitivity in the knee joints of CFA-induced arthritic mice in a TRPA1-dependent manner," *Arthritis Research & Therapy*, vol. 18, p. 7, 2016.
- [28] S.-J. Bae, J. Lim, S. Lee et al., "Augmented mechanical forces of the surface-modified nanoporous acupuncture needles elicit enhanced analgesic effects," *Frontiers in Neuroscience*, vol. 13, p. 652, 2019.
- [29] S. Kim, X. Zhang, S. C. O'Buckley et al., "Acupuncture resolves persistent pain and neuroinflammation in a mouse model of chronic overlapping pain conditions," *The Journal of Pain*, vol. 19, no. 12, pp. 1384.e1381–1384.e1314, 2016.
- [30] Z. Wang, J. Zhan, X. Wang et al., "Sodium hydrosulfide prevents hypoxia-induced behavioral impairment in neonatal mice," *Brain Research*, vol. 1538, pp. 126–134, 2013.
- [31] Y. Cui, X. Duan, H. Li et al., "Hydrogen sulfide ameliorates early brain injury following subarachnoid hemorrhage in rats," *Molecular Neurobiology*, vol. 53, no. 6, pp. 3646–3657, 2016.
- [32] V. M. Izaskun, N. Fabiana, D. S. Sara et al., "Striatal adenosine A_{2A} receptor expression is controlled by S-adenosyl-L-methionine-mediated methylation," *Purinergic Signalling*, vol. 10, no. 3, pp. 523–528, 2014.

- [33] H.-L. Zhao, P.-Y. Pan, Y. Yang et al., "Endogenous hydrogen sulphide attenuates NLRP3 inflammasome-mediated neuroinflammation by suppressing the P2X7 receptor after intracerebral haemorrhage in rats," *Journal of Neuroinflammation*, vol. 14, no. 1, p. 163, 2017.
- [34] F. Qi, Y. Zhou, Y. Xiao et al., "Promoter demethylation of cystathionine- β -synthetase gene contributes to inflammatory pain in rats," *Pain*, vol. 154, no. 1, pp. 34–45, 2013.
- [35] Y. Kimura, Y. Toyofuku, S. Koike et al., "Identification of H₂S₃ and H₂S produced by 3-mercaptopyruvate sulfurtransferase in the brain," *Scientific Reports*, vol. 5, p. 14774, 2015.
- [36] N. Shibuya, M. Tanaka, M. Yoshida et al., "3-Mercaptopyruvate sulfurtransferase produces hydrogen sulfide and bound sulfane sulfur in the brain," *Antioxidants & Redox Signaling*, vol. 11, no. 4, pp. 703–714, 2009.
- [37] H. Luo, P.-F. Wu, Q.-Q. Han et al., "Reactive sulfur species emerge as gliotransmitters to support memory via sulfuration-dependent gating of NR2A-containing n-methyl-D-aspartate subtype glutamate receptor function," *Antioxidants & Redox Signaling*, vol. 30, no. 16, pp. 1880–1899, 2019.
- [38] A. Asimakopoulou, P. Panopoulos, C. T. Chasapis et al., "Selectivity of commonly used pharmacological inhibitors for cystathionine β synthase (CBS) and cystathionine γ lyase (CSE)," *British Journal of Pharmacology*, vol. 169, no. 4, pp. 922–932, 2013.
- [39] K. Abe and H. Kimura, "The possible role of hydrogen sulfide as an endogenous neuromodulator," *The Journal of Neuroscience*, vol. 16, no. 3, pp. 1066–1071, 1996.
- [40] J. E. Roa-Coria, J. B. Pineda-Farias, P. Barragán-Iglesias et al., "Possible involvement of peripheral TRP channels in the hydrogen sulfide-induced hyperalgesia in diabetic rats," *BMC Neuroscience*, vol. 20, no. 1, p. 1, 2019.
- [41] S. Nishimura, O. Fukushima, H. Ishikura et al., "Hydrogen sulfide as a novel mediator for pancreatic pain in rodents," *Gut*, vol. 58, no. 6, pp. 762–770, 2009.
- [42] A. Kawabata, T. Ishiki, K. Nagasawa et al., "Hydrogen sulfide as a novel nociceptive messenger," *Pain*, vol. 132, no. 1–2, pp. 74–81, 2007.
- [43] Y. Maeda, Y. Aoki, F. Sekiguchi et al., "Hyperalgesia induced by spinal and peripheral hydrogen sulfide: evidence for involvement of Cav3.2 T-type calcium channels," *Pain*, vol. 142, no. 1–2, pp. 127–132, 2009.
- [44] J. L. Wallace, "Hydrogen sulfide-releasing anti-inflammatory drugs," *Trends in Pharmacological Sciences*, vol. 28, no. 10, pp. 501–505, 2007.
- [45] E. Distrutti, L. Sediari, A. Mencarelli et al., "Evidence that hydrogen sulfide exerts antinociceptive effects in the gastrointestinal tract by activating KATP channels," *Journal of Pharmacology and Experimental Therapeutics*, vol. 316, no. 1, pp. 325–335, 2006.
- [46] B. Yuan, W.-H. Tang, L.-J. Lu et al., "TLR4 upregulates CBS expression through NF- κ B activation in a rat model of irritable bowel syndrome with chronic visceral hypersensitivity," *World Journal of Gastroenterology*, vol. 21, no. 28, pp. 8615–8628, 2015.
- [47] E. J. Shim, Y. J. Seo, M. S. Kwon et al., "The intracerebroventricular kainic acid-induced damage affects animal nociceptive behavior," *Brain Research Bulletin*, vol. 73, no. 4–6, pp. 203–209, 2007.
- [48] A. A. Mutso, D. Radzicki, M. N. Baliki et al., "Abnormalities in hippocampal functioning with persistent pain," *Journal of Neuroscience*, vol. 32, no. 17, pp. 5747–5756, 2012.
- [49] J. Zheng, Y.-Y. Jiang, L.-C. Xu et al., "Adult hippocampal neurogenesis along the dorsoventral axis contributes differentially to environmental enrichment combined with voluntary exercise in alleviating chronic inflammatory pain in mice," *The Journal of Neuroscience*, vol. 37, no. 15, pp. 4145–4157, 2017.
- [50] X. Shan and W. D. Kruger, "Correction of disease-causing CBS mutations in yeast," *Nature Genetics*, vol. 19, no. 1, pp. 91–93, 1998.
- [51] T. Majtan and J. P. Kraus, "Folding and activity of mutant cystathionine β -synthase depends on the position and nature of the purification tag: characterization of the R266K CBS mutant," *Protein Expression and Purification*, vol. 82, no. 2, pp. 317–324, 2012.
- [52] C. Szabo and A. Papapetropoulos, "International union of basic and clinical pharmacology. CII: pharmacological modulation of H₂S levels: H₂S donors and H₂S biosynthesis inhibitors," *Pharmacological Reviews*, vol. 69, no. 4, pp. 497–564, 2017.
- [53] Y. Liu, L.-P. Zou, J.-B. Du, and V. Wong, "Electro-acupuncture protects against hypoxic-ischemic brain-damaged immature rat via hydrogen sulfide as a possible mediator," *Neuroscience Letters*, vol. 485, no. 1, pp. 74–78, 2010.
- [54] H. Shan, J. Qiu, P. Chang et al., "Exogenous hydrogen sulfide offers neuroprotection on intracerebral hemorrhage injury through modulating endogenous H(2)S metabolism in mice," *Frontiers in Cellular Neuroscience*, vol. 13, p. 349, 2019.
- [55] W. Niu, J. Wang, J. Qian et al., "Allosteric control of human cystathionine β -synthase activity by a redox active disulfide bond," *Journal of Biological Chemistry*, vol. 293, no. 7, pp. 2523–2533, 2018.
- [56] S. Pathirathna, D. F. Covey, S. M. Todorovic et al., "Differential effects of endogenous cysteine analogs on peripheral thermal nociception in intact rats," *Pain*, vol. 125, no. 1–2, pp. 53–64, 2006.
- [57] S.-P. Chen, Y. Kan, J.-L. Zhang et al., "Involvement of hippocampal acetylcholinergic receptors in electroacupuncture analgesia in neuropathic pain rats," *Behavioral and Brain Functions*, vol. 12, no. 1, p. 13, 2016.
- [58] J. Wang, Y. Gao, S. Chen et al., "The effect of repeated electroacupuncture analgesia on neurotrophic and cytokine factors in neuropathic pain rats," *Evidence-Based Complementary and Alternative Medicine*, vol. 2016, Article ID 8403064, 11 pages, 2016.
- [59] J.-S. Han, X.-H. Chen, Y. Yuan et al., "Transcutaneous electrical nerve stimulation for treatment of spinal spasticity," *Chinese Medical Journal*, vol. 107, no. 1, pp. 6–11, 1994.
- [60] J.-S. Han, "Acupuncture: neuropeptide release produced by electrical stimulation of different frequencies," *Trends in Neurosciences*, vol. 26, no. 1, pp. 17–22, 2003.
- [61] T. S. Hahm, "The effect of 2 Hz and 100 Hz electrical stimulation of acupoint on ankle sprain in rats," *Journal of Korean Medical Science*, vol. 22, no. 2, pp. 347–351, 2007.
- [62] R.-X. Zhang, L. Lao, X. Wang et al., "Electroacupuncture attenuates inflammation in a rat model," *The Journal of Alternative and Complementary Medicine*, vol. 11, no. 1, pp. 135–142, 2005.
- [63] H. J. Moon, B.-S. Lim, D.-I. Lee et al., "Effects of electroacupuncture on oxaliplatin-induced neuropathic cold hypersensitivity in rats," *The Journal of Physiological Sciences*, vol. 64, no. 2, pp. 151–156, 2014.
- [64] K. Wang, R. Zhang, F. He et al., "Electroacupuncture frequency-related transcriptional response in rat arcuate nucleus revealed region-distinctive changes in response to low- and high-frequency electroacupuncture," *Journal of Neuroscience Research*, vol. 90, no. 7, pp. 1464–1473, 2012.

- [65] P.-R. Lv, Y.-S. Su, W. He et al., “Electroacupuncture alleviated referral hindpaw hyperalgesia via suppressing spinal long-term potentiation (LTP) in tnbs-induced colitis rats,” *Neural Plasticity*, vol. 2019, pp. 1–11, 2019.
- [66] K. A. Sluka, T. L. Lisi, and K. N. Westlund, “Increased release of serotonin in the spinal cord during low, but not high, frequency transcutaneous electric nerve stimulation in rats with joint inflammation,” *Archives of Physical Medicine and Rehabilitation*, vol. 87, no. 8, pp. 1137–1140, 2006.
- [67] Y. Liu, Z. Li, X. Shi et al., “Neuroprotection of up-regulated carbon monoxide by electrical acupuncture on perinatal hypoxic-ischemic brain damage in rats,” *Neurochemical Research*, vol. 39, no. 9, pp. 1724–1732, 2014.
- [68] S. W. Lee, Y.-S. Hu, L.-F. Hu et al., “Hydrogen sulphide regulates calcium homeostasis in microglial cells,” *Glia*, vol. 54, no. 2, pp. 116–124, 2005.
- [69] J. Gao, L. Zhang, Y. Wang et al., “Antiarrhythmic effect of acupuncture pretreatment in rats subjected to simulative global ischemia and reperfusion—involvement of adenylate cyclase, protein kinase A, and L-type Ca²⁺ channel,” *The Journal of Physiological Sciences*, vol. 58, no. 6, pp. 389–396, 2008.
- [70] Q. Wang, H. Zhu, K. Zou et al., “Sensitization of P2X₃ receptors by cystathionine β -synthetase mediates persistent pain hypersensitivity in a rat model of lumbar disc herniation,” *Molecular Pain*, vol. 11, p. 15, 2015.
- [71] T. Chen, Y. Xiong, M. Long et al., “Electro-acupuncture pretreatment at zusanli (ST36) acupoint attenuates lipopolysaccharide-induced inflammation in rats by inhibiting Ca²⁺ influx associated with cannabinoid CB₂ receptors,” *Inflammation*, vol. 42, no. 1, pp. 211–220, 2019.