

Review Article

Medicinal Plants Used in the Management of Sexual Dysfunction, Infertility and Improving Virility in the East African Community: A Systematic Review

Christine Kyarimpa ¹, Christine Betty Nagawa ², Timothy Omara ³, Silver Odongo ⁴, Patrick Ssebugere ⁴, Solomon Omwoma Lugasi ⁵ and Ivan Gumula¹

¹Department of Chemistry, Faculty of Science, Kyambogo University, P.O. Box 1, Kampala, Uganda

²Department of Forestry, Biodiversity and Tourism, College of Agricultural and Environmental Sciences, Makerere University, P.O. Box 7062, Kampala, Uganda

³Chemistry Division (Food Safety Laboratories), Testing Department, Standards Directorate, Uganda National Bureau of Standards, P.O. Box 6329, Kampala, Uganda

⁴Department of Chemistry, College of Natural Sciences, Makerere University, P.O. Box 7062, Kampala, Uganda

⁵Department of Physical Sciences, Jaramogi Oginga Odinga University of Science and Technology, P.O. Box 210, Bondo 40601, Kenya

Correspondence should be addressed to Timothy Omara; prof.timo2018@mu.ac.ke

Received 15 February 2023; Revised 13 July 2023; Accepted 27 July 2023; Published 12 August 2023

Academic Editor: Chunpeng (Craig) Wan

Copyright © 2023 Christine Kyarimpa et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Sexual disorders such as erectile dysfunction (ED), sterility, and sexual inappetence represent some of the complex reproductive challenges that require addressing the underlying causes. The aim of this paper was to systematically synthesize literature on the ethnobotany, phytochemistry, bioactivities, and safety of plants used as remedies for managing sexual dysfunction and infertility, and improving fertility and virility in the EAC. Through an extensive review conducted in multidisciplinary electronic databases, 171 plant species were identified to have been reported for the management of sexual inappetence (i.e., used as aphrodisiacs, 39.4%), ED (35.9%), infertility (18.7%), and increasing fertility (6.0%). The most used plants are *Mondia whitei*, *Acalypha villicaulis*, *Combretum illairii*, *Erythrina abyssinica*, *Pappea capensis*, *Rhus vulgaris*, and *Warburgia ugandensis* while roots (44.9%), leaves (21.8%), stem and root barks (16.7%) of shrubs (35%), trees (31%), herbs (26%), and climbers (8%) are the preferred organs for making decoctions (69%). The research strides to date indicate that *Citropsis articulata*, *Cola acuminata*, *Ekebergia capensis*, *Plumbago zeylanica*, *Tarenna graveolens*, *Urtica massaica*, and *Zingiber officinale* have been assessed for their bioactivity. The majority (71.4%) of the plants either increased testosterone levels and mounting frequency or elicited prosexual stimulatory effects in male rats. More studies investigating the relevant pharmacological activities (aphrodisiac, fertility, and phosphodiesterase-5 inhibitory activities), safety aspects, responsible compounds, and clinical studies are warranted to establish the pharmacological potential of the unstudied species and elucidate the mechanism of action of the bioactive compounds.

1. Introduction

One of the universal interests enshrined in sustainable development goal (SDG) 3 is good health and well-being. It is linked to and affects other global goals such as SDG 1 (poverty reduction), SDG 2 (end poverty), and SDG 4 (quality and equitable education) [1]. Critical analysis of the

global disease burden shows that one-third of the total world population has more than five ailments [2]. Accordingly, three in five of the global deaths are ascribed to at least one of the four main noncommunicable diseases (NCDs), namely, cancer, diabetes, cardiovascular, and chronic lung diseases [3–5]. Most global mortalities (up to 71%) are due to NCDs [6], and 77% of these occur in low- and middle-income

countries due to limited access to medical services and poverty [7]. Whereas the global focus is on the major NCDs, conditions such as sexual dysfunction, infertility, and anaphrodisia (sexual inappetence) represent some of the complex health challenges.

Sexual dysfunction refers to the inability to achieve a normal sexual intercourse. It includes orgasmic disorder, retrograded, retarded, premature ejaculation, and erectile dysfunction [8]. Male erectile dysfunction (ED) or impotence is the inability to achieve or maintain an erection sufficient for satisfactory sexual performance and vaginal intercourse, typically for a period of more than six months [9]. Though to different degrees, ED affects more than 52% of men in the age bracket of 40 and 70 years. Erectile dysfunction is linked with conditions such as diabetes, sedentary lifestyle, hypertension, obesity, hypercholesterolemia, and smoking [10–12].

On the other hand, infertility is a medical condition characterized by failure to establish a clinical pregnancy after one year of regular and unprotected sex [13]. Infertility affects more than 48 million couples worldwide. It can be from either one or both partners, but 50% of all cases are due to male infertility [14]. In women, it may be due to endometriosis (premature ovarian failure) and uterine disorders such as fibroids or thyroid diseases. In males, infertility is associated with defective sperm function, azoospermia, low sperm counts, varicocele, undescended testes, testicular cancer, and low testosterone levels [13, 14]. Other risk factors for infertility include diabetes, sexually transmitted diseases, stress, obesity, drug abuse, age, exposure to environmental toxins, radiotherapy, and other cancer treatments [15, 16].

Sexual inappetence is a common reproductive challenge that accompanies or is a direct consequence of ED and infertility [17, 18]. Sexual inappetence (anaphrodisia or lack of desire/libido) is one of the most common sexual dysfunctions of women. Together, ED, infertility, and sexual inappetence are among the relatively common fecundity challenges that affect couples medically, sexually, and psychologically [17, 18]. With medical advancements in assisted reproduction technologies, the use of synthetic agents such as phosphodiesterase type 5 inhibitors (in intracavernosal injection therapy for ED) and stem cell therapy (for infertility) has been encouraged [13]. However, limited access to medical services, long-term treatment tenure, and side effects of injectable fertility drugs have limited their acceptability among the general population [19–21].

For indigenous communities in developing countries, the use of natural products for prevention and the management of reproductive diseases and conditions are common. The East African Community (EAC) is one of the regions with distinguished ethnomedicinal knowledge and use of natural products [22–25]. The high reliance of these communities on herbal medicine is explained by the exceptionally rich cultural heritage, acceptance, availability, and perceived efficacy [26–28]. In this context, traditional medicine practitioners correlate sicknesses and other medical conditions with their possible causes [25]. For this

reason, herbal medications and posology are prescribed based on the supposed cause of the diseases. Critical cases, or those due to supernatural forces, are managed through diviners' interventions [26, 29]. Illnesses are thought to be induced by external polluting influences (e.g., consumption of tabooed foods [30], breaching of taboos, witchcraft-related rites, fetishes or social rules, and use of objects planted by ill people) that interfere with body physiology [26, 31–33]. Therefore, traditional management of diseases involves health practices, knowledge, and beliefs that utilize plants and animal- and mineral-based remedies, dispensing of ritually protective herbal medicines or performing rituals for placating spirits [26, 33]. These perceptions are similar to traditional medicine concepts in other parts of Africa [34].

In the EAC, chronic poverty and resource-constrained healthcare systems are common, and the use of herbal remedies for the treatment of sexual dysfunction (ED) and infertility, and enhancing fertility and virility has been sporadically mentioned in ethnobotanical studies. However, no study has systematically collated literature on these medicinal plants with in-depth description and analysis of their claimed efficacy, phytochemistry, and safety. The aim of this paper was, therefore, to systematically synthesize literature on ethnobotany, phytochemistry, bioactivities, and the safety profile of plants used as remedies for managing sexual dysfunction and infertility, and improving fertility and virility in the EAC. As part of an ongoing project, we aimed at identifying highly cited but unstudied species that could be assessed for their aphrodisiac, fertility and phosphodiesterase-5 inhibitory activities, bioactive phytochemicals, and toxicity profiles. This could open lead to the discovery of molecules that can be used in modern medicine.

2. Methods

2.1. Study Design, Literature Sources, and Systematic Search Procedures. The Preferred Reporting Items for the Systematic Reviews and Meta-Analyses (PRISMA) 2020 guidelines [35] were followed (Supplementary file 1). The protocol used was registered with the International Prospective Register of Systematic Reviews (PROSPERO) with registration number CRD42022373152 (https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42022373152). Nine multidisciplinary electronic databases (Scopus, Web of Science, PubMed, Science Direct, Google Scholar, Wiley Online Library, Taylor and Francis Online, Springer Link, and Scientific Electronic Library Online) and regional university repositories were searched to gather relevant records on ethnobotany, phytochemistry, biological activities, and toxicity of medicinal plants exploited for the management of sexual dysfunction and infertility, and improving fertility and virility in the EAC. The dates on which we last consulted the databases were 7th January 2023, 31st December 2022, 20th November 2022, 20th January 2023, 4th January 2023, 17th January 2023, 11th November 2022, 10th January 2023, 24th November 2022, 2nd December 2022, and 2nd January 2023, respectively.

The EAC was considered as the region encompassing Uganda, Kenya, Tanzania, Rwanda, Burundi, South Sudan, and Democratic Republic of Congo (DRC) from April 2022 [36]. The searches were performed in parallel using search strings specified for a comprehensive search that covered all fields in records but broadened the scope in PubMed advanced search. Within each axis, keywords were combined with the “OR” operator in the Boolean operator and then linked the two axes’ search techniques to the “AND” operator. The keywords used were “plant” “erectile dysfunction” “aphrodisiac” “infertility” OR “fertility” “virility” AND “Uganda” “Kenya” “Tanzania” “Rwanda” “Burundi” “South Sudan” “Democratic Republic of Congo.” For example, in Scopus, the search string used was ALL (plants, AND erectile AND dysfunction, AND aphrodisiac, AND uganda) AND (LIMIT-TO (AFFILCOUNTRY, “Uganda”)) OR ALL (plants, AND erectile AND dysfunction, AND aphrodisiac, AND Kenya) AND (LIMIT-TO (AFFILCOUNTRY, “Kenya”)) OR ALL (plants, AND erectile AND dysfunction, AND aphrodisiac, AND rwanda) AND (LIMIT-TO (AFFILCOUNTRY, “Rwanda”)) OR ALL (plants, AND erectile AND dysfunction, AND aphrodisiac, AND burundi) AND (LIMIT-TO (AFFILCOUNTRY, “Burundi”)) OR ALL (plants, AND erectile AND dysfunction, AND aphrodisiac, AND south Sudan) AND (LIMIT-TO (AFFILCOUNTRY, “South Sudan”)) OR ALL (plants, AND erectile AND dysfunction, AND aphrodisiac, AND Democratic Republic of Congo) AND (LIMIT-TO (AFFILCOUNTRY, “Democratic Republic of Congo”)) OR ALL (plants, AND erectile AND dysfunction, AND aphrodisiac, AND Tanzania) AND (LIMIT-TO (AFFILCOUNTRY, “Tanzania”)) OR ALL (plant, AND infertility AND fertility AND uganda) AND (LIMIT-TO (AFFILCOUNTRY, “Uganda”)) OR ALL (plant, AND infertility AND fertility AND Kenya) AND (LIMIT-TO (AFFILCOUNTRY, “Kenya”)) OR ALL (plant, AND infertility AND fertility AND tanzania) AND (LIMIT-TO (AFFILCOUNTRY, “Tanzania”)) OR ALL (plant, AND infertility AND fertility AND rwanda) AND (LIMIT-TO (AFFILCOUNTRY, “Rwanda”)) OR ALL (plant, AND infertility AND fertility AND burundi) AND (LIMIT-TO (AFFILCOUNTRY, “Burundi”)) OR ALL (plant, AND infertility AND fertility AND south Sudan) AND (LIMIT-TO (AFFILCOUNTRY, “South Sudan”)) OR ALL (plant, AND infertility AND fertility AND democratic republic of congo) AND (LIMIT-TO (AFFILCOUNTRY, “Democratic Republic of Congo”)).

In addition, reference lists of the retrieved studies were also manually searched to access additional articles which were screened for their eligibility for inclusion in the study. The literature search was performed between 1st June 2022 and 20th January 2023.

2.2. Study Selection. All search results were imported into EndNote® X9 (Thomson Reuters, Philadelphia, PA, USA), and duplicate reports were removed. The screening was done according to the title and abstract of the articles. This was

conducted independently by 4 authors (CK, CBN, TO, and SO). Two independent reviewers (TO and SO) screened the articles against inclusion criteria, and possible contradictions during article selection and/or extraction were obviated through discussions and consensus.

2.3. Inclusion and Exclusion Criteria. To refrain the authors from bias, (1) only full-text articles or reports published in or translated into English and French; (2) cross-sectional original papers or reports on the ethnobotany, phytochemistry, bioactivities, and clinical trials of plants used in the management of sexual dysfunction and infertility, and improving fertility and virility in EAC; and (3) reports published online until 20th January 2023 were included. Excluded studies were those that (1) provided no data; (2) were neither from EAC nor full-text articles; (3) reported on the use of plants for managing conditions such as menorrhagia, blocked fallopian tubes, inducing twin birth or birth to a particular sex of children; (4) narrative and systematic reviews, or reports not based on original data (expert opinions, editorials, and perspective papers).

2.4. Risk of Bias Assessment. Quality of the considered reports (risk of bias) was established following the Joanna Briggs Institute quality assessment tool [37]. Two authors (TO and SO) independently assessed the quality of the included studies. Variations in the final risk of bias assessment among them were declared by discussing the prespecified criteria. The evaluation tool consisted of seven parameters: (1) appropriate sampling design; (2) correct sampling technique; (3) acceptable sample size; (4) adequate study subject and location explanation; (5) appropriate data investigation; (6) use of valid methods for identification of plants and the conditions that they treat; and (7) use of appropriate statistical/ethnobotanical analysis indices. Because most studies met parameters 5 to 7 that were similar across them, we relied on parameters 1 to 4 to ascertain the risk of bias status. A study that did not meet each parameter was scored as 1 if not 0. The risks for biases were classified as either low (total score, 0-1), moderate (total score, 2), or high (total score, 3-4) [38].

2.5. Data Extraction. Data were collated in a predesigned Microsoft Excel 2019 standardized sheet. Information on the reported medicinal plants, such as botanical names (and synonyms), plant family, traditional name(s), growth habit, part(s), and their uses (conditions treated), mode of preparation and administration, isolated pure compounds, and relevant efficacy reports were extracted. For each dataset, the first author’s last name, year of publication, and country were also extracted. Missing information in some reports such as local names, growth forms, and misspelled botanical names was checked from Google and botanical databases (WFO Plant List, International Plant Names Index, and Tropicos).

2.6. Data Analysis. Descriptive statistical methods were used to analyze the collected data. Results were expressed as ranges, percentages, and frequencies and presented as tables and charts. These analyses were performed in Microsoft Excel 2019 for Windows (Microsoft Corporation, Washington, DC, USA).

3. Results and Discussion

3.1. Description of Included Studies. Our systematic search (Figure 1) retrieved no previous review on the subject in East Africa. From databases, registers, and other websites, the search returned 396 unique reports published between 1962 and 2022. The highest number of reports were from Scopus ($n=172$) and Google Scholar ($n=108$), followed by Web of Science ($n=44$), Science Direct ($n=42$), Wiley Online Library ($n=11$), Taylor & Francis Online ($n=7$), Springer Link ($n=4$), PubMed ($n=3$), regional university repositories ($n=3$), and SciELO ($n=2$). Of these, duplicates ($n=47$) were removed, and 349 unique articles were screened. A total of 235 articles were excluded after reading their titles and abstracts, while 47 others were excluded because they were not from any country within the EAC. Therefore, 67 records were assessed for their eligibility and inclusion in the study. Based on the inclusion and exclusion criteria, some full-text articles were excluded with reasons, namely, (i) articles not in English or French ($n=11$), (ii) review articles ($n=9$), and (iii) those that did not provide any data ($n=8$). A manual search resulted in 7 eligible articles. Thus, data were extracted from a total of 46 articles in this systematic review. Regarding the assessment of the risk of bias among studies, most reports were judged as having a low (47.8%) or moderate risk of bias (34.8%) (Supplementary file 2).

3.2. Inventory of Medicinal Plants Reported. This review identified 46 reports on plants used in the management and treatment of ED and infertility, and increasing fertility and sexual appetite (virility) in the EAC. Some of the sexual dysfunctions captured from herbalists include erectile disorders, pain during penetration, premature ejaculation, lack of sexual arousal, and short-lasting erections (among men) and lack of orgasm, dyspareunia, lack of sexual arousal, atrophic vaginitis, and short orgasms among women [39].

In total, 171 plant species from 59 botanical families have been reported for the management of sexual inappetence, i.e., used as aphrodisiacs (39.4%), ED (35.9%), infertility (18.7%), and increasing fertility (6.0%) (Table 1). The highest number of plants cited was from Kenya (96), followed by Uganda (66), Tanzania (24), Rwanda (1), and DRC (1). Burundi and South Sudan had no reports on plants in the category under scrutiny. It is not surprising that Kenya ranked the highest, as it is known to have diversified flora with over 7,000 plant species [23, 40]. This is also supported by the fact that most of the ethnobotanical reports reviewed ($n=25$) were from Kenya as compared to Uganda ($n=18$), Tanzania ($n=7$), Rwanda ($n=1$), and DRC ($n=1$).

Analysis of transregional distribution of the plants revealed that Uganda and Kenya shared 8 species and Tanzania and Kenya shared 6 species while Kenya and Rwanda shared one species (*Tagetes minuta* L.). Only one plant (*Pachycarpus robusta*) was cited to be used in Uganda, Kenya, and Tanzania [41]. The rest of the countries did not share any plant. Such marked divergence in the use of plants across the region could be due to their preference which is related to specific cultural beliefs and traditions or centred around human relationships [42–44].

The majority of the plants retrieved in this study were from families: Fabaceae (16.9%, 29 species), Euphorbiaceae (7.0%, 12 species), Asteraceae (5.8%, 10 species), Apocynaceae, Rubiaceae (5.3%, 9 species each), and Capparaceae (4.7%, 8 species) (Figure 2). Species from these botanical families have been reported to have aphrodisiac and fertility potential in ethnobotanical surveys from Ethiopia [45], Southern Africa [46], Iran [47], and India [48]. The dominance of families, especially Fabaceae and Asteraceae, is due to the extensive range of their distribution across global biomes [49]. Moreover, they contain phytochemicals such as phenolics, tannins, and alkaloids which are known to have therapeutic effects [50, 51].

At the genus level, the most represented genera were *Acacia* (6 species), *Combretum* (5 species), *Cassia* and *Tragia* (3 species each), *Abrus*, *Allium*, *Boscia*, *Cadaba*, *Cleome*, *Croton*, *Impatiens*, *Maytenus*, *Sonchus*, *Uvaria*, *Vachellia*, and *Vernonia* (2 species each). The commonly mentioned plants were *Mondia whitei* (12 times), *Warburgia ugandensis* (4 times), *Acalypha villicaulis*, *Combretum illairii*, *Erythrina abyssinica*, *Pappea capensis*, and *Rhus vulgaris* (3 times each). Some of the plants listed such as *Abrus precatorius*, *Allium sativum*, *Cola acuminata*, *Combretum hereroense*, *Mondia whitei*, *Plumbago zeylanica*, *Ricinus communis*, and *Syzygium guineense* are traditionally used for treating infertility and ED in South Africa [52], Ghana [53], Cameroon, Guinea, Gabon [54], Iran [47], Benin [55], and Ethiopia [45]. It is worth mentioning that organs of some of the highly cited species such as *Abrus precatorius* and *Erythrina abyssinica* are used in Uganda for rituals and ceremonies of love, weddings, and childbirth [56].

In regards to the treatment of infertility, most plant species recorded were indicated to be used for the treatment of female infertility (Table 2). The most cited species were *Erythrina abyssinica* and *Combretum illairii* (3 times each). Interestingly, some species (*Cadaba glandulosa*, *Cadaba farinose*, *Combretum illairii*, *Hoslundia opposita*, and *Allophylus pervilleria*) were shown to be used for the treatment of both female and male infertility, which could make them good candidates for further studies of their biological activities.

3.3. Growth Habit, Organs Used, Dosage Forms, and Posology of the Herbal Remedies. The plants occurred as shrubs (35%), trees (31%), herbs (26%), and climbers (8%) (Figure 3). Figure 4 illustrates which plant organs are widely used in




FIGURE 1: PRISMA flow diagram showing the retrieval and exclusion steps of the systematic review adapted from Page et al. [35].

preparation of the herbal remedies, that is, roots (44.9%), leaves (21.8%), and stem and root barks (16.7%). The frequent use of roots is unsustainable but may be linked to the fact that the conditions treated are internal to the body (are hidden), just as root structures are hidden in the ground. On the other hand, the relatively frequent use of leaves could be related to their availability and the fact that they are the photosynthetic sheet of plants that accumulate therapeutic phytochemicals [57].

This review noted a tendency of including more than one plant part and adjuvants in herbal remedies. For multiple plant parts, a total of 13 species were encountered to be used in combination with others. For example, in Kenya, decoction of *Uvaria leptoclodon*, *Boscia coriacea*, and *Combretum hereroense* roots is used for treating ED. For infertility in women, the roots are used with *Croton dichagamus* roots [58, 59]. Similarly, the decoction of *Markhamia zanzibarica* roots mixed with *Uvaria acuminata* roots is administered as an aphrodisiac. For infertility in

women, it is used with *Salvadora persica* and *Uvaria acuminata* roots [58, 59]. A striking example of using adjuvants is from Tanzania where roots of *Polygala aphrodisiaca* are cooked with a young cock while *Duosperma kilimandscharicum* leaf and root decoction are taken with goat blood or goat meat soup as an aphrodisiac [41]. The use of cow and goat milk for preparation of *Morus mesozygia* roots as an aphrodisiac was also documented in Kenya [60]. In Uganda, *Acanthus pubescens* leaves are taken in *tonto*, a traditional beer prepared from *Musa × paradisiaca* L. var. *sapientum* fruits [61]. The use of more than one plant organ and adjuvants as witnessed in this review are tailored to various reasons. For instance, it may be an obvious way of masking the toxicity of herbal remedies or hiding the secrecy of the formulations [62, 63].

The commonest method of preparing herbal remedies is decoctions (69%). This could be because decoction procedures allow for better extraction of the bioactive phytochemicals in plant matrices [64]. However, the plant organs

TABLE 1: Plants used in the management of sexual disorders, infertility, and improving sexual virility and fertility in the East African Community.

Plant family	Botanical name	Local name	Part used	Habit	Preparation mode	Administration	Use	Country	Reference
Acanthaceae	<i>Acanthus pubescens</i> (T. Thoms.) Engl.	Amatovu (Luganda)	Leaves	Shrub	Decoction	Oral in a local brew (<i>tonto</i>)	Aphrodisiac	Uganda	[61]
Acanthaceae	<i>Duosperma kilimandscharicum</i> (C. B. Clarke) Dayton	Not reported	Leaves, roots	Shrub	Decoction. Taken with goat blood or extract from goat meat	Oral	Aphrodisiac	Tanzania	[41]
Alliaceae	<i>Allium cepa</i> L	Katunguru (Runyankore)	Bulb, leaves, root tuber	Herb	Used directly, decoction	Oral in water, and in food or just chewed	ED	Uganda	[39, 65]
Alliaceae	<i>Allium sativum</i> L	Tungurusumu (Rukonjo)	Stem bulb, leaves, roots tuber	Herb	Used directly, decoction	Oral in water, and in food	ED	Uganda	[65]
Aloaceae	<i>Aloe volkensii</i> Engl.	Hargeis, D'aar (Orma)	Leaves	Herb	Decoction	Used to wash genital area thrice daily	Infertility	Kenya	[58]
Anacardiaceae	<i>Mangifera indica</i> L	Muyembe (Luganda)	Bark	Tree	Decoction	Oral	Infertility (women)	Uganda	[22]
Anacardiaceae	<i>Ozoroa insignis</i> ssp. <i>reticulata</i> (Baker f.) J. B. Gillett	Not reported	Roots	Tree	Decoction	Oral	Aphrodisiac	Tanzania	[41]
Anacardiaceae	<i>Rhus vulgaris</i> Meikle	Mukanja (Runyankore), Mukanza (Rukonjo), Musatsa (Wanga)	Bark, roots, leaves, whole plant	Shrub	Used directly, decoction	Oral, chewed	ED, aphrodisiac	Uganda, Kenya	[41, 60, 65]
Annonaceae	<i>Ovariendron anisatum</i>	Ndonga (Embu)	Whole plant	Herb	Decoction	Oral	ED, infertility	Kenya	[61, 62]
Annonaceae	<i>Uvaria acuminata</i> Oliv	Mundagoni, murori (Pokomo)	Roots	Shrub	Decoction. Used with <i>Markhamia zanzibarica</i>	Oral, a glass daily for 5 days	Aphrodisiac	Kenya	[59]
Annonaceae	<i>Uvaria leptoclada</i>	Sholole (Orma)	Roots	Shrub	Decoction. Used with <i>Boscia coriacea</i> and <i>Combretum hereroense</i> . For infertility, use with <i>Croton dichagamus</i>	Oral, half a glass daily for 5 days. For infertility, half glass thrice daily for 3 days	ED, infertility (women)	Kenya	[58, 59]
Apocynaceae	<i>Carissa spinarum</i> L. (Synonym: <i>Carissa edulis</i> Forsk. (Vahl))	Leketetwo (Marakwet), Logetetwa (Pokot), Omukuyomonza	Roots, bark	Shrub	Decoction with <i>Elaeodendron buchannanii</i> bark or powder mixed with <i>Tragia furialis</i>	Oral, taken as tea	Aphrodisiac	Tanzania, Kenya	[64]
Apocynaceae	<i>Acokanthera schimperi</i> (A.D.C) Schweinf	Not reported	Roots	Tree	Infusion of powder	Oral	Aphrodisiac	Kenya	[65]

TABLE 1: Continued.

Plant family	Botanical name	Local name	Part used	Habit	Preparation mode	Administration	Use	Country	Reference
Apocynaceae	<i>Landolphia swynnertonii</i>	Mokokwet (Marakwet)	Roots	Shrub	Decoction	Oral	Increasing fertility (women)	Kenya	[66]
Apocynaceae	<i>Mondia whitei</i> (Hook. f.) Skeels	Omulondo (Luganda), Omurondo (Runyankore), Mukombelo (Luhya)	Roots/root bark	Vine	Used directly (chew when raw or dry), decoction	Oral in tea and in food	ED, aphrodisiac, fertility enhancer	Uganda, Kenya	[22, 23, 27, 39, 41, 65, 67–72]
Apocynaceae	<i>Cryptolepis obtusa</i> N. E. Br	Not reported	Roots	Shrub	Decoction (Tanzania), used directly	Drunk, chewed	Aphrodisiac	Kenya, Tanzania	[73]
Apocynaceae	<i>Dregea rubicunda</i> K. Schum	Not reported	Roots	Climber	Used directly	Fresh root chewed	Aphrodisiac	Tanzania	[60]
Apocynaceae	<i>Pachycarpus robusta</i> (A. Rich.) Bullock	Not reported	Roots	Herb	Not specified	Not specified	Aphrodisiac	East Africa	[41]
Apocynaceae	<i>Parquetina nigrescens</i> Afzel	Not reported	Roots	Climber	Decoction	Oral, once in the evening	Aphrodisiac	Kenya	[60]
Apocynaceae	<i>Periploca linearifolia</i> Dill. & Rich	Sinendet (Nandi)	Roots/milky latex	Climber/liana	Decoction	Oral	Fertility	Kenya	[75]
Areaceae	<i>Phoenix reclinata</i> Jacq	Akakindo, Mukindo (Runyankore), Mukindu (Pokomo)	Roots, leaves	Shrub	Decoction	Oral, a glass daily for 3 days	ED, aphrodisiac	Uganda, Kenya	[59, 65]
Asparagaceae	<i>Chlorophytum comosum</i> (Thunb.) Jacques	Nalwebe (Lusoga)	Tuber	Herb	Not specified	Not specified	Infertility	Uganda	[31]
Asphodelaceae	<i>Aspidoglossum biflorum</i> E. Mey	Drege (Kiswahili)	Roots	Herb	Not specified	Not specified	Aphrodisiac	Tanzania	[41]
Asteraceae	<i>Bidens pilosa</i> L.	Mucege, Enyabarashana (Runyankore)	Shoot, flowers	Herb	Decoction of young flowers as tea	Oral, 500 ml daily for 2 days for ED	ED, increase fertility	Kenya, Uganda	[27, 76]
Asteraceae	<i>Lainaea cornuta</i> (Hochst. Ex Oliv. & Hiern) C. Jeffrey	Uthuunga (Kikamba)	Leaves, stems	Herb	Infusion	Oral	Infertility (women)	Kenya	[77]
Asteraceae	<i>Lactuca inermis</i> Forsk. (L. Capensis Thurnb.)	Not reported	Roots	Herb	Not specified	Not specified	Aphrodisiac	Uganda	[56]
Asteraceae	<i>Microglossa pyrifolia</i> (Lam.) Kuntze	Omube/Mkuraiju	Leaves	Herb	Not specified	Not specified	Aphrodisiac	Tanzania	[41]
Asteraceae	<i>Sonchus asper</i>	Iviuvu (Embu)	Whole plant	Herb	Decoction	Oral	ED	Kenya	[61]
Asteraceae	<i>Psiadia punctulata</i> (DC.) Vatke	Konocho (Marakwet), Shiro (Luhya)	Roots	Herb	Decoction	Oral	Aphrodisiac, sterility (men)	Kenya	[23]
Asteraceae	<i>Sonchus schweinfurthii</i> Oliv. & Hiern	Sungasunga	Roots	Herb	Decoction	Oral	Aphrodisiac	Tanzania	[78]

TABLE 1: Continued.

Plant family	Botanical name	Local name	Part used	Habit	Preparation mode	Administration	Use	Country	Reference
Asteraceae	<i>Tagetes minuta</i> L	Nyiramunukanabi (Kinyarwanda)	Shoot	Herb	Decoction	Oral	Increase fertility, ED	Kenya, Rwanda	[76, 79]
Asteraceae	<i>Vernonia cinerea</i> (L.) Less (or <i>Cyanthillium cinereum</i> (L.) H. Rob.)	Kayayana (Luganda)	Leaves, roots	Shrub	Used directly, decoction	Oral	ED	Uganda	[65]
Asteraceae	<i>Vernonia lasiopis</i> O Hoffm	Shiroho	Roots	Shrub	Decoction/infusion	Oral, infusion drunk twice a day	Aphrodisiac	Kenya	[60]
Balanitaceae	<i>Balanites aegyptiaca</i> Del	Ng'oswet (Marakwet)	Roots	Tree	Decoction	Oral	Increasing fertility (women)	Kenya	[66]
Balsaminaceae	<i>Impatiens</i> species	Entungwa baishajja (Runyankore)	Whole plant	Herb	Used directly (chew), decoction	Oral	ED	Uganda	[39, 65]
Balsaminaceae	<i>Impatiens tinctoria</i> A. Rich	Chemakalbayi	Roots	Herb	Decoction	Oral	Fertility	Kenya	[80]
Bignoniaceae	<i>Markhamia zanzibarica</i>	Mubwoka (Pokomo)	Roots	Tree	Decoction. Used with <i>U. acuminata</i> roots. For infertility, it is used with <i>Salvadora persica</i> and <i>Uvaria acuminata</i>	Oral, a glass (or twice for infertility) daily for 5 days	Aphrodisiac (men), infertility (women)	Kenya	[58, 59]
Bignoniaceae	<i>Spathodea campanulata</i> Buch. -Harm. ex DC	Kifabakazi	Bark	Tree	Decoction	Oral	Infertility (men)	Uganda	[22]
Boraginaceae	<i>Ehretia cymosa</i> Thonn	Morori (Marakwet), Ponponat (Pokot), Shekhutu (Luhyia)	Roots, leaves	Shrub	Decoction	Oral	Aphrodisiac	Kenya	[23, 75]
Boraginaceae	<i>Kigelia africana</i> Lam	Sausage tree (English)	Fruits, seeds	Tree	Decoction	Oral	Aphrodisiac	Tanzania	[81]
Canellaceae	<i>Warburgia ugandensis</i> Sprague	Mwiba (Runyaruguru), Mugeta (Embu)	Bark, leaves, roots	Tree	Decoction	Oral in tea, 1 spoonful thrice daily or in porridge; 250 ml drunk	ED	Uganda, Kenya	[27, 61, 65, 71]
Cannabaceae	<i>Cannabis sativa</i> L	Njaji, olusambya (Luganda), Njaga (Runyankore)	Leaves, roots	Shrub	Used directly (chew), decoction	Oral, inhaling fumes (smoking)	ED	Uganda	[65, 72]
Capparaceae	<i>Boscia coriacea</i> Pax	Kalkacha (Orma)	Roots	Shrub	Decoction with <i>U. leptoclados</i> and <i>C. hereroense</i> roots	Oral, half a glass daily for 5 days	ED	Kenya	[59]
Capparaceae	<i>Boscia solicifolia</i> Oliv	Chelel (Marakwet)	Roots	Tree	Decoction	Oral	Increases fertility (male and female)	Kenya	[66]

TABLE 1: Continued.

Plant family	Botanical name	Local name	Part used	Habit	Preparation mode	Administration	Use	Country	Reference
Capparaceae	<i>Cadaba glandulosa</i> Forssk	Alakal (Orma)	Roots	Shrub	Decoction	Oral, half a glass daily for 5 days	Infertility (men and women)	Kenya	[58, 59]
Capparaceae	<i>Cadaba farinose</i>	Kumis (Orma)	Roots	Shrub	Decoction	Oral, half a glass daily for 3 days	Infertility (men and women)	Kenya	[58, 59]
Capparaceae	<i>Capparis tomentosa</i> Lam	Kumbolwop kimaget (Marakwet)	Roots	Climber	Decoction	Oral	Increasing fertility	Kenya	[66]
Capparaceae	<i>Cleome gynandra</i> L	Esoby/Amarera (Runkonjo), Eshogi (Runyankore)	Leaves, roots, flowers	Herb	Used directly (chew), decoction	Oral or as food	ED, aphrodisiac	Uganda	[65, 82]
Capparaceae	<i>Cleome usambarica</i> Pax	Not reported	Roots	Herb	Infusion. Mixed with roots of <i>Macaranga usambarensis</i>	Oral, one cup is taken before food twice daily	Aphrodisiac	Kenya	[60]
Capparaceae	<i>Maerua triphylla</i> A. Rich	Chokotwa (Marakwet), Chokowa (Pokot), Olamalogi (Massai)	Stem bark, leaves	Tree	Infusion	Oral	Aphrodisiac	Kenya	[23, 28]
Celastraceae	<i>Catha edulis</i> Forsk	Mairungi (Runyankore)	Leaves, stem	Shrub	Used directly	Oral (chewed)	ED	Uganda	[39, 65]
Celastraceae	<i>Elaeodendron buchananii</i> Loes	Omuharanyi	Roots	Tree	Decoction or powder used in porridge	Oral	Aphrodisiac	Tanzania	[64]
Celastraceae	<i>Maytenus putterlickioides</i> (Loes.) Excell & Mendonca	Muthuthi	Roots	Shrub	Decoction	Oral	Aphrodisiac	Kenya	[60]
Celastraceae	<i>Maytenus senegalensis</i> (Lam.) Exell	Omuwaiswa (Lusoga)	Roots	Shrub	Not specified	Not specified	Infertility	Uganda	[31]
Celastraceae	<i>Pristimera andogensis</i> var. <i>volkensii</i> (Loes.) N. Hallé	Not reported	Roots	Climber	Infusion	Oral	Aphrodisiac	Kenya	[60]
Combretaceae	<i>Combretum constrictum</i> (Benth.) Laws	Not reported	Roots	Climber	Decoction with salt or used directly	Oral, a cup drunk twice a day; root chewed	Aphrodisiac	Kenya	[60]
Combretaceae	<i>Combretum hereroense</i> Schinz	Konkon (Orma)	Roots	Tree	Decoction. Used with <i>U. leptoclados</i>	Oral, a glass daily until effective	ED, infertility	Kenya	[59, 83]
Combretaceae	<i>Combretum illairii</i> Engl	Mshinda alume (Pokomo)	Root bark	Tree	Decoction. Used with <i>Grewia tenax</i> for men	Oral, a glass daily for 7 days (or 2-3 times daily for 14 days for infertility in women)	ED, infertility (men & women)	Kenya	[58, 59, 83]
Combretaceae	<i>Combretum molle</i> R. Br. ex G. Don	Omurama (Runyankore)	Leaves	Tree	Decoction	Drink 500 ml (adult) daily	ED	Uganda	[27]

TABLE 1: Continued.

Plant family	Botanical name	Local name	Part used	Habit	Preparation mode	Administration	Use	Country	Reference
Combretaceae	<i>Combretum pentagonum</i> Laws	Not reported	Roots	Climber	Decoction with salt or used directly	Oral, a cup drunk twice/thrice a day or root is chewed	Aphrodisiac	Kenya	[60]
Cucurbitaceae	<i>Cucurbita maxima</i>	Owica (Lango)	Leaves/seeds	Herb	Decoction/used directly	Stew eaten or raw seeds chewed twice daily	Aphrodisiac	Uganda	[82]
Ebenaceae	<i>Flueggea virosa</i> (Roxb. Ex Willb.) Voigt	Lukandwa/mukandula	Leaves	Shrub	Decoction	Oral	Infertility in women	Uganda	[22, 65]
Euphorbiaceae	<i>Acalypha villicaulis</i> Hochst. ex A. Rich	Kaisokampanga (Lusoga)	Roots	Shrub	Infusion	Oral	ED, aphrodisiac	Uganda, Kenya	[31, 60, 65]
Euphorbiaceae	<i>Cluitia abyssinica</i> Jaub & Spach	Kapkurelwo (Marakwet)	Roots	Shrub	Decoction	Oral	ED	Kenya	[84]
Euphorbiaceae	<i>Croton dichagamus</i>	Qashin a'adha, Muuqaadhi (Orma)	Roots	Tree	Decoction. Used with <i>Uvaria leptocladon</i> roots	Taken, half glass 3 times daily for 6 days	Infertility in women	Kenya	[58]
Euphorbiaceae	<i>Croton menyharthii</i> Pax	Mualikaji, Muyama (Pokomo)	Roots, leaves	Tree	Decoction	Oral, half glass 2-3 times daily for 5 days	Infertility in women	Kenya	[58]
Euphorbiaceae	<i>Erythrococca fischeri</i> Pax	Mboga (Pokot)	Roots	Shrub	Decoction	Oral	Infertility	Kenya	[23]
Euphorbiaceae	<i>Euphorbia candelabrum</i> Kotschy	Olpoongi	Roots	Tree	Not specified	Not specified	Infertility	Kenya	[70]
Euphorbiaceae	<i>Euphorbia tirucalli</i> L	Not reported	Juice	Tree	Not specified	Not specified	Aphrodisiac	Tanzania	[41]
Euphorbiaceae	<i>Flueggea virosa</i> (Willd.) Voigt	Omukarara (Runyaruguru), Omukalali (Rukonjo)	Leaves, roots	Shrub	Decoction	Oral	ED, infertility	Uganda	[22, 65]
Euphorbiaceae	<i>Ricinus communis</i> L	Omukaakale (Lusoga)	Leaves	Shrub	Not specified	Not specified	ED	Uganda	[31]
Euphorbiaceae	<i>Tragia bentharii</i> Baker	Kamyu (Luganda)	Roots	Herb	Decoction	Oral	ED	Uganda	[71]
Euphorbiaceae	<i>Tragia brevipes</i> Pax	Engenyi (Runyankore)	Leaves	Herb	Decoction	Oral	ED	Uganda	[65]
Euphorbiaceae	<i>Tragia furtialis</i> Boj	Mgonampili	Roots	Climber	Decoction. Mixed with <i>Elaeodendron buchananii</i> or <i>Spathodea campanulata</i> and <i>Carisa spinarum</i>	Oral	Aphrodisiac	Tanzania	[64]

TABLE 1: Continued.

Plant family	Botanical name	Local name	Part used	Habit	Preparation mode	Administration	Use	Country	Reference
Fabaceae	<i>Abrus precatorius</i>	Mudanda, muturituri, Mudwadwa (Pokomo)	Roots, leaves, seeds	Shrub	Used directly. Seed powder taken or with seed extract or powder of <i>Indigofera cordifolia</i> or stem powder of <i>Tinospora cordifolia</i> ; root also chewed	Oral	Aphrodisiac	Kenya, Tanzania	[59, 60]
Fabaceae	<i>Abrus schimperi</i> Hochst. Ex Benth	Not reported	Roots	Shrub	Decoction	Oral	Aphrodisiac	Tanzania	[41]
Fabaceae	<i>Acacia brevispica</i> Harms	Kiptare (marakwet), Kiptara (Pokot)	Roots	Tree	Decoction	Oral	Aphrodisiac	Kenya	[23]
Fabaceae	<i>Acacia abyssinica</i> Hochst ex.Benth	Munyinya (Runyankore)	Bark	Tree	Decoction	Oral	ED	Uganda	[39]
Fabaceae	<i>Acacia drepanolobium</i> Harms ex Sjöstedt	Eluai (Massai)	Stem bark	Shrub	Not specified	Not specified	For fertility	Kenya	[28]
Fabaceae	<i>Acacia nilotica</i> (L.) Delile	Ngogwa (Marakwet), Kopokwo (Pokot)	Leaves, bark, roots	Tree	Decoction	Oral	Aphrodisiac	Kenya	[23]
Fabaceae	<i>Acacia reficiens</i> subsp. Misera (Vatke) Brenan	Leina (Marakwet), Panyarit (Pokot), Olchurrai (Massai)	Root/stem bark	Tree	Decoction	Oral	Aphrodisiac	Kenya	[23, 67]
Fabaceae	<i>Acacia sieberiana</i> Scheele	Munyinya (Runyankore, Runyaruguru)	Bark	Tree	Decoction	Oral	ED	Uganda	[65]
Fabaceae	<i>Azelia africana</i> Pers	Eiya (Lugbara)	Bark	Tree	Decoction	Oral	Aphrodisiac	Uganda	[85]
Fabaceae	<i>Albizia coriaria</i> Welw ex Oliver	Omusa (Runyankore)	Leaves, stem	Tree	Decoction	Oral	Aphrodisiac	Uganda	[39]
Fabaceae	<i>Arachis hypogaea</i> L	Ebinyobwa (Runyankore), Binyebwa (Rukonjo)	Seeds	Herb	Used directly (eaten raw or roasted)	Oral	ED	Uganda	[27, 65, 72]
Fabaceae	<i>Caesalpinia volkensii</i> Harms	Mucuthi, Muvuthi (Embu), Mujuthi (Meru)	Roots	Shrub	Used directly (eaten raw or cooked), taken with palm wine	Oral	Aphrodisiac	Kenya	[62]
Fabaceae	<i>Cajanus cajan</i> (L.) Millsp	Entondiirwa (Runyankore)	Leaves	Shrub	Decoction	Oral, drink 250 ml	ED	Uganda	[27]
Fabaceae	<i>Cassia abbreviata</i>	Mubaraka wa guba (Pokomo)	Roots	Tree	Decoction with <i>Cissampelos micronata</i> roots	Oral, a glass 3 times daily for 4 days	ED	Kenya	[59]

TABLE 1: Continued.

Plant family	Botanical name	Local name	Part used	Habit	Preparation mode	Administration	Use	Country	Reference
Fabaceae	<i>Cassia didymobotrya</i> Fresen	Mugabagaba (Runyankore), Mukyora (Rumyuruguru), Mucora (Rukonjo)	Leaves, roots	Shrub	Used directly (chew), decoction	Oral	ED	Uganda	[65]
Fabaceae	<i>Cassia occidentalis</i> L	Mwitanzoka (Runyankore, Rukonjo)	Leaves, roots	Herb	Used directly (chew), decoction	Oral	ED	Uganda	[65]
Fabaceae	<i>Desmodium salicifolium</i> Poir. DC	Mkongorana	Leaves, roots	Shrub	Decoction. Mixed with <i>Elaeodendron</i> <i>buchananii</i> and <i>Tragia furialis</i>	Oral, a glass taken daily	Aphrodisiac	Tanzania	[64]
Fabaceae	<i>Dolichos compressus</i> Wilczec	Chebugaa	Roots	Herb	Decoction	Oral	Fertility	Kenya	[80]
Fabaceae	<i>Eriosema psoraleoides</i> G. Don. Lam	Orutandaigwa	Leaves, roots	Shrub	Decoction	Oral	Aphrodisiac	Tanzania	[64]
Fabaceae	<i>Eritada abyssinica</i> Steud. ex A. Rich		Stem, bark	Tree	Not specified	Not specified	Infertility	Kenya	[60]
Fabaceae	<i>Erythrina abyssinica</i> Lam. Ex DC	Jjiirikiti (Luganda), Omutembe (Kuria), Muhuti (Kikuyu), Oloponi	Bark (stem bark), roots, stem	Tree	Decoction	Oral, eaten	Infertility (in women)	Kenya, Uganda	[22, 70, 86]
Fabaceae	<i>Dichrostachys cinerea</i> (L.) Wight & Arn	Muremanjojo (Runyankore)	Bark	Tree	Decoction	Oral	ED	Uganda	[39, 65]
Fabaceae	<i>Macrotyloma axillare</i> (E. Mey.) Verdc	Akaihabukuru (Runyaruguru)	Leaves, roots	Herb	Decoction	Oral	ED	Uganda	[65]
Fabaceae	<i>Mucuna pruriens</i> (L.) DC	Mukuna	Seeds	Herb	Decoction (tea)	Oral	Aphrodisiac	Uganda	[85]
Fabaceae	<i>Prosopis juliflora</i>	Mathenge	Root bark	Tree	Decoction. Used with <i>Zanthoxylum</i> <i>usamel</i> root bark	Oral, one teaspoonful daily for 5 days	Infertility in women	Kenya	[58]
Fabaceae	<i>Senegalia brevispica</i> (Harms) Seigler & Ebinger	Not reported	Roots, stem	Tree	Not specified	Not specified	Aphrodisiac	Kenya	[87]
Fabaceae	<i>Vachellia nilotica</i> (L.) P. J. H. Hurter & Mabb	Not reported	Stem bark, roots	Tree	Decoction	Oral	Aphrodisiac	Tanzania	[41, 60]
Fabaceae	<i>Vachellia sieberiana</i> (DC.) Kyal. & Boatwr var. <i>vermoesii</i> (De Wild.) Keay & Brenan	Not reported	Roots	Tree	Decoction	Oral	Aphrodisiac	Tanzania	[41]
Fabaceae	<i>Vigna unguiculata</i>	Bojo (Lango)	Leaves	Herb	Decoction (stewed)	Oral	Aphrodisiac	Uganda	[82]
Flacourtiaceae	<i>Ocoba spinosa</i> Forssk	Ekaepulepu (Ateso)	Roots	Herb	Decoction	Oral	ED	Uganda	[88]

TABLE 1: Continued.

Plant family	Botanical name	Local name	Part used	Habit	Preparation mode	Administration	Use	Country	Reference
Flacourtiaceae	<i>Xylotheca tettensis</i> (Klotzsch) Gilg	Not reported	Roots	Shrub	Decoction or used directly	Oral, taken or chewed	Aphrodisiac	Kenya	[60]
Lamiaceae	<i>Becium obovatum</i> (E. Mey. Ex. Benth) N. E. Br	Not reported	Not specified	Herb	Decoction	Oral	Genital stimulant/depressant	Kenya	[89]
Lamiaceae	<i>Hostundia opposita</i> Vahl	Simbaywa (Pokot), Shikuma (Luhya), Mtserere (Giriama)	Roots, leaves	Shrub	Decoction	Oral, a glass 2-3 times daily for 14 days	Aphrodisiac, infertility (men and women)	Kenya	[23, 58, 59]
Lamiaceae	<i>Ocimum suave</i> Wild	Omujaaja (Runyangkore)	Leaves	Shrub	Decoction with rock salt	Oral, drink 500 ml	ED	Uganda	[27]
Lamiaceae	<i>Plectranthus barbatus</i> Andrews	Papaha (Pokomo), Kan'gurwet (Markwet)	Roots	Shrub	Decoction. Used with <i>C. rotundifolia</i> for first 4 days	Oral, half glass daily for 30 days	ED	Kenya	[59]
Loganiaceae	<i>Buddleia polystachya</i> Fres	Chorwet	Roots	Herb	Decoction	Oral	Fertility	Kenya	[80]
Lythraceae	<i>Punica granatum</i> L	Mukungumanga (Embu, mbeere), Kukumanga (Meru)	Seeds	Shrub	Decoction	Oral	ED, infertility	Kenya	[61, 62]
Malvaceae	<i>Adansonia digitata</i> L	Muramba (Embu), Mbamburi (Swahili)	Bark	Tree	Decoction	Oral	ED	Kenya	[61]
Malvaceae	<i>Dombeya burgessiae</i> Gerrard ex Harv	Mukusa (Luhya)	Bark	Tree	Used directly (chewed)	Oral	Aphrodisiac	Kenya	[23]
Malvaceae	<i>Hibiscus fuscus</i> Garcke	Cheptelia (Marakwet), Pkapuyan (Pokot)	Roots	Herb	Used directly (chewed)	Oral	Aphrodisiac	Kenya	[23]
Malvaceae	<i>Sida tenuicarpa</i> Vollesen	Keyeyo (Rukonjo)	Leaves	Herb	Decoction	Oral	ED	Uganda	[65]
Meliaceae	<i>Ekebergia capensis</i> Sparrm	Cape ash (English)	Stem bark	Tree	Decoction	Oral	ED	Uganda	[69]
Meliaceae	<i>Bersama abyssinica</i> Fresen	Kipset (Marakwet)	Roots, leaves, branches, bark	Tree	Decoction (roots), used directly (leaves, branches and bark)	Oral	Aphrodisiac, infertility (women and men)	Kenya	[23]
Meliaceae	<i>Xylocarpus benadrensis</i> Mattei	Not reported	Unripe fruits	Tree	Used directly (exudate used)	Oral	Aphrodisiac	Tanzania	[41]
Menispermaceae	<i>Cissampelos micronata</i> A. Rich	Chovi, kivila kya mani (Pokomo), kashikiropaka (Giriama)	Roots	Herb	Decoction. Used with <i>C. abbreviate</i>	Oral, half glass daily for 3 days	Aphrodisiac, infertility, azoospermia	Kenya	[59]
Moraceae	<i>Artocarpus integer</i> (Thunb.) Merr	Fenensi (Runyangkore)	Seeds	Tree	Decoction	Oral, taken as tea	ED	Uganda	[27]

TABLE 1: Continued.

Plant family	Botanical name	Local name	Part used	Habit	Preparation mode	Administration	Use	Country	Reference
Moraceae	<i>Ficus natalensis</i> Hochst	Ekitooma (Runyankore)	Roots, root bark	Tree	Decoction	Oral, drunk 250 ml daily or 100 ml thrice daily (fresh root bark)	ED	Uganda	[27]
Moraceae	<i>Morus mesozygia</i> Stapf	Not reported	Roots	Tree	Decoction in cow/goat milk	Oral	Aphrodisiac	Kenya	[60]
Moringaceae	<i>Moringa oleifera</i> Lam	Moringa (English)	Seeds, leaves	Tree	Decoction, teas, food condiment	Oral; seed powder as tea; eat leaves as sauce; drink 100 ml	ED	Uganda	[27]
Myricaceae	<i>Myrica salicifolia</i> Hochst. ex A.Rich	Mujeje (Runyankore)	Roots, bark	Shrub	Decoction	Oral	ED	Uganda	[65]
Myrtaceae	<i>Syzygium guineense</i> (Willd.) DC	Lamaiwo (Marakwet), Cheptimanwa (Pokot)	Leaves, bark	Tree	Used directly (sap used)	Oral	Aphrodisiac, infertility	Kenya	[23, 60]
Olacaceae	<i>Capparis sepiaria</i> var. <i>caffra</i>	Hamwalika (Pokomo), Mugwada paka (Giriama)	Root bark	Shrub	Decoction. used with <i>Grewia plagiophylla</i>	Oral, half glass daily for 10 days	Aphrodisiac	Kenya	[59]
Passifloraceae	<i>Adenia gummifera</i> (Harv.) Harms	Mujoka (Pokomo)	Roots/stem	Climber	Decoction	Oral, half glass daily for 3 days	Infertility in women	Kenya	[58]
Pedaliaceae	<i>Sesamum indicum</i> L	Not reported	Leaves	Herb	Not specified	Not specified	Aphrodisiac	Kenya	[41]
Phytolaccaceae	<i>Phytolacca dodecandra</i> L'Her	Muhoko (Runyankore), Ruhuko (Rukonjo)	Roots, leaves	Shrub	Used directly	Smear on ripe banana and roast	ED	Uganda	[65]
Piperaceae	<i>Piper umbellatum</i> L	Not reported	Roots	Climber	Decoction with <i>Aframomum</i> roots and strained	Oral, one cup taken daily	Aphrodisiac	Kenya	[60]
Plumbaginaceae	<i>Plumbago zeylanica</i> L	Not reported	Roots	Shrub	Decoction	Oral	ED	Uganda	[70]
Polygalaceae	<i>Polygala aphrodisiaca</i> Gürke	Not reported	Roots	Herb	Decoction, i.e., cooked with a young cock	Oral, eaten in food	Aphrodisiac	Tanzania	[41]
Polygalaceae	<i>Polygala sphenoptera</i> Fresen	Not reported	Roots	Herb	Infusion	Oral	Aphrodisiac	Kenya, Tanzania	[23, 41, 60]
Polygalaceae	<i>Securidaca longipedunculata</i> Fres	Omukondwa (Luganda)	Leaves, bark	Tree	Decoction	Oral	ED	Uganda	[72]
Polygonaceae	<i>Coffea</i> species	Mwani (Runyankore)	Seeds	Shrub	Roasted and chewed	Oral as a beverage	ED	Uganda	[65]
Polygonaceae	<i>Hallea rubrostipulata</i> (K. Schum.) J. F. Leroy	Muziiko (Runyankore)	Bark, roots	Tree	Decoction	Oral	ED	Uganda	[65]
Polygonaceae	<i>Rumex abyssinicus</i> Jacq	Mufumbagyesi (Runyankore), Kasekekambaju (Luganda)	Leaves, stem	Shrub	Used directly (chewed)	Oral	ED	Uganda	[65]
Polygonaceae	<i>Rumex usambarensis</i> (Dammer) Dammer	Kaseke kambajjo (Luganda)	Leaves	Herb	Decoction	Oral	Aphrodisiac	Uganda	[61]

TABLE 1: Continued.

Plant family	Botanical name	Local name	Part used	Habit	Preparation mode	Administration	Use	Country	Reference
Polygonaceae	<i>Tarenna graveolens</i> (S. Moore) Bremek	Munyamazi (Rukonjo, Runyaruguru)	Leaves, roots, bark	Shrub	Decoction	Oral	ED, aphrodisiac	Uganda, Kenya	[65, 92]
Pteridaceae	<i>Actinopteris semiflabellata</i> Pic. Serm	Mwii wa ivia (Kikamba)	Whole plant	Herb	Infusion	Oral	Infertility in women	Kenya	[77]
Ranunculaceae	<i>Clematis hirsuta</i> Guill. & Perr	Omunkaamba (Runyagakore)	Leaves	Vine	Decoction	Oral	ED	Uganda	[27]
Rhamnaceae	<i>Berchemia discolor</i> (Klotzsch) Hemsl	Muchukwo (Marakwet)	Roots	Tree	Decoction	Oral	ED	Kenya	[84]
Rubiaceae	<i>Coffea arabica</i> L	Mwani (Runyankore)	Seeds	Shrub	Roasted and chewed	Oral as a beverage	ED	Uganda	[39, 65]
Rubiaceae	<i>Coffea canephora</i> Pierre ex A. Froehner (synonym: <i>Coffea robusta</i>)	Omwaani (Runyankore)	Leaves, fruits, seeds, roots	Shrub	Decoction of leaves and fruits, used directly (chew seeds) or cook with food	Oral	ED, aphrodisiac	Uganda, Kenya	[27, 60, 72]
Rubiaceae	<i>Craterispermum schweinfurthii</i> Hiern	Not reported	Roots	Shrub	Decoction, used directly (chew)	Oral	Aphrodisiac	Kenya	[60]
Rubiaceae	<i>Heinsia crinita</i>	Not reported	Stem bark	Shrub	Not specified	Not specified	ED	DRC	[93]
Rubiaceae	<i>Molinda citrifolia</i> Benth	Muziiko (Runyankore)	Roots	Tree	Decoction	Oral	ED	Uganda	[39]
Rubiaceae	<i>Psychotria capensis</i> subsp. <i>riparia</i> (K. Schum. & K. Krause) Verdc	Not reported	Roots	Shrub	Infusion/decoction	Oral	Aphrodisiac	Kenya	[60]
Rubiaceae	<i>Psychotria cyathicalyx</i> E.M. A. Petit	Not reported	Roots	Shrub	Decoction	Oral	Aphrodisiac	Kenya	[60]
Rubiaceae	<i>Psychotria lauracea</i> (K. Schum.) E. M. A. Petit	Not reported	Roots	Shrub	Decoction	Oral	Aphrodisiac	Kenya	[60]
Rubiaceae	<i>Vangueria infausta</i> Burch	Tabirirwo (Marakwet), Komolwo (Pokot)	Roots	Shrub	Decoction	Oral	ED, infertility	Kenya	[23]
Rutaceae	<i>Citropsis articulata</i> Swingle & Kellerman	Katimbolo (Luganda), Muboro (Runyankore)	Roots, bark	Tree	Decoction, used directly (chew)	Oral as a beverage in tea	ED, aphrodisiac	Uganda	[39, 61, 65]
Rutaceae	<i>Citrus sinensis</i> (L) Osbeck	Mudimu (Giriama)	Roots/stem bark	Tree	Decoction. Mixed with <i>Acacia robusta</i> and <i>Cissus rotundifolia</i> roots	Oral, a glass 3 times daily for 3 days	Infertility in women	Kenya	[58]
Rutaceae	<i>Fagaropsis hildebrandtii</i> (Engl.) Milne-Redh	Muvindavindi (Kamba)	Leaves	Shrub	Decoction	Oral	Infertility	Kenya	[94]
Salicaceae	<i>Flacourtia indica</i> (Burm.f.) Merr	Tungururwo (Marakwet), Tingoswa (Pokot)	Roots	Tree	Decoction	Oral	Infertility	Kenya	[23]
Sapindaceae	<i>Allophylus pervilliera</i> (A.Rich) Engl	Mnyanga kitswa (Pokomo)	Roots	Shrub	Decoction	Oral, a glass daily for 7 days	Infertility (men and women)	Kenya	[58, 59]

TABLE 1: Continued.

Plant family	Botanical name	Local name	Part used	Habit	Preparation mode	Administration	Use	Country	Reference
Sapindaceae	<i>Cardiospermum halicacabum</i> L.	Akambula (Lusoga)	Leaves	Climber	Not specified	Not specified	Infertility	Uganda	[31]
Sapindaceae	<i>Pappea capensis</i> Eckl. & Zeyh. var. <i>radlkoferi</i> Schinz	Oltimigomi (Massai)	Bark	Tree	Decoction	Oral	Aphrodisiac	Kenya, Tanzania	[28, 41, 70]
Solanaceae	<i>Capsicum frutescens</i> L.	Kamurari (Luganda), Eshenda (Runyankore)	Fruits, leaves, bark	Herb	Used directly (chew), decoction	Orally in food	ED	Uganda	[65, 72]
Solanaceae	<i>Solanum incanum</i> L.	Labotwa (Marakwet), Lopotwo (Pokot), Maduranzura (Luhya)	Roots	Herb	Used directly (chew)	Oral	ED	Kenya	[23]
Solanaceae	<i>Solanum nigrum</i> L.	Managu (Embu)	Whole plant	Herb	Decoction	Oral	ED	Kenya	[61]
Sterculiaceae	<i>Cola acuminata</i> Schott & Endl.	Engongoli (Rukonjo, Runyaruguru)	Fruits	Tree	Roasted and chewed	Oral in tea, porridge, milk	ED	Uganda	[65]
Sterculiaceae	<i>Sterculia africana</i> (Lour.) Fior	Iilwo (Marakwet)	Seeds	Tree	Used directly (chewed)	Oral	ED	Kenya	[84]
Stilbaceae	<i>Nuxia floribunda</i> Benth	Mngogo	Roots	Tree	Decoction	Oral	Aphrodisiac	Tanzania	[78]
Tiliaceae	<i>Grewia plagiophylla</i> . K. Schum	Mkoi (Pokomo)	Root bark	Shrub	Decoction. Used with <i>C. sepiaria</i>	Oral, a glass daily for 10 days	Aphrodisiac	Kenya	[59]
Tiliaceae	<i>Grewia similis</i> K. Schum	Mukarara (Runyaruguru)	Leaves, bark	Shrub	Decoction	Oral	ED	Uganda	[65]
Tiliaceae	<i>Grewia tenax</i> (forssk.) Fiori	Deeka (Orma), Mubavubavu, mukawa wa guba (Pokomo)	Root bark	Shrub	Decoction. Used with <i>Combretum illairii</i>	Oral, a glass daily for 7 days	ED, aphrodisiac, infertility	Kenya	[58, 59]
Urticaceae	<i>Urtica massaica</i> Mildbr	Engenyeni (Runyankore)	Whole plant, roots	Herb	Decoction	Oral	ED, aphrodisiac	Uganda, Kenya	[65, 76]
Verbenaceae	<i>Clerodendrum myricoides</i> (Hocst.) Vatke	Munjuga iria	Roots, root bark	Shrub	Decoction	Oral	Aphrodisiac	Kenya	[41, 76]
Vitaceae	<i>Cissus rotundifolia</i> (forsk.)	Mkwembe, Maneke, Neke (Pokomo), Arma (Orma)	Roots	Tree	Decoction. Sometimes used with <i>P. barbatus</i> for the first 4 days	Oral, a glass daily for 7 days	ED, increasing female fertility	Kenya, Tanzania	[59, 66]
Vitaceae	<i>Cyphostemma adenocaulle</i> (Steud. ex A. Rich) Desc. ex. Wild & R. B. Drumm	Akabombo akatono	Bark	Herb	Decoction	Oral	ED	Uganda	[71]
Zingiberaceae	<i>Zingiber officinale</i> Roscoe	Ntanghazi (Runyankore)	Rhizome	Herb	Decoction	Oral in tea, milk, porridge	ED	Uganda	[65, 72]
Zygophyllaceae	<i>Tribulus terrestris</i> L.	Kilesan (Marakwet)	Whole plant	Herb	Used directly (chewed)	Oral	ED	Kenya	[84]

Note. ED: erectile dysfunction; languages: Luganda, Lusoga, Lango, Rukonjo, and Runyankore are spoken in Uganda; Marakwet, Luhya, Nandi, Kikamba, Pokot, Orma, Wanga, Pokomo, Massai, Giriama, and Swahili are spoken in Kenya and Kinyarwanda in Rwanda.




FIGURE 2: Major botanical families from which remedies used for treating sexual dysfunction and infertility and improving virility are obtained in the EAC.

may also be used directly, i.e., chewed raw (16%) or prepared as an infusion (5%) and taken (Figure 5). The remedies are administered orally, either by taking decoctions, infusions, and eating or chewing. Only one study reported inhalation of fumes from *Cannabis sativa* leaves for treatment of ED in Uganda [65]. *Aloe volkensii* (leaf decoction) in Kenya when utilized for treating infertility is used as a wash for genitals [58], hinting that internally mediated fertility effects would be unlikely when such herbal remedies are administered orally. While most of the plants had their method of preparation and routes of administration indicated in the use reports, up to 8% of the species identified did not have specifications of the method of preparation and administration of the herbal remedies.

3.4. Bioactivity and Phytochemistry of the Reported Plants.

To decipher the therapeutic mechanisms and compounds responsible for the bioactivities of the plants reported in EAC, a holistic review of their bioactivity related to the traditional claims and phytochemistry was undertaken. However, only five reports on bioactivity from EAC were encountered for seven plants reported in this study. In this context, the aqueous extract of *Citropsis articulata* root bark was reported to increase the *in vivo* levels of serum testosterone and mounting frequency in male rats [66, 67]. Joseph et al. [68] found that aqueous extract of *Cola acuminata* (fruits) and *Zingiber officinale* (rhizome) had no significant effect on mounting frequency and testosterone levels in rats. Aqueous extract of *Tarenna graveolens* roots increased testosterone levels but had no significant effect on mounting frequency while aqueous extract of *Urtica massaiica* leaves elicited no appreciable increase in mounting frequency and testosterone levels in male rats [68]. Other

reports were for ethanolic stem bark extract of *Ekebergia capensis* which alleviated sexual dysfunction by increasing the mounting frequency and testosterone levels of male rats to 2.38 ± 0.02 ng/ml, 7.68 ± 0.66 , and 14.5 ± 0.777 ng/mL at doses of 300, 400, and 500 mg/kg, respectively [69]. The latest report is on *Plumbago zeylanica*, whose aqueous root extract administered at 150, 300, and 450 mg/kg was found to elicit prosexual stimulatory effects in male rats [70]. Though some of these reports supported the traditional use of the medicinal plants, most studies performed preliminary phytochemical screening only but not isolation and structural elucidation of the responsible bioactive compounds. Ndukupi et al. [69], for example, found saponins and steroid glycosides as the major secondary metabolites in *Ekebergia capensis* stem bark. Traces of tannins, anthraquinones, alkaloids, carotenoids, flavonoids, and anthracyanosides were also detected. Some of these secondary metabolites (tannins, phlobatannins, glycosides, phenols, saponins, quinones, terpenoids, and steroids) were also detected in *Plumbago zeylanica* [70]. It is worth noting that none of these studies probed into the mechanism of action of the extracts.

We, therefore, performed further searches and retrieved other 9 species (along with *Zingiber officinale*) cited in the EAC that have been explored for their phytochemical profiles as well as aphrodisiac, procopulatory, and fertility effects (Table 3). One of the most studied plants in this context is *Allium cepa* (*A. cepa*) which is locally used in culinary recipes. It has been reported to improve copulatory behaviour in sexually experienced rats [71]. Malviya et al. [72] indicated that ethyl acetate fraction of *A. cepa* bulb at 200 mg/kg restored the mating behaviour (ejaculatory latency, postejaculatory interval, mount, intromission, and ejaculatory frequencies and mount and intromission latencies) of drug-mediated sexually dysfunctional male rats.

TABLE 2: Synopsis of the most used plant species for the treatment of infertility among men and women in the East African Community.

Medicinal plant	Parts used	Mode of preparation	Mode of administration	Group treated (country)	References
<i>Uvaria leptoclados</i>	Roots	Decoction	Oral, half glass thrice daily for 3 days	Women (Kenya)	[58, 59]
<i>Markhamia zanzibarica</i>	Roots	Decoction with roots of <i>Salvadora persica</i> and <i>Uvaria acuminata</i>	Oral	Women (Kenya)	[58, 59]
<i>Spathodea campanulata</i> Buch.-Harm. ex DC	Bark	Decoction	Oral	Men (Uganda)	[22]
<i>Mangifera indica</i> L	Bark	Decoction drunk	Oral	Women (Uganda)	[22]
<i>Flueggea virosa</i> (Roxb. Ex Willb.) Voigt	Leaves	Decoction drunk	Oral	Women (Uganda)	[22, 65]
<i>Erythrina abyssinica</i> Lam. Ex	Bark (stem bark), roots, stem	Decoction, eaten directly	Oral	Women (Uganda, Kenya)	[22, 70, 86]
<i>Cadaba glandulosa</i> Forssk	Roots	Decoction	Oral, half a glass daily for 5 days	Women and men (Kenya)	[58, 59]
<i>Cadaba farinose</i>	Roots	Decoction	Oral, half a glass daily for 3 days	Women and men (Kenya)	[58, 59]
<i>Combretum illairii</i> Engl	Root bark	Decoction used with <i>Grewia tenax</i> for men	Oral, a glass daily for 7 days (or 2-3 times daily for 14 days for infertility in women)	Women and men (Kenya)	[58, 59, 83]
<i>Hoslundia opposita</i> Vahl	Leaves	Decoction	Oral, a glass 2-3 times daily for 14 days	Women and men (Kenya)	[58, 59]
<i>Allophylus pervillieria</i> (A. Rich) Engl	Roots	Decoction	Oral, a glass daily for 7 days	Women and men (Kenya)	[58, 59]




FIGURE 3: Life form of plants used for the preparation of remedies used in the treatment of erectile dysfunction and infertility, and increasing fertility and virility in the EAC.




FIGURE 4: Plant organs used in herbal preparations for treating erectile dysfunction and infertility, and enhancing fertility and virility in the EAC.

Quercetin (1) (Figure 6), a flavonoid present in extracts of *A. cepa*, enhanced sperm motility through the regulation of protein kinase C-mediated activation of the human voltage-gated proton channel and could explain its therapeutic effect when used in the treatment of human infertility [14]. Similarly, S-allyl cysteine (2) isolated from *Allium sativum* restored erectile function in diabetic rats through inhibiting reactive oxygen species formation via modulation of nicotinamide adenine dinucleotide phosphate oxidase subunit expression in penile tissues [73].

The third highly investigated species is *Mondia whitei*. It has been found to increase sexual arousal and copulatory efficiency and improve sexual sensation in rats [74–78]. A follow-up study with a polyherbal formulation containing *Mondia whitei*, *Dracaena arborea*, and *Bridelia ferruginea*

deduced that the administration of the formula enhanced the sexual performances and increased the mounting and intromission frequencies of normal rats and prediabetic rats [79].

Zingiber officinale (ginger) is the most thoroughly studied plant cited in this report. A bioactive compound from this species (zingerone, 5) attenuated zearalenone-induced steroidogenesis impairment in TM3 Leydig cell lines [80] and elicited dose-dependent enhancement of fertility in male and female rats as witnessed by increments in gonadal weights and sperm counts [81]. A gingerol (6)-rich fraction of ginger at 50, 100, and 200 mg/kg when administered to male rats with carbendazim-induced toxicity led to increased sperm motility and count but attenuated sperm abnormality [82].




FIGURE 5: Methods of preparation of herbal remedies used in the treatment of erectile dysfunction and infertility, and enhancing fertility and virility in the EAC.

Herbal extracts from plants such as *Allium cepa*, *Allium sativum*, *Mondia whitei*, and *Zingiber officinale* improve semen quality and sperm parameters such as concentration, viability, motility, morphology, and DNA integrity through increment in gonadal hormone levels (testosterone and luteinising hormone), sequestering free radicals and enhanced production of nitric oxide [83–85]. Such studies substantiate that the traditional claims of using the plants in the treatment of sexual dysfunction in EAC may be credible.

The contraceptive effect observed in plants such as *Catha edulis* (cathinone) and *Cannabis sativa* seed extracts is supported by studies which instead link their use to ED [86]. Nevertheless, plant extracts from certain families have been shown to elicit contradictory effects in fertility studies. Such differential bioactivities are species-specific and may depend on the extraction method and solvents employed [87, 88].

3.5. Clinical Studies. Clinical evaluation of herbal products is a requirement before they are promoted and used. In this study, we did not find any clinical studies in EAC that was performed on the extracts or isolated compounds from the cited plants. Further searches for global reports indicated that *Zingiber officinale* is the only plant encountered in this study that have been subjected to clinical studies investigating its effect on male ED, female sexual function, and infertility [89]. For example, its capsules improved the sexual function and quality of life in four weeks of a randomized, double-blind clinical trial involving women of reproductive age ($n = 190$) [90]. Another randomized double-blind placebo-controlled trial found that 3-month oral treatment using 500 mg/powder/day reduced sperm DNA fragmentation in infertile men [91]. Such promising clinical results demonstrate the need for more clinical trials on species such as *Mondia whitei*, *Acalypha villicaulis*,

Combretum illairii, *Erythrina abyssinica*, *Pappea capensis*, *Rhus vulgaris*, and *Warburgia ugandensis* that are widely used in the region.

3.6. Adverse Side Effects and Toxicity of Medicinal Plants and Bioactive Phytochemicals Reported. Further analysis of reports considered in this systematic review showed that no ethnobotanical survey captured the side effects of herbal preparations used in the management of sexual dysfunction, infertility, and improving virility in the EAC. However, some of the plants such as *Abrus precatorius* (roots, leaves, and seeds) cited in the EAC are known to contain highly poisonous compounds (abrine, precatorine, and hypaphorine) [139]. It could be positioned that the preparation of remedies with more than one plant and plant part or with the addition of adjuvants may be a way of masking the toxicity of the medicinal plants [36, 93].

From available toxicological studies, extracts from six of the investigated bioactive species have been shown to be safe (Table 4). Four species (*Abrus precatorius*, *Catha edulis*, *Cannabis sativa*, and *Parquetina nigrescens*) have been indicated to elicit marked toxicity, indicating that their use may lead to adverse reactions in herbal medicine practice. For bioactive compounds identified in the listed species (Table 3), quercetin (1) is potentially cytotoxic and hepatotoxic at higher doses (100 to 2,000 mg/kg) [94, 95]. Similarly, cathinone (3) is a psychoactive compound that is toxic to sperm cells [96], and its abuse has been associated with fatal renal, hepatic, and cardiac injuries [97]. On the other hand, S-allyl cysteine (2) is considered to be safe, with very minor acute/subacute toxicity in mice and rats ($LD_{50} > 54.7$ mM/kg) when administered intraperitoneally [98]. Sesamine (4) is the major lignan in sesame seeds and has been confirmed to be safe. It attenuated reactive oxygen

TABLE 3: Bioactivity and phytochemical profile of some plants used in the treatment of ED and infertility, and enhancing virility and fertility in EAC.

Plant	Part used	Extract	Bioactivity/mechanism of action and active phytoconstituents
<i>Abrus precatorius</i>	Seeds	Methanol	Antifertility effect in rats [101].
<i>Allium cepa</i> L.	Bulb	Aqueous, ethyl acetate	Enhanced copulatory behaviour in male rats [71, 106]. Ethyl acetate fraction of extract at 200 mg/kg restored the mating behaviour of drug-mediated sexually dysfunction male rats [72]. Quercetin (1) isolated from the plant-enhanced sperm motility [14]. Increased weight of seminal vesicles and epididymides in male rats [72, 106–108]. S-allyl cysteine (2) isolated from this species-promoted fertility [73]. Reduced epididymal sperm count in rats (contraceptive effect) [87, 88, 103]. Improvement of sexual behaviour and increase in plasma testosterone levels [112–114]. Cathinone (3) is toxic to sperm cells [96].
<i>Allium sativum</i> L.	Bulb	Aqueous, petroleum ether	Increase sperm count, motility, and fertilization ability in African catfish increase in testicular weight, body weight, testosterone levels, and follicle-stimulating hormone [116, 117].
<i>Cannabis sativa</i>	Seeds	Ethanol	Increased sexual arousal, copulatory efficiency, sexual sensation [74–78] through the activation/stimulation of nitric oxide synthase activity resulting in the elevation of levels of cyclic guanosine monophosphate [83].
<i>Catha edulis</i> Forsk	Shoots, small branches	Chloroform: diethyl ether extract (1:3)	The extracts improved sexual activity, behaviour, and competence through improving sexual hormone secretion [102, 125].
<i>Kigelia africana</i>	Fruits	Powder used directly	It promotes body weight gain, seminal parameters, antioxidant action, and testosterone level [126]. Sesamin (4), a compound in this species, resisted cyclophosphamide-induced sperm nuclear maturity and DNA damage by increasing the expression levels of histones H2A and H2B in the testis [127].
<i>Mondia whitei</i>	Root bark, roots	Aqueous, hexane	Powder at 100 mg has been cited to elicit positive effects in folliculogenesis and implantation [128]. Zingerone (5) isolated from the plant extract normalized zearalenone-impaired steroidogenesis in TM3 cells [80]. Similarly, a gingerol (6)-rich fraction of ginger enhanced sperm motility and count but attenuated sperm abnormality in male rats with carbendazim-induced toxicity [82].
<i>Parquetina nigrescens</i>	Leaves	Aqueous	
<i>Sesamum indicum</i>	Seeds	Ethanol	
<i>Zingiber officinale</i>	Rhizome		




FIGURE 6: Some of the bioactive molecules characterized from extracts of plants reported in EAC for treatment of ED and infertility, and enhancing fertility and virility (based on studies outside the region). The numbers 1 to 6 refer to the molecules mentioned in Table 3.

TABLE 4: Toxicity profile of plants with reports of efficacy that is used in the treatment of ED and infertility, and enhancing virility and fertility in EAC.

Plants	Toxicity reports
<i>Abrus precatorius</i>	Seeds contain abrin, a toxalbumin with a human lethal dose of 0.1–1 $\mu\text{g}/\text{kg}$ [148]. Poisoning is characterized by severe vomiting and abdominal pain, bloody diarrhoea, convulsions, and alteration of sensorium with depression of central nervous system [149].
<i>Allium cepa</i> L	Oral administration of extracts to mice at 250 and 500 mg/kg/day for 30–90 days had no visible toxicity symptoms. An oral dose of 30 g/kg/day for 30 days resulted into hypothermia, tachypnea, tachycardia, piloerection, and polyuria in the treated mice [150].
<i>Allium sativum</i> L	Its bulb extract induced mild alterations at 300 mg/kg in mice, indicating that it is relatively safe [151].
<i>Cannabis sativa</i>	Cannabidiol (a major nonpsychotropic constituent of this species) in extracts of this species is potentially toxic through the inhibition of hepatic drug metabolism, alterations of <i>in vitro</i> cell viability, reduced fertilization capacity, and decreased activities of <i>p</i> -glycoprotein and other drug transporters [152].
<i>Catha edulis</i> Forsk	Crude khat can damage the liver and kidneys and modulate levels of liver enzymes, urea, creatinine, and electrolytes essential for liver and kidney functions [153].
<i>Kigelia africana</i>	Low to moderately toxic [154].
<i>Mondia whitei</i>	Low toxicity in mice exposed to the extract for 90 days [155].
<i>Parquetina nigrescens</i>	Toxic to rats at 100 and 300 mg/kg of methanol leaf and aerial part extract. Renal haemorrhage, inflammation, and hepatic inflammation were noted [156].
<i>Sesamum indicum</i>	Ethanol extract had low toxicity at 500 mg/kg body weight [157].
<i>Zingiber officinale</i>	Extract had no toxicity at 5,000 mg/kg body weight [158].

species and nitric oxide production in zebra fish (LD₅₀) [99]. Toxicity studies with zingerone (5) and gingerol (6) have shown that they are safe [100, 104].

4. Conclusion

The EAC has a rich ethnobotanical knowledge of herbal remedies for the management of sexual dysfunction and infertility, and improving fertility and virility. Though we retrieved 171 medicinal plants being used, most of the species have not been subjected to phytochemical and bioactivity studies that lend credence to traditional claims of using them. We recommend performing toxicity studies and clinical trials using compounds isolated from some of the investigated species. Five highly cited unstudied species from this review (*Acalypha villicaulis*, *Combretum illairii*, *Erythrina abyssinica*, *Pappea capensis*, *Rhus vulgaris*, and *Warburgia ugandensis*) have been selected for further investigation of their phytochemistry, aphrodisiac, fertility, and phosphodiesterase-5 inhibitory activities.

Data Availability

This is a systematic review article, and no raw experimental data were collected. All data generated or analyzed during this study are included in this article.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

Supplementary Materials

Supplementary file 1: PRISMA 2020 checklist for the systematic review of medicinal plants used in the management of sexual dysfunction and infertility, and improving virility in the East African Community. Supplementary file 2: Risk of bias assessment of studies included for systematic review on medicinal plants used in the management of sexual dysfunction and infertility, and improving virility in the East African Community. (*Supplementary Materials*)

References

- [1] B. Y. F. Fong and V. T. S. Law, *Sustainable Development Goal 3: Health and Well-Being of Ageing in Hong Kong*, Taylor & Francis, New York, NY, USA, 2022.
- [2] T. Vos, R. M. Barber, B. Bell et al., “Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013,” *The Lancet*, vol. 386, no. 9995, pp. 743–800, 2015.
- [3] C. Hajat and E. Stein, “The global burden of multiple chronic conditions: a narrative review,” *Preventive Medicine Reports*, vol. 12, pp. 284–293, 2018.
- [4] H. Wang, M. Naghavi, C. Allen et al., “Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015,” *The Lancet*, vol. 388, no. 10053, pp. 1459–1544, 2016.
- [5] Who, “WHO reveals leading causes of death and disability worldwide: 2000–2019,” 2020, <https://www.who.int/news/item/09-12-2020-who-reveals-leading-causes-of-death-and-disability-worldwide-2000-2019>.
- [6] B. Pham, R. Jorry, N. Abori, V. D. Silas, A. D. Okely, and W. Pomat, “Non-communicable diseases attributed mortality and associated sociodemographic factors in Papua New Guinea: evidence from the comprehensive health and epidemiological surveillance system,” *PLoS Global Public Health*, vol. 2, no. 3, 2022.
- [7] Who, “Noncommunicable diseases,” 2021, <https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases>.
- [8] J. Ali, S. H. Ansari, and S. Kotta, “Exploring scientifically proven herbal aphrodisiacs,” *Pharmacognosy Reviews*, vol. 7, no. 1, pp. 1–10, 2013.
- [9] A. N. Sidawy, “Erectile dysfunction,” in *Rutherford’s Vascular Surgery and Endovascular Therapy*, Springer, Berlin, Germany, 2019.
- [10] P. Bearely, E. A. Phillips, S. Pan et al., “Long-term intracavernosal injection therapy: treatment efficacy and patient satisfaction,” *International Journal of Impotence Research*, vol. 32, no. 3, pp. 345–351, 2020.
- [11] D. Mobley, M. Khera, and N. Baum, “Recent advances in the treatment of erectile dysfunction,” *Postgraduate Medical Journal*, vol. 93, no. 1105, pp. 679–685, 2017.
- [12] P. Grant, G. Jackson, I. Baig, and J. Quin, “Erectile dysfunction in general medicine,” *Clinical Medicine*, vol. 13, no. 2, pp. 136–140, 2013.
- [13] S. Saha, P. Roy, C. Corbitt, and S. S. Kakar, “Application of stem cell therapy for infertility,” *Cells*, vol. 10, no. 7, p. 1613, 2021.
- [14] M. Chae, S. J. Kang, K. P. Lee et al., “Onion (*Allium cepa* L.) peel extract (OPE) regulates human sperm motility via protein kinase C-mediated activation of the human voltage-gated proton channel,” *Andrology*, vol. 5, no. 5, pp. 979–989, 2017.
- [15] B. V. Rossi, M. Abusief, and S. A. Missmer, “Modifiable risk factors and infertility: what are the connections?” *American Journal of Lifestyle Medicine*, vol. 10, no. 4, pp. 220–231, 2016.
- [16] A. Ilacqua, G. Izzo, G. P. Emerenziani, C. Baldari, and A. Aversa, “Lifestyle and fertility: the influence of stress and quality of life on male fertility,” *Reproductive Biology and Endocrinology*, vol. 16, no. 1, p. 115, 2018.
- [17] G. Biviá-Roig, A. Boldó-Roda, R. Blasco-Sanz et al., “Impact of the COVID-19 pandemic on the lifestyles and quality of life of women with fertility problems: a cross-sectional study,” *Frontiers in Public Health*, vol. 9, Article ID 686115, 2021.
- [18] A. Starc, M. Trampuš, D. Pavan Jukić, C. Rotim, T. Jukić, and A. Polona Mivšek, “Infertility and sexual dysfunctions: a systematic literature review,” *Acta Clinica Croatica*, vol. 58, no. 3, pp. 508–515, 2019.
- [19] A. Skalkidou, T. N. Sergentanis, S. P. Gialamas et al., “Risk of endometrial cancer in women treated with ovary-stimulating drugs for subfertility,” *The Cochrane Database of Systematic Reviews*, vol. 3, no. 3, 2017.
- [20] A. Bakhtiari, Z. Basirat, and F. Nasiri-Amiri, “Sexual dysfunction in women undergoing fertility treatment in Iran: prevalence and associated risk factors,” *Journal of Reproduction & Infertility*, vol. 17, no. 1, pp. 26–33, 2016.
- [21] A. Anand Ganapathy, V. M. Hari Priya, and A. Kumaran, “Medicinal plants as a potential source of Phosphodiesterase-

- 5 inhibitors: a review," *Journal of Ethnopharmacology*, vol. 267, Article ID 113536, 2021.
- [22] P. Tugume, E. K. Kakudidi, M. Buyinza et al., "Ethnobotanical survey of medicinal plant species used by communities around Mabira central forest reserve, Uganda," *Journal of Ethnobiology and Ethnomedicine*, vol. 12, no. 1, p. 5, 2016.
- [23] Y. M. Mbuni, S. Wang, B. N. Mwangi et al., "Medicinal plants and their traditional uses in local communities around cherangani hills, western Kenya," *Plants*, vol. 9, no. 3, p. 331, 2020.
- [24] T. Omara, A. K. Kiprof, R. C. Ramkat et al., "Medicinal plants used in traditional management of cancer in Uganda: a review of ethnobotanical surveys, phytochemistry, and anticancer studies," *Evidence-Based Complementary and Alternative Medicine*, vol. 2020, no. 6, Article ID 3529081, 26 pages, 2020.
- [25] T. Omara, "East African quintessential plants claimed to be used as blood purifiers, cleansers, detoxifiers and tonics: an appraisal of ethnobotanical reports and correlation with reported bioactivities," *Bulletin of the National Research Centre*, vol. 45, no. 1, p. 171, 2021.
- [26] E. Fratkin, "Traditional medicine and concepts of healing among Samburu pastoralists of Kenya," *Journal of Ethnobiology*, vol. 16, no. 1, pp. 63–97, 1996.
- [27] H. Gumisiriza, G. Birungi, E. A. Olet, and C. D. Sesaaizi, "Medicinal plant species used by local communities around queen elizabeth national park, maramagambo central forest reserve and ihmbo central forest reserve, south western Uganda," *Journal of Ethnopharmacology*, vol. 239, Article ID 111926, 2019.
- [28] J. Kimondo, J. Miaron, P. Mutai, and P. Njogu, "Ethnobotanical survey of food and medicinal plants of the Ilkisonko Maasai community in Kenya," *Journal of Ethnopharmacology*, vol. 175, pp. 463–469, 2015.
- [29] I. Sindiga, "Indigenous medical knowledge of the Maasai," *Indigenous Knowledge and Development Monitor*, vol. 2, pp. 16–18, 1994.
- [30] H. Trant, "Food taboos in East Africa," *The Lancet*, vol. 264, no. 6840, pp. 703–705, 1954.
- [31] J. Kigenyi, "Coping with resource extinction: the case of medicinal plants in Kawete village, Iganga district, Uganda," *Culture and Environment in Africa Series*, Vol. 9, The Cologne African Studies Centre, Cologne, Germany, 2016.
- [32] S. C. Chhabra, R. L. A. Mahunnah, and E. N. Mshiu, "Plants used in traditional medicine in eastern Tanzania. VI. Angiosperms (sapotaceae to zingiberaceae)," *Journal of Ethnopharmacology*, vol. 39, no. 2, pp. 83–103, 1993.
- [33] S. H. Ngere, V. Akelo, K. Ondeng'e et al., "Traditional medicine beliefs and practices among caregivers of children under five years-the child health and mortality prevention surveillance (champs), western Kenya: a qualitative study," *PLoS One*, vol. 17, no. 11, 2022.
- [34] S. van Vuuren and L. Frank, "Review: southern African medicinal plants used as blood purifiers," *Journal of Ethnopharmacology*, vol. 249, Article ID 112434, 2020.
- [35] M. J. Page, J. E. McKenzie, P. M. Bossuyt et al., "The PRISMA 2020 statement: an updated guideline for reporting systematic reviews," *International Journal of Surgery*, vol. 88, p. 105906, 2021.
- [36] T. Omara, C. K. Nakiguli, R. A. Naiyl, F. A. Opondo, S. B. Otieno, M. L. Ndiege et al., "Medicinal plants used as snake venom antidotes in East African Community: review and assessment of scientific evidences," *Journal of Medicinal and Chemical Sciences*, vol. 4, no. 2, pp. 107–144, 2021.
- [37] Joanna Briggs Institute, "Checklist for prevalence studies," 2017, https://jbi.global/sites/default/files/2019-05/JBI_Critical_Appraisal-Checklist_for_Prevalence_Studies2017_0.pdf.
- [38] D. Atlaw, B. Sahiledengle, S. Degno et al., "Utilization of provider-initiated HIV testing and counselling in Ethiopia: a systematic review and meta-analysis," *Tropical Medicine and Health*, vol. 50, no. 1, p. 29, 2022.
- [39] K. N. Kapere, M. Maud Kamatenesi, T. A. Otwine, and N. Waisindye, "Community perceptions on the use of traditional medicine among people experiencing sexual dysfunctions in Greater Mbarara, western Uganda," *Journal of Pharmacognosy and Phytochemistry*, vol. 11, no. 2, pp. 46–59, 2022.
- [40] P. Ssegawa and J. M. Kasenene, "Medicinal plant diversity and uses in the Sango bay area, Southern Uganda," *Journal of Ethnopharmacology*, vol. 113, no. 3, pp. 521–540, 2007.
- [41] J. M. Watt and M. G. Breyer-Brandwijk, "The medicinal and poisonous plants of Southern and Eastern Africa being an account of their medicinal and other uses," *Chemical Composition, Pharmacological Effects and Toxicology in Man and Animal Livingstone*, Edinburgh, 1962.
- [42] M. Kamatenesi-Mugisha and H. Oryem-Origa, "Traditional herbal remedies used in the management of sexual impotence and erectile dysfunction in Western Uganda," *African Health Sciences*, vol. 5, no. 1, pp. 40–49, 2005.
- [43] C. K. Kaingu, J. A. Oduma, J. M. Mbaria, and S. G. Kiama, "Medicinal plants traditionally used for the management of female reproductive health dysfunction in Tana River County, Kenya," *Tang Humanitarias Medicine*, vol. 3, no. 2, pp. 17.1–17.10, 2013.
- [44] N. Shiracko, B. O. Owuor, M. M. Gakuubi, and W. Wanzala, "A survey of ethnobotany of the AbaWanga people in Kakamega County, western province of Kenya," *Indian Journal of Traditional Knowledge*, vol. 15, no. 1, pp. 93–102, 2016.
- [45] P. G. Kareru, G. M. Kenji, A. N. Gachanja, J. M. Keriko, G. Mungai, and G. Mungai, "Traditional medicines among the Embu and Mbeere peoples of Kenya," *African Journal of Traditional, Complementary, and Alternative Medicines: AJTCAM*, vol. 4, no. 1, pp. 75–86, 2006.
- [46] J. Muriuki, "Medicinal trees in smallholder agroforestry systems: assessing some factors influencing cultivation by farmers East of Mt Kenya," Dissertation, University of Natural Resources and Applied Life Sciences, Vienna, 2011.
- [47] C. K. Kaingu, J. M. Mbaria, J. Oduma, and G. S. Kiama, "Ethnobotanical survey of medicinal plants used for the management of male sexual dysfunction and infertility in tana river county, Kenya," *The Journal of Ethnobiology and Traditional Medicine*, vol. 119, pp. 453–463, 2013.
- [48] M. J. Moshi, D. F. Otieno, P. K. Mbabazi, and A. Weisheit, "Ethnomedicine of the kagera region, north western Tanzania. Part 2: the medicinal plants used in katoro ward, bukoba district," *Journal of Ethnobiology and Ethnomedicine*, vol. 6, no. 1, p. 19, 2010.
- [49] G. H. Schmelzer, *Medicinal Plant. Plant Resources of Tropical Africa (PROTA)*, Technical Centre for Agricultural and Rural Cooperation, Netherlands, 2008.
- [50] K. J. Cheruiyot, E. Njenga, C. Mutai, C. Bii, R. Korir, and E. Too, "Ethnobotanical survey and plant monographs of medicinal plants used among the Elgeyo community in Kenya," *Photon*, vol. 120, pp. 633–649, 2013.
- [51] J. O. Kokwaro, *Medicinal Plants of East Africa*. University of Nairobi Press, Nairobi, Kenya, 3rd edition, 2009.

- [52] K. Munguti, "Indigenous knowledge in the management of malaria and visceral leishmaniasis among the tugen of Kenya," *Indigenous Knowledge and Development Monitoring*, vol. 5, pp. 10–12, 1997.
- [53] J. G. Agea, B. Katongole, D. Waiswa, and G. N. Nabanoga, "Market survey of *Mondia whytei* (mulondo) roots in Kampala City, Uganda," *African Journal of Traditional, Complementary, and Alternative Medicines: AJTCAM*, vol. 5, no. 4, pp. 399–408, 2008.
- [54] J. K. Muthee, D. W. Gakuya, J. M. Mbaria, P. G. Kareru, C. M. Mulei, and F. K. Njonge, "Ethnobotanical study of antihelmintic and other medicinal plants traditionally used in Loitoktok district of Kenya," *Journal of Ethnopharmacology*, vol. 135, no. 1, pp. 15–21, 2011.
- [55] S. Asimwe, J. Namukobe, R. Byamukama, and B. Imalingat, "Ethnobotanical survey of medicinal plant species used by communities around Mabira and Mpanga Central Forest Reserves, Uganda," *Tropical Medicine and Health*, vol. 49, no. 1, p. 52, 2021.
- [56] L. Ssozi, B. Kabiito, A. Byaruhanga, and W. Kanata, "Documenting baganda ethno-medicine: a step towards preservation and conservation," *Journal of Applied and Advanced Research*, vol. 1, pp. 15–22, 2016.
- [57] G. H. Schmelzer and A. Gurib-Fakim, "Plant resources of tropical Africa," *Medicinal Plants 1*, PROTA Foundation, Wageningen, Netherlands, 2008.
- [58] J. O. Kokwaro, *Medicinal Plants of East Africa*, East African Literature Bureau, Nairobi, Kenya, 2nd edition, 1993.
- [59] P. Jeruto, C. Lukhoba, G. Ouma, D. Otieno, and C. Mutai, "An ethnobotanical study of medicinal plants used by the Nandi people in Kenya," *Journal of Ethnopharmacology*, vol. 116, no. 2, pp. 370–376, 2008.
- [60] G. N. Njoroge and R. W. Bussman, "Ethnotherapeutic management of sexually transmitted diseases (STDs) and reproductive health conditions in central province of Kenya," *Indian Journal of Traditional Medicine*, vol. 8, no. 2, pp. 255–261, 2009.
- [61] W. Wanzala, S. M. Syombua, and J. O. Alwala, "A survey of the applications and use of ethnomedicinal plants and plant products for healthcare from the Ukambani region in Eastern Kenya," *Indian Journal of Ethnopharmacology*, vol. 2, no. 2, pp. 6–58, 2016.
- [62] E. K. Kakudidi, "Cultural and social uses of plants from and around kibale national park, western Uganda," *African Journal of Ecology*, vol. 42, no. s1, pp. 114–118, 2004.
- [63] R. A. Kitula, "Use of medicinal plants for human health in udzungwa mountains forests: a case study of new dabaga ulongambi forest reserve, Tanzania," *Journal of Ethnobiology and Ethnomedicine*, vol. 3, no. 1, p. 7, 2007.
- [64] P. C. Rwangabo and P. C. Rwangabo, *La médecine traditionnelle au Rwanda*, vol. 258, Editions Karthala et ACCT, Paris, 1993.
- [65] S. V. Okello, R. O. Nyunja, G. W. Netondo, and J. C. Onyango, "Ethnobotanical study of medicinal plants used by Sabaots of Mt. Elgon Kenya," *African Journal of Traditional, Complementary, and Alternative Medicines: AJTCAM*, vol. 7, no. 1, pp. 1–10, 2009.
- [66] H. D. Neuwinger, *African Ethnobotany: Poisons and Drugs: Chemistry, Pharmacology, Toxicology*, Chapman & Hall, London, 1996.
- [67] R. Nakaziba, M. K. Anyolitho, S. B. Amanyanya et al., "Traditional medicinal vegetables in northern Uganda: an ethnobotanical survey," *International Journal of Food Science*, vol. 2021, Article ID 5588196, 18 pages, 2021.
- [68] R. Korir, C. Kimani, J. Gathirwa, M. Wambura, and C. Bii, "In vitro antimicrobial properties of methanol extracts of three medicinal plants from Kilifi district, Kenya," *African Journal of Health Sciences*, vol. 20, pp. 4–10, 2012.
- [69] W. Kipkore, B. Wanjohi, H. Rono, and G. Kigen, "A study of the medicinal plants used by the Marakwet community in Kenya," *Journal of Ethnobiology and Ethnomedicine*, vol. 10, no. 1, p. 24, 2014.
- [70] G. Anywar, E. Kakudidi, R. Byamukama, J. Mukonzo, A. Schubert, and H. Oryem-Origa, "Indigenous traditional knowledge of medicinal plants used by herbalists in treating opportunistic infections among people living with HIV/AIDS in Uganda," *Journal of Ethnopharmacology*, vol. 246, Article ID 112205, 2020.
- [71] J. Nankaya, J. Nampushi, S. Petenya, and H. Balslev, "Ethnomedicinal plants of the loita Maasai of Kenya," *Environment, Development and Sustainability*, vol. 22, no. 3, pp. 2569–2589, 2019.
- [72] A. S. Delbanco, N. D. Burgess, and A. Cuni-Sanchez, "Medicinal plant trade in Northern Kenya: economic importance, uses, and origin," *Economic Botany*, vol. 71, no. 1, pp. 13–31, 2017.
- [73] K. Philip, M. Elizabeth, P. Cheplogoi, and K. Samuel, "Ethnobotanical survey of antimalarial medicinal plants used in Butebo county, Eastern Uganda," *European Journal of Medicinal Plants*, vol. 21, no. 4, pp. 1–22, 2017.
- [74] D. O. Okach, A. R. O. Nyunja, and G. Opande, "Phytochemical screening of some wild plants from Lamiaceae and their role in traditional medicine in Uiri district–Kenya," *International Journal of Herbal Medicine*, vol. 1, no. 5, pp. 135–143, 2013.
- [75] J. G. Ndukui, H. Muwonge, and L. F. Sembajwe, "Aphrodisiac potential and phytochemical profile of *Ekebergia capensis* (Cape ash) in male albino rats," *Spatula DD*, vol. 2, no. 4, pp. 237–243, 2012.
- [76] P. Onen, D. Ocira, T. Omara, J. Nyeko, and A. Okwir, "Preliminary phytochemical screening of *Plumbago zeylanica* L. Roots and its aphrodisiac effect in male rats," *Asian Journal of Applied Chemistry Research*, vol. 8, no. 3, pp. 24–31, 2021.
- [77] E. Odongo, N. Mungai, P. Mutai, E. Karumi, J. Mwangi, and J. Omale, "Ethnobotanical survey of medicinal plants used in Kakamega County, western Kenya," *Applied Medical Research*, vol. 4, no. 1, pp. 22–40, 2018.
- [78] P. Tshisekedi Tshibangu, P. Mutwale Kapepula, M. Kabongo Kapinga et al., "Anti-plasmodial activity of *Heinsia crinita* (Rubiaceae) and identification of new iridoids," *Journal of Ethnopharmacology*, vol. 196, pp. 261–266, 2017.
- [79] W. Musila, D. Kisangau, and J. Muema, "Conservation status and use of medicinal plants by traditional medical practitioners in machakos district, Kenya," in *Indigenous Knowledge Conference Proceedings*, p. 22, Pennsylvania State University. Traditional Aspects of Health and Wellness, Pennsylvania, PA, USA, 2004.
- [80] MEWNR, "Kenya biodiversity atlas. Ministry of environment natural resources and regional development authorities," 2015, <https://webfiles.york.ac.uk/KITE/Kenya%20Atlas/Kenyas%20Natural%20Atlas%20-%20A%20Biodiversity%20Atlas.pdf>.
- [81] T. Omara, S. Kagoya, A. Openy et al., "Antivenin plants used for treatment of snakebites in Uganda: ethnobotanical reports and pharmacological evidences," *Tropical Medicine and Health*, vol. 48, no. 1, pp. 6–16, 2020.

- [82] K. Medius, K. Esther, B. C. Anthony, and H. Robert, "Medicinal plants and herbalist preferences around bwindi impenetrable national park," *Journal of Medicinal Plants Research*, vol. 11, no. 8, pp. 161–170, 2017.
- [83] T. Omara, "Plants used in antivenom therapy in rural Kenya: ethnobotany and future perspectives," *Journal of Toxicology*, vol. 2020, Article ID 1828521, 9 pages, 2020.
- [84] D. Asmerom, T. H. Kalay, T. Y. Araya, D. M. Desta, D. Z. Wondafrash, and G. G. Tafere, "Medicinal plants used for the treatment of erectile dysfunction in Ethiopia: a systematic review," *BioMed Research International*, vol. 2021, Article ID 6656406, 12 pages, 2021.
- [85] S. S. Semenya and M. J. Potgieter, "Ethnobotanical survey of medicinal plants used by Bapedi traditional healers to treat erectile dysfunction in the Limpopo Province, South Africa," *Journal of Medicinal Plants Research*, vol. 7, pp. 349–357, 2013.
- [86] M. Nimrouzi, A. M. Jaladat, and M. M. Zarshenas, "A panoramic view of medicinal plants traditionally applied for impotence and erectile dysfunction in Persian medicine," *Journal of Traditional and Complementary Medicine*, vol. 10, no. 1, pp. 7–12, 2020.
- [87] N. S. Chauhan, V. Sharma, V. K. Dixit, and M. Thakur, "A review on plants used for improvement of sexual performance and virility," *BioMed Research International*, vol. 2014, pp. 1–19, 2014.
- [88] A. A. Ajao, N. P. Sibiyi, and A. N. Moteetee, "Sexual prowess from nature: a systematic review of medicinal plants used as aphrodisiacs and sexual dysfunction in sub-Saharan Africa," *South African Journal of Botany*, vol. 122, pp. 342–359, 2019.
- [89] D. Tungmunthum, A. Thongboonyou, A. Pholboon, and A. Yangsabai, "Flavonoids and other phenolic compounds from medicinal plants for pharmaceutical and medical aspects: an overview," *Medicines*, vol. 5, no. 3, p. 93, 2018.
- [90] C. Tanase, S. Coşarçã, and D. L. Muntean, "A critical review of phenolic compounds extracted from the bark of woody vascular plants and their potential biological activity," *Molecules*, vol. 24, no. 6, p. 1182, 2019.
- [91] B. Van Wyk and P. Van Wyk, *Field Guide to Trees of Southern Africa*, Struik publishers Ltd, Pretoria, South Africa, 1997.
- [92] M. Ofosuhen, "Assessment of the effects of *Mondia whitei* extracts on some biochemical processes associated with penile erection," MPhil. Thesis, Department of Biochemistry, University of Ghana, Ghana, 2005.
- [93] J. Adjanohoum, N. Aboubaka, K. Dramane, N. E. Ebot, J. A. Ekpere, and E. G. Enow-Orock, *Traditional Medicine and Pharmacopoeia. Contribution to Ethnobotanical and Floristic Studies in Cameroon OUA/STRC*, Lagos, Nigeria, 1996.
- [94] E. Agbodjento, J. R. Klotoé, T. I. Sacramento et al., "Ethnobotanical knowledge of medicinal plants used in the treatment of male infertility in southern Benin," *Advances in Traditional Medicine*, vol. 21, no. 4, pp. 655–673, 2020.
- [95] R. Machado, I. Alves-Pereira, and R. Ferreira, "Plant growth, phytochemical accumulation and antioxidant activity of substrate-grown spinach," *Heliyon*, vol. 4, no. 8, 2018.
- [96] K. A. M. Kuria, S. De Coster, G. Muriuki et al., "Antimalarial activity of *Ajuga remota* Benth (Labiatae) and *Caesalpinia volkensii* Harms (Caesalpinaceae): in vitro confirmation of ethnopharmacological use," *Journal of Ethnopharmacology*, vol. 74, no. 2, pp. 141–148, 2001.
- [97] T. Omara, "Antimalarial plants used across Kenyan communities," *Evidence-Based Complementary and Alternative Medicine*, vol. 2020, Article ID 4538602, 31 pages, 2020.
- [98] Q. W. Zhang, L. G. Lin, and W. C. Ye, "Techniques for extraction and isolation of natural products: a comprehensive review," *Chinese Medicine*, vol. 13, no. 1, p. 20, 2018.
- [99] O. Joseph, E. A. Paul, A. Martin, K. T. Julius, O. C. E. Joseph, and G. A. Amon, "Effects of aqueous root bark extract of *Citropsis articulata* (Swingle Kellerman) on sexual function in male rats," *African Journal of Pharmacy and Pharmacology*, vol. 9, no. 29, pp. 723–729, 2015.
- [100] P. Vudriko, M. K. Baru, J. Kateregga, and J. G. Ndukupi, "Crude ethanolic leaf extracts of *Citropsis articulata*: a potential phytomedicine for treatment of male erectile dysfunction associated with testosterone deficiency," *International Journal of Basic & Clinical Pharmacology*, vol. 3, no. 1, pp. 120–123, 2014.
- [101] O. Joseph, J. K. Tanayen, K. Barbra et al., "Phytochemical and efficacy study on four herbs used in erectile dysfunction: *mondia whitei*, *Cola acuminata*, *Urtica massaiica*, and *Tarenna graveolens*," *African Journal of Pharmacy and Pharmacology*, vol. 10, no. 37, pp. 785–790, 2016.
- [102] Y. Gigani, A. Vekaria, and S. A. Amir, "Effect of *Abrus precatorius* and *Amaranthus spinosus* combination treatment on fertility in male rats," *Journal of Pharmacology and Pharmacotherapeutics*, vol. 3, no. 3, pp. 272–273, 2012.
- [103] M. Z. Allouh, H. M. Daradka, M. M. A. Barbarawi, and A. G. Mustafa, "Fresh onion juice enhanced copulatory behavior in male rats with and without paroxetine-induced sexual dysfunction," *Experimental Biology and Medicine*, vol. 239, no. 2, pp. 177–182, 2014.
- [104] O. I. Adeyemi, O. O. Ige, E. Agbede, and C. A. Adebajo, "The effect of tadalafil on the aphrodisiac properties of *Allium sativum* and *Allium cepa* using behavioural models in rats," *Nigerian Journal of Natural Products and Medicine*, vol. 21, pp. 14–20, 2017.
- [105] N. Malviya, S. Jain, V. B. Gupta, and S. Vyas, "Management of drug induced sexual dysfunction in male rats by ethyl acetate fraction of onion," *Acta Poloniae Pharmaceutica*, vol. 70, no. 2, pp. 317–322, 2013.
- [106] A. M. Al-Bekairi, A. H. Shah, and S. Qureshi, "Effect of *Allium sativum* on epididymal spermatozoa, estradiol-treated mice and general toxicity," *Journal of Ethnopharmacology*, vol. 29, no. 2, pp. 117–125, 1990.
- [107] V. W. Pooja, J. Saravanan, T. K. Praveen, R. Emdormi, and S. Deepa, "Evaluation of aphrodisiac activity of *Allium sativum* in male rats," *International Journal Of Scientific & Technology Research*, vol. 8, no. 10, pp. 1683–1686, 2019.
- [108] J. Yang, T. Wang, J. Yang et al., "S-allyl cysteine restores erectile function through inhibition of reactive oxygen species generation in diabetic rats," *Andrology*, vol. 1, no. 3, pp. 487–494, 2013.
- [109] M. R. Sailani and H. Moeini, "Effect of *Ruta graveolens* and *Cannabis sativa* alcoholic extract on spermatogenesis in the adult wistar male rats," *Indian Journal of Urology*, vol. 23, no. 3, pp. 257–260, 2007.
- [110] N. Lotfi, M. Khazaei, S. M. A. Shariatzadeh, M. Soleimani Mehranjani, and A. Ghanbari, "The effect of *Cannabis sativa* hydroalcoholic extract on sperm parameters and testis histology in rats," *International Journal of Morphology*, vol. 31, no. 1, pp. 82–86, 2013.
- [111] A. O. Ige, D. T. Oluwole, M. O. Olaoye, and E. O. Adewoye, "Testicular function following oral exposure to Tramadol and *Cannabis sativa* ethanol extracts in male Wistar rats,"

- Research Journal of Health Sciences*, vol. 8, no. 2, pp. 63–72, 2020.
- [112] M. Abdulwaheb, E. Makonnen, A. Debella, and D. Abebe, “Effect of *Catha edulis* foresk (khat) extracts on male rat sexual behavior,” *Journal of Ethnopharmacology*, vol. 110, no. 2, pp. 250–256, 2007.
- [113] S. A. Taha, A. M. Ageel, M. W. Islam, and O. T. Ginawi, “Effect of (-)-cathinone, a psychoactive alkaloid from khat (*Catha edulis* forsk.) and caffeine on sexual behaviour in rats,” *Pharmacological Research*, vol. 31, no. 5, pp. 299–303, 1995.
- [114] A. Mohammed and E. Engidawork, “Reproductive parameters are differentially altered following subchronic administration of *Catha edulis* F. (Khat) extract and cathinone in male rats,” *Journal of Ethnopharmacology*, vol. 134, no. 3, pp. 977–983, 2011.
- [115] M. W. Islam, M. Tariq, A. M. Ageel, F. El-Feraly, I. Al-Meshal, and I. Ashraf, “An evaluation of the male reproductive toxicity of cathinone,” *Toxicology*, vol. 60, no. 3, pp. 223–234, 1990.
- [116] O. O. Azu, F. I. O. Duru, A. A. Osinubi et al., “Histomorphometric effects of *Kigelia africana* (Bignoniaceae) fruit extract on the testis following short-term treatment with cisplatin in male Sprague–Dawley rats,” *Middle East Fertility Society Journal*, vol. 15, no. 3, pp. 200–208, 2010.
- [117] E. O. Adeparusi, A. A. Dada, and O. V. Alale, “Effects of medicinal plant (*kigelia africana*) on sperm quality of african catfish *Clarias gariepinus* (burchel, 1822) broodstock,” *Journal of Agricultural Science*, vol. 2, no. 1, pp. 193–199, 2010.
- [118] P. Watcho, P. Kamtchoung, S. D. Sokeng et al., “Androgenic effect of *Mondia whitei* roots in male rats,” *Asian Journal of Andrology*, vol. 6, no. 3, pp. 269–272, 2004.
- [119] P. Watcho, F. Zelefact, T. B. Nguelefack et al., “Effects of the aqueous and hexane extracts of *mondia whitei* on the sexual behaviour and some fertility parameters of sexually inexperienced male rats,” *African Journal of Traditional, Complementary, and Alternative Medicines: AJTCAM*, vol. 4, no. 1, pp. 37–46, 2006.
- [120] P. Watcho, D. Fotsing, F. Zelefact et al., “Effects of *Mondia whitei* extracts on the contractile responses of isolated rat vas deferens to potassium chloride and adrenaline,” *Indian Journal of Pharmacology*, vol. 38, no. 1, pp. 33–37, 2006.
- [121] P. Watcho, F. Zelefact, and T. B. Nguelefack, “Effects of the hexane extract of *mondia whitei* on the reproductive organs of male rat,” *African Journal of Traditional, Complementary and Alternative Medicines*, vol. 2, pp. 302–311, 2005.
- [122] G. M. Gundidza, V. M. Mmbengwa, M. L. Magwa et al., “Aphrodisiac properties of some Zimbabwean medicinal plants formulations,” *African Journal of Biotechnology*, vol. 8, no. 22, pp. 6402–6407, 2009.
- [123] O. Quasie, O. N. Martey, A. K. Nyarko, W. S. Gbewonyo, and L. K. Okine, “Modulation of penile erection in rabbits by *Mondia whitei*: possible mechanism of action,” *African Journal of Traditional, Complementary, and Alternative Medicines: AJTCAM*, vol. 7, no. 3, pp. 241–252, 2010.
- [124] O. T. Oyelowo, O. V. Fabiyi, O. M. Jimoh, and B. V. Owoyele, “Aphrodisiac and male sexual characteristics in albino rats treated with the aqueous extract of *Parquetina nigrescens* root,” *Nigerian Journal of Natural Products and Medicine*, vol. 16, pp. 18–25, 2012.
- [125] O. T. Kayode and M. T. Yakubu, “*Parquetina nigrescens* leaves: chemical profile and influence on the physical and biochemical indices of sexual activity of male Wistar rats,” *Journal of Integrative Medicine*, vol. 15, no. 1, pp. 64–76, 2017.
- [126] E. Ashamu, E. Salawu, O. Oyewo, A. Alhassan, O. Alamu, and A. Adegoke, “Efficacy of vitamin C and ethanolic extract of *Sesamum indicum* in promoting fertility in male Wistar rats,” *Journal of Human Reproductive Sciences*, vol. 3, no. 1, pp. 11–14, 2010.
- [127] D. M. Hai, J. W. Ren, Y. N. Chi et al., “Protective effects of sesamin on cytoxin-induced spermatogenesis dysfunction by regulating RNF8-ubH2A/ubH2B pathways in male mice,” *Frontiers in Pharmacology*, vol. 12, Article ID 708467, 2021.
- [128] B. Seven, H. Timur, M. N. Kalem, Z. Kalem, O. Han, and B. Bilezikci, “Ginger (*zingiber officinale*) might improve female fertility: a rat model,” *Journal of the Chinese Medical Association*, vol. 81, no. 10, pp. 905–911, 2018.
- [129] M. Shahidi, A. Moradi, and P. Dayati, “Zingerone attenuates zearalenone-induced steroidogenesis impairment and apoptosis in TM3 Leydig cell line,” *Toxicol*, vol. 211, pp. 50–60, 2022.
- [130] M. Salihu, B. O. Ajayi, I. A. Adedara, D. de Souza, J. B. T. Rocha, and E. O. Farombi, “6-Gingerol-rich fraction from *Zingiber officinale* ameliorates carbendazim-induced endocrine disruption and toxicity in testes and epididymis of rats,” *Andrologia*, vol. 49, no. 5, 2017.
- [131] P. Watcho, G. M. A. Tchuenchie, and D. P. B. Defo, “Sexual stimulant effects of the mixture of *Mondia whitei*, *Dracaena arborea*, and *Bridelia ferruginea* in normal and prediabetic male Wistar rats,” *Journal of Basic and Clinical Physiology and Pharmacology*, vol. 30, no. 4, 2019.
- [132] B. Ghazi Eid, A. Hanafy, A. Hasan, and T. Neamatalla, “Zingerone enhances fertility markers in both male and female rats and increases aryl hydrocarbon receptor expression,” *International Journal of Pharmacology*, vol. 16, no. 3, pp. 267–275, 2020.
- [133] S. A. Banihani, “Effect of ginger (*Zingiber officinale*) on semen quality,” *Andrologia*, vol. 51, no. 6, 2019.
- [134] S. A. Banihani, “Testosterone in males as enhanced by onion (*Allium cepa* L.),” *Biomolecules*, vol. 9, no. 2, p. 75, 2019.
- [135] D. Pizzol, J. Demurtas, B. Stubbs et al., “Relationship between Cannabis use and erectile dysfunction: a systematic review and meta-analysis,” *American Journal of Men’s Health*, vol. 13, no. 6, Article ID 155798831989246, 2019.
- [136] M. Gholami-Ahangan, M. Karimi-Dehkordi, A. Akbari Javar, M. Haj Salehi, and M. Ostadpoor, “A systematic review on the effect of Ginger (*Zingiber officinale*) on improvement of biological and fertility indices of sperm in laboratory animals, poultry and humans,” *Veterinary Medicine and Science*, vol. 7, no. 5, pp. 1959–1969, 2021.
- [137] Z. Afshar, S. Shahali, and H. Rastad, “Effects of oral ginger capsule on sexual function and sexual quality of life in women: a double-blinded, randomized, placebo-controlled trial,” *Sexologies*, vol. 31, no. 4, pp. 387–393, 2022.
- [138] J. Hosseini, A. Mardi Mamaghani, H. Hosseinfar, M. A. Sadighi Gilani, F. Dadkhah, and M. Sepidarkish, “The influence of ginger (*Zingiber officinale*) on human sperm quality and DNA fragmentation: a double-blind randomized clinical trial,” *International Journal of Reproductive Biomedicine*, vol. 14, no. 8, pp. 533–540, 2016.
- [139] V. Kuete, *Physical, Hematological, and Histopathological Signs of Toxicity Induced by African Medicinal Plants*, Toxicological Survey of African Medicinal Plants, Africa, 2014.
- [140] L. L. Chen, R. Verpoorte, H. R. Yen et al., “Effects of processing adjuvants on traditional Chinese herbs,” *Journal of Food and Drug Analysis*, vol. 26, no. 2, pp. S96–S114, 2018.

- [141] A. Karthikeyan and S. D. Amalnath, "Abrus precatorius poisoning: a retrospective study of 112 patients," *Indian Journal of Critical Care Medicine*, vol. 21, no. 4, pp. 224-225, 2017.
- [142] L. Reedman, R. D. Shih, and O. Hung, "Survival after an intentional ingestion of crushed abrus seeds," *The Western Journal of Emergency Medicine*, vol. 9, no. 3, pp. 157-159, 2008.
- [143] S. Alqasoumi, T. H. Khan, M. Al-Yahya, I. Al-Mofleh, and S. Rafatullah, "Effect of acute and chronic treatment of common spices in Swiss albino mice: a safety assessment study," *International Journal of Pharmacology*, vol. 8, no. 2, pp. 80-90, 2012.
- [144] B. Lawal, O. K. Shittu, F. I. Oibiokpa et al., "Antimicrobial evaluation, acute and sub-acute toxicity studies of *Allium sativum*," *Journal of Acute Disease*, vol. 5, no. 4, pp. 296-301, 2016.
- [145] M. Machado Bergamaschi, R. Helena Costa Queiroz, A. Waldo Zuardi, and J. Alexandre S Crippa, "Safety and side effects of cannabidiol, a *Cannabis sativa* constituent," *Current Drug Safety*, vol. 6, no. 4, pp. 237-249, 2011.
- [146] E. Muema, J. Mbaria, J. Nguta et al., "Toxicity and safety of khat (*Catha edulis*) consumption during pregnancy using olive baboons (*Papio anubis*) as experimental models: a prospective randomised study," *Greener Journal of Epidemiology and Public Health*, vol. 4, no. 3, pp. 061-070, 2016.
- [147] S. Atawodi and O. Olowoniyi, "Pharmacological and therapeutic activities of *kigelia africana* (lam.) benth," *Annual Research & Review in Biology*, vol. 5, pp. 1-17, 2015.
- [148] J. Oloro, J. K. Tanayen, K. Barbra et al., "Toxicity of four herbs used in erectile dysfunction; *Mondia whiteii*, *Cola acuminata*, *Urtica massaica*, and *Tarenna graveolensis* male rats," *African Journal of Pharmacy and Pharmacology*, vol. 9, no. 30, pp. 756-763, 2015.
- [149] L. Adu-Amoah, C. Agyare, E. Kisseih, P. G. Ayande, and K. B. Mensah, "Toxicity assessment of *Erythrophleum ivorense* and *Parquetina nigrescens*," *Toxicology Reports*, vol. 1, pp. 411-420, 2014.
- [150] K. C. Uruakpa, K. E. Obeten, M. Eru, and A. O. Igiri, "The toxicity study of ethanolic leaf extract of *Sesamum indicum* on the histomorphology of the liver of adult Wistar rats," *Global Journal of Biology, Agriculture and Health Sciences*, vol. 23, pp. 42-46, 2013.
- [151] T. Plengsuriyakarn, V. Viyanant, V. Eursitthichai et al., "Cytotoxicity, toxicity, and anticancer activity of *Zingiber officinale* Roscoe against cholangiocarcinoma," *Asian Pacific Journal of Cancer Prevention*, vol. 13, no. 9, pp. 4597-4606, 2012.
- [152] R. Chen, J. Lin, J. Hong et al., "Potential toxicity of quercetin: the repression of mitochondrial copy number via decreased POLG expression and excessive TFAM expression in irradiated murine bone marrow," *Toxicology Reports*, vol. 1, pp. 450-458, 2014.
- [153] P. Singh, S. Sharma, and S. K. Rath, "A versatile flavonoid Quercetin: study of its toxicity and differential gene expression in the liver of mice," *Phytomedicine Plus*, vol. 2, no. 1, Article ID 100148, 2022.
- [154] A. L. Riley, K. H. Nelson, P. To et al., "Abuse potential and toxicity of the synthetic cathinones (i.e., "Bath salts")," *Neuroscience & Biobehavioral Reviews*, vol. 110, pp. 150-173, 2020.
- [155] Y. Kodera, A. Suzuki, O. Imada et al., "Physical, chemical, and biological properties of s-allylcysteine, an amino acid derived from garlic," *Journal of Agricultural and Food Chemistry*, vol. 50, no. 3, pp. 622-632, 2002.
- [156] W. Kitipaspallop, P. Phuwapraisirisan, W. K. Kim, C. Chanchao, and W. Pimpong, "Sesamin lacks zebrafish embryotoxicity but exhibits evidence of anti-angiogenesis, anti-oxidant and anti-inflammatory activities," *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, vol. 269, Article ID 109637, 2023.
- [157] V. A. S. Jesudoss, S. V. A. Santiago, K. Venkatachalam, and P. Subramanian, "Zingerone (ginger extract): antioxidant potential for efficacy in gastrointestinal and liver disease," in *Gastrointestinal Tissue. Oxidative Stress and Dietary Anti-oxidants*, pp. 289-297, 2017.
- [158] N. Promdam and P. Panichayupakaranant, "[6]-Gingerol: a narrative review of its beneficial effect on human health," *Food Chemistry Advances*, vol. 1, Article ID 100043, 2022.