Case Report

Calcifying Bowel Inflammation: A Case Report

Jacques Klein,¹ Philippe Morel,¹ and Christian Toso²

¹Abdominal Surgery, Department of Surgery, University Hospitals of Geneva, rue Gabrielle-Perret-Gentil 4, 1211 Genève 14, Switzerland

²Abdominal and Transplant Surgery, Department of Surgery, University Hospitals of Geneva, rue Gabrielle-Perret-Gentil 4, 1211 Genève 14, Switzerland

Correspondence should be addressed to Jacques Klein, jacques.klein@hcuge.ch

Received 28 December 2009; Accepted 11 March 2010

Academic Editor: Karel R. Geboes

Copyright © 2010 Jacques Klein et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We report about a previously healthy 72 year-old woman, presented with 6 days of left lower quadrant abdominal pain and constipation. There was no report of fever, melena, hematochezia or change in appetite. The physical exam demonstrated a distended abdomen with palpable left lower quadrant pain, without guarding. CT showed images compatible with a sigmoid diverticulitis and a calcification of the sigmoid colon. After antibiotic treatment, a colonoscopy was performed which revealed the presence of a shell in the sigmoid colon. Our case illustrates the need for a colonoscopy following an attack of diverticulitis to look for a cancer or rarely a foreign body.

A previously healthy 72-year-old woman presented with 6 days of left lower quadrant abdominal pain and constipation. There were no report of fever, melena, hematochezia or change in appetite. The physical exam demonstrated a distended abdomen with palpable left lower quadrant pain, without guarding. The blood work revealed a normal white blood cell count but an increased C-reactive protein (110 mg/L).

CT showed images compatible with a sigmoid diverticulitis and a calcification of the sigmoid colon (Figure 1(a), arrow).

The patient was treated with IV antibiotics (Ceftriaxone 2 g daily and Metronidazole 500 mg three times daily), with an adequate resolution of her signs and symptoms.

A colonoscopy performed one week later revealed the presence of a shell in the sigmoid colon (at 25 cm from the anus), which was extracted without complication (Figures 1(b) and 1(c)). The patient subsequently confessed to have eaten shellfish in paella the week prior to admission.

Diverticulosis is a common illness in developed countries where it is present in 50% of people over the age of 60.

The sigmoid colon is the most commonly affected. Approximately 10% of patients, who present with diverticulosis, will develop diverticulitis, usually caused by a fecalith.

The literature reports a few cases of diverticulitis caused by the presence of a foreign body (generally chicken or fish bones) [1, 2]. In 88% of cases the patients are unconscious about the ingested foreign body; furthermore, in 76% of cases the foreign body passes spontaneously without complication [2, 3]. Though rarely, the actual incidence of diverticulitis caused by a foreign body is unknown. The predisposing risk factor for ingestion is the use of dentures which reduces the sensibility of the palate (72%); however, this was not the case for our patient [2]. About 10%–20% of all ingested foreign bodies must be removed endoscopically and 1%–14% need operative removal [3].

Our case illustrates the need for a colonoscopy following an attack of diverticulitis to look for a cancer or rarely a foreign body. The patient left hospital after completing antibiotic treatment without complication. She remained asymptomatic three months after discharge.

FIGURE 1

References

- [1] J. Lubel and M. Wiley, "Images of interest: foreign bodies and diverticulitis," *Journal of Gastroenterology and Hepatology*, vol. 20, no. 4, p. 649, 2005.
- [2] J. I. Rodríguez-Hermosa, A. Codina-Cazador, J. M. Sirvent, A. Martín, J. Gironès, and E. Garsot, "Surgically treated perforations of the gastrointestinal tract caused by ingested foreign bodies," *Colorectal Disease*, vol. 10, no. 7, pp. 701–707, 2008.
- [3] N. G. Velitchkov, G. I. Grigorov, J. E. Losanoff, and K. T. Kjossev, "Ingested foreign bodies of the gastrointestinal tract: retrospective analysis of 542 cases," *World Journal of Surgery*, vol. 20, no. 8, pp. 1001–1005, 1996.