Research Article

Dietary Factors in Relation to *Helicobacter pylori* Infection

Seyyed Ali Mard,¹ Hossein Khadem Haghighian,² Vahid Sebghatullahi,³ and Bijan Ahmadi⁴

¹Physiology Research Center (PRC) and Department of Physiology, Research Institute for Infectious Diseases of Digestive System, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz 61357-15794, Iran

²Department of Nutrition, Research Institute for Infectious Diseases of Digestive System, School of Paramedical, Ahvaz Jundishapur University of Medical Sciences, Ahvaz 61357-15794, Iran

³Research Institute for Infectious Diseases of Digestive System, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz 61357-15794, Iran

⁴Research Institute for Infectious Diseases of Digestive System, School of Medicine, Kerman University of Medical Sciences, Kerman 7618747653, Iran

Correspondence should be addressed to Hossein Khadem Haghighian; khademnut@yahoo.com

Received 6 September 2014; Revised 9 November 2014; Accepted 9 November 2014; Published 2 December 2014

Academic Editor: Sergio Morini

Copyright © 2014 Seyyed Ali Mard et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background and Aim. *Helicobacter pylori* (HP) and diet are both risk factors for gastric cancer. The aim of this study was to evaluate the *Helicobacter pylori* infection and dietary habits common in Khuzestan province. **Methods.** This cross-sectional study was conducted in 2011–2013 on 374 patients. Participants were interviewed using a food frequency questionnaire and tissue sample of the antrum was sent for pathology lab. The histopathological major variables were graded on a scale of 3 (mild, moderate, and severe) and data analyzed using nonparametric tests. **Results.** In this study, of 160 patients (43%) that were determined, 8.1 percent had severe contamination. Among dietary patterns, relationship between energy intake and carbohydrate with *H. pylori* was significant. A direct association was found between mean daily intakes of sausage ($P = 0.001$) and burgers ($P < 0.05$) with HP infection. Low intake of fresh vegetables and fruits was the most significant risk factors ($P < 0.05$). **Conclusion.** There is a possibility that some dietary factors such as consumption of fast foods and low intake of fresh vegetables may increase the chance of HP and severity of this infection.

1. Introduction

Helicobacter pylori is a spiral, gram negative, acid tolerant, microaerophilic bacterium that lives in the stomach and duodenum [1, 2]. In Iran, *H. pylori* infection is present in nearly 90% of adult population [3] and appears to occur early in life, with >50% of children infected before age of 15 [4]. Despite the fact that the incidence of and mortality from gastric cancer have declined markedly worldwide over the past decades, gastric cancer is still the second most common cause of cancer-related death in the world [5]. Epidemiological data suggest that environmental factors are the predominant cause of this disease. The most important factors thought to be responsible for GC development are diet and *Helicobacter pylori* infection [6].

Besides the fact that *H. pylori* was introduced as a class I carcinogen [7], the infection is difficult to cure and requires various combination therapies [8]. Previous epidemiological studies have suggested that not only *H. pylori* infection but also varieties of environmental factors are important risk factors for GC [9–12]. It is believed that dietary factors may contribute to the *H. pylori* infection [13]. Adequate nutritional status, especially high consumption of fruits, vegetables, and vitamins, appears to protect against the pathological consequences of *H. pylori* infection [14]. Knoop et al. stressed the role of vitamin C as a chemopreventive factor in *H. pylori* gastric disorders [15, 16].

Furthermore, which environmental factor is involved in the development of GC among persons infected with *H. pylori* has rarely been investigated [17]. Therefore, we also

investigated this point among *H. pylori*-infected subjects. The aim of this study was to evaluate whether there are differences in dietary habits between patients who were infected and subjects who were not infected.

2. Materials and Methods

The study was carried out in 2011–2013 in a group of patients from the Gastroenterological Unit of the Emam Khomeini Hospital in Ahvaz who had functional dyspepsia and had been referred for endoscopic examination of the upper digestive tract. To minimize any known confounding effects, the subjects with the following conditions have been excluded from the study: subjects with habits such as smoking, tobacco chewing, and alcohol consumption; on corticosteroid therapy at pharmacological levels for duration of more than 6 months; diabetes mellitus (except for easily controlled, non-insulin-dependent diabetes mellitus).

For all patients, energy and nutrient intakes were measured using a 3-day 24 h recall (two week days and one weekend day). Three-day food diary form contains foods at every meal, snack, and the amount consumed per person. A validated food frequency questionnaire [18] was also used to determine dietary habits during the previous year. The information was obtained through face to-face interviews, with standard food models, and a variety of measuring tools to evaluate intake. To help people to recall precise quantities of food eaten homemade dishes and cups were accounted. The participants were asked to indicate consumption of specified food in daily, weekly, monthly, and yearly bases. The selected frequency choice, given by the subjects for each food item from the food groups, was then converted to a weekly intake. All subjects were asked about the average frequency of intake and portion size of 125 items over the past year. The frequency of other food items was classified into 9 categories (never, 1–3/month, 1–2, 3–4, or 5–6/week, almost once/day, or 2–3, 4–6, or 7+ times/day).

Values of each food were converted to grams by using household measures. An interview on dietary habits was performed by a dietician. Nutrients were analyzed by Nutritionist IV software (N-Squared computing, Salem, OR, USA), which was modified for Iranian foods.

After upper GI endoscopy a tissue sample from antrum, body, and fundus in plate containing formalin buffer was sent to the pathology lab. All samples were examined by a pathologist. Contamination detection was performed with hematoxylin eosin (H&E). Semiquantitative method of scoring according to the Updated Sydney Classification System was undertaken. Statistical analysis was done using SPSS software (version 13). To correlate variables with normal and nonnormal distribution, Pearson's and Spearman's correlations, respectively, were used. All tests were two-tailed, and $P < 0.05$ was the significance threshold.

3. Result

In this study, 374 patients were evaluated, 182 patients (48%) of them were females and 192 (52%) were males. Based

TABLE 1: Nutrient intakes of healthy group and patient group per day.

Dietary factors	Healthy subjects group Mean $\pm$ SD	Patient group Mean $\pm$ SD	Pv
Calories (kcal/d)	2110 $\pm$ 514	2491 $\pm$ 415	0.01
Carbohydrate (g/d)	300.01 $\pm$ 101.4	384.9 $\pm$ 109.02	0.02
Protein (g/d)	79.6 $\pm$ 21.3	81.4 $\pm$ 27.3	0.837
Fat (g/d)	65.4 $\pm$ 28.9	96.6 $\pm$ 31.5	0.279
PUFA (g/d)	24.31 $\pm$ 12.9	10.51 $\pm$ 11.7	0.02

TABLE 2: Food frequency data on meat samples from healthy group and patient group (per a week).

Meat	Healthy subjects group mean (minimum and maximum)	Patient group mean (minimum and maximum)	P
Mutton	1.25 (0.02, 4.7)	1.27 (0.01, 5)	0.714
Veal	2.23 (0.27, 7)	2.1 (0.2, 8.1)	0.317
Chicken	3 (0.26, 7.1)	3 (0.23, 7)	0.894
Fish	3 (0.47, 7)	1 (0.02, 4)	0.001
Egg	2 (0.04, 7)	2 (0.07, 8)	0.511
Sausage	1 (0.04, 3)	3 (0.08, 9.2)	0.001
Hamburger	0.56 (0.02, 3)	2 (0.08, 7)	0.001

on pathological studies on *H. pylori* infection, 214 healthy subjects (57%) and 160 patients (43%) were diagnosed that 8% of them had severe contamination and 37.5% and 54.4% had moderate and mild contamination, respectively.

As it is shown in Table 1, there was a significant correlation between energy and carbohydrate intake with *H. pylori* infection. ($P = 0.01, 0.02$) There was also a negative and significant correlation between daily intake of fish ($P = 0.001$), olive oil ($P = 0.002$) and honey ($P = 0.002$), and peas and beans ($P = 0.03$) with *H. pylori* infection.

Table 2 shows that there was positive relation between sausages ($P = 0.001$), hamburgers ($P = 0.002$), fat mayonnaise ($P = 0.002$), and soft drinks ($P = 0.001$) with *H. pylori* infection.

Taking weekly tomatoes ($P = 0.001$), onions ($P = 0.002$), green pepper ($P = 0.01$), apple ($P = 0.002$), and citrus fruits ($P = 0.001$) was significantly lower than in healthy individuals (Table 3).

Among the micronutrients reported, there were significant differences in intake of vitamin C and folate between the healthy subjects and patient groups (Table 4).

Subjects with *H. pylori* infection were categorized into 3 groups: severe, moderate, and mild contamination (on Revised Sydney System).

As it is shown in Table 5, there were negative and significant correlation and significant difference in tomatoes, onions, green pepper, apple, citrus fruits, fish, olive oil, and honey and peas intake with severity of *H. pylori* infection.

TABLE 3: Food frequency data of Fruits and vegetables intakes of Healthy group and Patient group (per a week).

Vegetables	Healthy subjects group mean (minimum and maximum)	Patient group mean (minimum and maximum)	Pv
Carrot	2 (6, 0.32)	1 (0.7, 2.2)	0.232
Tomato	6 (0.12, 21)	3 (0.15, 20.8)	0.001
Onion	7 (0.46, 21)	3 (0.23, 8)	0.002
Green pepper	3 (0.04, 7)	1 (0.05, 6.09)	0.01
Apple	4 (0.46, 10.2)	1 (0.11, 3)	0.002
Citrus fruits	4 (0.1, 9)	2 (0.03, 6)	0.001

4. Discussion

Epidemiological studies have shown that *H. pylori* is probably one of the most common bacterial infections throughout the world, involving 30% of the population living in developed countries and up to 80%–90% of the population in developing countries [19].

The treatment of *H. pylori* is difficult, requires a two-week application of at least three medicines (proton pump inhibitors and two antibiotics) simultaneously, proves successful in only 80%–90% of cases, and is connected with the risk of adverse effects of therapy with antibiotics (15%–30% of the treated) [20, 21].

A synergistic interaction between *H. pylori* infection and diet has been suggested [22]. In this research, the dietary factors were evaluated. The aim of this retrospective study was to investigate potential differences in the dietary habits of patients with *H. pylori* infection (group I) and in the control not-infected group (group II). We demonstrated a significantly higher consumption of fruit, vegetables, and vitamin C among persons who were not infected. Vitamin C is highly concentrated in stomach mucosa and gastric juice and probably lowers the risk of gastric cancer and influences the course of *H. pylori* infection through a number of mechanisms [23, 24]. It has a positive impact on the stimulation and activity of granulocytes, macrophages, and lymphocytes and the production of immunoglobulin. The direct inhibitory impact of this vitamin on the growth of *H. pylori* is now being examined. Jarosz et al. [13] showed that four-week treatment of *H. pylori* infected patients with chronic gastritis with a high dose of vitamin C caused *H. pylori* eradication in 30% of cases. In those patients, a highly significant rise in gastric juice total vitamin C concentration was demonstrated, which persisted for at least four weeks after treatment. Ruiz et al. [25] found a causal association between *H. pylori* infection and low ascorbic acid levels in the gastric juice. Their findings supported two hypotheses that explain this phenomenon: increased oxidation and decreased secretion of ascorbic acid.

Also diet may provide noxious agents that contribute to *H. pylori* pathogenicity or protective agents that hamper its activity as related to the appearance of cancer diseases. Production of nitrosylating species plays a fundamental role in

TABLE 4: Nutrient intakes of healthy group and patient group per day.

Micronutrient	Healthy subjects group	Patient group	Pv
Folate (μg)	392 ± 170.03	302 ± 131	0.01
Vitamin A (mg)	530 ± 491.1	517 ± 483	0.912
Vitamin C (mg)	88 ± 71.02	58 ± 53.9	0.02
Vitamin E (mg)	5.98 ± 1.03	6.74 ± 0.85	0.421
Zn (mg)	4.5 ± 3.9	4.2 ± 3.7	0.781

this regard. Observational cohort studies should ideally provide much more reliable evidence. Analysis of the data substantially confirmed the significant increased risk of developing GC due to high intake of total carbohydrates, processed meat, refined grains, and saturated fat [26, 27]. Prospective Investigation into Cancer and Nutrition study reported a significant increase of noncardiac cancer risk associated with intake of total meat, red meat, and processed meat [28].

Previous studies have reported the protection role of allium vegetables (onion) in gastric cancer [27–29]. In this study, we also observed negative significant relation between onion consumption and *H. pylori* infection.

In this study there was positive relation between carbohydrate, sausages ($P = 0.001$), hamburgers ($P = 0.002$), fat mayonnaise ($P = 0.002$), and soft drinks intake ($P = 0.001$) with *H. pylori* infection.

A high salt concentration in the stomach destroys the mucosal barrier, favors colonization by *H. pylori*, and leads to inflammation and damage-causing gastritis and diffuse erosion [30]. Processed meats have high salt concentration that contributes to *H. pylori* pathogenicity.

We observed a significant difference in intake of unsaturated fatty acid between two groups. It is thought that unsaturated fatty acid may inhibit gastric carcinogenesis. This inhibitory role was experimentally shown in mice [31]. An inverse association was reported between consumption of unsaturated fat and *H. pylori* with a significant dose dependency [32].

Although information on a potentially protective effect of vegetable oil consumption (unsaturated fat) or regarding specific types of unsaturated fatty acids and gastric cancer is very limited, but, in some studies, protective effect of vegetable oil consumption in gastric cancer has been reported [33].

Some studies of healthy adults demonstrated subclinical deficiency of some B vitamins and high prevalence of *H. pylori* infection [34].

H. pylori stimulates the macrophage system through the l-arginine/nitric oxide (NO) pathway [35]. As a consequence, chronic *H. pylori* infection of the human stomach may increase endogenous NO formation, yielding, after oxidation, the nitrosating agents N_2O_3 and N_2O_4 , which can produce nitrosamines or cause other types of DNA damage [36]. Nitrosating organisms are capable of catalysing a reaction between nitrite and other organic nitrogen compounds present in the gastric juice to form potent genotoxic N-nitroso compounds (NOC) [37]. Nitrite levels in the gastric juice are increased during hypochlorhydria, which typically occurs during *H. pylori*-induced atrophic gastritis. This may

TABLE 5: Nutrient intakes of subject without infection and patients with different infection.

Food item	Without infection	Mild infection	Moderate infection	Severe infection	Spearman's rho	Pv
Calories (kcal/d)	2110 ± 0.514	2230 ± 371.51	2532 ± 421.83	2711 ± 451.65	0.363	0.04
Carbohydrate (g/d)	300.01 ± 101.14	343.76 ± 97.36	390.3 ± 110.5	417 ± 118.1	0.327	0.037
Tomato*	6 ± 4.64	3.26 ± 1.53	3.04 ± 1.42	2.68 ± 1.26	-0.358	0.033
Onion*	7 ± 4.6	3.80 ± 1.292	3.54 ± 1.203	3.12 ± 1.06	-0.312	0.038
Green pepper*	3 ± 0.88	1.62 ± 0.47	1.51 ± 0.44	1.33 ± 0.39	-0.371	0.039
Apple*	4 ± 2.33	±1.5 ± 0.57	1.42 ± 0.53	1.22 ± 0.46	-0.311	0.027
Citrus fruits*	4 ± 2.133	1.90 ± 0.72	1.80 ± 0.68	1.55 ± 0.58	-0.382	0.042
Pea*	0.2 ± 0.77	0.17 ± 0.16	0.16 ± 0.1	0.13 ± 0.08	-0.527	0.003
Fish*	3 ± 1.93	2.55 ± .012	2.4 ± 0.128	1.95 ± 0.09	-0.348	0.001
Sausage*	1 ± 0.38	0.85 ± 0.32	1.65 ± .062	2.3 ± 0.87	0.339	0.011
Hamburgers*	0.56 ± 0.37	0.47 ± 0.31	0.91 ± 0.6	1.27 ± 0.83	0.312	0.021
Fat mayonnaise*	1 ± 0.51	0.83 ± 0.41	1.61 ± 0.8	2.25 ± 1.12	0.325	0.013
Olive and olive oil*	3 ± 1.02	2.49 ± 0.84	2.34 ± 0.79	1.92 ± 0.65	-0.351	0.031
Soft drink*	0.9 ± 0.51	0.74 ± 0.41	1.442 ± 0.81	2.012 ± 1.14	0.321	0.017
Honey*	2 ± 0.88	1.64 ± 0.72	1.56 ± 0.68	1.26 ± 0.55	-0.329	0.003

* Per a week.

result from bacteria within the stomach generating nitrite from dietary nitrate [38].

In present study among the micronutrients reported, there were significant differences in intake of folate between the healthy subjects and patient groups.

It has been established that chronic *H. pylori* infection causes atrophic gastritis and decreased absorption of vitamin folic acid in patients with this condition. Because methylation of homocysteine to methionine requires folate, an increased plasma homocysteine level is found in *H. pylori*-infected patients [14]. Our study has several potential limitations. First, we did not collect information about lifestyle. Second, our results could also be affected by measurement error in dietary intake, a common limitation of cross-sectional studies. However, prospective cohorts study is required, which evaluates the interaction between dietary habits and *H. pylori* infection. Third limitation was that level of education, socioeconomic, and race of subjects are not considered.

5. Conclusion

In summary, the present study on food interaction and *H. pylori*-infection displayed that a diet rich in fruit and vegetables and poor in meat, fat, and salt has a good prophylactic potential for cancer. Fruits and vegetables are sources (among other protective agents) of ascorbic acid, which were associated with a high effect against the progression of gastric carcinogenesis.

Thus, "diet for cancer prevention" can be recommended as a general role of well-being and can represent the basis for a rational health policy.

Conflict of Interests

None of the authors have any conflict of interests associated with this study.

References

- [1] M. J. Blaser and J. C. Atherton, "Helicobacter pylori persistence: biology and disease," *The Journal of Clinical Investigation*, vol. 113, no. 3, pp. 321–333, 2004.
- [2] C. Montecucco and R. Rappuoli, "Living dangerously: how *Helicobacter pylori* survives in the human stomach," *Nature Reviews Molecular Cell Biology*, vol. 2, no. 6, pp. 457–466, 2001.
- [3] R. Malekzadeh, M. Sotoudeh, M. H. Derakhshan et al., "Prevalence of gastric precancerous lesions in Ardabil, a high incidence province for gastric adenocarcinoma in the northwest of Iran," *Journal of Clinical Pathology*, vol. 57, no. 1, pp. 37–42, 2004.
- [4] J. Mikaeili, R. Malekzadeh, A. Z. B. Ziad et al., "Prevalence of *Helicobacter pylori* in two Iranian provinces with high and low incidence of gastric carcinoma," *Archives of Iranian Medicine*, vol. 3, no. 1, 2000.
- [5] D. M. Parkin, F. Bray, J. Ferlay, and P. Pisani, "Global cancer statistics, 2002," *A Cancer Journal for Clinicians*, vol. 55, no. 2, pp. 74–108, 2005.
- [6] A. Rocco and G. Nardone, "Diet, *H. pylori* infection and gastric cancer: evidence and controversies," *World Journal of Gastroenterology*, vol. 13, no. 21, pp. 2901–2912, 2007.
- [7] Working Group Report on Schistosomes, *Liver Flukes and Helicobacter pylori*, Lyon, France, 1994.
- [8] N. E. Daryani, M. Taher, and S. Shirzad, "Helicobacter pylori infection: a review," *Iranian Journal of Clinical Infectious Diseases*, vol. 6, no. 1, pp. 56–64, 2011.
- [9] Y. Watanabe, J. H. Kurata, S. Mizuno et al., "Helicobacter pylori infection and gastric cancer (A nested case-control study in a rural area of Japan)," *Digestive Diseases and Sciences*, vol. 42, no. 7, pp. 1383–1387, 1997.
- [10] K. Ozasa, J. H. Kurata, A. Higashi et al., "Helicobacter pylori infection and atrophic gastritis: a nested case-control study in a rural town in Japan," *Digestive Diseases and Sciences*, vol. 44, no. 2, pp. 253–256, 1999.
- [11] K. Shibata, M. Moriyama, T. Fukushima, H. Ume, M. Miyazaki, and N. Yamaguchi, "Relation of *Helicobacter pylori* infection

- and lifestyle to the risk of chronic atrophic gastritis: a cross-sectional study in Japan," *Journal of Epidemiology*, vol. 12, no. 2, pp. 105–111, 2002.
- [12] Y. Kuwahara, S. Kono, H. Eguchi, H. Hamada, K. Shinchi, and K. Imanishi, "Relationship between serologically diagnosed chronic atrophic gastritis, *Helicobacter pylori*, and environmental factors in Japanese men," *Scandinavian Journal of Gastroenterology*, vol. 35, no. 5, pp. 476–481, 2000.
- [13] M. Jarosz, E. Rychlik, H. Siuba et al., "Dietary and socio-economic factors in relation to *Helicobacter pylori* re-infection," *World Journal of Gastroenterology*, vol. 15, no. 9, pp. 1119–1125, 2009.
- [14] A. Izzotti, P. Durando, F. Ansaldi, F. Gianiorio, and A. Pulliero, "Interaction between *Helicobacter pylori*, diet, and genetic polymorphisms as related to non-cancer diseases," *Mutation Research: Fundamental and Molecular Mechanisms of Mutagenesis*, vol. 667, no. 1-2, pp. 142–157, 2009.
- [15] K. T. B. Knuops, L. C. P. G. M. De Groot, D. Kromhout et al., "Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women: the HALE project," *Journal of the American Medical Association*, vol. 292, no. 12, pp. 1433–1439, 2004.
- [16] E. Riboli, N. Slimani, and R. Kaaks, "Identifiability of food components for cancer chemoprevention," *IARC Scientific Publications*, no. 139, pp. 23–31, 1996.
- [17] A. Montani, S. Sasazuki, M. Inoue, K. Higuchi, T. Arakawa, and S. Tsugane, "Food/nutrient intake and risk of atrophic gastritis among the *Helicobacter pylori*-infected population of northeastern Japan," *Cancer Science*, vol. 94, no. 4, pp. 372–377, 2003.
- [18] A. F. Malekshah, M. Kimiagar, M. Saadatian-Elahi et al., "Validity and reliability of a new food frequency questionnaire compared to 24h recalls and biochemical measurements: pilot phase of Golestan cohort study of esophageal cancer," *European Journal of Clinical Nutrition*, vol. 60, no. 8, pp. 971–977, 2006.
- [19] J. Salgueiro, M. Zubillaga, C. Goldman et al., "Review article: is there a link between micronutrient malnutrition and *Helicobacter pylori* infection?" *Alimentary Pharmacology and Therapeutics*, vol. 20, no. 10, pp. 1029–1034, 2004.
- [20] P. Malfertheiner, F. Megraud, C. O'Morain et al., "Current concepts in the management of *Helicobacter pylori* infection: the Maastricht III Consensus Report," *Gut*, vol. 56, no. 6, pp. 772–781, 2007.
- [21] J. Dzieniszewski and M. Jarosz, "Guidelines in the medical treatment of *Helicobacter pylori* infection," *Journal of Physiology and Pharmacology*, vol. 57, no. 3, pp. 143–154, 2006.
- [22] N. Yamaguchi and T. Kakizoe, "Synergistic interaction between *Helicobacter pylori* gastritis and diet in gastric cancer," *The Lancet Oncology*, vol. 2, no. 2, pp. 88–94, 2001.
- [23] M. Jarosz, J. Dzieniszewski, E. Dabrowska-Ufniaz, M. Wartanowicz, S. Ziemiński, and P. Reed I, "Effects of high dose vitamin C treatment on *Helicobacter pylori* infection and total vitamin C concentration in gastric juice," *European Journal of Cancer Prevention*, vol. 7, no. 6, pp. 449–454, 1998.
- [24] L.-Q. Shi and R.-L. Zheng, "DNA damage and oxidative stress induced by *Helicobacter pylori* in gastric epithelial cells: protection by vitamin C and sodium selenite," *Pharmazie*, vol. 61, no. 7, pp. 631–637, 2006.
- [25] B. Ruiz, J. C. Rood, E. T. H. Fonstam et al., "Vitamin C concentration in gastric juice before and after anti-*Helicobacter pylori* treatment," *The American Journal of Gastroenterology*, vol. 89, no. 4, pp. 533–539, 1994.
- [26] R. W. Kneller, J. K. McLaughlin, E. Bjelke et al., "A cohort study of stomach cancer in a high-risk American population," *Cancer*, vol. 68, no. 3, pp. 672–678, 1991.
- [27] M. C. J. F. Jansen, H. B. Bueno-de-Mesquita, L. Räsänen et al., "Consumption of plant foods and stomach cancer mortality in the seven countries study. Is grain consumption a risk factor?" *Nutrition and Cancer*, vol. 34, no. 1, pp. 49–55, 1999.
- [28] C. A. González, P. Jakszyn, G. Pera et al., "Meat intake and risk of stomach and esophageal adenocarcinoma within the European Prospective Investigation Into Cancer and Nutrition (EPIC)," *Journal of the National Cancer Institute*, vol. 98, no. 5, pp. 345–354, 2006.
- [29] N. Muñoz, M. Plummer, J. Vivas et al., "A case-control study of gastric cancer in Venezuela," *International Journal of Cancer*, vol. 93, no. 3, pp. 417–423, 2001.
- [30] M. L. Caruso and L. Fucci, "Histological identification of *Helicobacter pylori* in early and advanced gastric cancer," *Journal of Clinical Gastroenterology*, vol. 12, no. 5, pp. 601–602, 1990.
- [31] R. A. Karmali, J. Marsh, and C. Fuchs, "Effect of omega-3 fatty acids on growth of a rat mammary tumor," *Journal of the National Cancer Institute*, vol. 73, no. 2, pp. 457–461, 1984.
- [32] L. López-Carrillo, M. López-Cervantes, M. H. Ward, J. Bravo-Alvarado, and A. Ramírez-Espitia, "Nutrient intake and gastric cancer in Mexico," *International Journal of Cancer*, vol. 83, no. 5, pp. 601–605, 1999.
- [33] E. Buiatti, D. Palli, A. Decarli et al., "A case-control study of gastric cancer and diet in Italy," *International Journal of Cancer*, vol. 44, no. 4, pp. 611–616, 1989.
- [34] Y. L. Domínguez, M. Hernández, C. M. Matos, and D. Zhou, "Is B vitamins deficiency associated with prevalence of Alzheimer's disease in Cuban elderly?" *Nutrition and Health*, vol. 18, no. 2, pp. 103–118, 2006.
- [35] K. B. Shapiro and J. H. Hotchkiss, "Induction of nitric oxide synthesis in murine macrophages by *Helicobacter pylori*," *Cancer Letters*, vol. 102, no. 1-2, pp. 49–56, 1996.
- [36] R. H. Liu and J. H. Hotchkiss, "Potential genotoxicity of chronically elevated nitric oxide: a review," *Mutation Research/Reviews in Genetic Toxicology*, vol. 339, no. 2, pp. 73–89, 1995.
- [37] D. Ziebarth, B. Spiegelhalter, and H. Bartsch, "N-nitrosation of medicinal drugs catalysed by bacteria from human saliva and gastro-intestinal tract, including *Helicobacter pylori*," *Carcinogenesis*, vol. 18, no. 2, pp. 383–389, 1997.
- [38] C. Mowat, A. Carswell, A. Wirz, and K. E. L. McColl, "Omeprazole and dietary nitrate independently affect levels of vitamin C and nitrite in gastric juice," *Gastroenterology*, vol. 116, no. 4, pp. 813–822, 1999.