Research Article

Expression of *Helicobacter pylori* *hspA* Gene in *Lactococcus lactis* NICE System and Experimental Study on Its Immunoreactivity

Xiao-Juan Zhang,¹ Shu-Ying Feng,² Zhi-Tao Li,² and Yan-Ming Feng³

¹Department of Pathogen Biology, Medical College, Henan University of Science and Technology, Luoyang 471003, China

²Department of Immunology, Medical College, Henan University of Science and Technology, Luoyang 471003, China

³Department of Epidemiology, Medical College, Henan University of Science and Technology, Luoyang 471003, China

Correspondence should be addressed to Xiao-Juan Zhang; gjfzxj0921@gmail.com

Received 5 May 2014; Accepted 3 March 2015

Academic Editor: Spiros D. Ladas

Copyright © 2015 Xiao-Juan Zhang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Aim. The aim of this study was to develop an oral *Lactococcus lactis* (*L. lactis*) vaccine against *Helicobacter pylori* (*H. pylori*). **Methods.** After *L. lactis* NZ3900/pNZ8110-*hspA* was constructed, growth curves were plotted to study whether the growth of recombinant *L. lactis* was affected after *hspA* was cloned into *L. lactis* and whether the growth of empty bacteria, empty plasmid bacteria, and recombinant *L. lactis* was affected by different concentrations of Nisin; SDS-PAGE and Western blot were adopted, respectively, to detect the HspA expressed by recombinant *L. lactis* and its immunoreactivity. **Results.** There was no effect observed from the growth curve after exogenous gene *hspA* was cloned into *L. lactis* NZ3900; different concentrations of Nisin did not affect the growth of NZ3900 and NZ3900/pNZ8110, while different concentrations of Nisin inhibited the growth of NZ3900/pNZ8110-*hspA* except 10 ng/mL Nisin. No HspA strip was observed from SDS-PAGE. Western blot analysis showed that HspA expressed by recombinant bacteria had favorable immunoreactivity. **Conclusion.** The growth of recombinant *L. lactis* was suppressed even though a small amount of HspA had been induced to express. Therefore recombinant *L. lactis* only express HspA which was not suitable to be oral vaccine against *Helicobacter pylori*.

1. Introduction

Helicobacter pylori (*H. pylori*) is a spiral and motile Gram-negative bacterium, which colonizes the human stomach mucosa and is the main cause of a range of gastric and duodenal diseases including chronic gastritis, stomach and duodenal ulcers, MALT lymphoma, and gastric adenocarcinoma, and it is considered as the second most frequent cause of cancer death worldwide [1, 2]. *H. pylori* is also the pathogenic microorganism which has relationship with diabetes mellitus, coronary artery disease, thrombocytopenic purpura, and so on [3–5]. Heat shock proteins (Hsp) of *H. pylori* play important roles in *H. pylori* adhesion to gastric epithelial, assembling of urease enzyme subunit and stimulating the body to produce an immune response; at the same time they are the main immunogens of *H. pylori*. HspA is the main functional unit of Hsp and it locates bacterial

surface, is relatively conservative, and has immunogenicity, so it is an important candidate vaccine antigen of *H. pylori*.

Lactococcus lactis (*L. lactis*) is a Gram-positive, non-invasive, nonpathogenic, and food-grade bacterium and is generally recognized as safe (GRAS), which has been used in the food industry for production of fermented milk products for many years. Because of its genetic accessibility and easy maneuverability, *L. lactis* has been extensively used as a live delivery vehicle of heterologous proteins [6–8]. Nisin-controlled expression (NICE) system is the most effective food-grade inducible expression system. As induction signal, Nisin has been successfully applied to a variety of strains, including *Lactococcus*, *Lactobacillus*, *Streptococcus*, and *Enterococcus* [9, 10].

To develop an oral vaccine against *H. pylori* in the present study, *hspA* gene was cloned from *H. pylori* MEL-Hp27 and sequenced. Then *hspA* gene was inserted into the vector

pNZ8110 and electrotransformed into *L. lactis* NZ3900. The immunoreactivity of HspA expressed by recombinant *Lactococcus* bacteria was identified by western blot analysis.

2. Materials and Methods

2.1. Microorganism, Plasmid, and Growth Conditions. The strain of *H. pylori* MEL-Hp27 was separated from a clinical atrophic gastritis patient of Zhengzhou city, China. *L. lactis* NZ3900 and the *E. coli*-*L. lactis* shuttle vector pNZ8110 were purchased from NIZO food research. *L. lactis* NZ3900 was grown in M17 medium (Difco, USA) supplemented with 0.5% glucose (GM17) at 30°C. MC1061 was grown in LB broth (Oxoid, UK) at 37°C. When necessary, antibiotics were added to culture medium at the following concentrations: ampicillin (100 µg/mL) and chloramphenicol (10 µg/mL).

2.2. PCR of *hspA* Gene. The *hspA* gene was amplified by PCR in the presence of MEL-Hp27 genomic DNA and NcoI-forward primer 5'-catgccatgggcatgaagtttctacca-3' and XbaI-reverse primer 5'-cggctctagattagtggtttttgtgatc-3'; NcoI/XbaI restriction endonuclease sites were underlined. To construct the translational fusion expression, the four base pairs ATGG of NcoI restriction endonuclease sites under the SD domain of pNZ8110 *nisA* promoter were added to the forward primer. Two base pairs GC were added to keep the reading frame invariant. PCR was carried out under the following conditions: 30 cycles of denaturation at 94°C for 1 min, annealing at 55°C for 1 min, and extension at 72°C for 1 min.

2.3. Construction of the Recombinant NZ3900/pNZ8110-*hspA*. The *hspA* gene was inserted into the *Escherichia coli*- (*E. coli*-) *Lactococcus lactis* (*L. lactis*) shuttle vector pNZ8110 and the recombinant plasmid was transformed into MC1061 using the method of calcium chloride. The transformants were selected on LB plates containing 10 µg/mL chloramphenicol. Identification of the transformants was operated by PCR of the *hspA* gene, digested by NcoI and XbaI, and subjected to DNA sequencing. Then the plasmid of pNZ8110-*hspA* was electrotransformed into *L. lactis* NZ3900 and the transformants were selected on GM17 plates containing 10 µg/mL chloramphenicol.

2.4. Influence of *hspA* Gene on the Growth of Recombinant *Lactococcus lactis*. NZ3900 was transferred into the 10 mL GM17 liquid medium according to 1:25 (v/v). NZ3900/pNZ8110 and NZ3900/pNZ8110-*hspA* were transferred, respectively, into the 10 mL GM17 liquid medium containing 10 µg/mL chloramphenicol according to 1:25 (v/v). Each tube was mixed at interval of 1 hour to measure the absorbance A600 of bacteria. Growth curves of NZ3900, NZ3900/pNZ8110, and NZ3900/pNZ8110-*hspA* were plotted with incubation time as the abscissa and A600 as the vertical axis.

2.5. Influence of Nisin Concentration on the Growth of Recombinant *Lactococcus lactis*. NZ3900/pNZ8110-*hspA* was transferred, respectively, into 7 tubes of 10 mL GM17 liquid medium containing 10 µg/mL chloramphenicol according to 1:25 (v/v); then Nisin stock solution was added to a final

FIGURE 1: PCR product of MEL-Hp *hspA* gene. (1), (2) PCR product of *hspA* gene from Hp genome; (3) DNA ladder.

concentration of 0 ng/mL, 10 ng/mL, 20 ng/mL, 40 ng/mL, 60 ng/mL, 80 ng/mL, and 100 ng/mL. Each tube was mixed at interval of 1 hour to measure the absorbance A600 of bacteria. Growth curves of NZ3900/pNZ8110-*hspA* were plotted with incubation time as the abscissa and A600 as the vertical axis. Both NZ3900 and NZ3900/pNZ8110 were operated as controls at the same time.

2.6. Expression of HspA Protein in Recombinant *Lactococcus lactis*. NZ3900/pNZ8110-*hspA* was overnight cultured, then 400 µL of which was transferred, respectively, into 2 tubes of 10 mL GM17 liquid medium containing 10 µg/mL chloramphenicol. After NZ3900/pNZ8110-*hspA* was cultured for 4 hours, Nisin stock solution was added, respectively, to the final concentrations 10 ng/mL and 20 ng/mL to induce the expression of HspA. NZ3900/pNZ8110-*hspA* was sonicated after 4 hours and supernatant protein was collected and quantified using Bradford amount. HspA expressed in *Lactococcus lactis* was detected by SDS-PAGE and Tricine SDS-PAGE.

2.7. Detection of the Immunoreactivity of HspA Expressed by Recombinant *Lactococcus lactis*. HspA expressed in recombinant *Lactococcus lactis* was electrophoresed on a 15% SDS-PAGE and then electrotransferred to the cellulose nitrate membrane. The membrane was probed with a primary polyclonal mice antiserum (diluted 1:50) and the membrane incubated at 37°C for 2 h, washed three times with TBS/Tween-20 (TBST) for 15 min, and then incubated with the horseradish peroxidase-linked secondary antibody (diluted 1:1000). Protein detection was visualized using the DAB western blot detection system (Tiangen, China).

3. Results

3.1. PCR Results of *hspA* Gene from MEL-Hp27. Figure 1 showed that the *hspA* gene (Genbank accession number KJ567076) had been cloned successfully, which contained 363 base pairs. The sequencing result showed that the sequences of *hspA* gene in this experiment were correct compared with the sequences of MEL-Hp27 *hspA* (Genbank accession

FIGURE 2: Identification of MC1061/pNZ8110-*hspA* by PCR. (1) Negative control. (2), (3), (4), and (5) PCR product of *hspA* from MC1061/pNZ8110-*hspA*. (6) DNA ladder.

number AY295084) except the additional six base pairs, as explained in Section 2.2. The sequence identification result of MEL-HP27 *hspA* gene is as follows.

```

atgggcatgaagtttc taccattagg agaaagggtc tttagtagaaa
gactgaaga agagaacaaa
accagttcag gcatcatcat ccctgataac gctaaagaaa agcctttaat
gggcgtagtc
aaagcgggta gccataaaat cagcgagggt tgcaaatcgg ttaaa-
gaagg cgaatgtgac
gcttttgca aatacaaaagg cgcagaaatc gttttagacg gcgttgaata
catgttgcta
gagctagaag acattctagg tattgtgggc tcaggctctt gttgtcatac
aaatagtcac
gaccataaac atgctaaga gcatgaagct tgctgtcatg atca-
caaaaa acactaa

```

3.2. Identification of Recombinant MC1061/pNZ8110-*hspA* and NZ3900/pNZ8110-*hspA*. To express the HspA protein in *L. lactis* transformant, the *hspA* gene was inserted into the *E. coli*-*L. lactis* shuttle vector pNZ8110 and the recombinant plasmid was transformed into the *E. coli* MC1061. The recombinant pNZ8110-*hspA* was detected to be recombined properly with PCR (Figure 2) and digestion experiments (Figure 3). Then the recombinant pNZ8110-*hspA* was electrotransformed into *L. lactis* NZ3900. NZ3900/pNZ8110-*hspA* was identified by PCR and digestion experiments (not shown).

3.3. Influence of *hspA* Gene on the Growth of Recombinant *Lactococcus lactis*. From the growth curves (Figure 4), it could be seen that the proliferation of NZ3900, NZ3900/pNZ8110, and NZ3900/pNZ8110-*hspA* was very slow within the former 2 hours; then they entered the logarithmic growth phase within the following 6 hours. After about 8 hours, the growth of bacteria got up to the plateau phase and the A₆₀₀ absorbance value got to 2.0 or so. The proliferation of NZ3900/pNZ8110 and NZ3900/pNZ8110-*hspA* was slower than NZ3900 in the logarithmic growth period, which may be due to the result

FIGURE 3: Identification of the recombinant vector of pNZ8110-*hspA* by restriction enzyme digestion. (1) pNZ8110-*hspA* digested by *Xba*I. (2) pNZ8110-*hspA* digested by *Nco*I and *Xba*I. (3) pNZ8110 digested by *Xba*I. (4) PCR product of *hspA*. (5) DNA ladder.

FIGURE 4: The growth curves of NZ3900, NZ3900/pNZ8110, and NZ3900/pNZ8110-*hspA*.

of plasmid replication, indicating that insertion of *hspA* gene did not impact the growth of recombinant *Lactococcus lactis*.

3.4. Influence of Nisin Concentration on the Growth of Recombinant *Lactococcus lactis*. It can be seen from Figures 5(a), 5(b), and 5(c) that the growth of NZ3900/pNZ8110-*hspA* was inhibited noticeably by different concentrations of Nisin. The A₆₀₀ absorbance value of NZ3900/pNZ8110-*hspA* got to 1.989 in no Nisin group, while in the 100 ng/mL Nisin group it only got to 1.125. The growth curve showed that different concentrations of Nisin did not affect the growth of blank control (NZ3900) and negative control (NZ3900/pNZ8110).

3.5. Expression of HspA in Recombinant *Lactococcus lactis*. The expected molecular weight bands could not be observed on the SDS-PAGE gel or Tricine SDS-PAGE gel after 10 ng/mL

FIGURE 5: (a) Influence of Nisin concentration on NZ3900. (b) Influence of Nisin concentration on NZ3900/pNZ8110. (c) Influence of Nisin concentration on NZ3900/pNZ8110-hspA.

or 20 ng/mL Nisin induction for 4 hours, indicating that no HspA protein was expressed or the expression level was too low to be detected. The expected molecular weight bands could not be observed in blank control (NZ3900) and negative control (NZ3900/pNZ8110).

3.6. Identification of the Immunoreactivity of HspA Expressed by Recombinant *Lactococcus lactis*. It was presented in Figure 6 that the HspA expressed by *L. lactis* transformant could be recognized by anti-Hp serum. This result indicated that the HspA protein expressed by *L. lactis* transformant had favorable immunoreactivity.

4. Discussion

H. pylori is now recognized as the most widespread human pathogen. Approximately half of the world's population is infected [11, 12]. A great interest in developing a vaccination method to prevent infection by *H. pylori* has been raised, given that immunization is always considered as the most economic and efficient means in such a field, especially for the developing countries. Many researchers had made a substantial contribution to *H. pylori* vaccine such as protein vaccine or *Salmonella* vector vaccine, while these vaccines could not be optimistical because of the difficult maneuverability or gastrointestinal toxicity or insecurity [13, 14].

FIGURE 6: Western blot reaction of HspA protein expressed by *Lactococcus lactis* with polyclonal antibody serum to *H. pylori*. (1) Negative control. (2), (3) Western blot reaction of HspA protein to polyclonal antibody serum to *H. pylori*. (4) Prestained protein marker.

L. lactis has a number of prominent advantages that make it a promising candidate for large scale gene expression: the bacterium is food-grade, and plasmid selection mechanisms are available that are food-grade and self-cloning. Furthermore, no endotoxins or inclusion bodies are formed and sophisticated genetic tools enable easy genetic handling. Finally, simple, nonaerated fermentation makes direct scale-up from 1-L scale to 1000-L scale possible. The commonly used regulated expression system of *L. lactis* is the Nisin-controlled gene expression (NICE) system [15–18]. In recent years, *L. lactis* has been used to express *H. pylori* antigen protein as live bacteria vehicle. The 3'-region (1152 bp) of the *cag7* gene of *Helicobacter pylori* 51 strain had been expressed in *Lactococcus lactis* ssp. MG1363 [19]. The results showed that the transformant could produce the Cag7-ct383 protein and the Cag7-ct383 protein level in the *L. lactis* transformant reached a maximum at the early stationary phase without extracellular secretion. The oral administration of the *L. lactis* transformant into mice generated anti-Cag7 antibody in serum in five of five mice. These results suggest that *L. lactis* transformant expressing Cag7-ct383 protein may be applicable as an oral vaccine to induce mucosal and systemic immunity to *H. pylori*.

NICE system was used in our experiments to express the MEL-Hp27 UreB and HspA protein in a food-grade delivery vehicle *L. lactis* NZ3900. The results showed that the maximum yield of the UreB protein was 27.26 μ g per milliliter medium and the percentage of UreB to cell extracts of the *L. lactis* transformant got to the peak at 20.19%, which was substantial quantities of antigen expressed in *L. lactis* [20]. In this study, it could be seen that the insertion of *hspA* did not impact the growth of NZ3900/pNZ8110-*hspA* and the bacterial density of NZ3900/pNZ8110-*hspA* could get the same value as NZ3900 and NZ3900/pNZ8110 when they grew into the plateau phase; different concentrations of Nisin did not affect the growth of NZ3900 and NZ3900/pNZ8110, while different concentrations of Nisin significantly inhibited the growth of NZ3900/pNZ8110-*hspA* except 10 ng/mL Nisin; the rest of concentrations of Nisin significantly inhibited the growth of NZ3900/pNZ8110-*hspA*. The expected molecular weight bands could not be observed on the SDS-PAGE gel or Tricine SDS-PAGE gel; western blot identification results showed that HspA expressed by NZ3900/pNZ8110-*hspA* had good immunoreactivity. From the growth curves, it could be inferred that it was not because of Nisin inhibiting the growth

of NZ3900/pNZ8110-*hspA* as inhibitory but because HspA had not been induced to be expressed effectively in 10 ng/mL Nisin group, while, in higher concentrations of Nisin group, a small amount of exogenous expression of HspA inhibited the growth of NZ3900/pNZ8110-*hspA* itself. The mechanism deserves to be further investigated.

This study established the experimental foundation for constructing the food-grade vaccine to *H. pylori* infection.

Conflict of Interests

The authors declare no conflict of interests.

References

- [1] S. P. Vyas and V. Sihorkar, "Exploring novel vaccines against *Helicobacter pylori*: protective and therapeutic immunization," *Journal of Clinical Pharmacy and Therapeutics*, vol. 24, no. 4, pp. 259–272, 1999.
- [2] V. De Francesco, A. Zullo, C. Hassan, F. Giorgio, R. Rosania, and E. Ierardi, "Mechanisms of *Helicobacter pylori* antibiotic resistance: an updated appraisal," *World Journal of Gastrointestinal Pathophysiology*, vol. 2, no. 3, pp. 35–41, 2011.
- [3] A. Pietroiusti, M. Giuliano, A. Magrini, A. Bergamaschi, and A. Galante, "Cytotoxin-associated gene A strains of *Helicobacter pylori* represent a risk factor for the development of microalbuminuria in type 2 diabetes," *Diabetes Care*, vol. 29, no. 6, pp. 1399–1401, 2006.
- [4] K.-S. Wu, C.-C. Hsiao, H.-R. Yu, E.-Y. Huang, W.-L. Mai, and J.-M. Sheen, "*Helicobacter pylori* infection and childhood idiopathic thrombocytopenic purpura," *Acta Paediatrica Taiwanica*, vol. 48, no. 5, pp. 263–266, 2007.
- [5] F. Franceschi, G. Niccoli, G. Ferrante et al., "CagA antigen of *Helicobacter pylori* and coronary instability: insight from a clinico-pathological study and a meta-analysis of 4241 cases," *Atherosclerosis*, vol. 202, no. 2, pp. 535–542, 2009.
- [6] P. M. Norton, J. M. Wells, H. W. G. Brown, A. M. Macpherson, and R. W. F. Le Page, "Protection against tetanus toxin in mice nasally immunized with recombinant *Lactococcus lactis* expressing tetanus toxin fragment C," *Vaccine*, vol. 15, no. 6-7, pp. 616–619, 1997.
- [7] L. G. Bermúdez-Humarán, N. G. Cortes-Perez, F. Lefèvre et al., "A novel mucosal vaccine based on live *Lactococci* expressing E7 antigen and IL-12 induces systemic and mucosal immune responses and protects mice against human papillomavirus type 16-induced tumors," *The Journal of Immunology*, vol. 175, no. 11, pp. 7297–7302, 2005.
- [8] R. Ramasamy, S. Yasawardena, A. Zomer, G. Venema, J. Kok, and K. Leenhouts, "Immunogenicity of a malaria parasite antigen displayed by *Lactococcus lactis* in oral immunisations," *Vaccine*, vol. 24, no. 18, pp. 3900–3908, 2006.
- [9] S. Pavan, P. Hols, J. Delcour et al., "Adaptation of the nisin-controlled expression system in *Lactobacillus plantarum*: a tool to study *in vivo* biological effects," *Applied and Environmental Microbiology*, vol. 66, no. 10, pp. 4427–4432, 2000.
- [10] E. M. Bryan, T. Bae, M. Kleerebezem, and G. M. Dunne, "Improved vectors for nisin-controlled expression in gram-positive bacteria," *Plasmid*, vol. 44, no. 2, pp. 183–190, 2000.
- [11] S. J. Czinn, "*Helicobacter pylori* infection: detection, investigation, and management," *Journal of Pediatrics*, vol. 146, no. 3, pp. S21–S26, 2005.

- [12] G. Rossi, P. Ruggiero, S. Peppoloni et al., "Therapeutic vaccination against *Helicobacter pylori* in the beagle dog experimental model: safety, immunogenicity, and efficacy," *Infection and Immunity*, vol. 72, no. 6, pp. 3252–3259, 2004.
- [13] M. Pizza, M. M. Giuliani, M. R. Fontana et al., "Mucosal vaccines: non toxic derivatives of LT and CT as mucosal adjuvants," *Vaccine*, vol. 19, no. 17–19, pp. 2534–2541, 2001.
- [14] C. Xu, Z.-S. Li, Y.-D. Du et al., "Construction of a recombinant attenuated *Salmonella typhimurium* DNA vaccine carrying *Helicobacter pylori* hpaA," *World Journal of Gastroenterology*, vol. 11, no. 1, pp. 114–117, 2005.
- [15] I. Mierau and M. Kleerebezem, "10 years of the nisin-controlled gene expression system (NICE) in *Lactococcus lactis*," *Applied Microbiology and Biotechnology*, vol. 68, no. 6, pp. 705–717, 2005.
- [16] T. Maischberger, I. Mierau, C. K. Peterbauer, J. Hugenholtz, and D. Haltrich, "High-level expression of *Lactobacillus* β -galactosidases in *Lactococcus lactis* using the food-grade, nisin-controlled expression system NICE," *Journal of Agricultural and Food Chemistry*, vol. 58, no. 4, pp. 2279–2287, 2010.
- [17] A. Berlec and B. Štrukelj, "Large increase in brazzein expression achieved by changing the plasmid/strain combination of the NICE system in *Lactococcus lactis*," *Letters in Applied Microbiology*, vol. 48, no. 6, pp. 750–755, 2009.
- [18] I. Mierau, P. Leij, I. van Swam et al., "Industrial-scale production and purification of a heterologous protein in *Lactococcus lactis* using the nisin-controlled gene expression system NICE: the case of lysostaphin," *Microbial Cell Factories*, vol. 4, article 15, 2005.
- [19] S. J. Kim, J. Y. Lee, D. Y. Jun et al., "Oral administration of *Lactococcus lactis* expressing *Helicobacter pylori* Cag7-ct383 protein induces systemic anti-Cag7 immune response in mice," *FEMS Immunology and Medical Microbiology*, vol. 57, no. 3, pp. 257–268, 2009.
- [20] X. J. Zhang, G. C. Duan, R. G. Zhang, and Q. Fan, "Optimized expression of *Helicobacter pylori ureB* gene in the *Lactococcus lactis* nisin-controlled gene expression (NICE) system and experimental study of its immunoreactivity," *Current Microbiology*, vol. 58, no. 4, pp. 308–314, 2009.