PREVENTION OF ENDOTOXAEMIA IN OBSTRUCTIVE JAUNDICE — A COMPARATIVE STUDY OF BILE SALTS

J.A. PAIN* and M.E. BAILEY

Department of Surgery, Royal Surrey County Hospital, Guildford, Surrey, UK

Systemic endotoxaemia is associated with postoperative renal dysfunction in obstructive jaundice, and can be prevented by the pre-operative administration of certain bile salts. In order to find the most effective bile salt for use in this condition, a comparison of the anti-endotoxic activities of different bile salts was performed. Bile salts were incubated *in vitro* with endotoxin and the resultant endotoxin level was measured with a quantitative limulus assay. The *in vivo* effects of different oral bile salts on the intestinal absorption of radiolabelled endotoxin from rats with obstructive jaundice were compared. The *in vitro* and *in vivo* anti-endotoxic activities of bile salts related to their known detergent activities. Deoxycholic acid and its conjugates were the most effective and should be the bile salts of choice for further clinical evaluation in obstructive jaundice in man.

KEY WORDS: Bile salts, endotoxin, limulus test, bile-duct obstruction — extra hepatic.

INTRODUCTION

Postoperative acute renal failure occurs in 9% of patients with obstructive jaundice¹. There is considerable evidence to implicate the presence of endotoxins within the systemic circulation in the development of this renal dysfunction^{2–4}. Systemic endotoxaemia is believed to result from an increased absorption of intestinal endotoxin into the portal circulation^{4,5} and its impaired removal by the Kupffer cells within the liver^{6,7}.

The presence of bile within the intestinal lumen reduces endotoxin absorption^{8,9}. Oral administration of bile salts abolishes endotoxaemia in rats with bile-duct ligation⁴. In patients with obstructive jaundice, oral sodium deoxycholate prevents both portal and systemic endotoxaemia and protects postoperative renal function¹⁰. However, chenodeoxycholic acid is less effective in these roles¹¹ and ursodeoxycholic acid has been found to confer no benefit¹². An *in vitro* and *in vivo* comparison of bile salts has been performed to find which bile salt should be the most effective anti-endotoxaemic agent for use in obstructive jaundice.

* Correspondence to: Mr. J.A. Pain, FRCS, Department of Surgery, Mayday Hospital, Thornton Heath, Croydon CR4 7YE, UK.

METHODS

In Vitro Study

Different bile salts and their conjugates were compared as to their ability to inhibit endotoxin activity *in vitro*. Bile salts were incubated with endotoxin and the resultant endotoxin level was measured.

The bile salts (200 mg, >98% purity, Sigma) were dissolved in 99% ethanol (4 ml), and then added to sodium phosphate buffer 86 ml, pH 7.0). Various concentration of the bile-salt/buffer solution were mixed with *Escherichia coli* endotoxin (20 µg, 0111B4, Difco) in water (0.2 ml), made up to 2 ml with sodium phosphate buffer and incubated at 37°C for 6 h. The mixtures were then assayed for endotoxin by means of a quantitative chromogenic limulus assay⁵. The assay had a lower limit of sensitivity of 1pg/ml endotoxin in water, with a coefficient of variation of 4% with 50pg/ml endotoxin in water. All tests were performed in triplicate with appropriate controls.

In Vivo Study

Different bile salts and their conjugates were compared in their ability to prevent the absorption of radiolabelled endotoxin from the intestines of rats with obstructive jaundice.

Endotoxin was labelled with Cr⁵¹ according to the method of Braude *et al.*¹³ Endotoxin, E.coli 0111B4 (700 mg, Sigma) was suspended in phosphate chloride buffer (140 ml, pH 7). The endotoxin suspension was incubated at 37°C for 24 h with radiolabelled sodium chromate (300 MBq, Amersham International). After incubation the endotoxin/sodium chromate was dialysed against de-ionized water, weakly buffered with phosphate chloride, at room temperature, until there was no longer any significant loss of radioactivity into the dialysate. The radiolabelled endotoxin was stored at 4°C in gas-evacuated vials and, prior to use, tested for unbound chromium by further dialysis.

Seventy-four male Wistar rats (weight 270–300gm) underwent common bile-duct (CBD) ligation and division or sham operation on Day 0 and were treated as shown in Figure 1. The extrapancreatic part of the CBD was mobilized and divided between silk ligatures, and in the sham operation a silk ligature was placed around the CBD and then removed.

On Day 12, 63 rats (seven groups of nine) received one of the following bile salts in an oral dose of 6mg/100gm body weight every 8 h for two days: sodium deoxycholate, cholic acid, chenodeoxycholic acid, ursodeoxycholic acid, sodium taurocholate, sodium taurodeoxycholate, and glycodeoxycholic acid. A dose of 6mg bile salt/100 gm body weight was chosen since preliminary experiments had shown that this dose of sodium deoxycholate had prevented the absorption of more than 60% of orally administered radiolabelled endotoxin. On Day 14, 4 h after the final dose of bile salts, radiolabelled endotoxin (2mg/100gm body weight) was given via a gavage tube to all rats.

Four hours later the rats were killed by cervical dislocation, and immediate laparotomy was performed. Ligatures were placed around the lower oesophagus and distal rectum, and the intestines including the stomach were removed. The attached mesentery was discarded and the intestines were counted for radioactivity to calculate the percentage radioactivity remaining in the intestines. From three control jaundiced rats the livers, lungs, spleens, kidneys, and samples of blood and urine were also measured for radioactivity.

Figure 1 Design of animal experiment.

Statistics

Comparisons between groups were made with the Student's t-test.

RESULTS

In Vitro Study

Controls in which only buffer was added to endotoxin showed no significant loss of detectable endotoxin after incubation. All bile salts tested reduced the detectable endotoxin level. The dihydroxy bile salts, glycodeoxycholic acid and deoxycholic acid, proved to have the greatest *in vitro* anti-endotoxic action. Only 400 µg/ml of these bile salts was required to reduce the detectable endotoxin level from 10 µg/ml to <50 pg/ml (20 000-fold decrease). The mean activities of other bile salts expressed as a percentage of that of deoxycholic acid are shown in Table 1.

In Vivo Study

The rats each received 1.7 ml radiolabelled endotoxin, which produced a mean 149 781 counts per minute. No diarrheogenic effect of the bile salts was evident. Two rats in the chenodeoxycholic-acid-treated group and one in the taurocholate-treated group died during the experiment and are not included in the results. In the sham-operated (non-jaundiced) rats less than 10% (mean intestinal radioactivity count $137\,079 \pm 7373$) of the radioactive label was absorbed, whereas in the jaundiced rats 65% (mean intestinal radioactivity count $52\,425 \pm 7193$) was absorbed ($t = 19.2$, d.f. = 9, $P < 0.001$). The mean radioactivity counts per minute of the tissue and body fluids taken from the three jaundiced control rats were liver 13 259, spleen 982, kidney 4873, urine 2240, packed cell blood 1871 per ml, plasma 724 per ml.

All bile salts tested, except ursodeoxycholic acid, produced a significant ($P < 0.05$)

Table 1 *In vitro* and *in vivo* comparison of bile salts. *In vitro* results expressed as bile-salt concentration required to reduce the detectable endotoxin level from 10 µg/ml to 50pg/ml. *In vivo* results expressed as mean $\pm$ s.d. residual intestinal endotoxin radioactivity counts. In bracket: percentage activity compared to glycodeoxycholic acid.

	<i>In Vitro</i>		<i>In Vivo</i>		
	µg/ml	%	$\times 10^3$	cpm	%
Secondary bile salts					
Glycodeoxycholic acid	400	(100)	114	± 12	(100)
Deoxycholic acid	400	(100)	116	± 18	(97)
Na Glycodeoxycholate	500	(80)			
Na Taurodeoxycholate	500	(80)	106	± 23	(90)
Chenodeoxycholic acid	600	(66)	105	± 16	(89)
Na Glycochenodeoxycholate	600	(66)			
Na Taurochenodeoxycholate	700	(57)			
Primary bile salts					
Cholic acid	1700	(23)	89	$\pm 23^a$	(79)
Glychocolic acid	1700	(23)			
Na Glycocholate	<2000	(>20)			
Na Taurocholate	<2000	(>20)	84	$\pm 19^b$	(74)
Tertiary bile salts					
Ursodeoxycholic acid	<2000	(>20)	70	$\pm 23^c$	(62)

Student's t-test ^a $P > 0.05$
^b $P > 0.01$
^c $P > 0.001$

reduction in the absorption of the radiolabelled endotoxin. Glycodeoxycholic acid was the most effective agent and reduced the percentage endotoxin absorption to a mean 24 ± 8 . The activities of other bile salts expressed as a percentage of that of glycodeoxycholic acid are shown in Table 1.

DISCUSSION

Bile salts are detergents and have a direct anti-endotoxic action *in vitro*, causing inactivation of the endotoxin molecule¹⁴⁻¹⁶. Endotoxin, which is aggregated into particles with a molecular weight of about 1 000 000, is disassociated by bile salts to form non-toxic subunits with molecular weights of about 20 000¹⁴.

The presence of bile within the intestinal lumen prevents endotoxin absorption. The clinical effects of endotoxaemia cannot be induced in healthy animals by the oral administration of endotoxin even if doses 500–3000 times the minimal lethal parenteral dose are given^{17,18}. Endotoxin absorption is increased in obstructive jaundice⁴ or after bile-duct cannulation and external biliary drainage⁸, but is reduced in obstructive jaundice if the endotoxin is incubated with a bile salt prior to its oral administration¹⁹, if oral bile salts are administered⁴, or if internal biliary drainage is performed⁹.

All bile salts tested had potent *in vitro* anti-endotoxic activities, and the most active were those of the dihydroxy bile acid, deoxycholic acid and its conjugates, and the least activity was shown by ursodeoxycholic acid. Lithocholic acid and its conjugates were not tested since they are toxic and would be unsuitable for use in

man. These results are consistent with previous studies examining the *in vitro* anti-endotoxic activity of bile salts. Iwasaki and Tanikawa²⁰ found that deoxycholic acid was the most effective bile salt compared to chenodeoxycholic acid, its conjugates and taurocholic acid. Liehr *et al.*²¹, comparing conjugates of bile salts, showed that those of deoxycholate had a greater anti-endotoxic activity than those of chenodeoxycholate, while those of cholate were the least effective. These previous studies measured endotoxin with a semiquantitative limulus assay based on a subjective assessment of the type of gel formed. The assay used in the present experiment was highly sensitive, reproducible and quantitative.

The comparative *in vitro* bile-salt activities related to their known critical micellar concentration^{22,23}, which supports the theory that it is the detergent properties of the bile salts that render endotoxins inactive.

The labelling of endotoxin with Cr⁵¹ was first described by Braude *et al.*¹³ Although there is uncertainty as to the exact location of the radiolabel within the endotoxin molecule, the attachment is very stable both *in vitro* and *in vivo*^{13,24–26}. Four hours after oral administration of the radiolabelled endotoxin to the jaundiced rats, some absorbed radioactivity was in the liver, but most was present in the plasma. This distribution is similar to that found after intravenous injection of Cr⁵¹-labelled endotoxin into animals¹³. Endotoxin binds to plasma high-density lipoproteins and circulates in this complex form until cleared from the circulation²⁷. In our study it is unlikely that the Cr⁵¹ present in the circulation was free, because if free Cr⁵¹ is injected intravenously the majority can subsequently be found within the red-cell layer and not within the plasma^{28,29}.

There was increased absorption of radiolabelled endotoxin in the bile-duct-ligated rat. This probably occurred due to a lack of intestinal bile rather than due to the hyperbilirubinaemic state, since bile-duct cannulation in rats also leads to an increased absorption of endotoxin^{8,9}. All of the bile salts tested, except for ursodeoxycholic acid, significantly reduced endotoxin absorption. As in the *in vitro* study deoxycholic acid and its conjugates were the most effective, and the bile-salt activities related to their known detergent activities^{20,21}. A greater difference between the bile-salt activities was shown by the *in vitro* test, but this probably related to the bile salts being incubated with the endotoxin for 6 h in controlled conditions, whereas in the *in vivo* test the endotoxin was given only 4 h before the animal was killed; also, numerous biological variables could effect the interaction of bile salts and endotoxin. The choice of the most suitable bile salt as a prophylactic agent against endotoxaemia in obstructive jaundice depends not only on the individual salt's activity as determined *in vitro* but also on its metabolism and action *in vivo*. It is probably unimportant whether a bile "acid" or bile "salt" is used, since most bile "salts" precipitate as bile "acids" in the stomach as their precipitation pH values are higher than the luminal pH³⁰.

In order to prevent endotoxaemia in obstructive jaundice it is recommended that deoxycholic acid or its conjugates be the agents of choice for further clinical evaluation.

References

1. Pain, J.A., Cahill, C.J. and Bailey, M.E. (1985) Perioperative complications in obstructive jaundice: therapeutic considerations. *British Journal of Surgery*, **72**, 942–945
2. Wardle, E.N. (1975) Endotoxaemia and the pathogenesis of acute renal failure. *Quarterly Journal of Medicine*, **44**, 389–398

3. Wilkinson, S.P., Moodie, H., Stamatakis, J.D., Kakkar, V.V. and Williams, R. (1976) Endotoxaemia and renal failure in cirrhosis and obstructive jaundice. *British Journal of Medicine*, **2**, 1415–1418
4. Bailey, M.E. (1976) Endotoxin, bile salts and renal function in obstructive jaundice. *British Journal of Surgery*, **63**, 774–778
5. Pain, J.A. and Bailey, M.E. (1987) Measurement of operative endotoxin plasma levels in jaundiced and non-jaundiced patients. *European Surgical Research*, **19**, 207–216
6. Drivas, G., James, O. and Wardle, N. (1976) Study of reticuloendothelial phagocytic capacity in patients with cholestasis. *British Medical Journal*, **i**, 1568–1569
7. Pain, J.A. (1987) Reticuloendothelial function in obstructive jaundice. *British Journal of Surgery*, **74**, 1091–1094
8. Koscar, L.T., Bertok, L. and Varteresz, V. (1969) Effect of bile acids on the intestinal absorption of endoxin in rats. *Journal of Bacteriology*, **100**, 220–223
9. Gouma, D.J., Coelho, J.C.U., Fisher, J.D., Schlegel, J.F., Li, Y.F. and Moody, F.G. (1986) Endotoxemia after relief of biliary obstruction by internal and external drainage in rats. *The American Journal of Surgery*, **151**, 476–479
10. Cahill, C.J. (1983) Prevention of post-operative renal failure in patients with obstructive jaundice — the role of bile salts. *British Journal of Surgery*, **70**, 590–593
11. Cahill, C.J., Pain, J.A. and Bailey, M.E. (1987) Bile salts, endotoxin and renal function in obstructive jaundice. *Surg. Gynecol. Obstet.*, **165**, 519–522
12. Thompson, J.N., Cohen, J., Blenkarn, J.I., McConnell, J.S., Barr, J. and Blumgart, L.H. (1986) A randomized clinical trial of oral ursodeoxycholic acid in obstructive jaundice. *British Journal of Surgery*, **73**, 634–636
13. Braude, A.I., Carey, F.J. and Zalesky, M. (1955) Studies with radioactive endotoxin. *Journal of Clinical Investigation*, **34**, 850–857
14. Ribi, E., Anacker, R.L., Brown, R., Haskins, W.T., Malmgren, B., Milner, K.C. and Rudbach, J.A. (1966) Reaction of endotoxin and surfactants. I. Physical and biological properties of endotoxin treated with sodium deoxycholate. *Journal of Bacteriology*, **92**, 1493–1509
15. Rudbach, J.A., Anacker, R.L., Haskins, W.T., Johnson, A.G., Milner, K.C. and Ribi, E. (1966) Physical aspects of reversible inactivation of endotoxin. *Annals of New York Academy of Science*, **133**, 629–643
16. Shands, J.W. and Chun, P.W. (1980) The dispersion of Gram-negative lipopolysaccharide by deoxycholate. *Journal of Biological Chemistry*, **255**, 1221–1226
17. Berdzi, I., Bertok, L., Baintner, K. and Veress, (1968) Failure of oral *Escherichia coli* endotoxin to induce either tolerance or toxic symptoms in rats. *Journal of Pathology and Bacteriology*, **96**, 481–486
18. Bertok, L. (1977) Physico-chemical defense of vertebrate organisms: the role of bile acids in defense against bacterial endotoxins. *Perspectives in Biology and Medicine*, **21**, 70–76
19. Filkins, J.P. (1971) Decreased shock lethality in rats with surfactant treated endotoxin. *Proceedings of the Society of Experimental Biology and Medicine*, **136**, 466–468
20. Iwasaki, M. and Tanikawa, K. (1971) Liver diseases and endotoxin. Part I. Effect of bile acid on endotoxin. *Nippon Shokakibyo Gakkai Zasshi*, **78**, 1232–1240
21. Liehr, H., Grun, M., Brunswig, D. and Sautter, T. (1976) Endotoxinamiebei leberzirrhose. *Zeitschrift für Gastroenterologie*, **14**, 14–23
22. Hofmann, A.F. (1963) The function of bile salts in fat absorption. *Biochemistry Journal*, **89**, 57–68
23. Hisadone, T., Nakama, T., Itoh, H. and Furusawa, T. (1980) Physical-chemical properties of chenodeoxycholic acid and ursodeoxycholic acid. *Gastroenterology Japan*, **15**, 257–263
24. Chedid, L., Skarnes, R.C. and Parant, M. (1963) Characterization of a Cr⁵¹-labeled endotoxin and its identification in plasma and urine after parenteral administration. *Journal of Experimental Medicine*, **117**, 561–571
25. Daniele, R., Singh, H., Appert, R.E., Pairent, F.W. and Howard, J.M. (1970) Lymphatic absorption of intraperitoneal endotoxin in the dog. *Surgery*, **67**, 484–487
26. Nolan, J.P., Hare, D.K., McDavitt, J.J. and Ali, M.V. (1977) *In vitro* studies of intestinal endotoxin absorption. I. Kinetics of absorption in the isolated everted gut sac. *Gastroenterology*, **72**, 434–439
27. Ulevitch, R.J., Johnstone, A.R. and Weinstein, D.B. (1981) New function for high density lipoproteins: isolation and characterization of a bacterial lipopolysaccharide–high density lipoprotein complex formed in rabbit plasma. *Journal of Clinical Investigation*, **67**, 827–837
28. Braude, A.I., Carey, F.J. and Zalesky, M. (1955) Studies with radioactive endotoxin. II. Correlation of physiological effects with distribution of radioactivity in rabbits injected with lethal doses of *E. coli* endotoxin labelled with radioactive sodium chromate. *Journal of Clinical Investigation*, **34**, 858–866

29. Manzo, L., Di Nucci, A., Edel, J., Gregotti, C. and Sabbioni, E (1983) Biliary and gastrointestinal excretion of chromium after administration of Cr-III and Cr-IV in rats. *Research Communications in Biochemical Pathology*, **42**, 113–125
30. Igimi, H. and Carey, M.C. (1980) pH-solubility relations of chenodeoxycholic and ursodeoxycholic acids: physical-chemical basis for dissimilar solution and membrane phenomena. *Journal of Lipid Research*, **21**, 72–90

Accepted by S. Bengmark on 14 March 1988