Images in Infectious Diseases in Obstetrics and Gynecology

Section Editor: David E. Soper, M.D.

Vaginal *Lactobacillus* Phage Plaques and Electron Micrograph

Lin Tao and Sylvia I. Pavlova

Department of Oral Biology, College of Dentistry, University
of Illinois at Chicago, Chicago, IL

Normally, lactobacilli are a dominant species in the vagina and play an important role in maintaining vaginal

health by producing hydrogen peroxide and lactic acid to inhibit other bacteria, such as pathogenic anaerobes. When bacterial vaginosis occurs, however, vaginal lactobacilli decrease or disappear for unknown reasons. Recently, we have observed that bacteriophages or viruses can infect vaginal lactobacilli. Here we present two figures to show these phages: (A) Photograph of *Lactobacillus* phage drop assay. Phages were induced from lactobacilli by mitomycin C and dropped onto a *Lactobacillus* indicator strain on soft agar plates. Note: Each of the small clear plaques in or near a lysis zone is caused by single phage. (B) Electron micrograph of vaginal *Lactobacillus* phage (bar = 100 nm). © 1999 Wiley-Liss, Inc.

Images in Infectious Diseases in Obstetrics and Gynecology presents clinically important visual images that a practitioner in women's health might encounter. If you have a high-quality color or black-and-white photograph or slide representing such an image that you would like considered for publication, send it with a descriptive legend to David E. Soper, MD, Department of Obstetrics and Gynecology, Rush-Presbyterian-St. Luke's Medical Center, 1653 West Congress Parkway, Chicago, IL 60612-3833.

Images is made possible through an educational grant from Pfizer, Inc.