Comparison of Gram stain and Pap smear procedures in the diagnosis of bacterial vaginosis

Enver Vardar¹, Izzet Maral², Murat Inal², Özgül Özgüder¹,
Funda Tasli¹ and Hakan Postaci¹

¹Pathology Department, SSK Izmir Training Hospital,

²SSK Ege Maternity Hospital, Izmir, Turkey

Objective: The purpose of this study was to examine the characteristics of Gram stain versus Pap smear in diagnosis of bacterial vaginosis (BV).

Methods: One-thousand and sixty women were enrolled in this study. All cases with symptoms of BV were determined by Amsel's criteria, which were accepted as the gold standard for diagnosis of BV. Pap smear and Gram stain evaluations were compared according to Amsel's criteria, without viewing the clinical results of the patients. Gram stain and Pap smear results were determined as negative or positive according to Amsel's criteria. Sensitivity, specificity and positive predictive values were calculated.

Results: After accepting the cases that were diagnosed as BV according to Amsel's criteria as reference cases, the sensitivity of the Gram stain method was calculated as 97% and the sensitivity of the Pap smear method as 93%. Similar specificity rates were obtained with both methods in diagnosis of BV related to the clinical results. There were no statistically significant differences in diagnosis of BV between these two groups.

Conclusion: If Amsel's criteria are accepted as the gold standard for diagnosis of BV, Gram stain and Pap smear methods will give similar results in diagnosis.

Key words: BACTERIAL VAGINOSIS; GRAM STAIN; PAP SMEAR

The most frequently encountered complaint in gynecological out-patient clinics is vaginal discharge. Normal vaginal mucosa does not have any glandular structure. Usually there is a biological balance in the microorganisms living in vaginal mucosa. The most important role in the continuity of this balance – and in preventing the growth of pathogenic microorganisms – is that of *Lactobacillus* species. Lactobacilli produce an acidic medium in the vagina via hydrogen peroxide (H₂O₂), which transforms glycogen present in vaginal epithelium to lactic acid¹. The acidic medium produced by lactobacilli suppresses the growth of other microorganisms. If the balance of vaginal flora is altered

against lactobacilli, a clinical picture of itching, pain and vaginal discharge or smell will be observed. In bacterial vaginosis (BV), microorganisms that are short, rod-like and mostly anaerobic, with variable Gram stain properties – such as *Gardnerella vaginalis*, *Bacteriodes*, β -streptococci and the *Mobilincus*–*Falcivibrio* group – replace the usual vaginal flora formed by lactobacilli^{1,2}.

The aim of the presented study is to compare the efficacy of Gram stain and Pap smear with that of Amsel's criteria in the diagnosis of BV – which is continually gaining importance in relation to morbidity and mortality problems – in cervico-vaginal specimens obtained in routine controls.

MATERIALS AND METHODS

Routine gynecologic examinations of 1060 patients aged 18–50 years, who were referred to the SSK Ege Maternity Hospital between July and December 1996 with complaints of bad-smelling vaginal discharge, itching and pain, were performed and histories of the patients were taken. Women using oral contraceptives and vaginal spermicides, those who used antibiotics or vaginal suppositories for any reason and those who performed vaginal lavage after coitus were excluded from the study (172 women).

Vaginal inspections were carried out using a dry, sterile speculum without applying antiseptic cleansing liquids and the characteristics of the vaginal discharge were evaluated. The pH of the vaginal discharge was evaluated with litmus paper (Merck KGaA Acilit pH, Darmstadt, Germany). An amine test was performed to the appropriately taken vaginal discharge sample with 10% KOH. Samples were taken from the vaginal lateral wall and posterior fornix with brushes, and spread out on three preparations. One of these preparations was examined immediately by light microscope; appropriate fixation, Gram stain and classic Papanicolaou (Pap) stain were applied to the other preparations.

The evaluation of the results obtained from Pap and Gram stains was based on positivity by Amsel's criteria. For a positive diagnosis of BV, the presence of at least three of the following four criteria of Amsel is required: (1) thin, homogeneous gray-white discharge on the vaginal wall, that can be easily sampled; (2) vaginal-discharge pH above 4.5; (3) observation of 'clue' cells in fresh vaginal smear; and (4) amine smell similar to that of stale fish, obtained by dripping 10% KOH on the vaginal discharge².

The aim of the study did not include establishing the prevalence of BV in the population.

In evaluation of Pap stain under $\times 400$ magnification, according to the method of Davis and colleagues³, cases with the presence of micro-organisms other than lactobacilli in a thin film form, the presence of clue cells – which are formed by the covering of the cytoplasmic membranes of squamous cells by non-*Lactobacillus* micro-organisms – and a predominance of non-*Lactobacillus* bacteria in most areas, were accepted as

BV-positive cases. Davis and colleagues reported the condition of absence of lactobacilli as a fourth diagnostic criterion, but the absence of lactobacilli is modified as predominance of non-*Lactobacillus* bacteria over lactobacilli in the present study.

In evaluation of Gram stain under $\times 400$ magnification, according to Thomason and colleagues' 'clue cell' method⁴ cases with the presence of non-*Lactobacillus* bacteria in most areas and the presence of clue cells in at least two of 20 areas in large magnification, were accepted as positive. Cells having only lactobacilli in their cytoplasm were accepted as negative, whereas the covering of cytoplasmic membranes by non-*Lactobacillus* bacteria was used to identify clue cells.

Cytologic evaluation related to each case was performed, without any data obtained via Amsel's criteria. Specificity, sensitivity and predictive values of the results obtained from Pap and Gram stains were calculated by using the results obtained according to Amsel's criteria as reference values.

RESULTS

The mean age of subjects was 31.7 years (range 18–51 years). Approximately forty-seven percent of all cases (416/888) were pregnant at the time of examination. In the pregnant group, the mean pregnancy number of the 416 cases with a determined pregnancy number was 3.4, and the mean parity of cases with a determined parity was 2.2.

BV was found by Amsel's criteria in 260 (29.3%) of the 888 cases. In BV-positive cases, the most frequently applied contraception methods of the non-pregnant women were intra-uterine device (IUD) (37%), coitus interruptus (17%) and condom (14%); 19% were not using any contraception.

All of the women were suffering from vaginal discharge, which was a yellow-gray discharge in 78% of cases, white in 17% and yellow-green in 4%. Itching was observed in 88% and bad smell was observed in 96%. Post-coital odor was observed in 97%.

Two hundred and forty-two cases were determined as BV-positive by Pap smear, compared with 251 BV-positive cases determined by Gram stain (Tables 1 and 2). Sensitivity and specificity were 93% and 94% respectively with Pap smear and 97%

Table 1 Distribution of positive cases according to diagnosis method

Diagnosis method	BV +ve cases (260 according to Amsel)
Gram	
True (+) cases	251
False (-) cases	9
Pap	
True (+) cases	242
False (-) cases	18

BV, bacterial vaginosis

Table 2 Distribution of negative cases according to diagnosis method

Diagnosis method	BV -ve cases (628 according to Amsel)
Gram	
True (-) cases	593
False (+) cases	35
Pap	
True (-) cases	588
False (+) cases	40

BV, bacterial vaginosis

and 94%, respectively with Gram stain. Positive predictive values were determined as 86% for the Pap-smear method and 88% for the Gram stain method. The comparison of results obtained from both groups with Amsel's criteria did not yield any significant difference.

DISCUSSION

BV is the most frequent cause of vaginitis, and is characterized by increase in growth of anaerobic and aerobic microorganisms due to an unbalanced eco-system in the vagina². Gardner and Dukes⁵ were the first to report *Haemophilus vaginalis* as a cause of nonspecific vaginitis in 1955. It is also reported that there are high sensitivity and specifity values of Gram stain in diagnosis of BV, besides Amsel's criteria^{2,4,6-12}.

The clinico-pathologic condition characterized by redness in the vaginal wall, bad-odored discharge and the presence of clue cells in

cervicovaginal specimens, which results from the transformation of the acidic vaginal pH to an alkaline pH via metabolic activity of the afore-defined bacteria is known as BV⁶.

It is also reported that BV causes premature rupture of membranes, preterm delivery and endometritis and, according to some authors, BV has possible carcinogenic effects due to abnormal vaginal cytology and mosaic colposcopic pattern in the cervix¹³⁻¹⁷.

Various methods have been recommended for the evaluation of preparations for BV by Gram stain. Spiegel and colleagues¹⁸ counted a total of 20 areas in × 1000 magnification in which they reported the diagnosis of BV, in correlation with the number of microorganisms present in each large magnification area. Nugent and colleagues¹⁹ determined a total score over six as BV, by taking *Lactobacillus Gardnerella* and other microorganisms into account in a total of 20 large magnification areas in × 1000 magnification. In the present study, the clue cell method of Thomason and colleagues⁴ has been used.

Platz-Christensen and colleagues⁹ ascertained the sensitivity of the Pap smear method as 88%, the specifity as 97% and positive predictive value as 97%, following a study comparing Pap smear and Gram stain methods in the diagnosis of BV. These rates were 100%, 97%, and 94% respectively for the Gram stain method⁹. In our study, the observed sensitivity and specificity were 93% and 94% respectively for Pap smear, and 97% and 94% respectively for Gram stain. The positive predictive values were determined as 86% for Pap smear and 88% for Gram stain. The aforementioned rates are similar to rates determined by Platz-Christensen and colleagues⁹. But the specificity rate of our study is smaller than the that determined by Platz-Christensen and colleagues. Also, Platz-Christensen and colleagues reported the sensitivity as 90% and specificity as 97% after the comparison of Pap smear and clinical criteria, in a similar study performed in 1989²⁰.

Davis and colleagues³ reported the sensitivity and specificity rates of Pap smear as 55 and 98% respectively, in a study in which the Gram stain results were accepted as the diagnostic standard. They reached their hypothesis because of lower rates of sensitivity. Davis and colleagues claimed

that the fact that Pap smear screening is a routine procedure and cannot be always evaluated by cytopathologists was the cause of lower rates of sensitivity. But the Pap smear evaluation criteria given to the cytotechnicians, despite a long period of training may explain the different results in Davis and colleagues' study.

Platz-Christensen and colleagues obtained similar results in two studies performed 6 years apart^{9,20}. In addition, Prey²¹ reported the necessity of the presence of coccobacilli-type micro-organisms for the diagnosis of BV by Pap smear. Bedrossian and colleagues²² reported that Pap smear gives results as good as Gram stain in the diagnosis of BV and cervicitis. They also reported that the exclusion of Gram stain from the routine examination method might decrease the economic cost. Lamont and colleagues reported the sensitivity and specificity as 81 and 91% respectively, in Pap smears evaluated by cytotechnicians, but these

rates were 87 and 97% respectively in Pap smears evaluated by only one cytopathologist⁸.

Long Gram stain evaluation under $\times 1000$ magnification, proposed by Nugent¹⁹ and Spiegel¹⁸ despite Pap smear evaluation at $\times 400$ magnification, will cause a significant loss of time especially in countries like Turkey, where cytotechnologists are not employed or do not exist. This time-loss will lead to an increase in the cost per preparation, and possibly a decrease in the time that can be used for the determination of malignancies – especially of cervix carcinoma – earlier in Pap smear determinations, if the chance of the Pap smear picking up other infection types is also taken into account. The difference between Gram stain and Pap smear methods, especially in specificity in BV diagnosis, can be neglected. If there is a negligible difference between Gram stain and Pap smear methods it is better to use Pap smears in routine gynecologic cytologic examinations.

REFERENCES

1. Klebanoff SJ, Hillier SL, Eschelbach DA, Waltersdorff AM. Control of the microbial flora of the vagina by H₂O₂-generating lactobacilli. *J Infect Dis* 1991;164:94–100
2. Amsel R, Totten PA, Spiegel CA, et al. Non-specific vaginitis. Diagnostic criteria and microbial and epidemiologic associations. *Am J Med* 1983;74:14–22
3. Davis JD, Connor EE, Clark P, et al. Correlation between cytologic results and Gram stain as diagnostic tests for bacterial vaginosis. *Am J Obstet Gynecol* 1997;177:532–5
4. Thomason JL, Anderson RJ, Gelbart SM, et al. Simplified Gram stain interpretive method for diagnosis bacterial vaginosis. *Am J Obstet Gynecol* 1992;167:16–19
5. Gardner H, Dukes CD. *Haemophilus vaginalis*, a newly defined specific infection previously classified 'nonspecific vaginosis'. *Am J Obstet Gynecol* 1955;69:962–76
6. Priestley CJ, Kinghorn GR. Bacterial vaginosis. *Br J Clin Pract* 1996;50:331–4
7. Thomason JL, Gelbart SM, Anderson RJ, et al. Statistical evaluation of diagnostic criteria for bacterial vaginosis. *Am J Obstet Gynecol* 1990;162:155–60
8. Lamont RF, Hudson EA, Hay PE, et al. A comparison of the use of Papanicolaou-stained cervical cytological smears with Gram-stained vaginal smears for the diagnosis of bacterial vaginosis in early pregnancy. *Int J STD AIDS* 1999;10:93–7
9. Platz-Christensen JJ, Larsson PG, Sundstrom E, Wiquist N. Detection of bacterial vaginosis in wet mount, Papanicolaou-stained vaginal smears and in Gram-stained smears. *Acta Obstet Gynecol Scand* 1995;74:67–70
10. Saharan SP, Surve C, Raut V, Bhattacharya M. Diagnosis and prevalence of bacterial vaginosis. *J Postgrad Med* 1993;39:72–3
11. Eschenbach DA, Hillier S, Critchlow C. Diagnosis and clinical manifestations of bacterial vaginosis. *Am J Obstet Gynecol* 1988;158:819–28
12. Joesoef MR, Hillier SL, Josodiwondo S, Linnan M. Reproducibility of a scoring system for Gram stain diagnosis of bacterial vaginosis. *J Clin Microbiol* 1991;29:1266–71
13. Paavonen J, Heinonen PK, Aine R, et al. Prevalence of nonspecific vaginitis and other cervicovaginal infections during the third trimester of pregnancy. *Sex Trans Dis* 1986;13:5–8
14. Goncales-Falco J, Jurado M, Madambra AM, Oriol A. *Gardnerella vaginalis* and mosaic colposcopic

- pattern of the cervix: casual or causal association? *Gynecol Obstet Invest* 1986;121:108–10
15. Pavic N. Is there a local production of nitrosamines by the vaginal microflora in anaerobic vaginosis/trichomoniasis? *Med Hypotheses* 1984;15:433–6
 16. Hauth JC, Goldenberg RL, Andrews WW, et al. Reduced incidence of preterm delivery with metronidazole and erythromycin in women with bacterial vaginosis. *N Engl J Med* 1995;333:1732–6
 17. Hillier SL, Nugent RP, Eschenbach DA, et al. Association between bacterial vaginosis and preterm delivery of low-birth-weight infant. *N Engl J Med* 1995;333:1737–42
 18. Spiegel CA, Amsel R, Holmes KK. Diagnosis of bacterial vaginosis by direct Gram stain of vaginal fluid. *J Clin Microbiol* 1983;18:170–7
 19. Nugent RP, Krohn MA, Hillier SL. Reliability of diagnosing bacterial vaginosis is improved by a standardized method of Gram stain interpretation. *J Clin Microbiol* 1991;29:297–301
 20. Platz-Christensen JJ, Larsson PG, Sundstrom E, Bondeson L. Detection of bacterial vaginosis in Papanicolaou smears. *Am J Obstet Gynecol* 1989;160:132–3
 21. Prey M. Routine Pap smears for the diagnosis of bacterial vaginosis. *Diagn Cytopathol* 1999;21:10–3
 22. Bedrossian UK, Fairfax MR, Ayers M. Pap smear follow-up of possible role of mucopurulent exudate as a prognosticator of a negative pregnancy outcome. *Diagn Cytopathol* 1999;21:4–6
-

RECEIVED 07/06/00; ACCEPTED 04/01/02