Research Article

Alzheimer's Disease: APP, Gamma Secretase, APOE, CLU, CR1, PICALM, ABCA7, BIN1, CD2AP, CD33, EPHA1, and MS4A2, and Their Relationships with Herpes Simplex, *C. Pneumoniae*, Other Suspect Pathogens, and the Immune System

Chris Carter

PolygenicPathways, Flat 2, 40 Baldslow Road, Hastings, East Sussex TN34 2EY, UK

Correspondence should be addressed to Chris Carter, chris_car@yahoo.com

Received 3 August 2011; Accepted 2 September 2011

Academic Editor: Francesco Panza

Copyright © 2011 Chris Carter. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Alzheimer's disease susceptibility genes, APP and gamma-secretase, are involved in the herpes simplex life cycle, and that of other suspect pathogens (*C. pneumoniae*, *H. pylori*, *C. neoformans*, *B. burgdorferi*, *P. gingivalis*) or immune defence. Such pathogens promote beta-amyloid deposition and *tau* phosphorylation and may thus be causative agents, whose effects are conditioned by genes. The antimicrobial effects of beta-amyloid, the localisation of APP/gamma-secretase in immunocompetent dendritic cells, and gamma secretase cleavage of numerous pathogen receptors suggest that this network is concerned with pathogen disposal, effects which may be abrogated by the presence of beta-amyloid autoantibodies in the elderly. These autoantibodies, as well as those to nerve growth factor and *tau*, also observed in Alzheimer's disease, may well be antibodies to pathogens, due to homology between human autoantigens and pathogen proteins. NGF or *tau* antibodies promote beta-amyloid deposition, neurofibrillary tangles, or cholinergic neuronal loss, and, with other autoantibodies, such as anti-ATPase, are potential agents of destruction, whose formation is dictated by sequence homology between pathogen and human proteins, and thus by pathogen strain and human genes. Pathogen elimination in the ageing population and removal of culpable autoantibodies might reduce the incidence and offer hope for a cure in this affliction.

1. Introduction

Hundreds of genes have been implicated in Alzheimer's disease, many of which can be grouped into discrete signalling networks and pathways relevant to the various subpathologies, risk factors, and biochemistry of Alzheimer's disease. Many of the environmental risk factors associated with Alzheimer's disease, including infectious agents (herpes simplex, *chlamydia pneumoniae*, and *Borrelia burgdorferi*) as well as Vitamin A deficiency, hypercholesterolaemia, hyperhomocysteinaemia or folate deficiency, oestrogen depletion, cerebral nerve growth factor (NGF) deprivation, diabetes, cerebral hypoperfusion (leading to hypoxia and hypoglycaemia) or are able to promote cerebral beta-amyloid deposition (in the absence of any particular gene variant) in animal models [1]. KEGG pathway and other analyses of the multiple genes implicated in Alzheimer's disease have shown that

subsets of susceptibility genes can be grouped into networks that are relevant to each of these amyloidogenic pathways (e.g., bacterial and viral entry pathways [1, 2], cholesterol/lipoprotein function [3, 4], growth factor signalling [5], folate and homocysteine pathways [6], insulin signalling [7], and steroid or Vitamin A metabolism [8, 9]). A large number of genes are also related to the immune network [10] (see <http://www.polygenicpathways.co.uk/alzkegg.htm> and a recent review for further details [1]). These gene subsets are thus related to multiple external factors that are each able to promote beta-amyloid deposition, suggesting that certain genes are related to the causes of Alzheimer's disease, (agents able to provoke beta-amyloid deposition) rather than (and as well as) to the underlying pathology of the disease itself.

Several studies have implicated the herpes simplex virus in the aetiology of Alzheimer's disease [11–13]. Viral DNA is found in amyloid plaques [14], which are also heavily

enriched in proteins used by the virus during its life cycle, as well as in proteins related to the immune network [15], and Immunoglobulin IgM, but not IgG seropositivity for herpes simplex is predictive of the subsequent development of Alzheimer's disease [16]. IgM seropositivity is indicative of viral reactivation which again can be induced by several of the risk factors relevant to Alzheimer's disease and its underlying genetic pathways (e.g., NGF deprivation, 17-beta oestradiol, hypoxia, or fever and interleukin 6 activation, with the latter being common and general consequences of infection [1]).

Along with herpes simplex, a number of other pathogens have been implicated in Alzheimer's disease and its associated pathologies. The viral, bacterial, spirochete, and fungal pathogens implicated in dementia or Alzheimer's disease are referenced at (<http://www.polygenicpathways.co.uk/alzenvrisk.htm>) and include HHV-6, *Chlamydia pneumoniae*, *Helicobacter pylori*, periodontal pathogens involved in gum disease [17], *Borrelia burgdorferi*, and *Cryptococcus neoformans*. HIV-1 is also able to provoke dementia with Alzheimer's disease pathology [18]. Of these, *H. pylori* eradication has been reported to improve performance and increase lifespan in Alzheimer's disease patients [19], while two case reports indicated virtually complete recovery from long-term (3 years) misdiagnosed dementia/Alzheimer's disease following antifungal treatment for *C. neoformans* infection [20, 21]. Many of these pathogens including herpes simplex, HHV-6, *C. Pneumoniae*, *H. pylori* and the periodontal pathogen, *P. Gingivalis*, have also been implicated in atherosclerosis [22–25], while *C. neoformans* infection in rabbits induces an increase in neutrophil superoxide production, plasma lipid peroxidation, and an increase in inflammatory cells, forerunners of atherosclerosis [26]. Atherosclerosis of the carotid arteries, or of the circle of Willis and leptomeningeal arteries, is a significant predictor of risk in dementia or Alzheimer's disease and correlates with Alzheimer's disease pathology [27, 28]. Cerebral hypoperfusion (hypoglycaemia, hypoxia, ischaemia, or carotid occlusion) or other factors linked to atherosclerosis (e.g., high cholesterol or homocysteine levels) are also able, *per se*, to induce cerebral beta-amyloid deposition in animal models (see above).

Genomewide association studies (GWAS) have now identified a subset of genes which, along with **APOE4** [29], contribute a high proportion of genetic risk. These include clusterin (**CLU**), phosphatidylinositol-binding clathrin assembly protein (**PICALM**) and complement receptor 1 (**CR1**) as well as the ATP cassette transporter **ABCA7**, Bridging integrator **BIN1**, a CD2-associated protein (**CD2AP**), **CD33**, ephrin A1 (**EPHA1**), and a membrane-spanning 4-domains, subfamily A (MS4A) cluster recently honed down to **MS4A2**, although other genes within this cluster may also be relevant [30, 31].

As discussed below, the major Alzheimer's disease genes implicated by the recent GWAS data, as well as **APP** and **gamma secretase**, and previous GWAS results are majoritarily involved in pathogen entry and defence, particularly in relation to herpes simplex, but also to other relevant pathogens, and in the immune network. This suggests that genes, pathogens, and the immune system act together to cause

Alzheimer's disease, and that a focus on pathogen detection and elimination should be a priority in the ageing at risk population.

2. Methods

The genes identified in a number of recent genomewide association studies are available at the GWAS repository at the National Human Genome Research Institute <http://www.genome.gov/gwastudies/> [32] and, along with pre-GWAS genes and environmental risk factors, at <http://www.polygenicpathways.co.uk/alzenvrisk.htm>. The genes returned from very large sample sets ($N > 10,000$) include **ABCA7**, **APOE**, **BIN1**, **CD2AP**, **CD33**, **CLU**, **CR1**, **EPHA1**, **MS4A2**, **MS4A4A**, **MS4A4E**, **MS4A6A**, and **PICALM** whose properties in relation to diverse pathogens were identified by literature survey. While it is recognised that such genes, particularly **APOE**, **ABCA7**, **CR1**, and **clusterin**, which are involved in lipoprotein function and/or amyloid processing (see below), may exert effects on other relevant branches of Alzheimer's disease pathophysiology, the focus of this paper is on pathogens and the immune system, which appear to be the common factors integrating this network. Throughout the text, these and other genes implicated in Alzheimer's disease from the GWAS and pre-GWAS era are highlighted in bold and appended to the various processes in which they are involved (derived from a KEGG pathway analysis of these genes <http://www.polygenicpathways.co.uk/alzkegg.htm>) Herpes simplex binding proteins, and key interactors, currently numbering over 450, are stocked and referenced at <http://www.polygenicpathways.co.uk/herpeshost.html>. KEGG pathway analysis of this interactome is provided at <http://www.polygenicpathways.co.uk/HERPESKEGG.htm>. Expression data are provided in Figure 1 and are also hyperlinked to the BioGPS webserver <http://www.biogps.gnf.org/>, which provides general gene information and mRNA expression profiles for most human genes, based on custom arrays from 79 human issues [33, 34]. Predicted B-cell epitopes from human beta-amyloid (1–42), nerve growth factor (NP_002497.2), or the microtubule protein, **tau** (NP_001116538.2) were identified using the BepiPred server <http://www.cbs.dtu.dk/services/BepiPred/> [35] and their sequences compared with pathogen proteomes (*Borrelia burgdorferi*, *C. neoformans*, *Helicobacter pylori*, herpes viruses HSV-1, HSV-2, HHV-6, and the cytomegalovirus (HHV-5)) using the NCBI BLAST server (Protein versus protein: BlastP) [36].

3. Results

3.1. The Complement System (ABCA7, CR1, CLU, CD2AP, and Beta-Amyloid) Figure 2. Complement receptor 1 (highly expressed in myeloid **CD33+** cells (bone marrow) <http://www.biogps.org/#goto=genereport&id=1378/>) is a receptor for herpes simplex, adenovirus 5, the influenza virus and HIV-1, as well as for a number of other pathogens, including *P. gingivalis*, *C. neoformans*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, and the malaria parasite,

FIGURE 1: Continued.

FIGURE 1: Continued.

FIGURE 1: Continued.

FIGURE 1: Continued.

FIGURE 1: The mRNA distribution of the major genes derived from GWAS in Alzheimer's disease, as well as that of APP and gamma-secretase components. Data are from the BioGps website.

FIGURE 2: Portions of the complement cascade in relation to beta-amyloid processing: Alzheimer's disease susceptibility genes returned from very large genomewide association studies are in black, and those from the pre-GWAS era in grey. Binding interactions are indicated by linked diamonds and other effects by arrows. C. Neo: *Cryptococcus neoformans*; C. Pneu: *chlamydia pneumoniae*; H. Pyl: *Helicobacter pylori*; P. Ging: *Porphyromonas gingivalis*; C1–C9: complement components. MAC: membrane attack complex. P-choline: phosphatidylcholine; black star : Herpes simplex: see text for details.

Plasmodium falciparum [37–43] and is a general clearance receptor for complement opsonised pathogens [44]. **Clusterin**, predominantly expressed in brain, liver, and testis, (<http://www.biogps.org/#goto=genereport&id=1191/>) is a ligand for the lipoprotein receptor, megalin (**LRP2**) that is involved in beta-amyloid clearance, and also a complement inhibitor that prevents the formation of the membrane attack complex, a channel that is inserted into pathogen membranes, killing them by lysis [45]. This complex is also seen in Alzheimer's disease neurones [46, 47]. The herpes simplex virus interacts with other members of the complement cascade, by binding to the complement component and **CR1** ligand, C3 and its derivatives and to CD59, a further inhibitor of the formation of the complement membrane attack complex (see review) [48]. *C. pneumoniae* interacts with this pathway by binding to properdin (CFP), a protein that stabilises the complement C3 and C5 convertase and contributes to the formation of the membrane attack complex [49]. CD59 is also incorporated into chlamydial inclusion bodies [50]. Complement component C3 binds to melanins derived from *C. neoformans* [51] and cryptococcal capsules bind to C3 and activate the alternative complement pathway [52]. Complement component C3 also binds to the bacterial surface of *H. pylori*, and the complement pathway

is involved in bactericidal effects against this pathogen [53]. *P. gingivalis* also uses complement receptor 3 (an integrin complex of integrin, alpha M/integrin, beta 2 (ITGAM/ITGB2)) for entry [54], and herpes simplex glycoprotein C also binds to this complex [55] as does *C. neoformans* [56], while ITGB2 is involved in *C. pneumoniae* entry in human coronary artery endothelial cells [57]. This macrophage complement receptor, also known as MAC-1, generally mediates the phagocytosis of pathogens coated with complement C3 derivatives [58]. T. C3 also binds to *P. gingivalis* although the pathogen has devised an elegant escape strategy involving digestion of complement components C3, C4, and C5 by bacterial secreted proteases, known as gingipains [59].

The complement inhibitor CD59 is also a ligand for CD2, and CD59 activation of this receptor, presumably involving **CD2AP**, activates T cell receptor signalling resulting in the secretion of interleukins (**IL1A**, **IL2** and **IL6**) and granulocyte macrophage colony stimulating factor (CSF2) [60, 61].

ABCA7 plays a role in the complement-mediated activation of phagocytosis in macrophages. Complement component C1q, which binds to IgM or IgG complexed antigens (relevant to most pathogens), binds to macrophage calreticulin and **LRP1** and C1q binding to macrophages markedly increased the expression of both **LRP1** and **ABCA7**, effects

FIGURE 3: A schematic representation of clathrin-mediated endocytosis and intracellular transport pathways. Alzheimer's disease susceptibility genes returned from very large genome-wide association studies are in black, and those from the pre-GWAS era in grey. Binding interactions are indicated by linked diamonds and other effects by arrows. C. Pneu: *chlamydia pneumoniae*; H. Pyl: *Helicobacter pylori*; P. Ging: *Porphyromonas gingivalis*; black star : Herpes simplex: see text for details.

which enhance the phagocytic abilities of macrophages [62]. C1q also binds to complement receptor CR1, an effect involved in the immune clearance of opsonised pathogens [63]. C1q also binds to beta-amyloid and is involved in amyloid-related complement activation [64].

3.2. Clathrin-Mediated Endocytosis (BIN1, CLU, CD2AP, PICALM) Figure 3. Mammalian surface receptors are endocytosed, via clathrin-dependent or independent processes (KEGG: ADRB1, ADRB2, BIN1, CAV1, CD2AP, CLU, DNM2, HLA-A, HSPA1B, LDLR, NTRK1, PICALM) and either recycled or tagged for destruction by the ubiquitin/proteasome system (KEGG: UBD, UBE2I, UBQLN1, UCHL1) or by lysosomes (KEGG: ABCA2, ARSA, ARSB, CTSD, CTSS, NPC1, NPC2, LIPA). Early endosomes receive traffic from the cell surface, which is transferred to late endosomes for traffic to lysosomes. Late endosomes also receive traffic from the trans-Golgi network used to synthesise proteins and from phagocytic pathways (KEGG: CTSS, DLD, DLST, DNM2, GAB2, HLA-A, HLA-DRB1, MPO, NOS1, OLR1, PIK3R1, PSK1, TAP2, TLR2, TLR4). Endosomal traffic moves along the microtubule (GSK3B, MAPT, TTLL7) or actin/myosin (MYH8, MYH13) networks via dynein/dynactin (DNM2, DNMBP) or kinesin (KIF18B, KIF20B, KNS2), related motors and Rho GTPases, and vacuolar sorting proteins (SORCS1, SORCS2, SORCS3, SORL1) *inter alia* [65]. These processes are usurped by many viruses

and other pathogens to gain access to cells and to various intracellular compartments, while the lysosomal or proteasomal pathways may be used to destroy pathogen proteins [66].

Clathrin-mediated endocytosis is one of several processes used by *Helicobacter pylori*, herpes simplex, and many other viral, bacterial and fungal pathogens to gain entry to cells [67–69].

PICALM, expressed primarily in myeloid and dendritic cells of the immune network <http://www.biogps.org/#goto=genereport&id=8301/>, plays a key role in clathrin-related endocytosis, binding to clathrin heavy chains (CLTC and CLTCL1), and recruiting the clathrin and adaptor protein 2 (AP-2) to the plasma membrane. The AP-2 complex is a heterotetramer consisting of permutations of two large adaptins (alpha (AP2A1, AP2A2)) or beta (AP2B1), a medium adaptin (AP1M1, AP1M2), and a small adaptin (sigma AP2S1). **PICALM** controls the endocytosis of the cation-independent mannose-6-phosphate IGF2 receptor (IGF2R) [70], one used by Herpes simplex for entry and cell-to-cell transmission [71] and by *C. pneumoniae* for cellular entry [57]. IGF2R is also a component of late endosomes disrupted by the *Helicobacter pylori* VacA cytotoxin [72]. The mannose-6-phosphate receptor binds to **clusterin**. **PICALM** also binds to a nuclear exportin crm-1 (XPO1) used by the herpes simplex virus during its life cycle [48].

Gamma-adaptins (GGA, GGA2, GGA3) bind to clathrins and mannose-6-phosphate receptors and regulate protein

traffic between the Golgi network and the lysosome and the sorting of mannose-6-phosphate receptors (IGF2R and M6PR) at the trans-Golgi network [73]. This network is also related to important Alzheimer's disease susceptibility genes as the interactions culled from NCBI gene show that GGA1 binds to the sortilin-related receptor, **SORL1**, and the APP cleaving beta-secretase **BACE2**, while GGA2 binds to the beta-secretases **BACE1** and **BACE2**, **SORL1** and the prolyl-isomerase **PIN1**.

CD2AP, primarily expressed in dendritic cells and B lymphoblasts <http://www.biogps.org/#goto=genereport&id=23607/>, is a scaffolding molecule that regulates the actin cytoskeleton and is primarily associated with the T-lymphocyte marker protein CD2. CD2 stimulates T cell activation and is involved in the creation of contacts between antigen presenting cells and T cells (the immunological synapse), effects mediated via **CD2AP** and clathrin [74]. **CD2AP** is also involved in the entry of the helicobacter vacuolating toxin VacA and connects the actin cytoskeleton to early endosomes containing VacA [75]. CD2 is cleaved by gingipain proteases from *P. gingivalis* [76].

CD2AP also binds to the actin-bonding protein, cortactin (CTTN), a protein that is exploited by several bacteria (*Escherichia coli*, *Shigella*, *Neisseria*, *Rickettsia*, *Chlamydia*, *Staphylococcus*, *Cryptosporidium*, and *Helicobacter pylori*), fungi (*Candida Albicans*), and viruses (Vaccinia) enabling them to modify the actin cytoskeleton, which they use for transport [77–79]. **CD2AP** has not been specifically associated with herpes simplex, although the actin cytoskeleton is exploited by this and many other viruses [80].

CD2AP also associated with E-Cadherin, (CDH1) [81]. The ectodomain of E-cadherin is involved in bacterial adherence to mammalian cells [82]. E-Cadherin binds to the *H. pylori* toxin CagA [83] and is also cleaved by the *Helicobacter pylori* protein HtRA allowing the pathogen to invade the intracellular compartment [84]. CDH1 and CDH5 expressions are increased by *C. pneumoniae* infection of human brain microvascular endothelial cells, contributing to vascular permeability changes and atherosclerosis [85].

Bridging integrator 1 (**BIN1**), also known as amphiphysin 2, is primarily expressed in the pineal and skeletal muscle, or otherwise ubiquitously <http://www.biogps.org/#goto=genereport&id=274/>. It is also involved in the clathrin-mediated endocytosis machinery [86] and binds to dynamin that regulate the clathrin network [87] including DNM1 and the herpes simplex binding partner **DNM2** [88] and to clathrins and the alpha adaptins, AP2A1 and AP2A2 [89]. **BIN1** also participates in phagocytosis in macrophages and is associated, but only transiently, with early phagosomes; however, it is retained on vacuoles containing *Chlamydia pneumoniae*, an effect that reduces the ability of the macrophage system to kill the bacteria via nitric oxide generation. Macrophages expressing a dominant negative **BIN1** internalise *C. pneumoniae*, but do not allow their killing [90]. **BIN1** also binds to a number of alpha integrins (ITGA1, ITGA3, and ITGA6) [91]: integrins are used for attachment by many viruses, bacteria, and fungi and may serve as pattern recognition receptors regulating the immune response [92]. Individual integrins bind to many

others, forming heteromeric complexes; for example, ITGA1 binds to ITGA3 or ITGA6, while ITGA3 binds to ITGB1 (a receptor for the *H. pylori* protein CagA [93]), ITGB4, or ITGB5, and ITGA6 binds to ITGB1 and ITGB4 (data from NCBI gene).

3.3. *The Immune Network* (APOE, **BIN1**, **CD2AP**, **CD33**, **MS4A2**) (Figure 4). **CD33**, mainly expressed in myeloid cells, monocytes, and dendritic cells (<http://www.biogps.org/#goto=genereport&id=945/>), is a member of the sialic acid binding Immunoglobulin g-like lectin (SIGLEC) family. **CD33**-related SIGLECs regulate adaptive immune responses and are also important as macrophage pattern recognition receptors for sialylated pathogens, including enveloped viruses [94]. **CD33** binds to alpha2-3- or alpha2-6-linked sialic acids (N-acetyl neuraminic acid) [95]. These particular sialic acids are expressed on the surface envelope glycoproteins (B, D, and H) of the herpes simplex virion, and these residues are required for viral entry into cells [96]. N-acetyl neuraminic acid is expressed by *C. neoformans*, is involved in fungal adhesion to macrophages [97], and is also a component of the cell wall of *B. burgdorferi* [98], while *Helicobacter pylori* adhesins also bind to this particular form of sialic acid [99, 100] as does *P. gingivalis* [101].

BIN1, as well as its relationship to the clathrin mediated endocytosis machinery, also regulates the expression of indoleamine 2,3-dioxygenase (IDO1), an enzyme that catalyzes the first rate-limiting step in tryptophan metabolism to N-formyl-kynurenine [102]. IDO1 upregulation is an important defence mechanism against pathogenic bacteria, many of which are unable to synthesise tryptophan. Their survival is compromised by the diversion of tryptophan metabolism to kynurenines [103]. This IDO1 response is also deleterious to other pathogens and parasites, including *T. gondii*, and to a number of viruses, including herpes simplex and other herpes viruses [104]. IDO1 protein expression is localised to plaques and tangles in the Alzheimer's disease brain. IDO1 activation can lead to the production of toxic tryptophan derivatives such as 3-hydroxyanthranilic acid or the N-methyl-D aspartate receptor agonist and excitatory neurotoxin, quinolinic acid [105] (**GRIN2B**, **GRIN3A**). Plasma tryptophan levels are also lower in the ageing population and in Alzheimer's disease, a pattern accompanied by immune activation, and by increased concentrations of quinolinic acid [106, 107].

MS4A2, expressed mainly in the tonsils, lymph nodes, B cells, and dendritic cells <http://www.biogps.org/#goto=genereport&id=931/>, is a component of the immunoglobulin E (IgE) receptor, which is involved in allergic responses in which allergens bound to receptor bound IgE result in the activation of allergic mediators such as histamine [108]. Mice immunised with inactivated herpes simplex develop IgE-specific antibodies to the virus [109]. High levels of IgE are also observed in man following recurrent herpes simplex infection [110] and human IgE antibodies are also known to interact with herpes family viruses including HSV-1 and 2 and the Epstein-Barr and cytomegalovirus [111] and also to *C. pneumoniae*, *H. pylori*, and *B. burgdorferi*

EPHA1 is an ephrin receptor, primarily expressed in the liver and otherwise ubiquitously (<http://www.biogps.org/#goto=genereport&id=2041/>). Only three protein/protein interactions for **EPHA1** are reported in the NCBI gene interaction section, including its ligand EFNA1, the anaplastic lymphoma receptor tyrosine kinase (ALK), and a SMAD-specific E3 ubiquitin protein ligase 2 (SMURF2). EFNA1 is one of several proteins identified as being important in the entry of *C. pneumoniae* into human coronary artery endothelial cells [57]. SMURF2 is known to bind to the VP22 tegument protein of herpes simplex [119] and plays a role in clathrin-mediated endocytosis and the subsequent ubiquitin-related proteasomal degradation of **TGF beta** receptors, to which it binds [120]. **Clusterin** is a ligand for **TGF beta** receptors (TGFBRI/TGFBRII) [121]. **TGF beta** signalling exerts immunosuppressive effects and inhibits host immunosurveillance and the recruitment of immunocompetent cells by chemokines [122]. ALK is ubiquitously expressed (<http://www.biogps.org/#goto=genereport&id=238/>). It plays a role in neural development, and its expression decreases with age [123]. ALK is best

3.4. *Lipoprotein Related (APOE, ABCA7, CLU)* (Figures 2 and 5). **ABCA7** is an ATP-binding cassette transporter, predominantly localised in the pineal gland and cells of the immune network (T cells, natural killer cells, and dendritic cells <http://www.biogps.org/#goto=genereport&id=10347/>). The lipoproteins **APOA1** and **APOE** are substrates for **ABCA7**, and in cultured HEK-293 cells, plasma membrane-situated **ABCA7** increases the efflux of phosphatidylcholine and sphingomyelin efflux to **APOA1** and **APOE**, with no effect on cholesterol efflux [125]. However, cholesterol efflux to lipid-laden **APOE**, but not to lipid free **APOE**, is increased by **ABCA7** expression in HEK-293 cells [126]. Sphingomyelin is enriched in extracellular herpes simplex viral membranes: this sphingomyelin, together with phosphatidylserine, is collected by the viral envelope during viral passage from the nuclear membrane to the exocytosis pathway [127]. Herpes viral infection leads to an increased incorporation of phosphate into membrane sphingomyelin of the host [128]. Inhibition of sphingomyelinase has also been shown to markedly reduce herpes simplex viral reproduction [129].

and also inhibits the antifungal effects of neutrophils against *C. neoformans* infection. Sphingomyelin is a receptor for the *Helicobacter* toxin VacA [130] and is also incorporated into inclusion bodies in *C. pneumoniae*-infected cells [131]. Phosphatidylcholine plays an important role in the fusion of herpes simplex glycoproteins B and H with the host cell lipid membrane, a process used in viral entry [132]. Phosphatidylcholine is also able to trigger capsular enlargement in *C. neoformans* infection [133].

ABCA7 expression increases the extracellular surface deposition of ceramide (derived from sphingomyelin) [134]. Ceramide, a potent activator of apoptosis, as well as its downstream target, caspase 3 (CASP3) are both able to reactivate the herpes simplex virus from latency [135]. Ceramide is also incorporated into *C. pneumoniae* inclusions, an effect that may play a role in the antiapoptotic effects of this bacterium [136]. **APOA1** exerts antiviral effects against herpes simplex and inhibits viral entry into cells as well as viral-induced cell fusion and intercellular spread [137]. In macrophages, **ABCA7** is expressed intracellularly and does not participate in cholesterol or phospholipid efflux, instead playing a role in the phagocytosis of apoptotic cells, an important general defence mechanism against invading pathogens [62, 138].

3.4.1. Apolipoprotein E. Possession of the **APOE4** allele facilitates the entry and transmission of herpes simplex in mice models [139]. In man, **APOE** is also involved in hepatitis C, HIV-1, and herpes simplex infectivity [140–143], and **APOE4** facilitates the binding of *C. pneumoniae* elementary bodies to host cells [144].

APOE mRNA is primarily expressed in the liver, adipocytes; kidney and brain, with very low expression in the peripheral immune network (<http://www.biogps.org/#goto=genereport&id=348/>) but nevertheless plays an important role in the immune system. For example, the presence of the **APOE4** allele is associated with an enhanced macrophage inflammatory response, and cytokine responses to the intracerebral injection of lipopolysaccharide are increased in **APOE4** transgenic mice, which also exhibit increased microglial activation. The anti-inflammatory effects of 17-beta-oestradiol on microglia are also reduced in such animals [145, 146]. C-reactive protein (**CRP**) levels are also decreased in **APOE4** carriers [145–147]. **CRP** is an acute phase protein that binds to phosphocholine on dead or dying cells and on bacteria, subsequently activating the complement pathway [148]. Resistance to infection (*Klebsiella pneumoniae*) or endotoxaemia is also decreased in **APOE** knockout mice [149].

In addition, atherosclerosis is induced or worsened by infection with a number of relevant pathogens (Cytomegalovirus, herpes simplex, *Helicobacter pylori*, influenza, *C. pneumoniae* or *P. gingivalis*) in **APOE** knockout mice [150–156]. *Helicobacter pylori* is able to promote atherosclerosis in heterozygous **APOE** (+/–) **LDLR** (+/–) mice, which is associated with an immune response to the bacterial heat shock protein hsp60 [157].

3.5. Other GWAS Genes (Figure 4). Prior to the very large GWAS collaboration, several other genes had been identified

FIGURE 5: Gamma-secretase-mediated cleavage of viral and pathogen receptors. **Gamma secretase** substrates are indicated by the square boxes and their pathogen ligands in the oval boxes. Cyto: cytomegalovirus; C. Neo: *Cryptococcus neoformans*; Adeno: adenovirus; EBV: Epstein-Barr virus; Flu: influenza A.; HepC: hepatitis C; HepE: hepatitis E; HHV-6: human herpesvirus 6; HIV-1: human immunodeficiency virus; HSV-1: herpes simplex; Lassa: Lassa fever virus; LCMV: Lymphocytic choriomeningitis virus; Leprosy: *mycobacterium leprae*; Meas: measles; Neiss: *Neisseria*; Papill: papillomavirus; P. Ging: *Porphyromonas gingivalis*; Pox: Vaccinia and other pox viruses; Rhino: rhinoviruses; Staph: *Staphylococcus aureus*; Strep: *streptococcus*; P. serine: phosphatidylserine; Alzheimer's disease susceptibility genes returned from very large genomewide association studies, and APP, are in black, as are the pathogens implicated in Alzheimer's disease. Genes from the pre-GWAS era are in grey.

in smaller genomewide studies (**APOC1**, **CELF2**, **DISC1**, **FAM113B**, **GAB2**, **MTHFD1L**, **PAX2**, **PCDH11X**, **PVRL2**, **RFC3**, **SASH1**, **TOMM40**, **TTL7**, and **ZNF224**). **PVRL2** is a receptor for herpes simplex (HSV-1 and HSV-2) [158], and the mitochondrial translocator, **TOMM40**, a receptor for certain chlamydial species [159]. The replication factor **RFC3** is part of a complex necessary for human DNA polymerase activity, a process exploited by many viruses including herpes simplex, whose virion component ICP34.5 binds to proliferating cell nuclear antigen (PCNA), an **RFC3** binding partner and also a cofactor for DNA polymerase [160]. **ZNF224** is a transcriptional repressor binding to the protein arginine methyltransferase, **PRMT5** [161]. Protein arginine methylation is important in viral infection and replication, as well as in cytokine signalling, and a related arginine methyltransferase, **PRMT1**, regulates herpes simplex replication via methylation of the ICP27 viral gene [162].

DISC1 is a component of the microtubule-associated dynein motor complex used in viral traffic [163]; **TTL7** (tubulin tyrosine ligase-like family, member 7) also regulates tubulin phosphorylation [164] and can again be related to viral traffic along the microtubule network (see below). **CELF2** (also known as CUGBP2) is a member of the APOBEC1 cytidine deaminase mRNA editing complex that also controls herpes simplex viral replication [165]. **GAB2**

is a member of the GRB2-associated binding protein family which act as adapter hubs transmitting signalling via cytokine and growth factor receptors, and T- and B-cell antigen receptors (definition from NCBI gene), while **PAX2** inhibits the expression of the antimicrobial peptide beta defensin (DEFB1) [166], a gene associated with HSV-1 and cytomegalovirus seropositivity in children with acute lymphoblastic leukaemia [167], as well as with *H. pylori* or chlamydial infections [168, 169], also endowed with antimicrobial activity against *C. neoformans* and other pathogens [170]. **MTHD1L** (methylenetetrahydrofolate dehydrogenase (NADP+ dependent) 1-like) is involved in the mitochondrial synthesis of tetrahydrofolate which in turn is important in the de novo synthesis of purines and thymidylate and in the regeneration of methionine from homocysteine (definition from NCBI gene). Many pathogens, including herpes simplex, express thymidylate kinases, which are important for viral replication and a target for acyclovir [171]. Hyperhomocysteinaemia correlates with *C. pneumoniae* IgG immunoreactivity in carotid artery atherosclerosis [172] and is also associated with *H. pylori* infection in the context of atherosclerosis [173]. The apolipoprotein **APOC1** is a component of high-density lipoprotein: herpes simplex is present in all lipoprotein blood fractions in blood (VLDL, LDL and HDL) and the lipid component of these lipoproteins binds to viral glycoprotein B [174] (c.f. **APOA1**, **APOA4**, **APOA5**, **APOC1**, **APOC2**, **APOC3**, **APOC4**, **APOD**, and **APOE**). No immediately apparent pathogen-relevant interactions were found for **FAM113B** (expressed exclusively in T cells, dendritic cells, and natural killer cells <http://www.biogps.org/#goto=genereport&id=91523/>), **PCDH11X** (which is ubiquitously expressed <http://www.biogps.org/#goto=genereport&id=27328/>), or **SASH1** (primarily expressed in the brain and lung although also in other tissues, including the immune network <http://www.biogps.org/#goto=genereport&id=23328/>) although the pathogen/immune theme is clearly carried through, particularly in relation to herpes simplex, in this second rank of Alzheimer's disease susceptibility genes.

3.6. Beta Amyloid Processing (Figure 2). **APOE**, **clusterin**, and **complement receptor 1** play key roles in beta amyloid clearance as do two further herpes simplex binding proteins **APOA1**, and alpha-2 macroglobulin (**A2M**). This is primarily mediated via lipoprotein receptors. **A2M**, or **APOE**-bound $A\beta$, is cleared by the lipoprotein receptor **LRP1**, while **LRP2** (megalin) clears clusterin-bound $A\beta$. **LRP8** is a receptor for both **APOE** and **clusterin**. **APOA1** is also involved in beta-amyloid clearance via its transporter **ABCA1**. The role of **ABCA7** has not been examined, although **APOA1** is also a ligand for this transporter (see above). The Varicella Zoster and herpes simplex glycoprotein E binding protein, **insulin-degrading enzyme**, are also involved in beta-amyloid degradation, as is caspase-3 which is activated by the herpes simplex viral US3 kinase. The HSV-1 binding protein, complement C3 is also a ligand for **LRP1** and **LRP8**, both of which play a role in C3 cellular uptake. Beta amyloid in the bloodstream is processed by its binding to complement C3, which subsequently binds to

complement receptor 1 on erythrocytes. The effects above are referenced in a recent review [48].

Clathrin-dependent endocytosis is also involved in the internalisation and recycling of neuronal **APP**, a procedure necessary for the subsequent cleavage of **APP** and the generation of beta-amyloid [175, 176], and in the neuronal [177], but not the microglial uptake of both soluble and aggregated beta-amyloids, the latter representing an important route of disposal [178]. However, while knockdown of the clathrin assembly protein AP180 in a neuronal cell line does reduce beta-amyloid generation, **PICALM** knockdown does not [179]. The accumulation of beta-amyloid in the brain interstitial space, related to the prior endocytosis of **APP**, is clathrin dependent [180]. Clathrin-mediated endocytosis is relevant to many receptors, including members of the lipoprotein family (**LRP1**, **LRP2**, **LRP8**, **LDLR**, **VLDLR**) [181], all of which are involved in beta-amyloid clearance, as well as in cholesterol and lipoprotein physiology [182].

ABCA7 plays a role in beta-amyloid secretion, which is increased in Chinese hamster ovary cells expressing **APP** and **ABCA7**. This was related to an effect on **APP** intracellular retention, rather than on secretase-mediated proteolysis of **APP** [126].

Gamma-secretase cleaves **APP**, and other gamma-secretase substrates also play key roles in **APP** processing (**ADAM10**) [183], lipid and cholesterol function (**LRP1**, **LDLR**, **VLDLR**), and other processes relevant to Alzheimer's disease, for example, **NOTCH** signalling [184].

No other immediately apparent relationships with beta-amyloid could be found by literature survey for **CD2AP**, **CD33**, and **EPHA1** or for **MS4A**-related proteins, although such are not precluded.

3.6.1. The Microtubule Network and Tau Phosphorylation. Many pathogens, including herpes simplex, *helicobacter*, *chlamydiae*, *P. gingivalis*, and *C. neoformans* [185–188], use the microtubule network that serves as a useful railway track between various cellular compartments, and may hijack dynein and kinesin motors for this purpose (see <http://www.polygenicpathways.co.uk/herpeshost.html> for herpes simplex). **Tau** (**MAPT**) stabilises microtubules by interacting with tubulins and promoting microtubule assembly [189]. When **tau** is phosphorylated, by any of several kinases, microtubules become disorganised. **Tau** hyperphosphorylation and neurofibrillary tangles are among the core pathologies of Alzheimer's disease [190] and can be promoted by herpes simplex infection [191]. In relation to viral/human protein homology, herpes simplex proteins are homologous to a number of kinases known to phosphorylate **tau** (**GSK3A**, **GSK3B**, **MAPK1**, and **CAMK2B**) suggesting that **tau** phosphorylation could be a direct result of a viral kinase [192].

3.6.2. APP and Gamma Secretase. **APP** plays a key role in the herpes simplex life cycle and is involved in its intracellular transport [193], an effect likely related to the ability of both **APP** and the herpes simplex protein, **US11**, to bind to the **APP** and kinesin binding protein **APPBP2** (also known as **pat1**) [194, 195].

The AntiMicrobial Effects of Beta-Amyloid. Beta-amyloid is an antimicrobial peptide with broad spectrum activity against a variety of yeasts and bacteria, effects that were attenuated by anti-A β antibodies [196]. Beta-amyloid also has antiviral effects and, like acyclovir, attenuates the stimulatory effects of herpes simplex on miRNA-146a levels in neuronal cells [197]. Beta-amyloid also activates innate immune responses via the activation of pattern recognition receptors, such as Toll receptors (**TLR2**, **TLR4**), which are also involved in beta-amyloid clearance [198, 199]. The antimicrobial, antiviral, and immunostimulant properties of beta amyloid are, however, likely to be abrogated by the presence of beta-amyloid autoantibodies in the sera of the ageing population and in Alzheimer's disease [200]. As immunogenic regions of beta-amyloid are homologous to similar regions within proteins expressed by all of the principal pathogens discussed in this paper, such antibodies are likely to be derived from antibodies raised to numerous pathogens (see below).

Gamma Secretase: Localisation to Dendritic Cells and Cleavage of Pathogen Receptors. **Gamma secretase** is constituted of four components: the presenilins (**PSEN1** or **PSEN2**), anterior pharynx-defective-1 (**APH1A**), the Presenilin enhancer-2 (**PSENEN**), and nicastrin (**NCSTN**) [201]. While all components are expressed in cerebral tissue, the major focus of distribution is within cells of the immune network; dendritic cells, myeloid cells, and monocytes for **PSEN1** <http://www.biogps.org/#goto=genereport&id=5663/>; dendritic cells and natural killer cells for **PSENEN** <http://www.biogps.org/#goto=genereport&id=55851/>, dendritic cells and myeloid cells for nicastrin <http://www.biogps.org/#goto=genereport&id=23385/> and B cells, dendritic cells, natural killer cells, and myeloid cells for **APH1A**. The substrate, **APP**, is the only gene in this set that appears to be preferentially distributed in brain compartments, but, as **with gamma-secretase** components, it is also highly expressed in dendritic cells of the immune system (<http://www.biogps.org/#goto=genereport&id=351/>). The primary function of such cells is to process antigens and present them to B cells and T cells. They scout for and recognise pathogens via the agency of numerous pattern recognition receptors, for example, Toll receptors (**TLR2**, **TLR4**), or viral DNA sensors, expressed on their surface [202, 203].

As well as cleaving **APP**, **gamma secretase** is involved in the proteolysis of at least three herpes simplex receptors, nectin 1 alpha (PVRL1) [204], and syndecans (SDC1, SDC2) [205]. SDC1 is also a receptor for HIV-1, Hepatitis E, and the human papillomavirus [206–209], while SDC3, also a **gamma secretase** substrate, is an HIV-1 and papillomavirus receptor [210, 211].

Several other **gamma secretase** substrates (reviewed by Lleó and Saura [201]) also function as viral/pathogen receptors, including ADAM10, a receptor for the cytotoxin *Staphylococcus aureus* alpha-haemolysin [212], CD44, an entry receptor for *C. neoformans* [213], CD46, a receptor for adenoviruses, measles virus, human herpes virus 6 (HHV-6), Streptococci, and Neisseria [214]; CD46 is also cleaved

by a protease secreted by *P. gingivalis* [215]. Desmoglein-2 is an adenovirus receptor [216], while rhinovirus receptors include the lipoprotein receptors **LRP1**, **LDLR**, and **VLDLR** [217, 218]. **LDLR** is also a hepatitis C receptor [219]. NOTCH1 and NOTCH4 are activated by the Epstein-Barr virus [220, 221], while ERBB4 is a receptor for vaccinia and other pox viruses [222]. The low affinity nerve growth factor (NGFR) is a rabies virus receptor, [223], Ephrin B2, (EFNB2) a Nipah virus and Hendra virus receptor [224] and sialophorin (SPN), a receptor for the influenza A, and both human and simian immunodeficiency viruses [225, 226] and for the *C. neoformans* virulence factor, galactoxylomannan [227]. Fractalkine (CX3CL1) binds to the cytomegalovirus chemokine receptor, US28 [228], while the chemokine CXCL16 is a scavenger receptor for phosphatidylserine and oxidized low density lipoprotein [229]: phosphatidylserine is the major lipid membrane component in most bacteria [230]. Dystroglycan is a receptor for the Lymphocytic choriomeningitis and Lassa fever viruses [231, 232] and also for *Mycobacterium leprae* (the Leprosy pathogen) [233].

The antimicrobial and immunostimulant effects of beta-amyloid, the cleavage of a number of pathogen receptors by **gamma secretase**, and the concentration of both **APP** and **gamma-secretase** components in dendritic cells suggest that a major function of this key group, implicated in both familial and late onset Alzheimer's disease, is dedicated to pathogen defence, and that increased beta-amyloid generation is primarily a defence mechanism to rid the body (and brain) of invading pathogens: This scenario is supported by the ability of herpes simplex, *C. pneumoniae*, and *B. burgdorferi* to increase beta-amyloid deposition [234–236]. One might expect many other pathogens to increase beta-amyloid deposition, and that, as has been noted in atherosclerosis, (a component of Alzheimer's disease pathology), the final extent of risk may depend upon the overall pathogen burden, rather than upon any specific pathogen [237].

3.6.3. Autoantibodies Derived from Pathogens as Contributory Causative Agents.. Viruses and bacteria express proteins containing short contiguous amino acid stretches (pentapeptides or more, or longer gapped consensi) that are identical to those in human proteins: these pathogen/human consensi number in millions and concern all human proteins [238–241].

Autoantibodies, which are observed in many, if not most human diseases, are often regarded as an epiphenomenon of little consequence. However, they can traverse the blood brain barrier [242] (which is compromised in Alzheimer's disease [243]) and are also able to enter cells, essentially by hitching a ride on viruses, via high affinity IgG receptors (Fc gamma receptors) (**FCER1G**) in the case of the rhinovirus, or the SARS coronavirus, or via the tripartite motif protein, TRIM21, in the case of adenoviruses, where they are able to activate an intracellular immune attack. It would appear that the cellular entry of antibody laden viruses is diverted from their usually preferred receptors towards those used by antibodies [244–246]. This may be relevant to the **MS4A** family. Fc gamma receptors are localised in microglia and

astrocytes in the brain and their expression is upregulated by blood brain barrier disruption [247], while TRIM21 appears to be exclusively localised in peripheral immunocompetent cells (<http://www.biogps.org/#goto=genereport&id=6737/>).

This ability places autoantibodies in a rather more sinister context, as their targeting of extracellular and intracellular human proteins would be expected to effect protein knockdown, a strictly immunopharmacological effect, as well as immune attack.

In multiple sclerosis, schizophrenia, and cystic fibrosis, as well as in Alzheimer's disease, numerous autoantigens targeted by the autoantibodies reported in these conditions, contain peptide sequences identical to those in the pathogens also implicated in the disease. Such regions of homology are focalised within epitope regions of the human autoantigen [192, 248–250]. In Parkinson's disease, antibodies to the Epstein-Barr virus, which has been implicated in postencephalitic adult and juvenile Parkinsonism, are also known to cross-react with synuclein, a key protein involved in neurodegeneration in this disorder [251–253]. In addition, 22 autoantigens reported in HIV-1/AIDS contain HIV-1/human matching sequences [254], supporting the contention that autoantibodies are in many cases antibodies initially raised to pathogens, which because of this homology, then target their human homologues. It has been argued that slightly dissimilar, rather than exact matches, are the more malignant in terms of autoimmunity, being less likely to be regarded as self, while the antibodies would retain low affinity for human counterparts [254, 255].

Autoantibody production would also be sustained, even after pathogen elimination, by continued encounter of the human homologue. The production of autoantibodies must be dependent upon the extent of pathogen/human matching, and thus by genes which determine human protein sequences. These pathogen/human matches are also highly and significantly enriched in the products of susceptibility genes implicated in Alzheimer's disease, multiple sclerosis, and schizophrenia [192, 248, 250]. Many genes related to Alzheimer's disease, including those described above, are involved in the immune network [10, 256], and the propensity for developing autoantibodies to particular proteins is also genetically determined and inherited [257]. Thus, despite the fact that all human proteins likely possess pathogen homologues, whether or not autoantibodies will be produced will depend on the extent of human/pathogen matching (determined by human genes and the strain of pathogen encountered), on whether the pathogen protein is deemed as self or nonself (a factor determined soon after birth) and on other genetic factors related to the immune network, and autoimmunity. Somatic hypermutation, that drives the creation of multiple antibodies and which selects against those reacting to self, is disrupted in autoimmune disorders [258]. These links suggest an interplay, applicable to many diseases, where susceptibility gene products, risk promoting pathogens and autoimmunity can all be related via protein sequence homology.

It has also been noted that autoantigens have a tendency to relate to proteins known to bind to dermatan sulphate, a component of dead cells [259] and a constituent of

glycosaminoglycan receptors for many bacteria and viruses [260].

3.6.4. Sequence Comparisons: Beta-Amyloid, NGF, and Tau versus Pathogen Proteins. All three of these proteins are autoantigens in Alzheimer's disease and were chosen for analysis because of the ability of their antibodies to promote features of Alzheimer's disease, *in vivo*. In mice, immunisation with neuronal **tau** produces neurofibrillary tangle-like structures, axonal damage, and gliosis, as in Alzheimer's disease [261]. In addition, in transgenic mice, initially expressing NGF antibodies only in lymphocytes, NGF antibodies subsequently enter the brain provoking extensive cortical degeneration, cholinergic neuronal loss, **tau** hyperphosphorylation, and beta-amyloid deposition [262]. Beta-amyloid autoantibodies are also able to promote meningoencephalitis, both in laboratory models and in clinical trials [263, 264]. Beta-amyloid plaques contain numerous inflammatory proteins, and even without meningoencephalitis, these are commonly found within the walls of meningeal and medium-sized cortical arteries in Alzheimer's disease [265].

Almost the entire length of the **tau** protein (638/776 amino acids = 82.2%) was predicted as immunogenic (B cell epitope), as defined by the server set cutoff index of 0.35. For the analysis in Table 1, only regions of the **tau** protein with an immunogenicity index >2.5 were examined for homology. For other proteins (NGF and beta-amyloid), the analysis concerned immunogenic regions above the cutoff value of 0.35.

All pathogens express proteins with homology to each autoantigen, specifically within predicted B-cell epitope regions of the human autoantigen (Table 1), suggesting that antibodies raised to any could be responsible for targeting these human proteins, under the appropriate circumstances. Perversely, the successful elimination of the pathogen via antibody production could set in motion the very autoimmune responses that may be crucial to the development of Alzheimer's disease, a Pyrrhic victory, which would also be promulgated by any further encounter with these very common pathogens, or by structurally related proteins from other pathogens, as well as by continual encounter of the human autoantigen homologue (see Section 3.6.5).

As well as autoantibodies to these three proteins, a number of autoantibodies targeting highly relevant proteins have been reported in Alzheimer's disease. These have functional effects on their target proteins and include antibodies that block the activity of ATP synthase, induce apoptosis, and increase the cellular uptake of high density lipoprotein [266, 267], antibodies to cholinergic neurones that cause immunolysis of brain synaptosomes [268], anti-brain antibodies that enhance intraneuronal beta-amyloid deposition [269] as well as antibodies to the nicotinic receptor **CHRNA7**, that displace its ligand alpha-bungarotoxin [270]: autoantibodies to the receptor for advanced glycation products (**AGER**) [271], to the antimicrobial peptide **S100B**, have also been reported [272].

3.6.5. Population Genetics: Susceptibility Genes Related to the Cause of the Disease, rather than to the Disease Itself. Using

TABLE 1: Viral, bacterial, and fungal protein homology with beta-amyloid (top), NGF (middle), or *tau* (bottom); Predicted B cell epitope segments were compared with *Borrelia burgdorferi*, *C. neoformans*, *C. pneumoniae*, *H. pylori*, *P. gingivalis* and herpes viruses (HSV1, HHV6, and cytomegalovirus) proteomes by BLAST analysis. The B cell antigenicity indices are shown, and those above the server set threshold of 0.35 are in *italics*. The first column shows the amino acid number within the peptide or protein sequence, and the second shows the amino acid pertaining to that position. The alignments with pathogen proteins are shown. Spaces represent nonidentical amino acids and + signs amino acids with similar physicochemical properties. Only highly antigenic regions of pentapeptides or more were processed. The VGGVV sequence, antibodies to which label beta-amyloid in brain tissue, despite relatively low antigenicity, has already been reported to be identical to proteins expressed by 69 viruses including HSV-1, HSV-2, and HHV6.

(a)

Position B-Amy	Amino acid	B-cell index	Alignments
1	D	<i>0.41</i>	<i>C. neoformans</i> +AE HDSG+ <i>Borrelia burgdorferi</i> DAE F H+SG EV <i>H. pylori</i> DA FRH HSV-1 +AE RH HHV-6 D FR DS <i>P. gingivalis</i> +AEFR <i>C. pneumoniae</i> DA EFRHD and +AEFR +SG
2	A	<i>0.35</i>	<i>C. neoformans</i> AEFR D GY+V <i>H. pylori</i> AEF D S YE and AE+RH+ <i>Borrelia burgdorferi</i> AEF H+ Cytomagalovirus AEFR HD HSV-1 AE R SG HHV-6 AE+ HD <i>P. gingivalis</i> A+F H+S and AEFR <i>C. pneumoniae</i> AEF DSG
3	E	<i>0.62</i>	<i>H. pylori</i> EFRHD HHV-6 EF DSG <i>Borrelia Burgdorferi</i> EFR DS <i>C. neoformans</i> EF R DS YE <i>P. gingivalis</i> E R DSGY V <i>C. pneumoniae</i> EF SGYEV
4	F	<i>0.73</i>	<i>C. neoformans</i> FRHDS <i>Borrelia burgdorferi</i> +RH SGY++ and F H+SG <i>H. pylori</i> F HD EV Cytomegalovirus FR SGY <i>P. gingivalis</i> +RHDS <i>C. Pneumonia</i> F H+SGY
5	R	<i>0.85</i>	<i>C. neoformans</i> R D GYEV <i>H. pylori</i> RHDS Y V and R SGYE <i>Borrelia burgdorferi</i> RH+ GY Cytomegalovirus RHD YE and RHD SG HSV-1 RH SG HHV-6 RHDS <i>P. gingivalis</i> R+DS Y+
6	H	<i>0.57</i>	<i>C. neoformans</i> HDSGY <i>H. pylori</i> <i>H. pylori</i> +DSGY and HD G EV and HD EV <i>Borrelia burgdorferi</i> H+SG Y+V HSV-1 HDSG <i>P. gingivalis</i> HDSG <i>C. pneumoniae</i> ++SGY+V
7	D	<i>0.69</i>	<i>C. neoformans</i> DSGY+V <i>H. pylori</i> +SG+EV HSV-1 DSGY <i>P. gingivalis</i> DSG+EV <i>C. pneumoniae</i> DSGY V
8	S	<i>0.38</i>	<i>P. gingivalis</i> SGYEV <i>H. pylori</i> SGYE <i>C. neoformans</i> SGY++ <i>C. pneumoniae</i> SG+EV
9	G	<i>0.63</i>	<i>H. pylori</i> GYEVH <i>Borrelia burgdorferi</i> GYE V KL+ <i>C. neoformans</i> GYE LV and GY++ + LV <i>P. gingivalis</i> GYEV <i>C. pneumoniae</i> GYEV and GY HH
10	Y	<i>0.56</i>	<i>H. pylori</i> YE HH and YE+ HQ and Y++H Q and YE HHQ Cytomegalovirus YEVH <i>Borrelia Burgdorferi</i> YE+ KL <i>C. neoformans</i> YE + QK FC <i>P. gingivalis</i> Y++H H+K <i>C. pneumoniae</i> Y+V +Q LV
11	E	<i>0.58</i>	<i>H. pylori</i> EV +QK Cytomegalovirus EV HQ L <i>Borrelia Burgdorferi</i> EV +KL <i>C. neoformans</i> EV Q LV <i>P. gingivalis</i> EV KLV <i>C. pneumoniae</i> EV QKLV
12	V	<i>0.35</i>	<i>H. pylori</i> . +H QK Cytomegalovirus V HQ LV HHV-6 VH QK+V <i>Borrelia Burgdorferi</i> VH KL <i>C. neoformans</i> +HH LV <i>P. gingivalis</i> VH + LV <i>C. pneumoniae</i> V HQKL
13	H	-0.17	<i>H. pylori</i> and <i>C. pneumoniae</i> HHQK Cytomegalovirus HH KL <i>P. gingivalis</i> HH KL
14	H	-0.66	<i>Borrelia Burgdorferi</i> HQKL+ <i>C. pneumoniae</i> and HSV-1 HQKL <i>P. gingivalis</i> +QKLV
15	Q	-1.03	<i>C. neoformans</i> and <i>P. gingivalis</i> and <i>C. pneumoniae</i> QKLV
16	K	-1.47	<i>H. pylori</i> : <i>Cryptococcus neoformans</i> <i>Borrelia burgdorferi</i> <i>Chlamydomytila pneumoniae</i> KLVFF Human herpesvirus 1 KLVF
17	L	-1.34	Human herpesvirus 5: Human herpesvirus 6 LVFF
18	V	-1.20	
19	F	-0.93	
20	F	-0.98	
21	A	-0.82	
22	E	-0.31	
23	D	0.23	
24	V	<i>0.81</i>	<i>Borrelia burgdorferi</i> <i>Cryptococcus neoformans</i> <i>Porphyromonas gingivalis</i> VGSNK Cytomegalovirus +GSNK <i>Helicobacter pylori</i> <i>Chlamydomytila pneumoniae</i> VGSN
25	G	<i>1.24</i>	<i>H. pylori</i> <i>Chlamydomytila pneumoniae</i> GSNK
26	S	<i>1.22</i>	
27	N	<i>0.90</i>	
28	K	<i>0.36</i>	

(a) Continued.

Position B-Amy	Amino acid	B-cell index	Alignments
29	G	0.30	
30	A	−0.24	
31	I	−0.58	
32	I	−1.00	
33	G	−1.14	
34	L	−1.19	
35	M	−1.23	
36	V	−1.16	69 viruses/phages VGGVV
37	G	−0.97	
38	G	−1.02	
39	V	−0.63	
40	V	−0.45	
41	I	−0.80	
42	A	−1.06	

(b)

NGF position	Amino acid	B-cell index	Alignment
18	A	0.59	<i>C. neoformans</i> AEPHS
19	E	1.16	<i>P. gingivalis</i> EPHSES—NVP Cytomegalovirus EPHS+S
20	P	1.32	<i>C. neoformans</i> P+S NVPAG and PHSES and P SESNV
21	H	1.78	<i>Borrelia burgdorferi</i> HSESN <i>C. neoformans</i> HSES VP and HSESN P H +P+
22	S	1.64	<i>C. neoformans</i> S+S VPAG T P <i>Borrelia burgdorferi</i> burgdorferi <i>C. neoformans</i> Chlamydomphila pneumoniae <i>P. gingivalis</i> SESNV <i>P. gingivalis</i> S+SNVP <i>C. pneumoniae</i> SESNV A
23	E	1.68	<i>C. neoformans</i> ESNVP and ESNV AG
24	S	1.51	<i>C. neoformans</i> SNVPA
25	N	1.61	<i>C. neoformans</i> Cytomegalovirus <i>P. gingivalis</i> NVPAG <i>C. neoformans</i> NV TIPQA <i>P. gingivalis</i> +VPAG HT
26	V	1.45	<i>C. neoformans</i> <i>P. gingivalis</i> VPAGH <i>C. neoformans</i> VP AGHT <i>C. neoformans</i> VPAG TI and V AGHT+
27	P	1.36	<i>C. neoformans</i> HSV-1 PAGHT <i>C. neoformans</i> PAGHT P <i>C. neoformans</i> PAG TIP
28	A	1.04	<i>C. neoformans</i> <i>H. pylori</i> AGHTI <i>C. neoformans</i> AG H IPQA and AGHT PQ and AGHT+P and AG TIP HSV-1 HSV-2 AGH PQ and AGHT QA
29	G	1.00	<i>C. neoformans</i> GHTI Q and GHT PQ and G TIPQ
30	H	0.93	<i>C. pneumoniae</i> HTI QA <i>C. neoformans</i> HT PQA and HTIP A
31	T	1.05	<i>C. neoformans</i> <i>P. gingivalis</i> TIPQA
32	I	0.95	
33	P	0.64	
34	Q	0.41	
35	A	0.46	
51	A	0.38	<i>C. neoformans</i> AR SAPA and AR APAA and A SAPAA and ARSA AA and ARSAP A and ARS PAA and AR—SAPAA
52	R	0.54	HSV-1 RSAPAA <i>Borrelia burgdorferi</i> burgdorferi RSA AA
53	S	0.91	<i>C. neoformans</i> <i>C. pneumoniae</i> Cytomegalovirus HSV-1 HSV-2 <i>P. gingivalis</i> SAPAA
54	A	0.89	
55	P	0.72	
56	A	0.69	
57	A	0.53	
64	A	0.46	<i>C. neoformans</i> AG TRNI and AGQT RN and AGQTR <i>P. gingivalis</i> AGQTR
65	G	0.59	<i>C. neoformans</i> + RNITV and GQTRN

(b) Continued.

NGF position	Amino acid	B-cell index	Alignment
66	Q	0.42	<i>P. gingivalis</i> Borrelia burgdorferi QTRNI Cytomegalovirus QTRN—IT
67	T	0.43	<i>C. neoformans</i> <i>P. gingivalis</i> TRNIT
68	R	0.49	<i>C. neoformans</i> RNIT DP and RNITV
69	N	0.76	<i>C. neoformans</i> <i>C. pneumoniae</i> <i>H. pylori</i> NITVD
70	I	0.66	
71	T	0.56	
90	S	0.40	<i>C. neoformans</i> STQPPR and STQPP AA and STQPP EA <i>C. pneumoniae</i> STQ PRE <i>C. pneumoniae</i> Cytomegalovirus STQPP HSV-1 STQ PR
91	T	0.63	<i>C. neoformans</i> TQP REA and TQPPR
92	Q	1.06	<i>C. neoformans</i> QPPRE
93	P	1.49	<i>C. neoformans</i> PP REAA <i>C. Neoformans</i> <i>P. gingivalis</i> PPREA
94	P	1.91	<i>C. neoformans</i> HSV-2 <i>H. pylori</i> PPREA
95	R	2.13	<i>C. neoformans</i> <i>P. gingivalis</i> REAAD and REAA TQ and RE ADT <i>C. neoformans</i> REAA DT and R AADT and R+AADT and READD HSV-1 REAA T Borrelia burgdorferi burgdorferi Cytomagalovirus RE ADT Cytomagalovirus REAA TQ <i>P. gingivalis</i> REA +TQ
96	E	2.13	<i>P. gingivalis</i> EAADTQ and EA TQDLD <i>C. neoformans</i> EAADT+ and EAAD QD and EAAD Q and EAA TQ
97	A	1.92	Borrelia burgdorferi burgdorferi <i>C. neoformans</i> AA TQD Borrelia burgdorferi burgdorferi <i>P. gingivalis</i> AAD QD HSV-2 AADT D HHV-6 AAD +DL <i>C. neoformans</i> AADT D and AA TQD and A DTQD and A TQDL and AADTQ and AADT+ D+D <i>P. gingivalis</i> ADTQ L <i>H. pylori</i> AADTQ
98	A	1.9	Borrelia burgdorferi ADT DLD <i>C. neoformans</i> AD QDLD and AD QDL <i>P. gingivalis</i> ADT DL and AD QDL <i>H. pylori</i> +TQDLD
99	D	1.39	<i>C. neoformans</i> DT DLD and D QDLD <i>C. pneumoniae</i> DTQDL
100	T	1.25	<i>C. neoformans</i> <i>H. pylori</i> <i>P. gingivalis</i> TQDLD
101	Q	0.66	Borrelia burgdorferi burgdorferi <i>C. neoformans</i> <i>H. pylori</i> <i>P. gingivalis</i> QDLDF <i>H. pylori</i> QD DFEV
102	D	0.73	Borrelia burgdorferi burgdorferi <i>C. neoformans</i> <i>H. pylori</i> DLD FE <i>C. neoformans</i> DLD EVG and DLDF VG
103	L	0.54	<i>C. neoformans</i> LD EVGG and LD FE GG and LD FEV and LDF VGG <i>H. pylori</i> LDF EVGG HSV-1 HSV-2 L+ EVGG
104	D	0.45	Borrelia burgdorferi burgdorferi DFEVG <i>C. neoformans</i> DF VGGA and D EVGGA
105	F	0.42	<i>C. neoformans</i> FEVGG Cytomegalovirus FE GGAA
106	E	0.49	<i>C. neoformans</i> EVGGAA and EVGGA P and EVGGA and EV GAAP and E+GGAAP <i>H. pylori</i> EVGGA Borrelia burgdorferi burgdorferi E+GG A PF+
107	V	0.39	<i>P. gingivalis</i> VGGAAP and VGGAA <i>C. neoformans</i> VGGAA PF and VGGAA NR and VGGAA and VG GAAP <i>C. pneumoniae</i> VGGA AP <i>H. pylori</i> VG GAAP and VGG APF and VGGAA <i>P. gingivalis</i> VGGAA
108	G	0.74	<i>C. neoformans</i> <i>C. pneumoniae</i> HSV-2 GGAAP
109	G	0.44	<i>C. neoformans</i> GA APFN T and GAAPF
110	A	0.95	<i>C. pneumoniae</i> AAPFN Borrelia burgdorferi AAP+NR HSV-1 AAP RT
111	A	0.83	<i>C. neoformans</i> APFN T RS <i>C. neoformans</i> <i>H. pylori</i> APFN TH <i>C. pneumoniae</i> AP N RT R RSSS
112	P	0.99	<i>C. neoformans</i> PFNRT and PF RTH Borrelia burgdorferi burgdorferi PF NRT
113	F	0.84	<i>C. neoformans</i> FNRT SKR
114	N	0.75	<i>C. neoformans</i> NR RSKRS S and NRT R RSSS Cytomegalovirus N T RSKRS
115	R	0.63	<i>C. neoformans</i> RT R SKRS and RTHRS Borrelia burgdorferi burgdorferi RTHRS
116	T	0.63	<i>C. neoformans</i> THRS RS and THRSK
117	H	0.64	<i>C. neoformans</i> HRS RSS and HRSKR <i>C. Pneumonie</i> HRSKR

(b) Continued.

NGF position	Amino acid	B-cell index	Alignment
118	R	0.94	<i>C. neoformans</i> RSKRSS and RS RSSS and RS KRSS and RS KRSS and RSK SSS and RSKRS S <i>P. gingivalis</i> RSKR S and RSKRS <i>Borrelia burgdorferi burgdorferi</i> RSKRS
119	S	0.88	<i>C. neoformans</i> SKRSSS and SKRSS <i>C. pneumoniae</i> <i>H. pylori</i> HSV-1 HSV-2 SKRSS
120	K	1.12	<i>C. neoformans</i> Cytomegalovirus HHV-6 <i>H. pylori</i> KRSSS
121	R	1.13	
122	S	1.12	
123	S	0.83	
124	S	0.50	
144	G	0.52	<i>C. neoformans</i> GDKTTA and GDKTT
145	D	0.54	<i>H. pylori</i> DK TATDI <i>Borrelia burgdorferi burgdorferi</i> <i>C. neoformans</i> <i>C. pneumoniae</i> <i>H. pylori</i> DKT TA HSV-1 +KTT TD
146	K	0.84	CF. <i>Pneumoniae</i> KTTAT+ <i>C. neoformans</i> <i>H. pylori</i> KTTAT
147	T	1.27	<i>C. neoformans</i> HSV-1 TTATDI <i>C. neoformans</i> <i>P. gingivalis</i> TTATD
148	T	1.22	<i>C. neoformans</i> TATDIK <i>Borrelia burgdorferi burgdorferi</i> <i>C. neoformans</i> <i>C. pneumoniae</i> <i>H. pylori</i> TATDI
149	A	1.21	<i>C. neoformans</i> HHV-6 HHV-6B <i>P. gingivalis</i> ATDIK
150	T	1.08	
151	D	1.01	
152	I	0.97	
153	K	0.61	
179	C	0.87	<i>C. neoformans</i> CR PNPV <i>C. pneumoniae</i> CRDPN P+ S RGI
180	R	1.40	<i>C. pneumoniae</i> RDPNPV <i>Borrelia burgdorferi burgdorferi</i> RD NP VDS <i>C. neoformans</i> RDP PVDS and RDPNP+ and RDPNP HHV-6 HHV-6B RDPNP HSV-1 HSV-2 RDPN V
181	D	1.36	<i>Borrelia burgdorferi burgdorferi</i> D NPVD and DPN VD <i>C. neoformans</i> DPNPV and DP PVD and DP PVDS and DPN VDS <i>C. pneumoniae</i> DPN VD HSV-1 DPNP S HSV-2 +P PVDS
182	P	1.69	<i>C. neoformans</i> PN VDSG and PNP DSG and PNPVD <i>P. gingivalis</i> PNPV+S and P PVDS
183	N	1.72	<i>H. pylori</i> NPVD G
184	P	1.85	<i>C. neoformans</i> PV DSGCR and P PVDS RG and PVDSG Cytomagalovirus P DSG RGI <i>P. gingivalis</i> PV DSGC
185	V	1.74	HSV-2 VDSG RG HSV-1 +DSG RG <i>C. neoformans</i> VDSG R
186	D	1.51	<i>C. neoformans</i> DSGC GI and DSG RG
187	S	1.28	<i>Borrelia burgdorferi burgdorferi</i> <i>C. neoformans</i> SGCRG and SG RGI and S CRGI
188	G	0.79	<i>C. neoformans</i> GC IDSKH W and GCRG D <i>P. gingivalis</i> GCRGI and GCRG+
189	C	0.79	<i>P. gingivalis</i> CRGID <i>C. neoformans</i> C GIDS
190	R	0.85	<i>C. neoformans</i> R IDSK and RGIDS and RGID K <i>H. pylori</i> RGIDS HSV-1 HSV-2 RG+DS <i>H. pylori</i> RG DSK <i>Borrelia burgdorferi burgdorferi</i> RGID K
191	G	0.65	<i>C. neoformans</i> GIDS HW and GIDSK and GIDS H <i>Borrelia burgdorferi burgdorferi</i> <i>H. pylori</i> <i>P. gingivalis</i> GIDSK <i>P. gingivalis</i> GID KH and G DSKH <i>H. pylori</i> GI SKH
192	I	0.60	<i>Borrelia burgdorferi burgdorferi</i> IDSKH <i>C. neoformans</i> <i>P. gingivalis</i> IDSK W
193	D	0.3	<i>C. neoformans</i> DSKH W <i>Borrelia burgdorferi burgdorferi</i> DSK WN
194	S	0.40	<i>C. neoformans</i> SKHW+
195	K	0.44	
197	W	0.41	
212	T	0.33	<i>C. neoformans</i> TMDGKQ and TMDGK
213	M	0.31	<i>P. gingivalis</i> MDGKQ
214	D	0.39	<i>C. neoformans</i> DGKQAA and DGKQA <i>C. pneumoniae</i> DGK AA

(b) Continued.

NGF position	Amino acid	B-cell index	Alignment
215	G	0.48	<i>C. neoformans</i> GKQAA
216	K	0.53	
217	Q	0.45	

(c)

Tau position	Amino acid	B-cell index	Alignment
53	E	2.516	<i>C. neoformans</i> (GSK3)EDGSEEP S and E+G EEPG and +DGS+EP S and ED GSEE GS and EDGS++ GS and EDGSE and +D EEPG and EDG S PGS Cytomegalovirus EDG EEP and +DG EE and ED GSEE <i>P. gingivalis</i> EDGSEE and E+G SEEP and EDGS EE HHV-6 HHV-6B EDGS EE <i>C. pneumoniae</i> E GSEE <i>Borrelia burgdorferi</i> E GSEE HHV-6 E GSEE
54	D	2.568	<i>C. neoformans</i> DGSEEP and DGS+EPG and DGSEE G <i>P. gingivalis</i> DGS EPGS and DGSEE and DGS EP and DGSE EP and DGSE P Cytomegalovirus DGS EP and DG EE G <i>C. pneumoniae</i> DGSE GS and +G EE GS <i>Borrelia burgdorferi</i> DGSE+ <i>P. gingivalis</i> +GSEE
55	G	2.621	<i>C. neoformans</i> GSEEP and G EEPG and G +EPGS and GS EEPG and GSEE G Cytomegalovirus GS+EP S <i>P. gingivalis</i> GS EPG <i>H. pylori</i> GSEE G <i>C. pneumoniae</i> GSEE GS
56	S	2.732	<i>C. neoformans</i> SEEPG and SEE GS and S EPGS and SEEP S <i>C. pneumoniae</i> SEEPG and SEE GS and SE PGS <i>P. gingivalis</i> SEE GS and S EPGS
57	E	2.774	<i>C. neoformans</i> EEPGS and +EPGS and EEPG E and EE GSE <i>C. pneumoniae</i> EEPGS HSV-1 HSV-2 EEPG
58	E	2.678	HHV-6 HHV-6B EPGSE <i>C. neoformans</i> EPGSE and EPGS+ <i>P. gingivalis</i> EPGS+
59	P	2.564	
60	G	2.544	
61	S	2.556	
62	E	2.287	
171	S	2.315	<i>Borrelia burgdorferi</i> SG GPED <i>C. neoformans</i> SG GPEDT and SGT PE and SGTGP and SG GPE and SGTG E and SG GP+ <i>P. gingivalis</i> SGTG E <i>C. pneumoniae</i> SGTG PE Cytomegalovirus SGTGP+
172	G	2.54	<i>C. neoformans</i> GT PED and GTG ED and GTGPE and G GPED and GTGP D and G EDTE HSV-1 GTGPE and GTGP D and GTGP+D HSV-2 GTGP D and G GPED <i>C. pneumoniae</i> GTGPE <i>H. pylori</i> GTGP D
173	T	2.731	HHV-6 TGPEDT and TG PEDT <i>C. neoformans</i> TG PEDT and TG EDT and T PEDT and TGPED <i>C. pneumoniae</i> TGPED <i>H. pylori</i> TG EDT <i>Borrelia burgdorferi</i> TGPE T+
174	G	2.709	<i>C. neoformans</i> GPEDT and GPED TE and GP DTE and GPED E HHV-6 HHV-6B TG PEDT
175	P	2.807	<i>C. neoformans</i> <i>H. pylori</i> <i>P. gingivalis</i> PEDTE
176	E	2.7	
177	D	2.563	
178	T	2.397	
179	E	2.225	
228	S	2.396	HSV-1 SP DSPP and SPQ SP <i>C. neoformans</i> SPQDS and S QDSP and SP DSPP and SP DSPP and SPQD PP and PQ DSPPS and SPQ PPSK and SPQ SP and SP DSP and SPQ SP Cytomegalovirus SP DSPP <i>P. gingivalis</i> SP DSP <i>Borrelia burgdorferi</i> SP++SPP and SP D PSK
229	P	2.744	<i>C. neoformans</i> PQ SPPS and P DSPPS and PQDSP and P+DSPP and PQ S PPSK and PQ SPP and PQD PP and P DSPP <i>P. gingivalis</i> PQDS P HSV-2 P DSPP HHV-6 PQ+ PP and PQ PP K <i>Borrelia burgdorferi</i> PQ+ P SK
230	Q	2.763	<i>C. neoformans</i> QDSPP and ++ PPSK and QDS PS and Q SPPS <i>P. gingivalis</i> QD PPS <i>C. pneumoniae</i> QDS PS
231	D	2.929	<i>C. neoformans</i> DSP PSK and DSPPS and DSP SK and D PPSK and +SPPS <i>P. gingivalis</i> DSP SK
232	S	2.882	<i>C. neoformans</i> SPPSK

(c) Continued.

Tau position	Amino acid	B-cell index	Alignment
233	P	2.75	
234	P	2.794	
235	S	2.68	
242	D	2.387	<i>C. neoformans</i> DGRPP <i>C. pneumoniae</i> DG PPQ HSV-1 DG PPQ HSV-2 DGRPP and DG PP+ Cytomegalovirus DG R PQ and DG PP and +GRPP <i>Borrelia burgdorferi</i> DG PP
243	G	2.49	<i>C. neoformans</i> GRPPQ
244	R	2.56	
245	P	2.42	
246	P	2.164	
247	Q	2.004	
331	P	2.259	<i>C. neoformans</i> PGEG PE and P EGPEA and PGEGP and PGEG EA and P EGPEA and PGEG E Cytomegalovirus PGEGP EA and PG GPE <i>C. pneumoniae</i> PGE PEA and PGEGP <i>H. pylori</i> PG GPE HSV-1 HHV-6 PG GP A
332	G	2.681	<i>C. neoformans</i> GEGPE and GEG EA and GE PEA and G GPEA and GEGP A and GEG EA and G+GPEA <i>C. pneumoniae</i> GE PEA <i>P. gingivalis</i> GEGPE and GEG EA
333	E	2.715	<i>C. neoformans</i> <i>P. gingivalis</i> EGPEA HSV-2 +GPEA
334	G	2.634	
335	P	2.592	
336	E	2.463	
337	A	2.405	
414	H	1.819	HSV-1 HSV-2 HPTPG <i>C. neoformans</i> HPT GSS and HP PGSS and HPTP SS and HP P SS <i>C. pneumoniae</i> HP PGS and +PT SS
415	P	2.274	Cytomegalovirus P PGSS and PTPG SS <i>C. pneumoniae</i> PT GSS and PTPG SS
416	T	2.503	
417	P	2.717	
418	G	2.469	
419	S	2.054	
420	S	1.76	
493	P	2.43	HSV-1 P APKTP and P APKTP and PPAP PP and PPA PP HSV-2 PPAP TPP Cytomegalovirus P APKT and PPAP PP and PP P PPS and PP KTPP and PP A TPPS and PPA TPP <i>C. neoformans</i> PPAP TPPS and PPAP KTP S and PP PKTP and PP PKTP and PPAPKT and PPAP KT <i>P. gingivalis</i> PPAPK P <i>C. pneumoniae</i> PPAPK and PP PKT and PP AP PP and PP TPP <i>Borrelia burgdorferi</i> PAP T PS and PPA KTP and PP P TPP
494	P	2.77	HSV-1 PAP KTP and PA KTPP HSV-2 PAPK PP HHV-6 PAP PPS <i>C. neoformans</i> PAPKTP S and PAPK PP <i>P. gingivalis</i> PAPKT <i>C. pneumoniae</i> PAP TP and PAPK P
495	A	2.94	<i>C. neoformans</i> AP PPS <i>P. gingivalis</i> A KTPPS HHV-6 APKTP
496	P	2.89	<i>C. neoformans</i> PKTPPS and PKTP PS and PK TPPS <i>P. gingivalis</i> PKT PS
497	K	2.95	<i>C. neoformans</i> KTPPS
498	T	2.98	
499	P	2.69	
500	P	2.51	
501	S	2.343	
502	S	2.123	

a classical example from the field of population genetics and Darwinism, the light coloured genes of the peppered moth favour its selective predation by *many different* birds when it alights on dark trees covered with soot pollution, while darker melanised forms are selectively targeted on lighter coloured trees [273]. The coloration susceptibility genes,

or the variety of tree (risk factors), do not kill the moth but allow *several* causes to do so. The causes can hide in plain sight, as epidemiological studies, as applied to human diseases, could conclude that the birds are not killing the moths, as they are always present, on both sets of trees, in both genetic conditions, whether the moths are alive or

dead (c.f. the ubiquitous *C. neoformans* and many other common pathogens). Many of the pathogens implicated in Alzheimer's disease (herpes simplex, *Borrelia burgdorferi* and *C. pneumoniae*), and several other risk factors (cholesterol, homocysteine, diabetes, or vitamin A or nerve growth factor deficiency) are able to promote cerebral beta-amyloid deposition in animal models, without the aid of any gene variant. Subsets of susceptibility genes can be related to each of these amyloidogenic pathways [1]. In the case of genomewide association studies, the genes returned, as well as **APP**, beta amyloid, and **gamma secretase**, seem intimately concerned with pathogen life cycles and defence and the immune network. This suggests that the diverse microbial risk factors as well as other dietary and environmental factors implicated in Alzheimer's disease are in fact causative agents, whose deleterious effects are conditioned by susceptibility genes. As the environmental risk factors are amenable to therapy, while the susceptibility gene products have so far proved largely resistant, this suggests numerous ways with which to tackle the problem of Alzheimer's disease.

Herpes Simplex Reactivation. Alzheimer's disease plaques and tangles are highly enriched in human proteins used the herpes simplex during its life cycle, as well as in many immune-related proteins, suggesting that immune attack on a reactivated virus, diverted to neurones, which contain the complement membrane attack complex, may be ultimately responsible for the extensive neuronal destruction seen in Alzheimer's disease [15]. The herpes simplex virus establishes latency in neurones, existing as a dormant form where only the latency transcript is expressed. A number of factors again related to susceptibility genes and environmental risk factors in Alzheimer's disease can be related to herpes simplex latency and reactivation. The cerebral herpes simplex viral load is decreased in **APOE** knockout mice, while the viral load is much increased in **APOE4** transgenic mice, compared to **APOE3** mice, suggesting that **APOE4** favours the establishment of a greater number of latent viruses in cerebral tissue [274]. In neuroblastoma cells, this latent form may even exert beneficial effects, blocking apoptosis and promoting neurite extension via **AKT1** upregulation [275]. However, this latent form can be reactivated by NGF deprivation [276], and NGF promotes viral latency via the TrkA receptor **NTRK1** [277], the expression of which is reduced in the Alzheimer's disease brain [278]. (Relevant pathways and genes: neurotrophin signalling (**GSK3B**, **NTRK1**, **NTRK2**, **PIK3R1**, and **SOS2**). Vitamin A supplementation in rats increases the cerebral levels of both NGF and **BDNF** [279], while oestrogen deficiency lowers cerebral NGF levels, an effect reversed by 17-beta oestradiol [280], which is, however, also able to reactivate the virus, via oestrogen receptor alpha (**ESR1**) [281]. High levels of total oestradiol have been reported as a risk factor for Alzheimer's disease in both women and men [282, 283]. Vitamin A related genes include **APOE4**, the isoform least able to transport the vitamin A precursor retinyl palmitate, **A2M** **ABCA1** **ALB** **ALDH2** **APOA1** **CHD4** **CLU** **CYP46A1** **ESR1** **GSTM1** **GSTP1** **HSPG2** **KLF5** **LIPA**

LPL **LRAT** **LRP2** **LRPAP1** **MEF2A** **NPAS2** **NR1H2** **PARP1** **PIN1** **POU2F1** **PPARA** **PPARG** **RXRA** **THRA** **TTR** **UBQLN1** **VDR** and many others controlled by retinoid receptor element (reviewed in [126]).

The virus can also be reactivated by heat [284], **IL6** [285], or **TNF** [286]. **IL6** plasma and CSF levels have been reported to be increased in Alzheimer's disease and the secretion of **IL6** from monocytes is increased [287–289]. **IL6** plasma levels are raised by infection with *C. pneumoniae* [290] or *Helicobacter pylori* [291], and **IL6** production in monocytes is stimulated by *C. neoformans* [292]. Stress-activated corticosterone is also able to reactivate the virus [293] (pathway = steroid hormone biosynthesis: **COMT**, **CYP19A1**, **HSD11B1**). Cortisol levels are increased in the ageing population and in Alzheimer's disease [294, 295]. A number of related stressors including adrenaline [296], or downstream effectors such as cyclic AMP, protein kinase A, or C activation [297] are also able to reactivate the virus. Herpes simplex reactivation is blocked by beta-receptor antagonism (**ADRB1**) [296, 298]. Glucocorticoids and prostaglandins are also able to reactivate the virus [299]: cyclooxygenase inhibitors (**PTGS2**), celecoxib and indomethacin block viral reactivation and nonsteroidal anti-inflammatory use has been associated with a lower incidence of Alzheimer's disease. Lysophosphatidic acid is also able to reactivate the virus [300] (**LPAR5**). Hypoxia also increases the replication of herpes simplex [301]: transient cerebral ischaemia also lowers NGF levels [302], and these effects are relevant to the cerebral hypoperfusion seen in ageing and in Alzheimer's disease [303]. Vitamin A and the retinoic acid isomers all-trans-, 9-cis-, and 13-cis-Retinoic Acid are all able to reduce viral replication [304]. Low Vitamin A levels are a problem in the ageing population and even in successfully ageing persons can be observed in 50% of the population over the age of 80–85 [305]. Low vitamin A levels are also a risk factor for Alzheimer's disease [306]. Mice deficient in the nitric oxide synthase **NOS2** are also more susceptible to herpes virus infection, and reactivation occurs is stimulated by caspase 3 activation (**CASP3**) [135].

Thus, fever or cytokine release induced by diverse infections might be expected to reactivate herpes simplex as would several of the conditions associated with Alzheimer's disease (cerebral hypoperfusion, low cerebral NGF levels, vitamin A deficiency, or high oestradiol levels).

3.6.6. The Microbiome in Alzheimer's Disease. These pathogens form a very small proportion of an extensive human microbiome comprising of trillions of bacteria, viruses, and other pathogens, whose influence on diseases is increasingly recognised [307]. Individual species may exert benevolent or malevolent effects which may well be dictated by the very extensive sequence homology between human and pathogen proteins which enables pathogens to intercalate with numerous important signalling networks, via competition with their human counterparts [239, 241, 248, 250, 254]. These networks are implicated in diseases, and their efficiency, as well as pathogen/host homology, is dictated by susceptibility genes. This host/pathogen matching covers

the entire human proteome and is represented by millions of short consensi of pentapeptides or more, not counting the longer gapped consensi. Clearly, powerful algorithms are needed to trawl human and pathogen proteomes and to link these homologies to susceptibility genes and to epitopes and autoantigens.

The principles discussed here may apply to many other, if not most, human diseases.

Protective Agents in Alzheimer's Disease. Statins [308], fish consumption [309], omega-3 polyunsaturated fatty acids [310], the Mediterranean diet [311], nonsteroidal anti-inflammatories [312], or the generally healthy life-style of nuns living in convents [313] have all been associated with a reduced incidence or severity of Alzheimer's disease, while homocysteine lowering B vitamins and folate have been shown to slow the rate of brain atrophy in cognitively impaired elderly patients [314]. A rather pessimistic study [315] recently stated that no firm conclusions could be drawn on the association of any modifiable factors with risk of Alzheimer's disease. While this may be a justifiable statistical conclusion from meta-analysis, this is not to reckon with the diversity of the underlying genomic platform of each individual or with the profusion of diverse interacting risk and protective factors (epistasis, gene/environment, and environment/environment interactions).

As with the risk factors, protective agents can be linked to genetic and pathological pathways involved in cholesterol and lipid function (statins, fish, and diet, e.g., ABCA1, ABCA2, ABCA7, ABCG1, ACADS, ALDH2, APP, APOA1, APOA4, APOA5, APOC1-4, APOE, CH25H, CYP46A1, DHCR24, FDPS, HMGCR, HMGCS2, HSD11B1, LIPC, LRP1, LRP2, LRP8, LDLR, LIPA, LPL, OLR1, PPARA, PPARG, PTPLA, VLDLR, NPC1, NPC2, SOAT1, SREBF1), homocysteine, methionine, and folate metabolism (folate and vitamins, e.g., BLMH, CBS, COMT, NAT2, MTHFD1L, MTHFR, MTR, MTRR, PON1, PON2, PON3, VDR) and inflammatory pathways (NSAID'S e.g., C4A, C4B, CD2AP, CD33, CFH, CCL2, CCL3, CCR2, CSF1, CLU, CR1, FAS, GSK3B, IL1A, IL1B, IL6, IL8, IL10, IL18, PLA2, SERPINA1, SERPINE1, SERPINF2, PTGS2, TGFBI, TNF). As with the risk factors, the success of such protective agents is likely to be determined by genes and other confounding factors. In this genomic era, affordable whole genome sequencing will soon be achieved, ushering in an age of more effective treatments tailored to individual genetic profiles. Alzheimer's disease is clearly multifactorial with many related genetic and environmental risk factors, several underlying pathologies, and several available protective strategies. A recent study has also shown that many other seemingly benign factors (eyesight, hearing, denture wearing, stomach, kidney, bladder or bowel problems, coughs, and colds) as well as high blood pressure and diabetes, constituting a frailty index, combine to markedly increase both the incidence and severity of Alzheimer's disease [316]. A further study, identifying physical inactivity, depression, smoking, mid-life hypertension or obesity, low education, and diabetes as key risk factors, has estimated that ~50% of Alzheimer's disease cases may be preventable [317].

Elimination of the risk factors, including the regular detection and elimination of pathogens in the elderly, adherence to sensible dietary and vitamin supplement recommendations, and the genetically tailored use of certain drug regimens are together likely to be able to markedly reduce the incidence of Alzheimer's disease. Using phage display, it is now possible to express peptide fragments of the entire human proteome in a phage library, and to use this to trap autoantibodies in blood or other bodily fluid samples. The antigen expressed by the labelled phage can be identified by high throughput sequencing [318]. Such technology is likely to be extremely useful in characterising biomarkers and pathological immune processes as well as potential pathogen/human cross-reactivity.

Proteome-wide characterisation of the autoantibodies relevant to Alzheimer's disease (a move from GWAS to PWAS) would also be very informative as selective autoantibody removal via affinity dialysis might well be expected to influence the severity and progression of Alzheimer's disease.

Since the submission of this paper, Miklosy has reported a highly significant association between spirochete infection and Alzheimer's disease. As well as *B. burgdorferi*, several periodontal pathogen treponemas species were detected in brain samples and the pathological features of Alzheimer's disease were reproduced by infection *in vitro* [319]. In addition, two groups have reported a very extensive repertoire of autoantigens in Alzheimer's disease, which can be characterised with a high degree of accuracy by a definitive immunosignature [320, 321]. The principles outlined above could thus be tested by analysis of pathogen/autoantigen cross-reactivity.

References

- [1] C. J. Carter, "The fox and the rabbits, environmental variables and population genetics. 1: replication problems in association studies and the untapped power of GWAS. 2: vitamin A deficiency, herpes simplex reactivation and other causes of Alzheimer's disease," *ISRN Neurology*, vol. 2011, Article ID 394678, 29 pages, 2011.
- [2] C. J. Carter, "Interactions between the products of the Herpes simplex genome and Alzheimer's disease susceptibility genes: relevance to pathological-signalling cascades," *Neurochemistry International*, vol. 52, no. 6, pp. 920–934, 2008.
- [3] C. J. Carter, "Convergence of genes implicated in Alzheimer's disease on the cerebral cholesterol shuttle: APP, cholesterol, lipoproteins, and atherosclerosis," *Neurochemistry International*, vol. 50, no. 1, pp. 12–38, 2007.
- [4] A. Papassotiropoulos, M. A. Wollmer, M. Tsolaki et al., "A cluster of cholesterol-related genes confers susceptibility for Alzheimer's disease," *Journal of Clinical Psychiatry*, vol. 66, no. 7, pp. 940–947, 2005.
- [5] R. Menon and C. Farina, "Shared molecular and functional frameworks among five complex human disorders: a comparative study on interactomes linked to susceptibility genes," *PLoS ONE*, vol. 6, no. 4, Article ID e18660, 2011.
- [6] A. C. Naj, G. W. Beecham, E. R. Martin et al., "Dementia revealed: novel chromosome 6 locus for Late-onset alzheimer disease provides genetic evidence for Folate-pathway abnormalities," *PLoS Genetics*, vol. 6, no. 9, Article ID e1001130, 2010.

- [7] D. Liolitsa, J. Powell, and S. Lovestone, "Genetic variability in the insulin signalling pathway may contribute to the risk of late onset Alzheimer's disease," *Journal of Neurology Neurosurgery and Psychiatry*, vol. 73, no. 3, pp. 261–266, 2002.
- [8] A. B. Goodman and A. B. Pardee, "Evidence for defective retinoid transport and function in late onset Alzheimer's disease," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 5, pp. 2901–2905, 2003.
- [9] V. Tseveleki, R. Rubio, S. S. Vamvakas et al., "Comparative gene expression analysis in mouse models for multiple sclerosis, Alzheimer's disease and stroke for identifying commonly regulated and disease-specific gene changes," *Genomics*, vol. 96, no. 2, pp. 82–91, 2010.
- [10] J. C. Lambert, B. Grenier-Boley, V. Chouraki et al., "Implication of the immune system in Alzheimer's disease: evidence from genome-wide pathway analysis," *Journal of Alzheimer's Disease*, vol. 20, no. 4, pp. 1107–1118, 2010.
- [11] J. M. Hill, Y. Zhao, C. Clement, D. M. Neumann, and W. J. Lukiw, "HSV-1 infection of human brain cells induces miRNA-146a and Alzheimer-type inflammatory signaling," *NeuroReport*, vol. 20, no. 16, pp. 1500–1505, 2009.
- [12] R. F. Itzhaki and M. A. Wozniak, "Herpes simplex virus type 1 in Alzheimer's disease: the enemy within," *Journal of Alzheimer's Disease*, vol. 13, no. 4, pp. 393–405, 2008.
- [13] R. B. Pyles, "The association of herpes simplex virus and Alzheimer's disease: a potential synthesis of genetic and environmental factors," *Herpes*, vol. 8, no. 3, pp. 64–68, 2001.
- [14] M. A. Wozniak, A. P. Mee, and R. F. Itzhaki, "Herpes simplex virus type 1 DNA is located within Alzheimer's disease amyloid plaques," *Journal of Pathology*, vol. 217, no. 1, pp. 131–138, 2009.
- [15] C. J. Carter, "Alzheimer's disease plaques and tangles: cemeteries of a Pyrrhic victory of the immune defence network against herpes simplex infection at the expense of complement and inflammation-mediated neuronal destruction," *Neurochemistry International*, vol. 58, no. 3, pp. 301–320, 2011.
- [16] L. Letenneur, K. Pérès, H. Fleury et al., "Seropositivity to Herpes Simplex Virus antibodies and risk of Alzheimer's disease: a population-based cohort study," *PLoS ONE*, vol. 3, no. 11, Article ID e3637, 2008.
- [17] A. R. Kamer, R. G. Craig, A. P. Dasanayake, M. Brys, L. Glodzik-Sobanska, and M. J. de Leon, "Inflammation and Alzheimer's disease: possible role of periodontal diseases," *Alzheimer's and Dementia*, vol. 4, no. 4, pp. 242–250, 2008.
- [18] M. M. Esiri, S. C. Biddolph, and C. S. Morris, "Prevalence of Alzheimer plaques in AIDS," *Journal of Neurology Neurosurgery and Psychiatry*, vol. 65, no. 1, pp. 29–33, 1998.
- [19] J. Kountouras, M. Boziki, E. Gavalas et al., "Five-year survival after *Helicobacter pylori* eradication in Alzheimer disease patients," *Cognitive and Behavioral Neurology*, vol. 23, no. 3, pp. 199–204, 2010.
- [20] T. A. Ala, R. C. Doss, and C. J. Sullivan, "Reversible dementia: a case of cryptococcal meningitis masquerading as Alzheimer's disease," *Journal of Alzheimer's Disease*, vol. 6, no. 5, pp. 503–508, 2004.
- [21] M. Hoffmann, J. Muniz, E. Carroll, and J. de Villasante, "Cryptococcal meningitis misdiagnosed as Alzheimer's disease: complete neurological and cognitive recovery with treatment," *Journal of Alzheimer's Disease*, vol. 16, no. 3, pp. 517–520, 2009.
- [22] I. Kaklikkaya, N. Kaklikkaya, I. Birincioglu, K. Buruk, and N. Turan, "Detection of human herpesvirus 6 DNA but not human herpesvirus 7 or 8 DNA in atherosclerotic and nonatherosclerotic vascular tissues," *Heart Surgery Forum*, vol. 13, no. 5, pp. E345–E349, 2010.
- [23] G. C. Makris, M. C. Makris, V. V. Wilmot, G. Geroulakos, and M. E. Falagas, "The role of infection in carotid plaque pathogenesis and stability: the clinical evidence," *Current Vascular Pharmacology*, vol. 8, no. 6, pp. 861–872, 2010.
- [24] A. Nazmi, A. V. Diez-Roux, N. S. Jenny, M. Y. Tsai, M. Szklo, and A. E. Aiello, "The influence of persistent pathogens on circulating levels of inflammatory markers: a cross-sectional analysis from the Multi-Ethnic Study of Atherosclerosis," *BMC Public Health*, vol. 10, article 706, 2010.
- [25] A. Taniguchi, F. Nishimura, Y. Murayama et al., "Porphyromonas gingivalis infection is associated with carotid atherosclerosis in non-obese Japanese type 2 diabetic patients," *Metabolism*, vol. 52, no. 2, pp. 142–145, 2003.
- [26] C. J. Hall, L. Bouhafs, U. Dizcalusy, and K. Sandstedt, "Cryptococcus neoformans causes lipid peroxidation; therefore it is a potential inducer of atherogenesis," *Mycologia*, vol. 102, no. 3, pp. 546–551, 2010.
- [27] A. E. Roher, C. Esh, A. Rahman, T. A. Kokjohn, and T. G. Beach, "Atherosclerosis of cerebral arteries in Alzheimer disease," *Stroke*, vol. 35, no. 11, pp. 2623–2627, 2004.
- [28] M. van Oijen, F. J. de Jong, J. C. Witteman, A. Hofman, P. J. Koudstaal, and M. M. B. Breteler, "Atherosclerosis and risk for dementia," *Annals of Neurology*, vol. 61, no. 5, pp. 403–410, 2007.
- [29] E. H. Corder, A. M. Saunders, W. J. Strittmatter et al., "Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families," *Science*, vol. 261, no. 5123, pp. 921–923, 1993.
- [30] C. Antúnez, M. Boada, A. González-Pérez et al., "The membrane-spanning 4-domains, subfamily A (MS4A) gene cluster contains a common variant associated with Alzheimer's disease," *Genome Medicine*, vol. 3, no. 5, article 33, 2011.
- [31] K. Morgan, "The three new pathways leading to Alzheimer's disease," *Neuropathology and Applied Neurobiology*, vol. 37, no. 4, pp. 353–357, 2011.
- [32] L. A. Hindorf, P. Sethupathy, H. A. Junkins et al., "A Catalog of Published Genome-Wide Association Studies," <http://www.genome.gov/gwastudies/>. Accessed [June 2011].
- [33] A. I. Su, T. Wiltshire, S. Batalov et al., "A gene atlas of the mouse and human protein-encoding transcriptomes," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 101, no. 16, pp. 6062–6067, 2004.
- [34] C. Wu, C. Orozco, J. Boyer et al., "BioGPS: an extensible and customizable portal for querying and organizing gene annotation resources," *Genome Biology*, vol. 10, no. 11, article R130, 2009.
- [35] J. E. Larsen, O. Lund, and M. Nielsen, "Improved method for predicting linear B-cell epitopes," *Immunome Research*, vol. 2, no. 2, 2006.
- [36] S. F. Altschul, T. L. Madden, A. A. Schäffer et al., "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs," *Nucleic Acids Research*, vol. 25, no. 17, pp. 3389–3402, 1997.
- [37] Z. Beck, B. K. Brown, G. R. Matyas, V. R. Polonis, M. Rao, and C. R. Alving, "Infection of human peripheral blood mononuclear cells by erythrocyte-bound HIV-1: effects of antibodies and complement," *Virology*, vol. 412, no. 2, pp. 441–447, 2011.

- [38] R. C. Carlisle, Y. Di, A. M. Cerny et al., "Human erythrocytes bind and inactivate type 5 adenovirus by presenting Cox-sackie virus-adenovirus receptor and complement receptor 1," *Blood*, vol. 113, no. 9, pp. 1909–1918, 2009.
- [39] E. Gyimesi, A. J. Bankovich, T. A. Schuman, J. B. Goldberg, M. A. Lindorfer, and R. P. Taylor, "Staphylococcus aureus bound to complement receptor 1 on human erythrocytes by bispecific monoclonal antibodies is phagocytosed by acceptor macrophages," *Immunology Letters*, vol. 95, no. 2, pp. 185–192, 2004.
- [40] Y. Hatano, S. Taniuchi, M. Masuda et al., "Phagocytosis of heat-killed Staphylococcus aureus by eosinophils: comparison with neutrophils," *APMIS*, vol. 117, no. 2, pp. 115–123, 2009.
- [41] J. Li, J. P. Wang, I. Ghiran et al., "Complement receptor 1 expression on mouse erythrocytes mediates clearance of Streptococcus pneumoniae by immune adherence," *Infection and Immunity*, vol. 78, no. 7, pp. 3129–3135, 2010.
- [42] C. Spadafora, G. A. Awandare, K. M. Kopydlowski et al., "Complement receptor 1 is a sialic acid-independent erythrocyte receptor of Plasmodium falciparum," *PLoS Pathogens*, vol. 6, no. 6, Article ID e1000968, 2010.
- [43] J. H. Powers, B. L. Buster, C. J. Reist et al., "Complement-independent binding of microorganisms to primate erythrocytes in vitro by cross-linked monoclonal antibodies via complement receptor 1," *Infection and Immunity*, vol. 63, no. 4, pp. 1329–1335, 1995.
- [44] A. Repik, S. E. Pincus, I. Ghiran et al., "A transgenic mouse model for studying the clearance of blood-borne pathogens via human complement receptor 1 (CR1)," *Clinical and Experimental Immunology*, vol. 140, no. 2, pp. 230–240, 2005.
- [45] M. Calero, T. Tokuda, A. Rostagno et al., "Functional and structural properties of lipid-associated apolipoprotein J (clusterin)," *Biochemical Journal*, vol. 344, no. 2, pp. 375–383, 1999.
- [46] S. Itagaki, H. Akiyama, H. Saito, and P. L. McGeer, "Ultrastructural localization of complement membrane attack complex (MAC)-like immunoreactivity in brains of patients with Alzheimer's disease," *Brain Research*, vol. 645, no. 1-2, pp. 78–84, 1994.
- [47] P. L. McGeer, H. Akiyama, S. Itagaki, and E. G. McGeer, "Activation of the classical complement pathway in brain tissue of Alzheimer patients," *Neuroscience Letters*, vol. 107, no. 1–3, pp. 341–346, 1989.
- [48] C. J. Carter, "APP, APOE, complement receptor 1, clusterin and PICALM and their involvement in the herpes simplex life cycle," *Neuroscience Letters*, vol. 483, no. 2, pp. 96–100, 2010.
- [49] C. Cortes, V. P. Ferreira, and M. K. Pangburn, "Native properdin binds to Chlamydia pneumoniae and promotes complement activation," *Infection and Immunity*, vol. 79, no. 2, pp. 724–731, 2011.
- [50] A. Hasegawa, L. F. Sogo, M. Tan, and C. Sütterlin, "Host complement regulatory protein CD59 is transported to the chlamydial inclusion by a golgi apparatus-independent pathway," *Infection and Immunity*, vol. 77, no. 4, pp. 1285–1292, 2009.
- [51] Á. L. Rosas, R. S. MacGill, J. D. Nosanchuk, T. R. Kozel, and A. Casadevall, "Activation of the alternative complement pathway by fungal melanins," *Clinical and Diagnostic Laboratory Immunology*, vol. 9, no. 1, pp. 144–148, 2002.
- [52] M. A. Gates-Hollingsworth and T. R. Kozel, "Phenotypic heterogeneity in expression of epitopes in the Cryptococcus neoformans capsule," *Molecular Microbiology*, vol. 74, no. 1, pp. 126–138, 2009.
- [53] G. Gonzalez-Valencia, G. I. Perez-Perez, R. G. Washburn, and M. J. Blaser, "Susceptibility of Helicobacter pylori to the bactericidal activity of human serum," *Helicobacter*, vol. 1, no. 1, pp. 28–33, 1996.
- [54] G. Hajishengallis, M. Wang, S. Liang et al., "Subversion of innate immunity by periodontopathic bacteria via exploitation of complement receptor-3," *Advances in Experimental Medicine and Biology*, vol. 632, pp. 203–219, 2008.
- [55] D. C. Altieri, O. R. Etingin, D. S. Fair et al., "Structurally homologous ligand binding of integrin Mac-1 and viral glycoprotein C receptors," *Science*, vol. 254, no. 5035, pp. 1200–1202, 1991.
- [56] G. J. Jones, J. C. Wiseman, K. J. Marr, S. Wei, J. Y. Djeu, and C. H. Mody, "In contrast to anti-tumor activity, YT cell and primary NK cell cytotoxicity for Cryptococcus neoformans bypasses LFA-1," *International Immunology*, vol. 21, no. 4, pp. 423–432, 2009.
- [57] A. Wang, S. C. Johnston, J. Chou, and D. Dean, "A systemic network for Chlamydia pneumoniae entry into human cells," *Journal of Bacteriology*, vol. 192, no. 11, pp. 2809–2815, 2010.
- [58] Z. Y. Huang, S. Hunter, P. Chien et al., "Interaction of two phagocytic host defense systems: Fcγ receptors and complement receptor 3," *Journal of Biological Chemistry*, vol. 286, no. 1, pp. 160–168, 2011.
- [59] K. Popadiak, J. Potempa, K. Riesbeck, and A. M. Blom, "Biphasic effect of gingipains from Porphyromonas gingivalis on the human complement system," *Journal of Immunology*, vol. 178, no. 11, pp. 7242–7250, 2007.
- [60] M. Deckert, M. Ticchioni, B. Mari, D. Mary, and A. Bernard, "The glycosylphosphatidylinositol-anchored CD59 protein stimulates both T cell receptor ζ/ZAP-70-dependent and -independent signaling pathways in T cells," *European Journal of Immunology*, vol. 25, no. 7, pp. 1815–1822, 1995.
- [61] S. Naderi, P. Hofmann, S. Seiter, W. Tilgen, H. Abken, and U. Reinhold, "CD2-mediated CD59 stimulation in keratinocytes results in secretion of IL-1α, IL-6, and GM-CSF: implications for the interaction of keratinocytes with intraepidermal T lymphocytes," *International Journal of Molecular Medicine*, vol. 3, no. 6, pp. 609–614, 1999.
- [62] A. W. Jehle, S. J. Gardai, S. Li et al., "ATP-binding cassette transporter A7 enhances phagocytosis of apoptotic cells and associated ERK signaling in macrophages," *Journal of Cell Biology*, vol. 174, no. 4, pp. 547–556, 2006.
- [63] S. W. Tas, L. B. Klickstein, S. F. Barbashov, and A. Nicholson-Weller, "C1q and C4b bind simultaneously to CR1 and additively support erythrocyte adhesion," *Journal of Immunology*, vol. 163, no. 9, pp. 5056–5063, 1999.
- [64] M. Sarvari, I. Vago, C. S. Weber et al., "Inhibition of C1q-beta-amyloid binding protects hippocampal cells against complement mediated toxicity," *Journal of Neuroimmunology*, vol. 137, no. 1-2, pp. 12–18, 2003.
- [65] B. Alberts, A. Johnson, J. Lewis, M. Raff, K. Roberts, and P. Walker, *Molecular Biology of the Cell*, Garland Science, New York, NY, USA, 2007.
- [66] S. J. Flint, L. W. Enquist, V. R. Racaniello, and A. M. Skalka, *Principles of Virology*, ASM Press, Herndon, Va, USA, 2008.
- [67] T. Gianni, G. Campadelli-Fiume, and L. Menotti, "Entry of herpes simplex virus mediated by chimeric forms of nectin1 retargeted to endosomes or to lipid rafts occurs through acidic endosomes," *Journal of Virology*, vol. 78, no. 22, pp. 12268–12276, 2004.

- [68] H. Parker, K. Chitcholtan, M. B. Hampton, and J. I. Keenan, "Uptake of *Helicobacter pylori* outer membrane vesicles by gastric epithelial cells," *Infection and Immunity*, vol. 78, no. 12, pp. 5054–5061, 2010.
- [69] J. Pizarro-Cerdá, M. Bonazzi, and P. Cossart, "Clathrin-mediated endocytosis: what works for small, also works for big," *BioEssays*, vol. 32, no. 6, pp. 496–504, 2010.
- [70] F. Tebar, S. K. Bohlander, and A. Sorkin, "Clathrin assembly lymphoid myeloid leukemia (CALM) protein: localization in endocytic-coated pits, interactions with clathrin, and the impact of overexpression on clathrin-mediated traffic," *Molecular Biology of the Cell*, vol. 10, no. 8, pp. 2687–2702, 1999.
- [71] C. R. Brunetti, R. L. Burke, B. Hoflack, T. Ludwig, K. S. Dingwell, and D. C. Johnson, "Role of mannose-6-phosphate receptors in herpes simplex virus entry into cells and cell-to-cell transmission," *Journal of Virology*, vol. 69, no. 6, pp. 3517–3528, 1995.
- [72] M. Molinari, C. Galli, N. Norais et al., "Vacuoles induced by *Helicobacter pylori* toxin contain both late endosomal and lysosomal markers," *Journal of Biological Chemistry*, vol. 272, no. 40, pp. 25339–25344, 1997.
- [73] K. Nakayama and S. Wakatsuki, "The structure and function of GGAs, the traffic controllers at the TGN sorting cross-roads," *Cell Structure and Function*, vol. 28, no. 5, pp. 431–442, 2003.
- [74] E. M. Rabin, K. Gordon, M. H. Knoppers et al., "Inhibition of T cell activation and adhesion functions by soluble CD2 protein," *Cellular Immunology*, vol. 149, no. 1, pp. 24–38, 1993.
- [75] N. C. Gauthier, P. Monzo, T. Gonzalez et al., "Early endosomes associated with dynamic F-actin structures are required for late trafficking of *H. pylori* VacA toxin," *Journal of Cell Biology*, vol. 177, no. 2, pp. 343–354, 2007.
- [76] P. L. W. Yun, A. A. Decarlo, C. C. Chapple, C. A. Collyer, and N. Hunter, "Binding of *Porphyromonas gingivalis* gingipains to human CD4⁺ T cells preferentially down-regulates surface CD2 and CD4 with little effect on co-stimulatory molecule expression," *Microbial Pathogenesis*, vol. 38, no. 2-3, pp. 85–96, 2005.
- [77] E. Moreno-Ruiz, M. Galán-Díez, W. Zhu et al., "Candida albicans internalization by host cells is mediated by a clathrin-dependent mechanism," *Cellular Microbiology*, vol. 11, no. 8, pp. 1179–1189, 2009.
- [78] M. Selbach and S. Backert, "Cortactin: an Achilles' heel of the actin cytoskeleton targeted by pathogens," *Trends in Microbiology*, vol. 13, no. 4, pp. 181–189, 2005.
- [79] N. Tegtmeyer, R. Wittelsberger, R. Hartig, S. Wessler, N. Martinez-Quiles, and S. Backert, "Serine phosphorylation of cortactin controls focal adhesion kinase activity and cell scattering induced by *Helicobacter pylori*," *Cell Host and Microbe*, vol. 9, no. 6, pp. 520–531, 2011.
- [80] S. Cudmore, I. Reckmann, and M. Way, "Viral manipulations of the actin cytoskeleton," *Trends in Microbiology*, vol. 5, no. 4, pp. 142–148, 1997.
- [81] H. Mustonen, A. Lepistö, S. Lehtonen, E. Lehtonen, P. Puolakkainen, and E. Kivilaakso, "CD2AP contributes to cell migration and adhesion in cultured gastric epithelium," *Biochemical and Biophysical Research Communications*, vol. 332, no. 2, pp. 426–432, 2005.
- [82] M. Lecuit, R. Hurme, J. Pizarro-Cerda, H. Ohayon, B. Geiger, and P. Cossart, "A role for α - and β -catenins in bacterial uptake," *Proceedings of the National Academy of Sciences*, vol. 97, no. 18, pp. 10008–10013, 2000.
- [83] S. Backert, N. Tegtmeyer, and M. Selbach, "The versatility of *Helicobacter pylori* caga effector protein functions: the master key hypothesis," *Helicobacter*, vol. 15, no. 3, pp. 163–176, 2010.
- [84] B. Hoy, M. Löwer, C. Weydig et al., "*Helicobacter pylori* HtrA is a new secreted virulence factor that cleaves E-cadherin to disrupt intercellular adhesion," *EMBO Reports*, vol. 11, no. 10, pp. 798–804, 2010.
- [85] A. MacIntyre, C. J. Hammond, C. S. Little, D. M. Appelt, and B. J. Balin, "Chlamydia pneumoniae infection alters the junctional complex proteins of human brain microvascular endothelial cells," *FEMS Microbiology Letters*, vol. 217, no. 2, pp. 167–172, 2002.
- [86] M. J. Taylor, D. Perrais, and C. J. Merrifield, "A high precision survey of the molecular dynamics of mammalian clathrin-mediated endocytosis," *PLoS Biology*, vol. 9, no. 3, Article ID e1000604, 2011.
- [87] A. S. Nicot, A. Toussaint, V. Tosch et al., "Mutations in amphiphysin 2 (BIN1) disrupt interaction with dynamin 2 and cause autosomal recessive centronuclear myopathy," *Nature Genetics*, vol. 39, no. 9, pp. 1134–1139, 2007.
- [88] K. Nishi and K. Saigo, "Cellular internalization of green fluorescent protein fused with herpes simplex virus protein VP22 via a lipid raft-mediated endocytic pathway independent of caveolae and Rho family GTPases but dependent on dynamin and Arf6," *Journal of Biological Chemistry*, vol. 282, no. 37, pp. 27503–27517, 2007.
- [89] H. T. McMahon, P. Wigge, and C. Smith, "Clathrin interacts specifically with amphiphysin and is displaced by dynamin," *FEBS Letters*, vol. 413, no. 2, pp. 319–322, 1997.
- [90] E. S. Gold, R. M. Simmons, T. W. Petersen, L. A. Campbell, C. C. Kuo, and A. Aderem, "Amphiphysin II is required for survival of *Chlamydia pneumoniae* in macrophages," *Journal of Experimental Medicine*, vol. 200, no. 5, pp. 581–586, 2004.
- [91] V. Wixler, E. Laplantine, D. Geerts et al., "Identification of novel interaction partners for the conserved membrane proximal region of α -integrin cytoplasmic domains," *FEBS Letters*, vol. 445, no. 2-3, pp. 351–355, 1999.
- [92] M. Ulanova, S. Gravelle, and R. Barnes, "The role of epithelial integrin receptors in recognition of pulmonary pathogens," *Journal of Innate Immunity*, vol. 1, no. 1, pp. 4–17, 2008.
- [93] N. Tegtmeyer, S. Wessler, and S. Backert, "Role of the cag-pathogenicity island encoded type IV secretion system in *Helicobacter pylori* pathogenesis," *FEBS Journal*, vol. 278, no. 8, pp. 1190–1202, 2011.
- [94] P. R. Crocker and P. Redelinghuys, "Siglecs as positive and negative regulators of the immune system," *Biochemical Society Transactions*, vol. 36, no. 6, pp. 1467–1471, 2008.
- [95] E. C. Brinkman-Van der Linden and A. Varki, "New aspects of siglec binding specificities, including the significance of fucosylation and of the sialyl-Tn epitope. Sialic acid-binding immunoglobulin superfamily lectins," *Journal of Biological Chemistry*, vol. 275, no. 12, pp. 8625–8632, 2000.
- [96] J. R. Teuton and C. R. Brandt, "Sialic acid on herpes simplex virus type 1 envelope glycoproteins is required for efficient infection of cells," *Journal of Virology*, vol. 81, no. 8, pp. 3731–3739, 2007.
- [97] M. L. Rodrigues, S. Rozental, J. N. Couceiro, J. Angluster, C. S. Alviano, and L. R. Travassos, "Identification of N-Acetylneuraminic acid and its 9-O-acetylated derivative on the cell surface of *Cryptococcus neoformans*. Influence on fungal phagocytosis," *Infection and Immunity*, vol. 65, no. 12, pp. 4937–4942, 1997.

- [98] D. Hulínská, P. Volf, and L. Grubhoffer, "Characterization of *Borrelia burgdorferi* glycoconjugates and surface carbohydrates," *Zentralblatt für Bakteriologie*, vol. 276, no. 4, pp. 473–480, 1992.
- [99] M. Aspholm, F. O. Olfat, J. Nordén et al., "SabA is the *H. pylori* hemagglutinin and is polymorphic in binding to sialylated glycans," *PLoS Pathogens*, vol. 2, no. 10, Article ID e110, 2006.
- [100] H. J. Bennett and I. S. Roberts, "Identification of a new sialic acid-binding protein in *Helicobacter pylori*," *FEMS Immunology and Medical Microbiology*, vol. 44, no. 2, pp. 163–169, 2005.
- [101] U. Hallén, A. E. Björkner, and E. C. Hallberg, "Binding of the periodontitis associated bacterium *Porphyromonas gingivalis* to glycoproteins from human epithelial cells," *Oral Microbiology and Immunology*, vol. 23, no. 5, pp. 367–371, 2008.
- [102] A. J. Muller, J. B. DuHadaway, P. S. Donover, E. Sutanto-Ward, and G. C. Prendergast, "Inhibition of indoleamine 2,3-dioxygenase, an immunoregulatory target of the cancer suppression gene Bin1, potentiates cancer chemotherapy," *Nature Medicine*, vol. 11, no. 3, pp. 312–319, 2005.
- [103] A. Müller, K. Heseler, S. K. Schmidt, K. Spekter, C. R. MacKenzie, and W. Däubener, "The missing link between indoleamine 2,3-dioxygenase mediated antibacterial and immunoregulatory effects," *Journal of Cellular and Molecular Medicine*, vol. 13, no. 6, pp. 1125–1135, 2009.
- [104] C. R. MacKenzie, K. Heseler, A. Müller, and W. Däubener, "Role of indoleamine 2,3-dioxygenase in antimicrobial defence and immuno-regulation: tryptophan depletion versus production of toxic kynurenines," *Current Drug Metabolism*, vol. 8, no. 3, pp. 237–244, 2007.
- [105] L. Capuron and A. H. Miller, "Immune system to brain signaling: neuropsychopharmacological implications," *Pharmacology and Therapeutics*, vol. 130, no. 2, pp. 226–238, 2011.
- [106] E. Gulaj, K. Pawlak, B. Bien, and D. Pawlak, "Kynurenine and its metabolites in Alzheimer's disease patients," *Advances in Medical Sciences*, vol. 55, no. 2, pp. 204–211, 2010.
- [107] B. Widner, F. Leblhuber, J. Walli, G. P. Tilz, U. Demel, and D. Fuchs, "Tryptophan degradation and immune activation in Alzheimer's disease," *Journal of Neural Transmission*, vol. 107, no. 3, pp. 343–353, 2000.
- [108] R. G. Hamilton, D. W. MacGlashan Jr., and S. S. Saini, "IgE antibody-specific activity in human allergic disease," *Immunologic Research*, vol. 47, no. 1–3, pp. 273–284, 2010.
- [109] S. Ida, R. P. Siraganian, and A. L. Notkins, "Cell-bound and circulating IgE antibody to herpes simplex virus," *Journal of General Virology*, vol. 64, no. 3, pp. 533–537, 1983.
- [110] J. Lagace-Simard, J. D. Portnoy, and M. A. Wainberg, "High levels of IgE in patients suffering from frequent recurrent herpes simplex lesions," *Journal of Allergy and Clinical Immunology*, vol. 77, no. 4, pp. 582–585, 1986.
- [111] E. Calenoff, J. C. Zhao, E. L. Derlacki et al., "Patients with Meniere's disease possess IgE reacting with herpes family viruses," *Archives of Otolaryngology: Head and Neck Surgery*, vol. 121, no. 8, pp. 861–864, 1995.
- [112] A. Aceti, D. Celestino, M. Caferro et al., "Basophil-bound and serum immunoglobulin E directed against *Helicobacter pylori* in patients with chronic gastritis," *Gastroenterology*, vol. 101, no. 1, pp. 131–137, 1991.
- [113] M. H. Bluth, J. Robin, M. Ruditsky et al., "IgE anti-*Borrelia burgdorferi* components (p18, p31, p34, p41, p45, p60) and increased blood CD8⁺CD60⁺ T cells in children with Lyme disease," *Scandinavian Journal of Immunology*, vol. 65, no. 4, pp. 376–382, 2007.
- [114] U. Emre, N. Sokolovskaya, P. M. Roblin, J. Schachter, and M. R. Hammerschlag, "Detection of anti-*Chlamydia pneumoniae* IgE in children with reactive airway disease," *Journal of Infectious Diseases*, vol. 172, no. 1, pp. 265–267, 1995.
- [115] M. Liutu, K. Kalimo, J. Uksila, and J. Savolainen, "Extraction of ige-binding components of *Helicobacter pylori* by immunoblotting analysis in chronic urticaria patients," *International Archives of Allergy and Immunology*, vol. 126, no. 3, pp. 213–217, 2001.
- [116] M. Feldmesser, A. Casadevall, Y. Kress, G. Spira, and A. Orlofsky, "Eosinophil-Cryptococcus neoformans interactions in vivo and in vitro," *Infection and Immunity*, vol. 65, no. 5, pp. 1899–1907, 1997.
- [117] K. Ishibashi, M. Suzuki, S. Sasaki, and M. Imai, "Identification of a new multigene four-transmembrane family (MS4A) related to CD20, HTm4 and β subunit of the high-affinity IgE receptor," *Gene*, vol. 264, no. 1, pp. 87–93, 2001.
- [118] H. Xu, Y. Yan, M. S. Williams et al., "MS4a4B, a CD20 Homologue in T Cells, Inhibits T Cell Propagation by Modulation of Cell Cycle," *PLoS ONE*, vol. 5, no. 11, Article ID e13780, 2010.
- [119] M. Li, L. Wang, X. Ren, and C. Zheng, "Host cell targets of tegument protein VP22 of herpes simplex virus 1," *Archives of Virology*, vol. 156, no. 6, pp. 1079–1084, 2011.
- [120] G. M. di Guglielmo, C. Le Roy, A. F. Goodfellow, and J. L. Wrana, "Distinct endocytic pathways regulate TGF- β receptor signalling and turnover," *Nature Cell Biology*, vol. 5, no. 5, pp. 410–421, 2003.
- [121] K. B. Reddy, M. C. Karode, A. K. Harmony, and P. H. Howe, "Interaction of transforming growth factor β receptors with apolipoprotein J/clusterin," *Biochemistry*, vol. 35, no. 1, pp. 309–314, 1996.
- [122] L. Yang, Y. Pang, and H. L. Moses, "TGF- β and immune cells: an important regulatory axis in the tumor microenvironment and progression," *Trends in Immunology*, vol. 31, no. 6, pp. 220–227, 2010.
- [123] C. J. Tartari, L. Scapozza, and C. Gambacorti-Passerini, "The ALK gene, an attractive target for inhibitor development," *Current Topics in Medicinal Chemistry*, vol. 11, no. 11, pp. 1406–1419, 2011.
- [124] L. Lamant, R. D. Gascoyne, M. M. Duplantier et al., "Non-muscle myosin heavy chain (MYH9): a new partner fused to ALK in anaplastic large cell lymphoma," *Genes Chromosomes and Cancer*, vol. 37, no. 4, pp. 427–432, 2003.
- [125] N. Wang, D. Lan, M. Gerbod-Giannone et al., "ATP-binding cassette transporter A7 (ABCA7) binds apolipoprotein A-I and mediates cellular phospholipid but not cholesterol efflux," *Journal of Biological Chemistry*, vol. 278, no. 44, pp. 42906–42912, 2003.
- [126] S. L. Chan, W. S. Kim, J. B. Kwok et al., "ATP-binding cassette transporter A7 regulates processing of amyloid precursor protein in vitro," *Journal of Neurochemistry*, vol. 106, no. 2, pp. 793–804, 2008.
- [127] I. L. Van Genderen, R. Brandimarti, M. R. Torrisi, G. Campadelli, and G. Van Meer, "The phospholipid composition of extracellular herpes simplex virions differs from that of host cell nuclei," *Virology*, vol. 200, no. 2, pp. 831–836, 1994.
- [128] W. L. Steinhart, C. M. Nicolet, and J. L. Howland, "Incorporation of ³²P-phosphate into membrane phospholipids during infection of cultured human fibroblasts by herpes

- simplex virus type 1," *Intervirology*, vol. 16, no. 2, pp. 80–85, 1981.
- [129] W. L. Steinhart, J. S. Busch, J. P. Oettgen, and J. L. Howland, "Sphingolipid metabolism during infection of human fibroblasts by herpes simplex virus type 1," *Intervirology*, vol. 21, no. 2, pp. 70–76, 1984.
- [130] V. R. Gupta, B. A. Wilson, and S. R. Blanke, "Sphingomyelin is important for the cellular entry and intracellular localization of *Helicobacter pylori* VacA," *Cellular Microbiology*, vol. 12, no. 10, pp. 1517–1533, 2010.
- [131] K. Wolf and T. Hackstadt, "Sphingomyelin trafficking in *Chlamydia pneumoniae*-infected cells," *Cellular Microbiology*, vol. 3, no. 3, pp. 145–152, 2001.
- [132] S. Galdiero, A. Falanga, G. Vitiello et al., "Role of membraneotropic sequences from herpes simplex virus type I glycoproteins B and H in the fusion process," *Biochimica et Biophysica Acta*, vol. 1798, no. 3, pp. 579–591, 2010.
- [133] C. J. Chrisman, P. Albuquerque, A. J. Guimaraes, E. Nieves, and A. Casadevall, "Phospholipids trigger *Cryptococcus neoformans* capsular enlargement during interactions with amoebae and macrophages," *PLoS Pathogens*, vol. 7, no. 5, Article ID e1002047, 2011.
- [134] D. Kielar, W. E. Kaminski, G. Liebisch et al., "Adenosine triphosphate binding cassette (ABC) transporters are expressed and regulated during terminal keratinocyte differentiation: a potential role for ABCA7 in epidermal lipid reorganization," *Journal of Investigative Dermatology*, vol. 121, no. 3, pp. 465–474, 2003.
- [135] E. A. Hunsperger and C. L. Wilcox, "Caspase-3-dependent reactivation of latent herpes simplex virus type 1 in sensory neuronal cultures," *Journal of NeuroVirology*, vol. 9, no. 3, pp. 390–398, 2003.
- [136] J. Marino, I. Stoeckli, M. Walch et al., "Chlamydia pneumoniae derived from inclusions late in the infectious cycle induce apoptosis in human aortic endothelial cells," *BMC Microbiology*, vol. 8, article 32, 2008.
- [137] R. V. Srinivas, Y. V. Venkatachalapathi, Z. Rui et al., "Inhibition of virus-induced cell fusion by apolipoprotein A-I and its amphipathic peptide analogs," *Journal of Cellular Biochemistry*, vol. 45, no. 2, pp. 224–237, 1991.
- [138] N. Tanaka, S. Abe-Dohmae, N. Iwamoto, and S. Yokoyama, "Roles of ATP-binding cassette transporter A7 in cholesterol homeostasis and host defense system," *Journal of Atherosclerosis and Thrombosis*, vol. 18, pp. 274–281, 2011.
- [139] J. S. Burgos, C. Ramirez, I. Sastre, and F. Valdivieso, "Apolipoprotein E genotype influences vertical transmission of herpes simplex virus type 1 in a gender specific manner," *Aging Cell*, vol. 6, no. 6, pp. 841–842, 2007.
- [140] W. Cun, J. Jiang, and G. Luo, "The C-terminal α -helix domain of apolipoprotein E is required for interaction with nonstructural protein 5A and assembly of hepatitis C virus," *Journal of Virology*, vol. 84, no. 21, pp. 11532–11541, 2010.
- [141] T. Hishiki, Y. Shimizu, R. Tobita et al., "Infectivity of hepatitis C virus is influenced by association with apolipoprotein E isoforms," *Journal of Virology*, vol. 84, no. 22, pp. 12048–12057, 2010.
- [142] J. Turchan-Cholewo, Y. Liu, S. Gartner et al., "Increased vulnerability of ApoE4 neurons to HIV proteins and opiates: protection by diosgenin and l-deprenyl," *Neurobiology of Disease*, vol. 23, no. 1, pp. 109–119, 2006.
- [143] W. R. Lin, M. A. Wozniak, M. M. Esiri, P. Klennerman, and R. F. Itzhaki, "Herpes simplex encephalitis: involvement of apolipoprotein E genotype," *Journal of Neurology Neurosurgery and Psychiatry*, vol. 70, no. 1, pp. 117–119, 2001.
- [144] H. C. Gérard, E. Fomicheva, J. A. Whittum-Hudson, and A. P. Hudson, "Apolipoprotein E4 enhances attachment of *Chlamydia pneumoniae* elementary bodies to host cells," *Microbial Pathogenesis*, vol. 44, no. 4, pp. 279–285, 2008.
- [145] C. M. Brown, E. Choi, Q. Xu, M. P. Vitek, and C. A. Colton, "The APOE4 genotype alters the response of microglia and macrophages to 17 β -estradiol," *Neurobiology of Aging*, vol. 29, no. 12, pp. 1783–1794, 2008.
- [146] M. P. Vitek, C. M. Brown, and C. A. Colton, "APOE genotype-specific differences in the innate immune response," *Neurobiology of Aging*, vol. 30, no. 9, pp. 1350–1360, 2009.
- [147] M. N. Haan, A. E. Aiello, N. A. West, and W. J. Jagust, "C-reactive protein and rate of dementia in carriers and non carriers of Apolipoprotein APOE4 genotype," *Neurobiology of Aging*, vol. 29, no. 12, pp. 1774–1782, 2008.
- [148] D. Thompson, M. B. Pepys, and S. P. Wood, "The physiological structure of human C-reactive protein and its complex with phosphocholine," *Structure*, vol. 7, no. 2, pp. 169–177, 1999.
- [149] N. de Bont, M. G. Netea, P. N. Demacker et al., "Apolipoprotein E knock-out mice are highly susceptible to endotoxemia and *Klebsiella pneumoniae* infection," *Journal of Lipid Research*, vol. 40, no. 4, pp. 680–685, 1999.
- [150] D. G. Alber, K. L. Powell, P. Vallance, D. A. Goodwin, and C. Grahame-Clarke, "Herpesvirus infection accelerates atherosclerosis in the apolipoprotein E-deficient mouse," *Circulation*, vol. 102, no. 7, pp. 779–785, 2000.
- [151] K. Ayada, K. Yokota, K. Kobayashi, Y. Shoenfeld, E. Matsuura, and K. Oguma, "Chronic infections and atherosclerosis," *Annals of the New York Academy of Sciences*, vol. 1108, pp. 594–602, 2007.
- [152] R. Ezzahiri, F. R. Stassen, H. A. Kurvers, M. M. van Pul, P. J. Kitslaar, and C. A. Bruggeman, "Chlamydia pneumoniae infection induces an unstable atherosclerotic plaque phenotype in LDL-receptor, ApoE double knockout mice," *European Journal of Vascular and Endovascular Surgery*, vol. 26, no. 1, pp. 88–95, 2003.
- [153] E. Hsich, Y. F. Zhou, B. Paigen, T. M. Johnson, M. S. Burnett, and S. E. Epstein, "Cytomegalovirus infection increases development of atherosclerosis in Apolipoprotein-E knockout mice," *Atherosclerosis*, vol. 156, no. 1, pp. 23–28, 2001.
- [154] L. Li, E. Messas, E. L. Batista Jr., R. A. Levine, and S. Amar, "Porphyromonas gingivalis infection accelerates the progression of atherosclerosis in a heterozygous apolipoprotein E-deficient murine model," *Circulation*, vol. 105, no. 7, pp. 861–867, 2002.
- [155] M. Naghavi, P. Wyde, S. Litovsky et al., "Influenza infection exerts prominent inflammatory and thrombotic effects on the atherosclerotic plaques of apolipoprotein E-deficient mice," *Circulation*, vol. 107, no. 5, pp. 762–768, 2003.
- [156] L. R. Portugal, L. R. Fernandes, G. C. Cesar et al., "Infection with *Toxoplasma gondii* increases atherosclerotic lesion in ApoE-deficient mice," *Infection and Immunity*, vol. 72, no. 6, pp. 3571–3576, 2004.
- [157] K. Ayada, K. Yokota, K. Hirai et al., "Regulation of cellular immunity prevents *Helicobacter pylori*-induced atherosclerosis," *Lupus*, vol. 18, no. 13, pp. 1154–1168, 2009.
- [158] M. S. Warner, R. J. Geraghty, W. M. Martinez et al., "A cell surface protein with herpesvirus entry activity (Hvab) confers susceptibility to infection by mutants of herpes simplex virus type 1, herpes simplex virus type 2, and

- pseudorabies virus," *Virology*, vol. 246, no. 1, pp. 179–189, 1998.
- [159] I. Derré, M. Pypaert, A. Dautry-Varsat, and H. Agaisse, "RNAi screen in *Drosophila* cells reveals the involvement of the tom complex in *Chlamydia* infection," *PLoS Pathogens*, vol. 3, no. 10, pp. 1446–1458, 2007.
- [160] J. Harland, P. Dunn, E. Cameron, J. Conner, and S. M. Brown, "The herpes simplex virus (HSV) protein ICP34.5 is a virion component that forms a DNA-binding complex with proliferating cell nuclear antigen and HSV replication proteins," *Journal of NeuroVirology*, vol. 9, no. 4, pp. 477–488, 2003.
- [161] A. Lupo, E. Cesaro, G. Montano, P. Izzo, and P. Costanzo, "ZNF224: structure and role of a multifunctional KRAB-ZFP protein," *International Journal of Biochemistry and Cell Biology*, vol. 43, no. 4, pp. 470–473, 2011.
- [162] J. Yu, B. Shin, E. S. Park et al., "Protein arginine methyltransferase 1 regulates herpes simplex virus replication through ICP27 RGG-box methylation," *Biochemical and Biophysical Research Communications*, vol. 391, no. 1, pp. 322–328, 2010.
- [163] A. Kamiya, K. Kubo, T. Tomoda et al., "A schizophrenia-associated mutation of DISC1 perturbs cerebral cortex development," *Nature Cell Biology*, vol. 7, no. 12, pp. 1067–1078, 2005.
- [164] C. Janke, K. Rogowski, D. Wloga et al., "Biochemistry: tubulin polyglutamylase enzymes are members of the TTL domain protein family," *Science*, vol. 308, no. 5729, pp. 1758–1762, 2005.
- [165] P. Gee, Y. Ando, H. Kitayama et al., "APOBEC1-mediated editing and attenuation of HSV-1 DNA implicates an antiviral role in neurons during encephalitis," *The Journal of Virology*, vol. 85, no. 19, pp. 9726–9736, 2011.
- [166] S. K. Bose, W. Gibson, R. S. Bullard, and C. D. Donald, "PAX2 oncogene negatively regulates the expression of the host defense peptide human beta defensin-1 in prostate cancer," *Molecular Immunology*, vol. 46, no. 6, pp. 1140–1148, 2009.
- [167] R. Tesse, N. Santoro, P. Giordano, F. Cardinale, D. D. Mattia, and L. Armenio, "Association between defb1 gene haplotype and herpes viruses seroprevalence in children with acute lymphoblastic leukemia," *Pediatric Hematology and Oncology*, vol. 26, no. 8, pp. 573–582, 2009.
- [168] A. K. Kocsis, Z. F. Kiss, L. Tiszlavicz, Z. Tiszlavicz, and Y. Mandi, "Potential role of human beta-defensin 1 in *Helicobacter pylori*-induced gastritis," *Scandinavian Journal of Gastroenterology*, vol. 44, pp. 289–295, 2009.
- [169] B. Wiechula, K. Cholewa, A. Ekiel, M. Romanik, H. Dolezych, and G. Martirosian, "HBD-1 and hBD-2 are expressed in cervico-vaginal lavage in female genital tract due to microbial infections," *Ginekologia Polska*, vol. 81, no. 4, pp. 268–271, 2010.
- [170] R. Circo, B. Skerlavaj, R. Gennaro, A. Amoroso, and M. Zanetti, "Structural and functional characterization of hBD-1(Ser35), a peptide deduced from a DEFB1 polymorphism," *Biochemical and Biophysical Research Communications*, vol. 293, no. 1, pp. 586–592, 2002.
- [171] P. Meyer, C. Caillat, D. Topalis, J. Balzarini, and D. Deville-Bonne, "Structural basis for the specificity of thymidylate kinases from human pathogens: implications for nucleotide analogues activation," *Nucleic Acids Symposium Series*, no. 53, p. 41, 2009.
- [172] Y. Sawayama, M. Tatsukawa, S. Maeda, H. Ohnishi, N. Furusyo, and J. Hayashi, "Association of hyperhomocysteinemia and *Chlamydia pneumoniae* infection with carotid atherosclerosis and coronary artery disease in Japanese patients," *Journal of Infection and Chemotherapy*, vol. 14, no. 3, pp. 232–237, 2008.
- [173] T. Matsui, "Helicobacter pylori and Arteriosclerosis," *Gan To Kagaku Ryoho*, vol. 38, no. 3, pp. 365–369, 2011.
- [174] H. P. Huemer, H. J. Menzel, D. Potratz et al., "Herpes simplex virus binds to human serum lipoprotein," *Intervirology*, vol. 29, no. 2, pp. 68–76, 1988.
- [175] S. I. Vieira, S. Rebelo, H. Esselmann et al., "Retrieval of the Alzheimer's amyloid precursor protein from the endosome to the TGN is S655 phosphorylation state-dependent and retromer-mediated," *Molecular Neurodegeneration*, vol. 5, no. 1, article 40, 2010.
- [176] N. R. Marquez-Sterling, A. C. Lo, S. S. Sisodia, and E. H. Koo, "Trafficking of cell-surface β -amyloid precursor protein: evidence that a sorting intermediate participates in synaptic vesicle recycling," *Journal of Neuroscience*, vol. 17, no. 1, pp. 140–151, 1997.
- [177] M. S. Song, G. B. Baker, K. G. Todd, and S. Kar, "Inhibition of beta-amyloid1-42 internalization attenuates neuronal death by stabilizing the endosomal-lysosomal system in rat cortical cultured neurons," *Neuroscience*, vol. 178, pp. 181–188, 2011.
- [178] S. Mandrekar, Q. Jiang, C. Y. Lee, J. Koenigsnecht-Talboo, D. M. Holtzman, and G. E. Landreth, "Microglia mediate the clearance of soluble $a\beta$ through fluid phase macropinocytosis," *Journal of Neuroscience*, vol. 29, no. 13, pp. 4252–4262, 2009.
- [179] F. Wu, Y. Matsuoka, M. P. Mattson, and P. J. Yao, "The clathrin assembly protein AP180 regulates the generation of amyloid- β peptide," *Biochemical and Biophysical Research Communications*, vol. 385, no. 2, pp. 247–250, 2009.
- [180] J. R. Cirrito, J. E. Kang, J. Lee et al., "Endocytosis is required for synaptic activity-dependent release of amyloid- β in vivo," *Neuron*, vol. 58, no. 1, pp. 42–51, 2008.
- [181] M. M. Hussain, "Structural, biochemical and signaling properties of the low-density lipoprotein receptor gene family," *Frontiers in Bioscience*, vol. 6, pp. D417–D428, 2001.
- [182] M. Dieckmann, M. F. Dietrich, and J. Herz, "Lipoprotein receptors—an evolutionarily ancient multifunctional receptor family," *Biological Chemistry*, vol. 391, no. 11, pp. 1341–1363, 2010.
- [183] M. Deuss, K. Reiss, and D. Hartmann, "Part-time α -secretases: the functional biology of ADAM 9, 10 and 17," *Current Alzheimer Research*, vol. 5, no. 2, pp. 187–201, 2008.
- [184] C. Sato, G. Zhao, and M. X. Ilagan, "An overview of notch signaling in adult tissue renewal and maintenance," *Current Alzheimer Research*. In press.
- [185] B. B. Fuchs, R. J. Tang, and E. Mylonakis, "The temperature-sensitive role of *Cryptococcus neoformans* ROM2 in cell morphogenesis," *PLoS ONE*, vol. 2, no. 4, article e368, 2007.
- [186] J. Mital and T. Hackstadt, "Diverse requirements for SRC-family tyrosine kinases distinguish chlamydial species," *MBio*, vol. 2, no. 2, 2011.
- [187] R. Pai, T. L. Cover, and A. S. Tarnawski, "Helicobacter pylori vacuolating cytotoxin (VacA) disorganizes the cytoskeletal architecture of gastric epithelial cells," *Biochemical and Biophysical Research Communications*, vol. 262, no. 1, pp. 245–250, 1999.
- [188] H. Takeuchi, N. Furuta, I. Morisaki, and A. Amano, "Exit of intracellular *Porphyromonas gingivalis* from gingival epithelial cells is mediated by endocytic recycling pathway," *Cellular Microbiology*, vol. 13, no. 5, pp. 677–691, 2011.
- [189] M. D. Weingarten, A. H. Lockwood, S. Y. Hwo, and M. W. Kirschner, "A protein factor essential for microtubule assembly," *Proceedings of the National Academy of Sciences of*

- the United States of America*, vol. 72, no. 5, pp. 1858–1862, 1975.
- [190] P. J. Dolan and G. V. Johnson, "The role of tau kinases in Alzheimer's disease," *Current Opinion in Drug Discovery and Development*, vol. 13, no. 5, pp. 595–603, 2010.
 - [191] Á. Zambrano, L. Solís, N. Salvadores, M. Cortés, R. Lerchundi, and C. Otth, "Neuronal cytoskeletal dynamic modification and neurodegeneration induced by infection with herpes simplex virus type 1," *Journal of Alzheimer's Disease*, vol. 14, no. 3, pp. 259–269, 2008.
 - [192] C. J. Carter, "Alzheimer's disease: a pathogenetic autoimmune disorder caused by herpes simplex in a gene-dependent manner," *International Journal of Alzheimer's Disease*, 2010, article 140539.
 - [193] S. B. Cheng, P. Ferland, P. Webster, and E. L. Bearer, "Herpes simplex virus dances with amyloid precursor protein while exiting the cell," *PLoS ONE*, vol. 6, no. 3, Article ID e17966, 2011.
 - [194] L. Benboudjema, M. Mulvey, Y. Gao, S. W. Pimplikar, and I. Mohr, "Association of the herpes simplex virus type 1 Us11 gene product with the cellular kinesin light-chain-related protein PAT1 results in the redistribution of both polypeptides," *Journal of Virology*, vol. 77, no. 17, pp. 9192–9203, 2003.
 - [195] P. Zheng, J. Eastman, P. S. Vande, and S. W. Pimplikar, "PAT1, a microtubule-interacting protein, recognizes the basolateral sorting signal of amyloid precursor protein," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 95, no. 25, pp. 14745–14750, 1998.
 - [196] S. J. Soscia, J. E. Kirby, K. J. Washicosky et al., "The Alzheimer's disease-associated amyloid β -protein is an antimicrobial peptide," *PLoS ONE*, vol. 5, no. 3, Article ID e9505, 2010.
 - [197] W. J. Lukiw, J. G. Cui, L. Y. Yuan et al., "Acyclovir or A β 42 peptides attenuate HSV-1-induced miRNA-146a levels in human primary brain cells," *NeuroReport*, vol. 21, no. 14, pp. 922–927, 2010.
 - [198] A. Salminen, J. Ojala, A. Kauppinen, K. Kaarniranta, and T. Suuronen, "Inflammation in Alzheimer's disease: amyloid- β oligomers trigger innate immunity defence via pattern recognition receptors," *Progress in Neurobiology*, vol. 87, no. 3, pp. 181–194, 2009.
 - [199] K. Tahara, H. D. Kim, J. J. Jin, J. A. Maxwell, L. Li, and K. Fukuchi, "Role of toll-like receptor signalling in A β uptake and clearance," *Brain*, vol. 129, no. 11, pp. 3006–3019, 2006.
 - [200] J.-H. Sohn, J. O. So, H. J. Hong et al., "Identification of autoantibody against beta-amyloid peptide in the serum of elderly," *Frontiers in Bioscience*, vol. 14, no. 10, pp. 3879–3883, 2009.
 - [201] A. Lleó and C. A. Saura, " γ -secretase substrates and their implications for drug development in Alzheimer's disease," *Current Topics in Medicinal Chemistry*, vol. 11, no. 12, pp. 1513–1527, 2011.
 - [202] O. Takeuchi and S. Akira, "Innate immunity to virus infection," *Immunological Reviews*, vol. 227, no. 1, pp. 75–86, 2009.
 - [203] I. Zanoni and F. Granucci, "Regulation of antigen uptake, migration, and lifespan of dendritic cell by Toll-like receptors," *Journal of Molecular Medicine*, vol. 88, no. 9, pp. 873–880, 2010.
 - [204] D. Y. Kim, L. A. Ingano, and D. M. Kovacs, "Nectin-1 α , an immunoglobulin-like receptor involved in the formation of synapses, is a substrate for presenilin/ γ -secretase-like cleavage," *Journal of Biological Chemistry*, vol. 277, no. 51, pp. 49976–49981, 2002.
 - [205] M. L. Hemming, J. E. Elias, S. P. Gygi, and D. J. Selkoe, "Proteomic profiling of gamma-secretase substrates and mapping of substrate requirements," *PLoS Biology*, vol. 6, no. 10, p. e257, 2008.
 - [206] S. Bacsa, G. Karasneh, S. Dosa, J. Liu, T. Valyi-Nagy, and D. Shukla, "Syndecan-1 and syndecan-2 play key roles in herpes simplex virus type-1 infection," *Journal of General Virology*, vol. 92, no. 4, pp. 733–743, 2011.
 - [207] M. Kalia, V. Chandra, S. A. Rahman, D. Sehgal, and S. Jameel, "Heparan sulfate proteoglycans are required for cellular binding of the hepatitis E virus ORF2 capsid protein and for viral infection," *Journal of Virology*, vol. 83, no. 24, pp. 12714–12724, 2009.
 - [208] S. Shafit-Keramat, A. Handisurya, E. Kriehuber, G. Meneguzzi, K. Slupetzky, and R. Kirnbauer, "Different heparan sulfate proteoglycans serve as cellular receptors for human papillomaviruses," *Journal of Virology*, vol. 77, no. 24, pp. 13125–13135, 2003.
 - [209] A. J. Smith, T. W. Schacker, C. S. Reilly, and A. T. Haase, "A role for syndecan-1 and claudin-2 in microbial translocation during HIV-1 infection," *Journal of Acquired Immune Deficiency Syndromes*, vol. 55, no. 3, pp. 306–315, 2010.
 - [210] L. de Witte, M. Bobardt, U. Chatterji et al., "Syndecan-3 is a dendritic cell-specific attachment receptor for HIV-1," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 49, pp. 19464–19469, 2007.
 - [211] L. de Witte, Y. Zoughlami, B. Aengeneyndt et al., "Binding of human papilloma virus L1 virus-like particles to dendritic cells is mediated through heparan sulfates and induces immune activation," *Immunobiology*, vol. 212, no. 9–10, pp. 679–691, 2008.
 - [212] G. A. Wilke and W. J. Bubeck, "Role of a disintegrin and metalloprotease 10 in *Staphylococcus aureus* α -hemolysin-mediated cellular injury," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 30, pp. 13473–13478, 2010.
 - [213] A. Jong, C. H. Wu, G. M. Shackleford et al., "Involvement of human CD44 during *Cryptococcus neoformans* infection of brain microvascular endothelial cells," *Cellular Microbiology*, vol. 10, no. 6, pp. 1313–1326, 2008.
 - [214] B. D. Persson, N. B. Schmitz, C. Santiago et al., "Structure of the extracellular portion of CD46 provides insights into its interactions with complement proteins and pathogens," *PLoS Pathogens*, vol. 6, no. 9, Article ID e01122, 2010.
 - [215] H. Mahtout, F. Chandad, J. M. Rojo, and D. Grenier, "Porphyromonas gingivalis mediates the shedding and proteolysis of complement regulatory protein CD46 expressed by oral epithelial cells," *Oral Microbiology and Immunology*, vol. 24, no. 5, pp. 396–400, 2009.
 - [216] H. Wang, Z.-Y. Li, Y. Liu et al., "Desmoglein 2 is a receptor for adenovirus serotypes 3, 7, 11 and 14," *Nature Medicine*, vol. 17, no. 1, pp. 96–104, 2011.
 - [217] F. Hofer, M. Gruenberger, H. Kowalski et al., "Members of the low density lipoprotein receptor family mediate cell entry of a minor-group common cold virus," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 91, no. 5, pp. 1839–1842, 1994.
 - [218] B. Ronacher, T. C. Marlovits, R. Moser, and D. Blaas, "Expression and folding of human very-low-density lipoprotein receptor fragments: neutralization capacity toward human rhinovirus HRV2," *Virology*, vol. 278, no. 2, pp. 541–550, 2000.

- [219] S. Jahan, S. Khaliq, B. Samreen et al., "Effect of combined siRNA of HCV E2 gene and HCV receptors against HCV," *Virology Journal*, vol. 8, article 295, 2011.
- [220] L. J. Anderson and R. Longnecker, "Epstein-Barr virus latent membrane protein 2A exploits Notch1 to alter B-cell identity in vivo," *Blood*, vol. 113, no. 1, pp. 108–116, 2009.
- [221] S. Kusano and N. Raab-Traub, "An Epstein-Barr virus protein interacts with Notch," *Journal of Virology*, vol. 75, no. 1, pp. 384–395, 2001.
- [222] E. Tzahar, J. D. Moyer, H. Waterman et al., "Pathogenic poxviruses reveal viral strategies to exploit the ErbB signaling network," *EMBO Journal*, vol. 17, no. 20, pp. 5948–5963, 1998.
- [223] C. Tuffereau, J. Bénéjean, D. Blondel, B. Kieffer, and A. Flament, "Low-affinity nerve-growth factor receptor (P75NTR) can serve as a receptor for rabies virus," *EMBO Journal*, vol. 17, no. 24, pp. 7250–7259, 1998.
- [224] T. A. Bowden, A. R. Aricescu, R. J. Gilbert, J. M. Grimes, E. Y. Jones, and D. I. Stuart, "Structural basis of Nipah and Hendra virus attachment to their cell-surface receptor ephrin-B2," *Nature Structural and Molecular Biology*, vol. 15, no. 6, pp. 567–572, 2008.
- [225] R. J. Orentas and J. E. Hildreth, "Association of host cell surface adhesion receptors and other membrane proteins with HIV and SIV," *AIDS Research and Human Retroviruses*, vol. 9, no. 11, pp. 1157–1165, 1993.
- [226] S. W. Rothwell and D. G. Wright, "Characterization of influenza A virus binding sites on human neutrophils," *Journal of Immunology*, vol. 152, no. 5, pp. 2358–2367, 1994.
- [227] E. Pericolini, E. Gabrielli, E. Cenci et al., "Involvement of glycoreceptors in galactoxylomannan-induced T cell death," *Journal of Immunology*, vol. 182, no. 10, pp. 6003–6010, 2009.
- [228] J. Vomaske, R. M. Melnychuk, P. P. Smith et al., "Differential ligand binding to a human cytomegalovirus chemokine receptor determines cell type-specific motility," *PLoS Pathogens*, vol. 5, no. 2, Article ID e1000304, 2009.
- [229] M. Minami, N. Kume, T. Shimaoka et al., "Expression of SR-PSOX, a novel cell-surface scavenger receptor for phosphatidylserine and oxidized LDL in human atherosclerotic lesions," *Arteriosclerosis, Thrombosis, and Vascular Biology*, vol. 21, no. 11, pp. 1796–1800, 2001.
- [230] O. Geiger, N. González-Silva, I. M. López-Lara, and C. Sohlenkamp, "Amino acid-containing membrane lipids in bacteria," *Progress in Lipid Research*, vol. 49, no. 1, pp. 46–60, 2010.
- [231] W. Cao, M. D. Henry, P. Borrow et al., "Identification of α -dystroglycan as a receptor for lymphocytic choriomeningitis virus and Lassa fever virus," *Science*, vol. 282, no. 5396, pp. 2079–2081, 1998.
- [232] N. Sevilla, S. Kunz, A. Holz et al., "Immunosuppression and resultant viral persistence by specific viral targeting of dendritic cells," *Journal of Experimental Medicine*, vol. 192, no. 9, pp. 1249–1260, 2000.
- [233] S. Sasaki, F. Takeshita, K. Okuda, and N. Ishii, "Mycobacterium leprae and leprosy: a compendium," *Microbiology and Immunology*, vol. 45, no. 11, pp. 729–736, 2001.
- [234] E. Boelen, F. R. Stassen, A. J. van der Ven et al., "Detection of amyloid beta aggregates in the brain of BALB/c mice after Chlamydia pneumoniae infection," *Acta Neuropathologica*, vol. 114, no. 3, pp. 255–261, 2007.
- [235] J. Miklossy, A. Kis, A. Radenovic et al., " β -amyloid deposition and Alzheimer's type changes induced by Borrelia spirochetes," *Neurobiology of Aging*, vol. 27, no. 2, pp. 228–236, 2006.
- [236] R. Piacentini, L. Civitelli, C. Ripoli et al., "HSV-1 promotes Ca^{2+} -mediated APP phosphorylation and $\text{A}\beta$ accumulation in rat cortical neurons," *Neurobiology of Aging*, vol. 32, no. 12, pp. 2323.e13–2323.e26, 2011.
- [237] M. S. Elkind, "Infectious burden: a new risk factor and treatment target for atherosclerosis," *Infectious Disorders*, vol. 10, no. 2, pp. 84–90, 2010.
- [238] D. Kanduc, "Describing the hexapeptide identity platform between the influenza A H5N1 and Homo sapiens proteomes," *Biologicals*, vol. 4, pp. 245–261, 2010.
- [239] D. Kanduc, A. Stufano, G. Lucchese, and A. Kusalik, "Massive peptide sharing between viral and human proteomes," *Peptides*, vol. 29, no. 10, pp. 1755–1766, 2008.
- [240] R. Ricco and D. Kanduc, "Hepatitis B virus and Homo sapiens proteome-wide analysis: a profusion of viral peptide overlaps in neuron-specific human proteins," *Biologics*, no. 4, pp. 75–81, 2010.
- [241] B. Trost, G. Lucchese, A. Stufano, M. Bickis, A. Kusalik, and D. Kanduc, "No human protein is exempt from bacterial motifs, not even one," *Self/Nonsense*, vol. 1, no. 4, pp. 328–334, 2010.
- [242] W. M. Pardridge, "Re-engineering biopharmaceuticals for delivery to brain with molecular Trojan horses," *Bioconjugate Chemistry*, vol. 19, no. 7, pp. 1327–1338, 2008.
- [243] P. Grammas, J. Martinez, and B. Miller, "Cerebral microvascular endothelium and the pathogenesis of neurodegenerative diseases," *Expert Reviews in Molecular Medicine*, vol. 13, Article ID e19, 2011.
- [244] G. Baravalle, M. Brabec, L. Snyers, D. Blaas, and R. Fuchs, "Human Rhinovirus Type 2-Antibody Complexes Enter and Infect Cells via Fc- γ Receptor IIB1," *Journal of Virology*, vol. 78, no. 6, pp. 2729–2737, 2004.
- [245] D. L. Mallery, W. A. McEwan, S. R. Bidgood, G. J. Towers, C. M. Johnson, and L. C. James, "Antibodies mediate intracellular immunity through tripartite motif-containing 21 (TRIM21)," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 46, pp. 19985–19990, 2010.
- [246] M. Jaume, M. S. Yip, C. Y. Cheung et al., "Anti-severe acute respiratory syndrome coronavirus spike antibodies trigger infection of human immune cells via a pH- and cysteine protease-independent Fc γ R pathway," *The Journal of Virology*, vol. 85, no. 20, pp. 10582–10597, 2011.
- [247] Y. N. Li, X. J. Qin, F. Kuang et al., "Alterations of Fc γ receptor I and Toll-like receptor 4 mediate the antiinflammatory actions of microglia and astrocytes after adrenaline-induced blood-brain barrier opening in rats," *Journal of Neuroscience Research*, vol. 86, no. 16, pp. 3556–3565, 2008.
- [248] C. J. Carter, "Epstein-Barr and other viral mimicry of autoantigens, myelin and vitamin D related proteins, and of EIF2B, the cause of vanishing white matter disease: massive mimicry of multiple sclerosis relevant proteins by the Synchococcus phage," *Immunopharmacology and Immunotoxicology*. In press.
- [249] C. J. Carter, "Pathogen and autoantigen homologous regions within the cystic fibrosis transmembrane conductance regulator (CFTR) protein suggest an autoimmune treatable component of cystic fibrosis," *FEMS Immunology & Medical Microbiology*, vol. 62, no. 2, pp. 197–214, 2011.
- [250] C. J. Carter, "Schizophrenia: a pathogenetic autoimmune disease caused by viruses and pathogens and dependent on genes," *Journal of Pathogens*, vol. 2011, Article ID 128318, 37 pages, 2011.

- [251] A. J. Espay and K. K. Henderson, "Postencephalitic parkinsonism and basal ganglia necrosis due to Epstein-Barr virus infection," *Neurology*, vol. 76, no. 17, pp. 1529–1530, 2011.
- [252] F. Roselli, I. Russo, A. Fraddosio et al., "Reversible Parkinsonian syndrome associated with anti-neuronal antibodies in acute EBV encephalitis: a case report," *Parkinsonism and Related Disorders*, vol. 12, no. 4, pp. 257–260, 2006.
- [253] J. Woulfe, H. Hoogendoorn, M. Tarnopolsky, and D. G. Munoz, "Monoclonal antibodies against Epstein-Barr virus cross-react with alpha-synuclein in human brain," *Neurology*, vol. 55, no. 9, pp. 1398–1401, 2000.
- [254] C. J. Carter, "Extensive viral mimicry of 22 AIDS-related autoantigens by HIV-1 proteins and pathway analysis of 561 viral/human homologues suggest an initial treatable autoimmune component of AIDS," *FEMS Immunology & Medical Microbiology*, vol. 63, no. 2, pp. 254–268, 2011.
- [255] D. Kanduc, "The self/nonself issue a confrontation between proteomes," *Self/Nonself*, vol. 1, no. 3, pp. 255–258, 2010.
- [256] L. Jones, P. A. Holmans, M. L. Hamshire et al., "Genetic evidence implicates the immune system and cholesterol metabolism in the aetiology of Alzheimer's disease," *PLoS ONE*, vol. 5, no. 11, Article ID e13950, 2010.
- [257] D. Phillips, L. Prentice, M. Upadhyaya et al., "Autosomal dominant inheritance of autoantibodies to thyroid peroxidase and thyroglobulin—studies in families not selected for autoimmune thyroid disease," *Journal of Clinical Endocrinology and Metabolism*, vol. 72, no. 5, pp. 973–975, 1991.
- [258] N. S. Zuckerman, H. Hazanov, M. Barak et al., "Somatic hypermutation and antigen-driven selection of B cells are altered in autoimmune diseases," *Journal of Autoimmunity*, vol. 35, no. 4, pp. 325–335, 2010.
- [259] J. H. Rho, W. Zhang, M. Murali, M. H. Roehrl, and J. Y. Wang, "Human proteins with affinity for dermatan sulfate have the propensity to become autoantigens," *American Journal of Pathology*, vol. 178, no. 5, pp. 2177–2190, 2011.
- [260] T. Wadström and Å. Ljungh, "Glycosaminoglycan-binding microbial proteins in tissue adhesion and invasion: key events in microbial pathogenicity," *Journal of Medical Microbiology*, vol. 48, no. 3, pp. 223–233, 1999.
- [261] H. Rosenmann, N. Grigoriadis, D. Karussis et al., "Tauopathy-like abnormalities and neurologic deficits in mice immunized with neuronal tau protein," *Archives of Neurology*, vol. 63, no. 10, pp. 1459–1467, 2006.
- [262] S. Capsoni, G. Ugolini, A. Comparini, F. Ruberti, N. Berardi, and A. Cattaneo, "Alzheimer-like neurodegeneration in aged antineurite growth factor transgenic mice," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 97, no. 12, pp. 6826–6831, 2000.
- [263] I. Ferrer, R. M. Boada, M. L. Sánchez Guerra, M. J. Rey, and F. Costa-Jussà, "Neuropathology and pathogenesis of encephalitis following amyloid- β immunization in Alzheimer's disease," *Brain Pathology*, vol. 14, no. 1, pp. 11–20, 2004.
- [264] R. Furlan, E. Brambilla, F. Sanvito et al., "Vaccination with amyloid- β peptide induces autoimmune encephalomyelitis in C57/BL6 mice," *Brain*, vol. 126, no. 2, pp. 285–291, 2003.
- [265] A. J. Rozemuller, W. A. van Gool, and P. Eikelenboom, "The neuroinflammatory response in plaques and amyloid angiopathy in Alzheimer's disease: therapeutic implications," *Current Drug Targets*, vol. 4, no. 3, pp. 223–233, 2005.
- [266] D. Vacirca, C. Barbat, B. Scazzocchio et al., "Anti-ATP synthase autoantibodies from patients with Alzheimer's disease reduce extracellular HDL level," *Journal of Alzheimer's Disease*, vol. 26, no. 3, pp. 441–445, 2011.
- [267] D. Vacirca, F. Delunardo, P. Matarrese et al., "Autoantibodies to the adenosine triphosphate synthase play a pathogenetic role in Alzheimer's disease," *Neurobiology of Aging*. In press.
- [268] P. Foley, H. F. Bradford, M. Docherty et al., "Evidence for the presence of antibodies to cholinergic neurons in the serum of patients with Alzheimer's disease," *Journal of Neurology*, vol. 235, no. 8, pp. 466–471, 1988.
- [269] R. G. Nagele, P. M. Clifford, G. Siu et al., "Brain-reactive autoantibodies prevalent in human sera increase intra-neuronal amyloid- γ 1-42 deposition," *Journal of Alzheimer's Disease*, vol. 25, no. 4, pp. 605–622, 2011.
- [270] L. Koval, O. Lykhmus, O. Kalashnyk et al., "The presence and origin of autoantibodies against α 4 and α 7 nicotinic acetylcholine receptors in the human blood: possible relevance to Alzheimer's pathology," *Journal of Alzheimer's Disease*, vol. 25, no. 4, pp. 747–761, 2011.
- [271] S. Mruthinti, J. J. Buccafusco, W. D. Hill et al., "Autoimmunity in Alzheimer's disease: increased levels of circulating IgGs binding A β and RAGE peptides," *Neurobiology of Aging*, vol. 25, no. 8, pp. 1023–1032, 2004.
- [272] M. A. Gruden, T. B. Davidova, M. Mališauskas et al., "Differential neuroimmune markers to the onset of Alzheimer's disease neurodegeneration and dementia: autoantibodies to A β (25–35) oligomers, S100b and neurotransmitters," *Journal of Neuroimmunology*, vol. 186, no. 1–2, pp. 181–192, 2007.
- [273] H. B. D. Kettlewell, "Selection experiments on industrial melanism in the *Lepidoptera*," *Heredity*, vol. 9, pp. 323–342, 1955.
- [274] J. S. Burgos, C. Ramirez, I. Sastre, and F. Valdivieso, "Effect of apolipoprotein E on the cerebral load of latent herpes simplex virus type 1 DNA," *Journal of Virology*, vol. 80, no. 11, pp. 5383–5387, 2006.
- [275] S. Li, D. Carpenter, C. Hsiang, S. L. Wechsler, and C. Jones, "Herpes simplex virus type 1 latency-associated transcript inhibits apoptosis and promotes neurite sprouting in neuroblastoma cells following serum starvation by maintaining protein kinase B (AKT) levels," *Journal of General Virology*, vol. 91, no. 4, pp. 858–866, 2010.
- [276] C. L. Wilcox and E. M. Johnson Jr., "Nerve growth factor deprivation results in the reactivation of latent herpes simplex virus in vitro," *Journal of Virology*, vol. 61, no. 7, pp. 2311–2315, 1987.
- [277] V. Camarena, M. Kobayashi, J. Y. Kim et al., "Nature and duration of growth factor signaling through receptor tyrosine kinases regulates HSV-1 latency in neurons," *Cell Host and Microbe*, vol. 8, no. 4, pp. 320–330, 2010.
- [278] S. E. Counts, M. Nadeem, J. Wu, S. D. Ginsberg, H. U. Saragovi, and E. J. Mufson, "Reduction of cortical TrkA but not p75NTR protein in early-stage Alzheimer's disease," *Annals of Neurology*, vol. 56, no. 4, pp. 520–531, 2004.
- [279] S. Kheirvari, K. Uezu, S. Yamamoto, and Y. Nakaya, "High-dose dietary supplementation of vitamin A induces brain-derived neurotrophic factor and nerve growth factor production in mice with simultaneous deficiency of vitamin A and zinc," *Nutritional Neuroscience*, vol. 11, no. 5, pp. 228–234, 2008.
- [280] B. Jiang, E. Y. Liao, L. M. Tan, R. C. Dai, Z. J. Xiao, and H. J. Liao, "Effects of long-term replacement therapy of compound nylestriol tablet or low-dose 17 beta-estradiol on the expression of nerve growth factor in OVX rat hippocampal formation," *Zhong Nan Da Xue Xue Bao. Yi Xue Ban*, vol. 29, no. 5, pp. 529–533, 2004.
- [281] R. D. Vicetti Miguel, B. S. Sheridan, S. A. K. Harvey, R. S. Schreiner, R. L. Hendricks, and T. L. Cherpes, "17- β estradiol

- promotion of herpes simplex virus type 1 reactivation is estrogen receptor dependent," *Journal of Virology*, vol. 84, no. 1, pp. 565–572, 2010.
- [282] M. I. Geerlings, L. J. Launer, F. H. de Jong et al., "Endogenous estradiol and risk of dementia in women and men: the Rotterdam study," *Annals of Neurology*, vol. 53, no. 5, pp. 607–615, 2003.
- [283] G. Ravaglia, P. Forti, F. Maioli et al., "Endogenous sex hormones as risk factors for dementia in elderly men and women," *Journals of Gerontology*, vol. 62, no. 9, pp. 1035–1041, 2007.
- [284] C. Clement, P. S. Bhattacharjee, H. E. Kaufman, and J. M. Hill, "Heat-induced reactivation of HSV-1 in latent mice: upregulation in the TG of CD83 and other immune response genes and their LAT-ICP0 locus," *Investigative Ophthalmology and Visual Science*, vol. 50, no. 6, pp. 2855–2861, 2009.
- [285] J. D. Kriesel, J. Ricigliano, S. L. Spruance, H. H. Garza Jr., and J. M. Hill, "Neuronal reactivation of herpes simplex virus may involve interleukin-6," *Journal of NeuroVirology*, vol. 3, no. 6, pp. 441–448, 1997.
- [286] I. Walev, J. Podlech, and D. Falke, "Enhancement by TNF- α of reactivation and replication of latent herpes simplex virus from trigeminal ganglia of mice," *Archives of Virology*, vol. 140, no. 6, pp. 987–992, 1995.
- [287] J. Kálmán, A. Juhász, G. Laird et al., "Serum interleukin-6 levels correlate with the severity of dementia in down syndrome and in Alzheimer's disease," *Acta Neurologica Scandinavica*, vol. 96, no. 4, pp. 236–240, 1997.
- [288] F. Shalit, B. Sredni, L. Stern, E. Kott, and M. Huberman, "Elevated interleukin-6 secretion levels by mononuclear cells of Alzheimer's patients," *Neuroscience Letters*, vol. 174, no. 2, pp. 130–132, 1994.
- [289] Y. X. Sun, L. Minthon, A. Wallmark, S. Warkentin, K. Blennow, and S. Janciauskiene, "Inflammatory markers in matched plasma and cerebrospinal fluid from patients with Alzheimer's disease," *Dementia and Geriatric Cognitive Disorders*, vol. 16, no. 3, pp. 136–144, 2003.
- [290] P. Kraggsbjerg, T. Vikerfors, and H. Holmberg, "Cytokine responses in patients with pneumonia caused by Chlamydia or Mycoplasma," *Respiration*, vol. 65, no. 4, pp. 299–303, 1998.
- [291] N. Mehmet, M. Refik, M. Harputluoglu, Y. Ersoy, N. E. Aydin, and B. Yildirim, "Serum and gastric fluid levels of cytokines and nitrates in gastric diseases infected with Helicobacter pylori," *New Microbiologica*, vol. 27, no. 2, pp. 139–148, 2004.
- [292] D. Delfino, L. Cianci, E. Lupis et al., "Interleukin-6 production by human monocytes stimulated with Cryptococcus neoformans components," *Infection and Immunity*, vol. 65, no. 6, pp. 2454–2456, 1997.
- [293] S. Noisakran, W. P. Halford, L. Veress, and D. J. Carr, "Role of the hypothalamic pituitary adrenal axis and IL-6 in stress-induced reactivation of latent herpes simplex virus type 1," *Journal of Immunology*, vol. 160, no. 11, pp. 5441–5447, 1998.
- [294] K. L. Davis, B. M. Davis, B. S. Greenwald et al., "Cortisol and Alzheimer's disease. I: basal studies," *American Journal of Psychiatry*, vol. 143, no. 3, pp. 300–305, 1986.
- [295] C. W. Huang, C. C. Lui, W. N. Chang, C. H. Lu, Y. L. Wang, and C. C. Chang, "Elevated basal cortisol level predicts lower hippocampal volume and cognitive decline in Alzheimer's disease," *Journal of Clinical Neuroscience*, vol. 16, no. 10, pp. 1283–1286, 2009.
- [296] A. Rodriguez, D. L. M. Sainz, J. Missry, and C. S. Foster, "The role of cyclic nucleotide mediators in latency and reactivation of HSV-1 infected neuroblastoma cells," *Eye*, vol. 5, no. 5, pp. 627–635, 1991.
- [297] R. L. Smith, L. I. Pizer, E. M. Johnson Jr., and C. L. Wilcox, "Activation of second-messenger pathways reactivates latent herpes simplex virus in neuronal cultures," *Virology*, vol. 188, no. 1, pp. 311–318, 1992.
- [298] B. M. Gebhardt and H. E. Kaufman, "Propranolol suppresses reactivation of herpesvirus," *Antiviral Research*, vol. 27, no. 3, pp. 255–261, 1995.
- [299] B. Sainz, J. M. Loutsch, M. E. Marquart, and J. M. Hill, "Stress-associated immunomodulation and herpes simplex virus infections," *Medical Hypotheses*, vol. 56, no. 3, pp. 348–356, 2001.
- [300] R. E. Martin, J. M. Loutsch, H. H. Garza, D. J. Boedeker, and J. M. Hill, "Iontophoresis of lysophosphatidic acid into rabbit cornea induces HSV-1 reactivation: evidence that neuronal signaling changes after infection," *Molecular Vision*, vol. 5, p. 36, 1999.
- [301] M. K. Aghi, T. C. Liu, S. Rabkin, and R. L. Martuza, "Hypoxia enhances the replication of oncolytic herpes simplex virus," *Molecular Therapy*, vol. 17, no. 1, pp. 51–56, 2009.
- [302] Y. Hashimoto, H. Kawatsura, Y. Shiga, S. Furukawa, and T. Shigeno, "Significance of nerve growth factor content levels after transient forebrain ischemia in gerbils," *Neuroscience Letters*, vol. 139, no. 1, pp. 45–46, 1992.
- [303] J. C. de la Torre, "The vascular hypothesis of Alzheimer's disease: bench to bedside and beyond," *Neurodegenerative Diseases*, vol. 7, no. 1–3, pp. 116–121, 2010.
- [304] C. E. Isaacs, R. Kascak, R. K. Pullarkat, W. Xu, and K. Schneidman, "Inhibition of herpes simplex virus replication by retinoic acid," *Antiviral Research*, vol. 33, no. 2, pp. 117–127, 1997.
- [305] E. D. Toffanello, E. M. Inelmen, N. Minicuci et al., "Ten-year trends in vitamin intake in free-living healthy elderly people: the risk of subclinical malnutrition," *Journal of Nutrition, Health and Aging*, vol. 15, pp. 99–103, 2010.
- [306] F. J. Jimenez-Jimenez, J. A. Molina, F. de Bustos et al., "Serum levels of β -carotene, α -carotene and vitamin A in patients with Alzheimer's disease," *European Journal of Neurology*, vol. 6, pp. 495–497, 1999.
- [307] D. N. Frank, W. Zhu, R. B. Sartor, and E. Li, "Investigating the biological and clinical significance of human dysbioses," *Trends in Microbiology*, vol. 19, no. 9, pp. 427–434, 2011.
- [308] B. Wolozin, W. Kellman, P. Ruosseau, G. G. Celesia, and G. Siegel, "Decreased prevalence of Alzheimer disease associated with 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors," *Archives of Neurology*, vol. 57, no. 10, pp. 1439–1443, 2000.
- [309] M. C. Morris, D. A. Evans, J. L. Bienias et al., "Consumption of fish and n-3 fatty acids and risk of incident Alzheimer disease," *Archives of Neurology*, vol. 60, no. 7, pp. 940–946, 2003.
- [310] L. B. Lopez, D. Kritz-Silverstein, and C. E. Barrett-Connor, "High dietary and plasma levels of the omega-3 fatty acid docosahexaenoic acid are associated with decreased dementia risk: the Rancho Bernardo study," *Journal of Nutrition, Health and Aging*, vol. 15, pp. 25–31, 2010.
- [311] V. Solfrizzi, F. Panza, and A. Capurso, "The role of diet in cognitive decline," *Journal of Neural Transmission*, vol. 110, no. 1, pp. 95–110, 2003.
- [312] P. L. McGeer, M. Schulzer, and E. G. McGeer, "Arthritis and anti-inflammatory agents as possible protective factors for Alzheimer's disease: a review of 17 epidemiologic studies," *Neurology*, vol. 47, no. 2, pp. 425–432, 1996.

- [313] D. A. Snowdon, "Healthy aging and dementia: findings from the nun study," *Annals of Internal Medicine*, vol. 139, no. 5, pp. 450–454, 2003.
- [314] A. D. Smith, S. M. Smith, C. A. de Jager et al., "Homocysteine-lowering by b vitamins slows the rate of accelerated brain atrophy in mild cognitive impairment: a randomized controlled trial," *PLoS ONE*, vol. 5, no. 9, Article ID e12244, pp. 1–10, 2010.
- [315] M. L. Daviglus, B. L. Plassman, A. Pirzada et al., "Risk factors and preventive interventions for Alzheimer disease: state of the science," *Archives of Neurology*, vol. 68, no. 9, pp. 1185–1190, 2011.
- [316] X. Song, A. Mitnitski, and K. Rockwood, "Nontraditional risk factors combine to predict Alzheimer disease and dementia," *Neurology*, vol. 77, no. 3, pp. 227–234, 2011.
- [317] D. E. Barnes and K. Yaffe, "The projected effect of risk factor reduction on Alzheimer's disease prevalence," *The Lancet Neurology*, vol. 10, no. 9, pp. 819–828, 2011.
- [318] H. B. Larman, Z. Zhao, U. Laserson et al., "Autoantigen discovery with a synthetic human peptidome," *Nature Biotechnology*, vol. 29, no. 6, pp. 535–541, 2011.
- [319] J. Miklosy, "Alzheimer's disease—a neurospirochetosis. Analysis of the evidence following Koch's and Hill's criteria," *Journal of Neuroinflammation*, vol. 8, pp. 90–96, 2011.
- [320] E. Nagele, M. Han, C. Demarshall, B. Belinka, and R. Nagele, "Diagnosis of Alzheimer's disease based on disease-specific autoantibody profiles in human sera," *PLoS ONE*, vol. 6, no. 8, Article ID e23112, 2011.
- [321] L. Restrepo, P. Stafford, D. M. Magee, and S. A. Johnston, "Application of immunosignatures to the assessment of Alzheimer's disease," *Annals of Neurology*, vol. 70, no. 2, pp. 286–295, 2011.