Research Article

Uniqueness of Meromorphic Functions Sharing Fixed Point

Subhas S. Bhoosnurmath and Veena L. Pujari

Department of Mathematics, Karnatak University, Dharwad 580003, India

Correspondence should be addressed to Subhas S. Bhoosnurmath; ssbmath@gmail.com

Received 28 September 2012; Revised 22 January 2013; Accepted 30 January 2013

Academic Editor: Alexandre Timonov

Copyright © 2013 S. S. Bhoosnurmath and V. L. Pujari. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study the uniqueness of meromorphic functions concerning differential polynomials sharing fixed point and obtain some significant results, which improve the results due to Lin and Yi (2004).

1. Introduction and Main Results

Let $f(z)$ be a nonconstant meromorphic function in the whole complex plane $\mathbb{C}$. We will use the following standard notations of value distribution theory: $T(r, f)$, $m(r, f)$, $N(r, f)$, $\overline{N}(r, f)$, $\dots$ (see [1, 2]). We denote by $S(r, f)$ any function satisfying

$$S(r, f) = o\{T(r, f)\}, \quad \text{as } r \rightarrow +\infty, \quad (1)$$

possibly outside of a set with finite measure.

Let a be a finite complex number and k a positive integer. We denote by $N_k(r, 1/(f-a))$ the counting function for the zeros of $f(z) - a$ in $|z| \leq r$ with multiplicity $\leq k$ and by $\overline{N}_k(r, 1/(f-a))$ the corresponding one for which multiplicity is not counted. Let $N_{(k)}(r, 1/(f-a))$ be the counting function for the zeros of $f(z) - a$ in $|z| \leq r$ with multiplicity $\geq k$ and $\overline{N}_{(k)}(r, 1/(f-a))$ the corresponding one for which multiplicity is not counted. Set

$$N_k\left(r, \frac{1}{f-a}\right) = \overline{N}\left(r, \frac{1}{f-a}\right) + \overline{N}_{(2)}\left(r, \frac{1}{f-a}\right) + \dots + \overline{N}_{(k)}\left(r, \frac{1}{f-a}\right). \quad (2)$$

Let $g(z)$ be a nonconstant meromorphic function. We denote by $\overline{N}_L(r, 1/(f-a))$ the counting function for a -points of both $f(z)$ and $g(z)$ about which $f(z)$ has larger multiplicity than $g(z)$, where multiplicity is not counted. Similarly, we have notation $\overline{N}_L(r, 1/(g-a))$.

We say that f and g share a CM (counting multiplicity) if $f-a$ and $g-a$ have same zeros with the same multiplicities. Similarly, we say that f and g share a IM (ignoring multiplicity) if $f-a$ and $g-a$ have same zeros with ignoring multiplicities.

In 2004, Lin and Yi [3] obtained the following results.

Theorem A. *Let f and g be two transcendental meromorphic functions, $n \geq 12$ an integer. If $f^n(f-1)f'$ and $g^n(g-1)g'$ share z CM, then either $f(z) \equiv g(z)$ or*

$$g = \frac{(n+2)(1-h^{n+1})}{(n+1)(1-h^{n+2})}, \quad f = \frac{(n+2)h(1-h^{n+1})}{(n+1)(1-h^{n+2})}, \quad (3)$$

where h is a nonconstant meromorphic function.

Theorem B. *Let f and g be two transcendental meromorphic functions, $n \geq 13$ an integer. If $f^n(f-1)^2 f'$ and $g^n(g-1)^2 g'$ share z CM, then $f(z) \equiv g(z)$.*

In this paper, we study the uniqueness problems of entire or meromorphic functions concerning differential polynomials sharing fixed point, which improves Theorems A and B.

1.1. Main Results

Theorem 1. *Let f and g be two nonconstant meromorphic functions, $n \geq 11$ a positive integer. If $f^n(f-1)f'$ and*

$g^n(g-1)g'$ share z CM, f and g share ∞ IM, then either $f(z) \equiv g(z)$ or

$$g = \frac{(n+2)(1-h^{n+1})}{(n+1)(1-h^{n+2})}, \quad f = \frac{(n+2)h(1-h^{n+1})}{(n+1)(1-h^{n+2})}, \quad (4)$$

where h is a nonconstant meromorphic function.

Theorem 2. Let f and g be two nonconstant meromorphic functions, $n \geq 12$ a positive integer. If $f^n(f-1)^2f'$ and $g^n(g-1)^2g'$ share z CM, f and g share ∞ IM, then $f(z) \equiv g(z)$.

Theorem 3. Let f and g be two nonconstant entire functions, $n \geq 7$ an integer. If $f^n(f-1)f'$ and $g^n(g-1)g'$ share z CM, then $f(z) \equiv g(z)$.

2. Some Lemmas

Lemma 4 (see [4]). Let f_1, f_2 , and f_3 be nonconstant meromorphic functions such that $f_1 + f_2 + f_3 = 1$. If f_1, f_2 , and f_3 are linearly independent, then

$$T(r, f_1) < \sum_{i=1}^3 N_2\left(r, \frac{1}{f_i}\right) + \sum_{i=1}^3 \bar{N}(r, f_i) + o(T(r)), \quad (5)$$

where $T(r) = \max_{1 \leq i \leq 3} \{T(r, f_i)\}$ and $r \notin E$.

Lemma 5 (see [1]). Let f_1 and f_2 be two nonconstant meromorphic functions. If $c_1f_1 + c_2f_2 = c_3$, where c_1, c_2 , and c_3 are non-zero constants, then

$$T(r, f_1) \leq \bar{N}(r, f_1) + \bar{N}\left(r, \frac{1}{f_1}\right) + \bar{N}\left(r, \frac{1}{f_2}\right) + S(r, f_1). \quad (6)$$

Lemmas 4 and 5 play a very important role in proving our theorems.

Lemma 6 (see [1]). Let f be a nonconstant meromorphic function and let k be a nonnegative integer, then

$$N\left(r, \frac{1}{f^{(k)}}\right) \leq N\left(r, \frac{1}{f}\right) + k\bar{N}(r, f) + S(r, f). \quad (7)$$

The following lemmas play a cardinal role in proving our results.

Lemma 7. Let f and g be nonconstant meromorphic functions. If $f^n(f-1)f'$ and $g^n(g-1)g'$ share z CM and $n > 6$, then

$$T(r, g) \leq \left(\frac{n+3}{n-6}\right)T(r, f) + \log r + S(r, g). \quad (8)$$

Proof. Applying Nevanlinna's second fundamental theorem to $g^n(g-1)g'$, we have

$$\begin{aligned} T(r, g^n(g-1)g') &\leq \bar{N}\left(r, g^n(g-1)g'\right) \\ &\quad + \bar{N}\left(r, \frac{1}{g^n(g-1)g'}\right) \\ &\quad + \bar{N}\left(r, \frac{1}{g^n(g-1)g'-z}\right) + S(r, g) \\ &\leq \bar{N}(r, g) + \bar{N}\left(r, \frac{1}{g}\right) \\ &\quad + \bar{N}\left(r, \frac{1}{g-1}\right) + \bar{N}\left(r, \frac{1}{g'}\right) \\ &\quad + \bar{N}\left(r, \frac{1}{f^n(f-1)f'-z}\right) + S(r, g). \end{aligned} \quad (9)$$

By the first fundamental theorem and (9), we have

$$\begin{aligned} (n+1)T(r, g) &\leq T(r, g^n(g-1)g') + S(r, g) \\ &\leq T(r, g^n(g-1)g') + T\left(r, \frac{1}{g'}\right) + S(r, g) \\ &\leq \bar{N}(r, g) + \bar{N}\left(r, \frac{1}{g}\right) \\ &\quad + \bar{N}\left(r, \frac{1}{g-1}\right) + \bar{N}\left(r, \frac{1}{g'}\right) \\ &\quad + \bar{N}\left(r, \frac{1}{f^n(f-1)f'-z}\right) \\ &\quad + T(r, g') + S(r, g). \end{aligned} \quad (10)$$

We know that

$$\begin{aligned} \bar{N}\left(r, \frac{1}{f^n(f-1)f'-z}\right) &\leq T\left(r, \frac{1}{f^n(f-1)f'-z}\right) \\ &= T(r, f^n(f-1)f'-z) + O(1) \\ &\leq T(r, f^n(f-1)f') \\ &\quad + T(r, z) + O(1) \\ &\leq nT(r, f) + T(r, f-1) \\ &\quad + T(r, f') + \log r + O(1) \\ &\leq (n+3)T(r, f) + \log r + O(1). \end{aligned} \quad (11)$$

Therefore using Lemma 6, (10) becomes

$$\begin{aligned}
 (n+1)T(r, g) &\leq \bar{N}(r, g) + \bar{N}\left(r, \frac{1}{g}\right) \\
 &\quad + \bar{N}\left(r, \frac{1}{g-1}\right) + \bar{N}\left(r, \frac{1}{g'}\right) \\
 &\quad + (n+3)T(r, f) + T(r, g') \\
 &\quad + \log r + S(r, g) \\
 &\leq 7T(r, g) + (n+3)T(r, f) + \log r + S(r, g)
 \end{aligned} \tag{12}$$

$$\implies (n-6)T(r, g) \leq (n+3)T(r, f) + \log r + S(r, g); \tag{13}$$

since $n > 6$, we have

$$T(r, g) \leq \left(\frac{n+3}{n-6}\right)T(r, f) + \log r + S(r, g) \tag{14}$$

This completes the proof of Lemma 7. $\square$

Lemma 8. Let f and g be nonconstant entire functions. If $f^n(f-1)f'$ and $g^n(g-1)g'$ share z CM and $n > 3$, then

$$T(r, g) \leq \left(\frac{n+2}{n-3}\right)T(r, f) + \log r + S(r, g). \tag{15}$$

Proof. Applying Nevanlinna's second fundamental theorem to $g^n(g-1)g'$, we have

$$\begin{aligned}
 T(r, g^n(g-1)g') &\leq \bar{N}(r, g^n(g-1)g') \\
 &\quad + \bar{N}\left(r, \frac{1}{g^n(g-1)g'}\right) \\
 &\quad + \bar{N}\left(r, \frac{1}{g^n(g-1)g' - z}\right) + S(r, g) \\
 &\leq \bar{N}(r, g) + \bar{N}\left(r, \frac{1}{g}\right) \\
 &\quad + \bar{N}\left(r, \frac{1}{g-1}\right) + \bar{N}\left(r, \frac{1}{g'}\right) \\
 &\quad + \bar{N}\left(r, \frac{1}{f^n(f-1)f' - z}\right) + S(r, g).
 \end{aligned} \tag{16}$$

Since g is an entire function, we have $\bar{N}(r, g) = 0$ and the above equation becomes

$$\begin{aligned}
 T(r, g^n(g-1)g') &\leq \bar{N}\left(r, \frac{1}{g}\right) + \bar{N}\left(r, \frac{1}{g-1}\right) + \bar{N}\left(r, \frac{1}{g'}\right) \\
 &\quad + \bar{N}\left(r, \frac{1}{f^n(f-1)f' - z}\right) + S(r, g).
 \end{aligned} \tag{17}$$

By the first fundamental theorem and (17), we have

$$\begin{aligned}
 (n+1)T(r, g) &\leq T(r, g^n(g-1)g') + S(r, g) \\
 &\leq T(r, g^n(g-1)g') + T\left(r, \frac{1}{g'}\right) + S(r, g) \\
 &\leq \bar{N}\left(r, \frac{1}{g}\right) + \bar{N}\left(r, \frac{1}{g-1}\right) + \bar{N}\left(r, \frac{1}{g'}\right) \\
 &\quad + \bar{N}\left(r, \frac{1}{f^n(f-1)f' - z}\right) \\
 &\quad + T(r, g') + S(r, g).
 \end{aligned} \tag{18}$$

We know that

$$\begin{aligned}
 \bar{N}\left(r, \frac{1}{f^n(f-1)f' - z}\right) &\leq T\left(r, \frac{1}{f^n(f-1)f' - z}\right) \\
 &= T(r, f^n(f-1)f' - z) + O(1) \\
 &\leq T(r, f^n(f-1)f') \\
 &\quad + T(r, z) + O(1) \\
 &\leq nT(r, f) + T(r, f-1) \\
 &\quad + T(r, f') + \log r + O(1) \\
 &\leq (n+2)T(r, f) + \log r + O(1).
 \end{aligned} \tag{19}$$

Therefore using Lemma 6, (18) becomes

$$\begin{aligned}
 (n+1)T(r, g) &\leq \bar{N}\left(r, \frac{1}{g}\right) + \bar{N}\left(r, \frac{1}{g-1}\right) + \bar{N}\left(r, \frac{1}{g'}\right) \\
 &\quad + (n+2)T(r, f) + T(r, g') \\
 &\quad + \log r + S(r, g) \\
 &\leq 4T(r, g) + (n+2)T(r, f) + \log r + S(r, g)
 \end{aligned} \tag{20}$$

or

$$(n-3)T(r, g) \leq (n+2)T(r, f) + \log r + S(r, g); \tag{21}$$

since $n > 3$, we have

$$T(r, g) \leq \left(\frac{n+2}{n-3}\right)T(r, f) + \log r + S(r, g). \tag{22}$$

This completes the proof of Lemma 8. $\square$

Lemma 9 (see [5]). Suppose that $f(z)$ is a meromorphic function in the complex plane and $P(f) = a_0 f^n + a_1 f^{n-1} + \dots + a_n$, where $a_0 (\neq 0)$, $a_1, \dots, a_n$ are small meromorphic functions of $f(z)$. Then

$$T(r, P(f)) = nT(r, f) + S(r, f). \tag{23}$$

Lemma 10 (see [6]). Let f_1, f_2 , and f_3 be three meromorphic functions satisfying $\sum_{j=1}^3 f_j = 1$, let $g_1 = -f_3/f_2$, $g_2 = 1/f_2$, and $g_3 = -f_1/f_2$. If f_1, f_2 , and f_3 are linearly independent, then g_1, g_2 , and g_3 are linearly independent.

3. Proof of Theorems

Proof of Theorem 1. By assumption, $f^n(f-1)f'$ and $g^n(g-1)g'$ share z CM, and f and g share ∞ IM. Let

$$H = \frac{f^n(f-1)f' - z}{g^n(g-1)g' - z}. \quad (24)$$

Then, H is a meromorphic function satisfying

$$\begin{aligned} T(r, H) &= T\left(r, \frac{f^n(f-1)f' - z}{g^n(g-1)g' - z}\right) \\ &\leq T(r, f^n(f-1)f' - z) \\ &\quad + T(r, g^n(g-1)g' - z) + O(1) \\ &\leq (n+3)(T(r, f) + T(r, g)) + O(\log r). \end{aligned} \quad (25)$$

Therefore,

$$T(r, H) = O(T(r, f) + T(r, g)). \quad (26)$$

From (24), we easily see that the zeros and poles of H are multiple and satisfy

$$\overline{N}(r, H) \leq \overline{N}_L(r, f), \quad \overline{N}\left(r, \frac{1}{H}\right) \leq \overline{N}_L(r, g). \quad (27)$$

Let

$$\begin{aligned} f_1 &= \frac{f^n(f-1)f'}{z}, & f_2 &= H, \\ f_3 &= \frac{-Hg^n(g-1)g'}{z}. \end{aligned} \quad (28)$$

Then, $f_1 + f_2 + f_3 = 1$ and $T(r)$ denote the maximum of $T(r, f_j)$, $j = 1, 2, 3$.

We have

$$\begin{aligned} T(r, f_1) &= O(T(r, f)), \\ T(r, f_2) &= O(T(r, f) + T(r, g)), \\ T(r, f_3) &= O(T(r, f) + T(r, g)). \end{aligned} \quad (29)$$

Therefore, $T(r) = O(T(r, f) + T(r, g))$, and thus

$$S(r, f) + S(r, g) = o(T(r)). \quad (30)$$

Now, we discuss the following three cases.

Case 1. Suppose that neither f_2 nor f_3 is a constant.

If f_1, f_2 , and f_3 are linearly independent, then by Lemma 4 and (28), we have

$$\begin{aligned} T(r, f_1) &< \sum_{i=1}^3 N_2\left(r, \frac{1}{f_i}\right) + \sum_{i=1}^3 \overline{N}(r, f_i) + o(T(r)) \\ &\leq N_2\left(r, \frac{1}{f_1}\right) + N_2\left(r, \frac{1}{f_2}\right) + N_2\left(r, \frac{1}{f_3}\right) \\ &\quad + \overline{N}(r, f_1) + \overline{N}(r, f_2) + \overline{N}(r, f_3) + o(T(r)) \\ &\leq N_2\left(r, \frac{z}{f^n(f-1)f'}\right) + N_2\left(r, \frac{1}{H}\right) \\ &\quad + N_2\left(r, \frac{z}{Hg^n(g-1)g'}\right) \\ &\quad + \overline{N}\left(r, \frac{f^n(f-1)f'}{z}\right) \\ &\quad + \overline{N}(r, H) + \overline{N}\left(r, \frac{Hg^n(g-1)g'}{z}\right) + o(T(r)) \\ &= N_2\left(r, \frac{1}{f^n(f-1)f'}\right) + N_2\left(r, \frac{1}{H}\right) \\ &\quad + N_2\left(r, \frac{1}{Hg^n(g-1)g'}\right) \\ &\quad + \overline{N}(r, f^n(f-1)f') \\ &\quad + \overline{N}(r, H) + \overline{N}(r, Hg^n(g-1)g') \\ &\quad + 2 \log r + o(T(r)). \end{aligned} \quad (31)$$

Using (27), we note that

$$\begin{aligned} &N_2\left(r, \frac{1}{Hg^n(g-1)g'}\right) \\ &\leq N_2\left(r, \frac{1}{H}\right) + N_2\left(r, \frac{1}{g^n(g-1)g'}\right) \\ &\leq 2\overline{N}\left(r, \frac{1}{H}\right) + N_2\left(r, \frac{1}{g^n(g-1)g'}\right) \\ &\leq 2\overline{N}_L(r, g) + N_2\left(r, \frac{1}{g^n(g-1)g'}\right). \end{aligned} \quad (32)$$

Since $\overline{N}_L(r, g) = 0$, we obtain that

$$N_2\left(r, \frac{1}{Hg^n(g-1)g'}\right) \leq N_2\left(r, \frac{1}{g^n(g-1)g'}\right), \quad (33)$$

$$\begin{aligned} \overline{N}(r, Hg^n(g-1)g') &\leq \overline{N}(r, H) + \overline{N}(r, g^n(g-1)g') \\ &\leq \overline{N}_L(r, f) + \overline{N}(r, g). \end{aligned} \quad (34)$$

But $\overline{N}_L(r, f) = 0$, so we get

$$\overline{N}\left(r, Hg^n(g-1)g'\right) \leq \overline{N}(r, g). \quad (35)$$

Using (33) and (35) in (31), we get

$$\begin{aligned} T(r, f_1) &\leq N_2\left(r, \frac{1}{f^n(f-1)f'}\right) + N_2\left(r, \frac{1}{H}\right) \\ &+ N_2\left(r, \frac{1}{g^n(g-1)g'}\right) + \overline{N}(r, f) \\ &+ \overline{N}(r, H) + \overline{N}(r, g) + 2\log r + o(T(r)). \end{aligned} \quad (36)$$

Since f and g share ∞ IM, we have $\overline{N}(r, f) = \overline{N}(r, g)$.
Using this with (27), we get

$$\begin{aligned} T(r, f_1) &\leq N_2\left(r, \frac{1}{f^n(f-1)f'}\right) + 2\overline{N}_L(r, g) \\ &+ N_2\left(r, \frac{1}{g^n(g-1)g'}\right) + \overline{N}_L(r, f) \\ &+ 2\overline{N}(r, f) + 2\log r + o(T(r)) \\ &\leq N\left(r, \frac{1}{f^n(f-1)f'}\right) \\ &- \left[N_{(3)}\left(r, \frac{1}{f^n(f-1)f'}\right) \right. \\ &\quad \left. - 2\overline{N}_{(3)}\left(r, \frac{1}{f^n(f-1)f'}\right) \right] \\ &+ N\left(r, \frac{1}{g^n(g-1)g'}\right) \\ &- \left[N_{(3)}\left(r, \frac{1}{g^n(g-1)g'}\right) \right. \\ &\quad \left. - 2\overline{N}_{(3)}\left(r, \frac{1}{g^n(g-1)g'}\right) \right] \\ &+ 2\overline{N}_L(r, g) + \overline{N}_L(r, f) + 2\overline{N}(r, f) \\ &+ 2\log r + o(T(r)). \end{aligned} \quad (37)$$

If z_0 is a zero of f with multiplicity p , then z_0 is a zero of $f^n(f-1)f'$ with multiplicity $np + p - 1 \geq 3$; we have

$$\begin{aligned} &\left[N_{(3)}\left(r, \frac{1}{f^n(f-1)f'}\right) - 2\overline{N}_{(3)}\left(r, \frac{1}{f^n(f-1)f'}\right) \right] \\ &\geq (n-2)N\left(r, \frac{1}{f}\right). \end{aligned} \quad (38)$$

Similarly,

$$\begin{aligned} &\left[N_{(3)}\left(r, \frac{1}{g^n(g-1)g'}\right) - 2\overline{N}_{(3)}\left(r, \frac{1}{g^n(g-1)g'}\right) \right] \\ &\geq (n-2)N\left(r, \frac{1}{g}\right). \end{aligned} \quad (39)$$

Let

$$f_1^* = \frac{f^{n+2}}{n+2} - \frac{f^{n+1}}{n+1}. \quad (40)$$

By Lemma 9, we have $T(r, f_1^*) = (n+2)T(r, f) + S(r, f)$.
Since $(f_1^*)' = zf_1$, we have

$$\begin{aligned} m\left(r, \frac{1}{f_1^*}\right) &\leq m\left(r, \frac{1}{zf_1}\right) + m\left(r, \frac{(f_1^*)'}{f_1^*}\right) \\ &\leq m\left(r, \frac{1}{f_1}\right) + \log r + S(r, f). \end{aligned} \quad (41)$$

By the first fundamental theorem, we have

$$\begin{aligned} T(r, f_1^*) &\leq T(r, f_1) + N\left(r, \frac{1}{f_1^*}\right) - N\left(r, \frac{1}{f_1}\right) \\ &+ \log r + S(r, f); \end{aligned} \quad (42)$$

we have

$$\begin{aligned} &N\left(r, \frac{1}{f_1^*}\right) \\ &= (n+1)N\left(r, \frac{1}{f}\right) + N\left(r, \frac{1}{f - (n+2)/(n+1)}\right). \end{aligned} \quad (43)$$

From (37)–(43), we get

$$\begin{aligned} T(r, f_1^*) &\leq N\left(r, \frac{1}{f^n(f-1)f'}\right) - (n-2)N\left(r, \frac{1}{f}\right) \\ &+ N\left(r, \frac{1}{g^n(g-1)g'}\right) - (n-2)N\left(r, \frac{1}{g}\right) \\ &+ 2\overline{N}_L(r, g) \\ &+ \overline{N}_L(r, f) + 2\overline{N}(r, f) + (n+1)N\left(r, \frac{1}{f}\right) \\ &+ N\left(r, \frac{1}{f - (n+2)/(n+1)}\right) \\ &- N\left(r, \frac{1}{f^n(f-1)f'}\right) + 3\log r + o(T(r)). \end{aligned} \quad (44)$$

Using Lemma 6, we get

$$\begin{aligned}
 T(r, f_1^*) &\leq 3N\left(r, \frac{1}{f}\right) + 3N\left(r, \frac{1}{g}\right) \\
 &\quad + 2\bar{N}(r, f) + \bar{N}(r, g) + 2\bar{N}_L(r, g) \\
 &\quad + \bar{N}_L(r, f) + N\left(r, \frac{1}{g-1}\right) \\
 &\quad + N\left(r, \frac{1}{f-(n+2)/(n+1)}\right) \\
 &\quad + 3\log r + o(T(r)) \\
 (n+2)T(r, f) &\leq 3N\left(r, \frac{1}{f}\right) + 3N\left(r, \frac{1}{g}\right) \\
 &\quad + 2\bar{N}(r, f) + \bar{N}(r, g) \\
 &\quad + 2\bar{N}_L(r, g) + \bar{N}_L(r, f) + N\left(r, \frac{1}{g-1}\right) \\
 &\quad + N\left(r, \frac{1}{f-(n+2)/(n+1)}\right) \\
 &\quad + 3\log r + o(T(r)).
 \end{aligned} \tag{45}$$

Let

$$\begin{aligned}
 g_1 &= -\frac{f_3}{f_2} = \frac{g^n(g-1)g'}{z}, \\
 g_2 &= \frac{1}{f_2} = \frac{1}{H}, \quad g_3 = -\frac{f_1}{f_2} = \frac{f^n(f-1)f'}{zH}.
 \end{aligned} \tag{46}$$

Then $g_1 + g_2 + g_3 = 1$. By Lemma 10, g_1, g_2 , and g_3 are linearly independent. In the same manner as above, we get

$$\begin{aligned}
 (n+2)T(r, g) &\leq 3N\left(r, \frac{1}{g}\right) + 3N\left(r, \frac{1}{f}\right) \\
 &\quad + 2\bar{N}(r, g) + \bar{N}(r, f) + 2\bar{N}_L(r, f) \\
 &\quad + \bar{N}_L(r, g) + N\left(r, \frac{1}{f-1}\right) \\
 &\quad + N\left(r, \frac{1}{g-(n+2)/(n+1)}\right) \\
 &\quad + 3\log r + o(T(r)).
 \end{aligned} \tag{47}$$

Note that

$$\bar{N}_L(r, f) + \bar{N}_L(r, g) \leq \bar{N}(r, f) = \bar{N}(r, g). \tag{48}$$

Adding (45) and (47) gives

$$\begin{aligned}
 (n+2)(T(r, f) + T(r, g)) &\leq 6\left(N\left(r, \frac{1}{f}\right) + N\left(r, \frac{1}{g}\right)\right) + 3(\bar{N}(r, f) + \bar{N}(r, g)) \\
 &\quad + (\bar{N}_L(r, f) + \bar{N}_L(r, g)) \\
 &\quad + \left(N\left(r, \frac{1}{f-1}\right) + N\left(r, \frac{1}{g-1}\right)\right) \\
 &\quad + N\left(r, \frac{1}{f-(n+2)/(n+1)}\right) \\
 &\quad + N\left(r, \frac{1}{g-(n+2)/(n+1)}\right) + 6\log r + o(T(r)) \\
 (n-6)(T(r, f) + T(r, g)) &\leq 3(\bar{N}(r, f) + \bar{N}(r, g)) \\
 &\quad + 3(\bar{N}_L(r, f) + \bar{N}_L(r, g)) + 6\log r + o(T(r)).
 \end{aligned} \tag{49}$$

Using (48), we get

$$\begin{aligned}
 (n-6)(T(r, f) + T(r, g)) &\leq 3\bar{N}(r, f) + 6\bar{N}(r, g) + 6\log r + o(T(r))
 \end{aligned} \tag{50}$$

or

$$\begin{aligned}
 (n-6)(T(r, f) + T(r, g)) &\leq 6\bar{N}(r, f) + 3\bar{N}(r, g) + 6\log r + o(T(r)).
 \end{aligned} \tag{51}$$

Combining (50) and (51), we get

$$\begin{aligned}
 (n-6)(T(r, f) + T(r, g)) &\leq \frac{9}{2}(\bar{N}(r, f) + \bar{N}(r, g)) \\
 &\quad + 6\log r + o(T(r)) \\
 \left(n - \frac{21}{2}\right)(T(r, f) + T(r, g)) &\leq 6\log r + o(T(r)).
 \end{aligned} \tag{52}$$

By $n \geq 11$ and (30), we get a contradiction. Thus f_1, f_2 , and f_3 are linearly dependent. Then, there exists three constants $(c_1, c_2, c_3) \neq (0, 0, 0)$ such that

$$c_1 f_1 + c_2 f_2 + c_3 f_3 = 0. \tag{53}$$

If $c_1 = 0$, from (53) $c_2 \neq 0, c_3 \neq 0$, and

$$f_3 = -\frac{c_2}{c_3} f_2; \tag{54}$$

$$\implies g^n(g-1)g' = \frac{c_2}{c_3} z. \tag{55}$$

On integrating, we get

$$\begin{aligned} \frac{g^{n+2}}{n+2} - \frac{g^{n+1}}{n+1} &= \frac{c_2}{c_3} \frac{z^2}{2} + k, \quad k \text{ is a constant,} \\ T\left(r, \frac{g^{n+2}}{n+2} - \frac{g^{n+1}}{n+1}\right) &\leq T(r, z^2) + O(1), \\ (n+2)T(r, g) &\leq 2 \log r + O(1); \end{aligned} \quad (56)$$

since $n \geq 11$, we get a contradiction.

Thus $c_1 \neq 0$, and by (53) we have

$$\begin{aligned} c_1 f_1 &= -c_2 f_2 - c_3 f_3 \\ \Rightarrow f_1 &= -\frac{c_2}{c_1} f_2 - \frac{c_3}{c_1} f_3. \end{aligned} \quad (57)$$

Substituting this in $f_1 + f_2 + f_3 = 1$, we get

$$-\frac{c_2}{c_1} f_2 - \frac{c_3}{c_1} f_3 + f_2 + f_3 = 1; \quad (58)$$

that is,

$$\left(1 - \frac{c_2}{c_1}\right) f_2 + \left(1 - \frac{c_3}{c_1}\right) f_3 = 1, \quad \text{where } c_1 \neq c_3, \quad c_2 \neq c_3. \quad (59)$$

From (28), we obtain

$$\begin{aligned} \left(1 - \frac{c_2}{c_1}\right) H + \left(1 - \frac{c_3}{c_1}\right) \left(\frac{-H g^n (g-1) g'}{z}\right) &= 1 \\ \Rightarrow \left(1 - \frac{c_2}{c_1}\right) - \left(1 - \frac{c_3}{c_1}\right) \left(\frac{g^n (g-1) g'}{z}\right) &= 1 \\ \Rightarrow \left(1 - \frac{c_3}{c_1}\right) \left(\frac{g^n (g-1) g'}{z}\right) + \frac{1}{H} &= \left(1 - \frac{c_2}{c_1}\right). \end{aligned} \quad (60)$$

Applying Lemma 5 to the above equation, we get

$$\begin{aligned} T\left(r, \frac{g^n (g-1) g'}{z}\right) &\leq \bar{N}\left(r, \frac{g^n (g-1) g'}{z}\right) \\ &\quad + \bar{N}\left(r, \frac{z}{g^n (g-1) g'}\right) \\ &\quad + \bar{N}(r, H) + S(r, g). \end{aligned} \quad (61)$$

Note that

$$\begin{aligned} T(r, g^n (g-1) g') &\leq T\left(r, \frac{g^n (g-1) g'}{z}\right) + T(r, z) \\ &\leq T\left(r, \frac{g^n (g-1) g'}{z}\right) + \log r. \end{aligned} \quad (62)$$

Using (61), we get

$$\begin{aligned} T(r, g^n (g-1) g') &\leq \bar{N}(r, g) + \bar{N}\left(r, \frac{1}{g^n (g-1) g'}\right) \\ &\quad + \bar{N}(r, g) + 2 \log r + S(r, g) \\ &= \bar{N}\left(r, \frac{1}{g^n (g-1) g'}\right) + 2\bar{N}(r, g) \\ &\quad + 2 \log r + S(r, g). \end{aligned} \quad (63)$$

By Lemmas 9 and 6 and (63), we have

$$\begin{aligned} (n+1)T(r, g) &= T(r, g^n (g-1)) + S(r, g) \\ &\leq T(r, g^n (g-1) g') + T\left(r, \frac{1}{g'}\right) + S(r, g) \\ &\leq \bar{N}\left(r, \frac{1}{g^n (g-1) g'}\right) + 2\bar{N}(r, g) \\ &\quad + 2 \log r + 2T(r, g) + S(r, g) \\ &\leq \bar{N}\left(r, \frac{1}{g}\right) + \bar{N}\left(r, \frac{1}{g-1}\right) + N\left(r, \frac{1}{g'}\right) \\ &\quad + 2\bar{N}(r, g) + 2T(r, g) + 2 \log r + S(r, g) \\ &\leq 8T(r, g) + 2 \log r + S(r, g) \\ &\Rightarrow (n-7)T(r, g) \leq 2 \log r + S(r, g); \end{aligned} \quad (64)$$

we obtain $n \leq 7$, which contradicts $n \geq 11$.

Case 2. Suppose that $f_2 = c$ ($\neq 0$), where c is a constant.

If $c \neq 1$, then we have

$$\begin{aligned} f_1 + f_2 + f_3 &= 1 \\ \Rightarrow \frac{f^n (f-1) f'}{z} + c - \frac{c g^n (g-1) g'}{z} &= 1; \end{aligned} \quad (65)$$

$$\Rightarrow \frac{f^n (f-1) f'}{z} - \frac{c g^n (g-1) g'}{z} = 1 - c. \quad (66)$$

Applying Lemma 5 to the above equation, we have

$$\begin{aligned} T\left(r, \frac{f^n (f-1) f'}{z}\right) &\leq \bar{N}\left(r, \frac{f^n (f-1) f'}{z}\right) \\ &\quad + \bar{N}\left(r, \frac{z}{g^n (g-1) g'}\right) \\ &\quad + \bar{N}\left(r, \frac{z}{f^n (f-1) f'}\right) + S(r, f) \end{aligned}$$

$$\begin{aligned}
&\leq \overline{N}(r, f) + \overline{N}\left(r, \frac{1}{g^n(g-1)g'}\right) \\
&\quad + \overline{N}\left(r, \frac{1}{f^n(f-1)f'}\right) \\
&\quad + \log r + S(r, f).
\end{aligned} \tag{67}$$

Note that

$$T(r, f^n(f-1)f') \leq T\left(r, \frac{f^n(f-1)f'}{z}\right) + \log r. \tag{68}$$

Therefore,

$$\begin{aligned}
T(r, f^n(f-1)f') &\leq \overline{N}(r, f) + \overline{N}\left(r, \frac{1}{g^n(g-1)g'}\right) \\
&\quad + \overline{N}\left(r, \frac{1}{f^n(f-1)f'}\right) \\
&\quad + 2 \log r + S(r, f).
\end{aligned} \tag{69}$$

Using Lemmas 9 and 6 and (69), we have

$$\begin{aligned}
(n+1)T(r, f) &= T(r, f^n(f-1)) + S(r, f) \\
&\leq T(r, f^n(f-1)f') \\
&\quad + T\left(r, \frac{1}{f'}\right) + S(r, f) \\
&\leq \overline{N}(r, f) + \overline{N}\left(r, \frac{1}{f^n(f-1)f'}\right) \\
&\quad + \overline{N}\left(r, \frac{1}{g^n(g-1)g'}\right) \\
&\quad + T(r, f') + 2 \log r + S(r, f) \\
&\leq \overline{N}(r, f) + \overline{N}\left(r, \frac{1}{f}\right) \\
&\quad + \overline{N}\left(r, \frac{1}{f-1}\right) + N\left(r, \frac{1}{f'}\right)
\end{aligned}$$

$$\begin{aligned}
&+ \overline{N}\left(r, \frac{1}{g}\right) + \overline{N}\left(r, \frac{1}{g-1}\right) \\
&+ N\left(r, \frac{1}{g'}\right) + 2T(r, f) \\
&+ 2 \log r + S(r, f) \\
&\leq 7T(r, f) + 4T(r, g) + 2 \log r + S(r, f) \\
&\implies (n-6)T(r, f) \leq 4T(r, g) + 2 \log r + S(r, f).
\end{aligned} \tag{70}$$

Using Lemma 7, we get

$$(n-6)T(r, f) \leq 4\left(\frac{n+3}{n-6}\right)T(r, f) + 2 \log r + S(r, f); \tag{71}$$

since $n \geq 11$, we get a contradiction.

Therefore $c = 1$, and by (27) and (24) we have

$$f^n(f-1)f' = g^n(g-1)g'. \tag{72}$$

On integrating, we get

$$\begin{aligned}
\frac{f^{n+2}}{n+2} - \frac{f^{n+1}}{n+1} &= \frac{g^{n+2}}{n+2} - \frac{g^{n+1}}{n+1} + k \\
F^* &= G^* + k, \quad \text{where } k \text{ is a constant.}
\end{aligned} \tag{73}$$

We claim that $k = 0$. Suppose that $k \neq 0$, then

$$\begin{aligned}
&\Theta(0, F^*) + \Theta(k, F^*) + \Theta(\infty, F^*) \\
&= \Theta(0, F^*) + \Theta(0, G^*) + \Theta(\infty, F^*).
\end{aligned} \tag{74}$$

We have

$$\begin{aligned}
&\overline{N}\left(r, \frac{1}{F^*}\right) \\
&= \overline{N}\left(r, \frac{1}{f}\right) + \overline{N}\left(r, \frac{1}{f - (n+2)/(n+1)}\right) \leq 2T(r, f).
\end{aligned} \tag{75}$$

Similarly,

$$\begin{aligned}
&\overline{N}\left(r, \frac{1}{G^*}\right) \leq 2T(r, g), \\
&\overline{N}(r, F^*) = \overline{N}(r, f) \leq T(r, f).
\end{aligned} \tag{76}$$

Using Lemma 9, we have

$$\begin{aligned}
T(r, F^*) &= (n+2)T(r, f) + S(r, f), \\
T(r, G^*) &= (n+2)T(r, g) + S(r, g).
\end{aligned} \tag{77}$$

Thus,

$$\Theta(0, F^*) = 1 - \lim_{r \rightarrow \infty} \frac{\overline{N}(r, 1/F^*)}{T(r, F^*)} \geq 1 - \frac{2}{n+2}. \tag{78}$$

Similarly,

$$\begin{aligned}\Theta(0, G^*) &\geq 1 - \frac{2}{n+2}, \\ \Theta(\infty, F^*) &= 1 - \lim_{r \rightarrow \infty} \frac{\bar{N}(r, F^*)}{T(r, F^*)} \geq 1 - \frac{1}{n+2}.\end{aligned}\quad (79)$$

Therefore, (74) becomes

$$\begin{aligned}\Theta(0, F^*) + \Theta(k, F^*) + \Theta(\infty, F^*) \\ \geq 2\left(1 - \frac{2}{n+2}\right) + 1 - \frac{1}{n+2} \\ = \frac{3n+1}{n+2} > 2 \quad \text{for } n \geq 11,\end{aligned}\quad (80)$$

which contradicts $\sum_{a \in \bar{\mathbb{C}}} \Theta(a, f) \leq 2$. Thus, we have

$$\frac{f^{n+2}}{n+2} - \frac{f^{n+1}}{n+1} = \frac{g^{n+2}}{n+2} - \frac{g^{n+1}}{n+1}.\quad (81)$$

Let $h = f/g$. If $h \neq 1$, then we easily obtain that

$$g = \frac{(n+2)(h^{n+1}-1)}{(n+1)(h^{n+2}-1)}, \quad f = \frac{(n+2)h(h^{n+1}-1)}{(n+1)(h^{n+2}-1)}.\quad (82)$$

If $h \equiv 1$, that is, $f \equiv g$.

Case 3. Suppose that $f_3 = c (\neq 0)$, where c is a constant.

If $c \neq 1$, then we have

$$\begin{aligned}f_1 + f_2 + f_3 &= 1 \\ f_1 + f_2 &= 1 - c \\ \frac{f^n(f-1)f'}{z} - \frac{cz}{g^n(g-1)g'} &= 1 - c.\end{aligned}\quad (83)$$

Applying Lemma 5 to the above equation, we have

$$\begin{aligned}T\left(r, \frac{f^n(f-1)f'}{z}\right) &\leq \bar{N}\left(r, \frac{f^n(f-1)f'}{z}\right) \\ &\quad + \bar{N}\left(r, \frac{z}{f^n(f-1)f'}\right) \\ &\quad + \bar{N}\left(r, \frac{g^n(g-1)g'}{z}\right) + S(r, f) \\ &\leq \bar{N}(r, f) + \bar{N}\left(r, \frac{1}{f^n(f-1)f'}\right) \\ &\quad + \bar{N}(r, g) + 2 \log r + S(r, f).\end{aligned}\quad (84)$$

Note that

$$\begin{aligned}T(r, f^n(f-1)f') &\leq T\left(r, \frac{f^n(f-1)f'}{z}\right) + T(r, z) \\ &\leq T\left(r, \frac{f^n(f-1)f'}{z}\right) + \log r.\end{aligned}\quad (85)$$

Therefore using (84), we have

$$\begin{aligned}T(r, f^n(f-1)f') &\leq \bar{N}(r, f) + \bar{N}\left(r, \frac{1}{f^n(f-1)f'}\right) \\ &\quad + \bar{N}(r, g) + 3 \log r + S(r, f).\end{aligned}\quad (86)$$

Using Lemmas 9 and 6 and (86), we have

$$\begin{aligned}(n+1)T(r, f) &= T(r, f^n(f-1)) + S(r, f) \\ &\leq T(r, f^n(f-1)f') \\ &\quad + T\left(r, \frac{1}{f'}\right) + S(r, f) \\ &\leq \bar{N}(r, f) + \bar{N}\left(r, \frac{1}{f^n(f-1)f'}\right) \\ &\quad + \bar{N}(r, g) + T(r, f') + 3 \log r + S(r, f) \\ &\leq \bar{N}(r, f) + \bar{N}\left(r, \frac{1}{f}\right) + \bar{N}\left(r, \frac{1}{f-1}\right) \\ &\quad + N\left(r, \frac{1}{f'}\right) + \bar{N}(r, g) + 2T(r, f) \\ &\quad + 3 \log r + S(r, f) \\ &\leq 7T(r, f) + T(r, g) + 3 \log r + S(r, f) \\ &\implies (n-6)T(r, f) \leq T(r, g) + 3 \log r + S(r, f).\end{aligned}\quad (87)$$

Using Lemma 7, we get

$$(n-6)T(r, f) \leq \frac{(n+3)}{(n-6)}T(r, f) + 3 \log r + S(r, f); \quad (88)$$

since $n \geq 11$, we get a contradiction. Thus, $c = 1$. Hence,

$$\begin{aligned}\frac{f^n(f-1)f'}{z} - \frac{z}{g^n(g-1)g'} &= 0 \\ \implies f^n(f-1)f'g^n(g-1)g' &= z^2.\end{aligned}\quad (89)$$

Let z_0 be a zero of f of order p . From (89), we know that z_0 is a pole of g . Suppose that z_0 is a pole of g of order q . From (89), we obtain

$$\begin{aligned}np + p - 1 &= nq + 2q + 1 \\ \implies (n+1)(p-q) &= q + 2,\end{aligned}\quad (90)$$

which implies that $p \geq q + 1$ and $q + 2 \geq n + 1$. Hence,

$$p \geq n.\quad (91)$$

Let z_1 be a zero of $(f-1)$ of order p_1 , then from (89) z_1 is a pole of g (say order q_1). By (89), we get

$$\begin{aligned} p_1 + p_1 - 1 &= nq_1 + 2q_1 + 1 \\ 2p_1 - 1 &\geq n + 3 \\ 2p_1 \geq n + 4 &\implies p_1 \geq \frac{n+4}{2}. \end{aligned} \quad (92)$$

Let z_2 be a zero of f' of order p_2 that is not zero of $f(f-1)$, then from (89), z_2 is a pole of g of order q_2 . Again by (89), we get

$$\begin{aligned} p_2 &= nq_2 + 2q_2 + 1 \\ \implies p_2 &\geq n + 3. \end{aligned} \quad (93)$$

In the same manner as above, we have similar results for the zeros of $g^n(g-1)g'$. From (89)–(93), we have

$$\overline{N}(r, f^n(f-1)f') = \overline{N}\left(r, \frac{z^2}{g^n(g-1)g'}\right); \quad (94)$$

that is,

$$\begin{aligned} \overline{N}(r, f) &\leq \overline{N}\left(r, \frac{1}{g}\right) + \overline{N}\left(r, \frac{1}{g-1}\right) + N\left(r, \frac{1}{g'}\right) \\ &\leq \frac{1}{11}N\left(r, \frac{1}{g}\right) + \frac{2}{15}N\left(r, \frac{1}{g-1}\right) + \frac{1}{14}N\left(r, \frac{1}{g'}\right) \\ &\leq \frac{1}{11}T(r, g) + \frac{2}{15}T(r, g) + \frac{2}{14}T(r, g) + S(r, g) \\ &= \left(\frac{1}{11} + \frac{2}{15} + \frac{2}{7}\right)T(r, g) + S(r, g) \\ \implies \overline{N}(r, f) &< \frac{2}{3}T(r, g) + S(r, g). \end{aligned} \quad (95)$$

By Nevanlinna's second fundamental theorem, we have from (91), (92), and (95) that

$$\begin{aligned} T(r, f) &\leq \overline{N}\left(r, \frac{1}{f}\right) + \overline{N}\left(r, \frac{1}{f-1}\right) + \overline{N}(r, f) + S(r, f) \\ &< \frac{1}{11}T(r, f) + \frac{2}{15}T(r, f) + \frac{2}{3}T(r, g) \\ &\quad + S(r, f) + S(r, g) \\ \implies T(r, f) &\leq \frac{37}{165}T(r, f) + \frac{2}{3}T(r, g) + S(r, f) + S(r, g). \end{aligned} \quad (96)$$

Similarly,

$$T(r, g) \leq \frac{37}{165}T(r, g) + \frac{2}{3}T(r, f) + S(r, f) + S(r, g). \quad (97)$$

From (96) and (97), we get

$$\begin{aligned} T(r, f) + T(r, g) &\leq \frac{37}{165}(T(r, f) + T(r, g)) \\ &\quad + \frac{2}{3}(T(r, f) + T(r, g)) + S(r, f) + S(r, g) \\ \frac{18}{165}(T(r, f) + T(r, g)) &\leq S(r, f) + S(r, g); \end{aligned} \quad (98)$$

since $n \geq 11$, we get a contradiction.

This completes the proof of Theorem 1. $\square$

Using the same argument as in the proof of Theorem 1, we can prove Theorem 2.

Proof of Theorem 3. By the assumption of the theorem, we know that either both f and g are two transcendental entire functions or both f and g are polynomials.

If f and g are transcendental entire functions, putting $\overline{N}(r, f) = 0$, $\overline{N}(r, g) = 0$ and using similar arguments as in the proof of Theorem 1, we easily obtain Theorem 3.

If f and g are polynomials, $f^n(f-1)f'$ and $g^n(g-1)g'$ share z CM, we get

$$f^n(f-1)f' - z = k(g^n(g-1)g' - z), \quad (99)$$

where k is a nonzero constant. Suppose that $k \neq 1$, (99) can be written as

$$\frac{f^n(f-1)f'}{z} - k \frac{g^n(g-1)g'}{z} = 1 - k. \quad (100)$$

Applying Lemma 5 to the above equation, we have

$$\begin{aligned} T\left(r, \frac{f^n(f-1)f'}{z}\right) &\leq \overline{N}\left(r, \frac{f^n(f-1)f'}{z}\right) \\ &\quad + \overline{N}\left(r, \frac{z}{g^n(g-1)g'}\right) \\ &\quad + \overline{N}\left(r, \frac{z}{f^n(f-1)f'}\right) + S(r, f). \end{aligned} \quad (101)$$

Since f is a polynomial, so it does not have any poles. Thus, we have,

$$\begin{aligned} T\left(r, \frac{f^n(f-1)f'}{z}\right) &\leq \overline{N}\left(r, \frac{1}{g^n(g-1)g'}\right) \\ &\quad + \overline{N}\left(r, \frac{1}{f^n(f-1)f'}\right) + 2 \log r + S(r, f). \end{aligned} \quad (102)$$

Note that

$$T(r, f^n(f-1)f') \leq T\left(r, \frac{f^n(f-1)f'}{z}\right) + \log r. \quad (103)$$

Therefore,

$$\begin{aligned} T(r, f^n(f-1)f') &\leq \overline{N}\left(r, \frac{1}{g^n(g-1)g'}\right) \\ &\quad + \overline{N}\left(r, \frac{1}{f^n(f-1)f'}\right) + 3\log r + S(r, f). \end{aligned} \quad (104)$$

Using Lemmas 9 and 6 and (104), we have

$$\begin{aligned} (n+1)T(r, f) &= T(r, f^n(f-1)) + S(r, f) \\ &\leq T(r, f^n(f-1)f') \\ &\quad + T\left(r, \frac{1}{f'}\right) + S(r, f) \\ &\leq \overline{N}\left(r, \frac{1}{f^n(f-1)f'}\right) \\ &\quad + \overline{N}\left(r, \frac{1}{g^n(g-1)g'}\right) \\ &\quad + T(r, f') + 3\log r + S(r, f) \\ &\leq \overline{N}\left(r, \frac{1}{f}\right) + \overline{N}\left(r, \frac{1}{f-1}\right) + N\left(r, \frac{1}{f'}\right) \\ &\quad + \overline{N}\left(r, \frac{1}{g}\right) + \overline{N}\left(r, \frac{1}{g-1}\right) \\ &\quad + N\left(r, \frac{1}{g'}\right) + T(r, f) \\ &\quad + 3\log r + S(r, f) \\ &\leq 4T(r, f) + 3T(r, g) + 3\log r + S(r, f) \\ &\Rightarrow (n-3)T(r, f) \leq 3T(r, g) + 3\log r + S(r, f). \end{aligned} \quad (105)$$

Using Lemma 8, we get

$$(n-3)T(r, f) \leq 3\left(\frac{n+2}{n-3}\right)T(r, f) + 3\log r + S(r, f); \quad (106)$$

since $n \geq 7$, we get a contradiction.

Therefore, $k = 1$; so (99) becomes

$$f^n(f-1)f' = g^n(g-1)g'. \quad (107)$$

On integrating, we get

$$\begin{aligned} \frac{f^{n+2}}{n+2} - \frac{f^{n+1}}{n+1} &= \frac{g^{n+2}}{n+2} - \frac{g^{n+1}}{n+1} + c \\ F^* &= G^* + c, \quad \text{where } c \text{ is a constant.} \end{aligned} \quad (108)$$

We claim that $c = 0$. Suppose that $c \neq 0$, then

$$\Theta(0, F^*) + \Theta(c, F^*) = \Theta(0, F^*) + \Theta(0, G^*). \quad (109)$$

We have

$$\begin{aligned} \overline{N}\left(r, \frac{1}{F^*}\right) &= \overline{N}\left(r, \frac{1}{f}\right) + \overline{N}\left(r, \frac{1}{f-(n+2)/(n+1)}\right) \leq 2T(r, f). \end{aligned} \quad (110)$$

Similarly,

$$\overline{N}\left(r, \frac{1}{G^*}\right) \leq 2T(r, g). \quad (111)$$

Using Lemma 9, we have

$$\begin{aligned} T(r, F^*) &= (n+2)T(r, f) + S(r, f), \\ T(r, G^*) &= (n+2)T(r, g) + S(r, g). \end{aligned} \quad (112)$$

Thus,

$$\Theta(0, F^*) = 1 - \lim_{r \rightarrow \infty} \frac{\overline{N}(r, 1/F^*)}{T(r, F^*)} \geq 1 - \frac{2}{n+2}. \quad (113)$$

Similarly,

$$\Theta(0, G^*) \geq 1 - \frac{2}{n+2}. \quad (114)$$

Therefore, (109) becomes

$$\begin{aligned} \Theta(0, F^*) + \Theta(c, F^*) &\geq 2\left(1 - \frac{2}{n+2}\right) \\ &= \frac{2n}{n+2} \geq \frac{14}{9} > 1 \quad \text{for } n \geq 7, \end{aligned} \quad (115)$$

which contradicts $\sum_{a \in \mathbb{C}} \Theta(a, f) \leq 1$. Thus, we have

$$\frac{f^{n+2}}{n+2} - \frac{f^{n+1}}{n+1} = \frac{g^{n+2}}{n+2} - \frac{g^{n+1}}{n+1}. \quad (116)$$

Let $h = f/g$. If $h \neq 1$, then by (116) we have

$$g = \frac{(n+2)(1+h+h^2+\dots+h^n)}{(n+1)(1+h+h^2+\dots+h^{n+1})}. \quad (117)$$

By Picard's theorem, $h(z)$ is a constant. Hence, g is a constant, which is a contradiction. Therefore, $h(z) \equiv 1$, that is, $f(z) \equiv g(z)$. $\square$

4. Remarks

If the condition “ $f^n(f-1)f'$ and $g^n(g-1)g'$ share z CM” is replaced by the condition “ $f^n(f-1)f'$ and $g^n(g-1)g'$ share $\alpha(z)$ CM,” where α is a meromorphic function such that $\alpha \neq 0, \infty$ and $T(r, \alpha) = o\{T(r, f), T(r, g)\}$; the conclusion of Theorems 1, 2, and 3 still holds. We, thus, obtain the following results.

Theorem 11. Let f and g be two nonconstant meromorphic functions, $n \geq 11$ a positive integer. If $f^n(f-1)f'$ and $g^n(g-1)g'$ share $\alpha(z)$ CM, f and g share ∞ IM, then either $f(z) \equiv g(z)$ or

$$g = \frac{(n+2)(1-h^{n+1})}{(n+1)(1-h^{n+2})}, \quad f = \frac{(n+2)h(1-h^{n+1})}{(n+1)(1-h^{n+2})}, \quad (118)$$

where h is a nonconstant meromorphic function.

Theorem 12. Let f and g be two nonconstant meromorphic functions, $n \geq 12$ a positive integer. If $f^n(f-1)^2f'$ and $g^n(g-1)^2g'$ share $\alpha(z)$ CM, f and g share ∞ IM, then $f(z) \equiv g(z)$.

Theorem 13. Let f and g be two nonconstant entire functions, $n \geq 7$ an integer. If $f^n(f-1)f'$ and $g^n(g-1)g'$ share $\alpha(z)$ CM, then $f(z) \equiv g(z)$.

Acknowledgments

The authors thank the referees for their valuable suggestions. This research work is supported by the Department of Science and Technology, Government of India, Ministry of Science and Technology, Technology Bhavan, New Delhi, India (no. SR/S4/MS: 520/08).

References

- [1] C.-C. Yang and H.-X. Yi, *Uniqueness Theory of Meromorphic Functions*, vol. 557 of *Mathematics and Its Applications*, Kluwer Academic, Dordrecht, The Netherlands, 2003.
- [2] L. Yang, *Value Distribution Theory*, Springer, Berlin, Germany, 1993.
- [3] W. Lin and H. Yi, "Uniqueness theorems for meromorphic functions concerning fixed-points," *Complex Variables*, vol. 49, no. 11, pp. 793–806, 2004.
- [4] J.-F. Xu, F. Lü, and H.-X. Yi, "Fixed-points and uniqueness of meromorphic functions," *Computers and Mathematics with Applications*, vol. 59, no. 1, pp. 9–17, 2010.
- [5] C. C. Yang, "On deficiencies of differential polynomials. II," *Mathematische Zeitschrift*, vol. 125, pp. 107–112, 1972.
- [6] H. X. Yi, "Meromorphic functions that share two or three values," *Kodai Mathematical Journal*, vol. 13, no. 3, pp. 363–372, 1990.