Research Article

Molecular Evolution of the Vertebrate FK506 Binding Protein 25

Fei Liu,^{1,2} Xiao-Long Wei,³ Hao Li,^{1,2} Ji-Fu Wei,² Yong-Qing Wang,² and Xiao-Jian Gong¹

¹ Department of Pharmacology, China Pharmaceutical University, Nanjing 210009, China

² Research Division of Clinical Pharmacology, The First Affiliated Hospital, Nanjing Medical University, 300 Guangzhou Road, Nanjing 210029, China

³ Department of Pathology, Cancer Hospital of Shantou University Medical College, Shantou, China

Correspondence should be addressed to Yong-Qing Wang; wyqjsh@hotmail.com and Xiao-Jian Gong; gongxj66@sina.com

Received 25 November 2013; Accepted 16 January 2014; Published 2 March 2014

Academic Editor: Huai-Rong Luo

Copyright © 2014 Fei Liu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

FK506 binding proteins (FKBPs) belong to immunophilins with peptidyl-prolyl isomerases (PPIases) activity. FKBP25 (also known as FKBP3) is one of the nuclear DNA-binding proteins in the FKBP family, which plays an important role in regulating transcription and chromatin structure. The calculation of nonsynonymous and synonymous substitution rates suggested that FKBP25 undergoes purifying selection throughout the whole vertebrate evolution. Moreover, the result of site-specific tests showed that no sites were detected under positive selection. Only one PPIase domain was detected by searching FKBP25 sequences at Pfam and SMART domain databases. Mammalian FKBP25 possess exon-intron conservation, although conservation in the whole vertebrate lineage is incomplete. The result of this study suggests that the purifying selection triggers FKBP25 evolutionary history, which allows us to discover the complete role of the PPIase domain in the interaction between FKBP25 and nuclear proteins. Moreover, intron alterations during FKBP25 evolution that regulate gene splicing may be involved in the purifying selection.

1. Introduction

Immunophilins include three families with peptidyl-prolyl isomerases (PPIases) activity, FK506 binding proteins (FKBPs), cyclophilins, and parvulins. FKBP25 is named for binding to the immunosuppressive drug FK506, characterized by one or more PPIase domains. The 15 identified members of human FKBP25 are divided into 4 groups: cytoplasmic, TPR domain, endoplasmic reticulum (ER), and nucleus. FKBP25 and FKBP133 locate in the nucleus, containing a single PPIase domain [1].

FKBP25 (also known as FKBP3) is the first mammalian FKBP with a calculated molecular mass of 25 kDa found in the nucleus, which plays a role in regulating transcription and chromatin structure. The FKBP25 comprises a conserved PPIase domain at its C-terminus with a 43% sequence identity to FKBP12 and a helix-loop-helix (HLH) motif at its unique hydrophilic N-terminal [2, 3]. This conserved PPIase domain functions in binding to the immunosuppressive agent FK506 or rapamycin. Unlike another FKBP, FKBP25 shows a strong affinity for binding rapamycin ($K_i = 0.9$ nM) over

FK506 ($K_i = 200$ nM) [4]. The FKBP25 was reported to be associated with nuclear proteins including transcription factor Yin-Yang1 (YY1), mouse double minute 2 (MDM2), and histone deacetylases (HDACs) [5]. FKBP25 binds to YY1 at N-terminal and increases its DNA-binding activity without the involvement of the FK506/rapamycin binding domain [6]. In addition, the level and activity of the tumor suppressor protein p53 are negatively regulated by MDM2. The HLH motif of FKBP25 mediates protein-protein interaction to enhance ubiquitination and degradation of oncogene MDM2, increasing the expression of tumor suppressor p53 and its downstream effector p21 [7]. Moreover, the protein-protein interaction contributes to form HDAC complexes, which is critical for the chromatin structure [2].

In 1992, Jin et al. reported the molecule cloning of human FKBP25 and performed a homology comparison between FKBP25 and FKBP12/FKBP13 [8]. Furthermore, Mas et al. showed the molecule cloning of mouse FKBP25 and expression pattern of FKBP25 gene during cerebral cortical neurogenesis [9]. However, the relationships between nuclear functions and evolution in FKBP25 are seldom reported.

TABLE 1: Site-specific tests for positive selection of FKBP25.

Species	Models	Estimates of parameters	lnL	2Δl	Positively selected sites
Vertebrate	M7	$p = 0.91900$ $q = 8.19764$	-5463.938465	0.003806	NA
	M8	$p_0 = 0.99999$ $p = 0.91899$ $q = 8.19758$ ($p_1 = 0.00001$) $w = 1.86072$	-5463.940368		None
Mammalian	M7	$p = 0.33823$ $q = 1.62046$	-2182.244789	0.000258	NA
	M8	$p_0 = 0.99999$ $p = 0.33824$ $q = 1.62055$ ($p_1 = 0.00001$) $w = 1.00000$	-2182.244918		None
Primate	M7	$p = 4.13016$ $q = 99.00000$	-997.077389	0.000102	NA
	M8	$p_0 = 0.99999$ $p = 4.12942$ $q = 99.00000$ ($p_1 = 0.00001$) $w = 1.00000$	-997.077440		None
Mammalian excluding primate	M7	$p = 0.28229$ $q = 1.41420$	-2242.306222	0.000160	NA
	M8	$p_0 = 0.99999$ $p = 0.28230$ $q = 1.41430$ ($p_1 = 0.00001$) $w = 1.00000$	-2242.306302		NS
Rodent	M7	$p = 0.13287$ $q = 1.19752$	-1372.902164	0.000058	NA
	M8	$p_0 = 0.99999$ $p = 0.13287$ $q = 1.19764$ ($p_1 = 0.00001$) $w = 1.00000$	-1372.902193		NS
Teleost	M7	$p = 0.38691$ $q = 4.30540$	-2354.923181	0.000408	NA
	M8	$p_0 = 0.99999$ $p = 0.38690$ $q = 4.30545$ ($p_1 = 0.00001$) $w = 3.90806$	-2354.923385		NS

lnL: the log-likelihood difference between the two models; 2Δl: twice the log-likelihood difference between the two models (In all the species, 2Δl < 9.21, the P-value is more than the significance level 0.05, indicating that M8 model is not better than M7 model); NA: not allowed; NS: not shown (it means the sites under positive selection but not reaching the significance level of 0.9).

In this study, we exhibit an evolutionary analysis not only on selective pressure but also on intron-exon conversion among vertebrate FKBP25 genes.

2. Materials and Methods

2.1. Sequence Data Collection. All the FKBP25 gene and amino acid sequences were obtained from the ENSEMBL (<http://www.ensembl.org/index.html>) [10], based on orthologous and paralogous relationships. The gained FKBP25 sequences were applied as queries to search known FKBP25 genes using BLAST at the National Center for Biotechnology Information (NCBI), in order to confirm whether their best hit was an FKBP25 gene [11].

Incomplete sequences of FKBP25 genes in four species (tree shrew, horse, platypus, and turkey) were retrieved from both ENSEMBL and NCBI. After eliminating these incomplete sequences, 28 sequences were applied for this study. The 28 sequences from 23 species comprised *human* (ENSG00000100442), *chimpanzee* (ENSPTRG00000006305), *gorilla* (ENSGGOG00000013322), *orangutan* (ENSPPYG00000005778), *macaque* (ENSMMUG00000016512), *marmoset* (ENSCJAG00000015972), *mouse* (ENSMUSG0000020949), *rat* (ENSRNOG00000004629), *guinea pig* (ENSCPOG00000001444), *rabbit1* (ENSOCUG00000007535), *rabbit2* (ENSOCUG00000026892), *dog1* (ENSCAFG00000014018), *dog2* (ENSCAFG00000014093), *dog3* (ENSCAFG00000024192), *dog4* (ENSCAFG00000000578), *cow* (ENSBTAG00000002610), *elephant1* (ENSLAFG000000003572), *elephant2* (ENSLAFG000000027553), *opossum* (ENSMODG00000007352), *chicken* (ENSGALG00000012466),

zebra finch (ENSTGUG00000013231), *anole lizard* (ENSACAG00000004080), *xenopus* (ENSXETG00000003052), *fugu* (ENSTRUG00000011887), *medaka* (ENSORLG00000015070), *stickleback* (ENSGACG00000012834), *tetraodon* (ENSTNIG00000010980), and *zebrafish* (ENS DARG00000079018).

2.2. Molecular Phylogenetic Analyses. The protein coding sequences of FKBP25 were aligned using CLUSTAL W program in MEGA 5.05. We constructed a maximum likelihood (ML) tree of FKBP25 amino acid sequences by MEGA 5.05 with the optimal model (Kimura 2-parameter model). The relative support of internal node was performed by bootstrap analyses with 1000 replications for ML reconstructions [12].

2.3. Selection Pressure Analyses. The numbers of nonsynonymous substitutions per nonsynonymous site (dN) and the numbers of synonymous substitutions per synonymous site (dS) were computed by MEGA 5.05 with the modified Nei-Gojobori method. The dN/dS <1, =1 and >1 demonstrate purifying selection, neutral selection, and positive selection, respectively [13]. The dN is the numbers of nonsynonymous substitutions per nonsynonymous site, and the dS is the numbers of synonymous substitutions per synonymous site. The transition/transversion ratio was 1.55 estimated using the ML method by MEGA 5.05 [14].

The FASTA format of FKBP25 sequences was converted to the PAML format using DAMBE software for subsequent site analyses [13]. The CODEML program implemented in the PAML 4.7 package was used to detect positive selection of individual sites. The site-specific model was exerted using

TABLE 2: Exon and intron lengths of FKBP25.

Species	Length (bp)															Total exons
	Exon1	Intron1	Exon2	Intron2	Exon3	Intron3	Exon4	Intron4	Exon5	Intron5	Exon6	Intron6	Exon7	Intron7	Exon8	
Human	108	3548	102	797	108	8173	136	530	68	2761	98	1775	55	—	—	675
Chimpanzee	108	3524	102	797	108	8898	136	530	68	2725	98	1789	55	—	—	675
Gorilla	108	3538	102	796	108	8214	136	530	68	2753	98	1778	55	—	—	675
Orangutan	108	3498	102	793	108	8395	136	533	68	2457	98	1432	55	—	—	675
Macaque	108	3496	102	786	108	8273	136	531	68	2845	98	1818	55	—	—	675
Marmoset	108	3592	102	780	108	5644	136	507	68	2537	98	2100	55	—	—	675
Mouse	108	3762	102	841	108	2224	136	837	68	1961	98	937	55	—	—	675
Rat	108	3528	102	816	108	2030	136	942	68	1667	98	1118	55	—	—	675
Guinea pig	108	3232	102	772	108	3600	136	1416	68	1346	98	1340	55	—	—	675
Rabbit1	108	2189	102	1082	108	4634	136	1115	68	1826	98	1266	55	—	—	675
Rabbit2	620	40	55	—	—	—	—	—	—	—	—	—	—	—	—	675
Dog1	108	2573	102	1076	108	2088	136	468	68	1823	98	1216	55	—	—	675
Dog2	296	13	229	4	129	—	—	—	—	—	—	—	—	—	—	654
Dog3	30	2	195	4	252	2	33	2	33	12	102	—	—	—	—	645
Dog4	427	190	248	—	—	—	—	—	—	—	—	—	—	—	—	675
Cow	108	2332	102	603	108	2835	136	484	68	1706	98	1309	55	—	—	675
Elephant1	108	3176	102	1089	108	4756	136	483	68	1580	98	1725	55	—	—	675
Elephant2	675	—	—	—	—	—	—	—	—	—	—	—	—	—	—	675
Opossum	108	2560	102	1484	108	2807	136	1051	68	1261	98	554	55	—	—	675
Chicken	111	76	102	75	114	408	136	1040	68	1011	98	829	55	—	—	684
Zebra finch	111	112	102	76	108	494	136	892	97	2054	16	69	53	49	55	678
Anole lizard	186	1699	108	1333	136	1078	68	824	98	610	55	—	—	—	—	651
Xenopus	111	2319	102	403	108	418	136	129	68	186	98	787	55	—	—	678
Fugu	105	375	102	78	105	65	136	82	68	68	98	106	55	—	—	669
Medaka	105	109	102	71	99	738	136	75	68	70	98	804	55	—	—	663
Stickleback	105	294	102	76	102	93	136	135	68	81	98	96	55	—	—	666
Tetraodon	105	305	102	80	102	75	136	91	68	70	98	75	55	—	—	666
Zebra fish	105	2527	102	447	16	1117	20	244	16	359	19	904	28	108	11	666
<hr/>																
Intron8	Exon9	Intron9	Exon10	Intron10	Exon11	Intron11	Exon12	Intron12	Exon13	Intron13	Exon14	Intron14	Exon15	Intron15		
1042	24	316	26	708	14	99	10	98	31	107	15	26	76	1784		
<hr/>																
Exon16	Intron16	Exon17														
98	118	55														

Human	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESI	78
Chimpanzee	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESI	78
Gorilla	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESI	78
Orangutan	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESI	78
Macaque	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESI	78
Marmoset	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESI	78
Mouse	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTETI	78
Rat	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTETI	78
Guinea	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESV	78
Rabbit1	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESV	78
Rabbit2	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESV	78
Dog1	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTESI	78
Dog2	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTESI	78
Dog3	MAAAD.PQNAW.VEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTET.L	76
Dog4	MVVAV.PQQAWTMEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTESI	78
Cow	MAAAV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTESI	78
Elephant1	MAAIV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESI	78
Elephant2	MAAIV.PQRAWVEQLRSEQLPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESI	78
Opossum	MAAA.GPQRTWSAEQLRSEALPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTESV	78
Chicken	MAAATAPACQFWSAEELRSEALPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESV	79
Zebra	MAAAGACQFWSAEELRSEALPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESV	79
Anole	MLYL.GVVVSRQPC.LRISFLV.SDFWFFQ.....FLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.SFKGSESP	70
Xenopus	MTAAEFAPAEWSNEQLRSDLPKDDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEETK.RFKGTESV	79
Medaka	MA...DEPMREWTDEQLRSEALPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTETP.L	76
Fugu	MA...EPTREWSDEQLRSDLPKDDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTESV	78
Stickleback	MA...DEPTREWSDEQLRSDLPKDDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTETP	77
Tetraodon	MAA...EPTREWSDEQLRSDLPKDDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTESV	77
Zebrafish	MAAA...EPQWSDQLRSEALPKKDIIFLQDHGSDSFLAEHKLGNIKNVAKTANNDHIVTAYNHLFEESK.RFKGTESV	76
Consensus	m 1 fl e llg nv kta k l a lf	
Human	SKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSAKKK	156
Chimpanzee	SKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSAKKK	156
Gorilla	SKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSAKKK	156
Orangutan	SKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSAKKK	156
Macaque	SKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSAKKK	156
Marmoset	SKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSAKKK	156
Mouse	SKVSEQVNVNVLNEDKE...KDSKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	156
Rat	SKVSEQVNVNVLNEDKE...KDSKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	156
Guinea	SKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	156
Rabbit1	TKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	156
Rabbit2	TKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	156
Dog1	SKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	156
Dog2	SKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	151
Dog3	SKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	154
Dog4	SKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	156
Cow	SKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	156
Elephant1	NKVSEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	156
Elephant2	NNVLEQVNVNVLNEDKE...KETKSEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	156
Opossum	AKVTEQVNVNVLNEDKE...KEVKEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	156
Chicken	EKAAEKAPKAAEAEAKG...KAVKAAEAEVEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	159
Zebra	EKAAEKAPKAAEAEAKG...KAVKAAEAEVEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	157
Anole	EELAEQVNVNVLNEDKE...KEARQEAIDEAGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	148
Xenopus	ETVTKDVRDINDGKE...KETKLEETLDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	157
Medaka	EELTEQVNVNVLNEDKE...KEVKA.EFVDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	152
Fugu	EEVTEQVNVNVLNEDKE...KEVKT.EFVDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	154
Stickleback	EEVTEQVNVNVLNEDKE...KEVKT.EFVDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	153
Tetraodon	EEVTEQVNVNVLNEDKE...KEVKT.EFVDEGPPHYTKSVLKGDKTNFPKKGTVVHCWYTGILQDGTVFDTNIQTSSKKK	153
Zebrafish	KSPTEQVNVNVLNEDKE...NIDTSQVYEDFENHTLNLGKYLNTFREAGTSSQWDSKSHQRSDLDVILSAKKK	153
Consensus	p k l k k n g d v c g d k k	
Human	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Chimpanzee	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Gorilla	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Orangutan	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Macaque	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Marmoset	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Mouse	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Rat	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Guinea	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Rabbit1	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Rabbit2	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Dog1	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Dog2	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	217
Dog3	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	214
Dog4	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Cow	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Elephant1	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Elephant2	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Opossum	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	224
Chicken	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	227
Zebra	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	225
Anole	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	216
Xenopus	KNAPFLSFRVGVGKVIKRGWDEALLTMSKGEKARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVDID	225
Medaka	RQTKFLSFRVGLGRVIRGWDEALLTMSKGETARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVSD	220
Fugu	RQTKFLSFRVGLGRVIRGWDEALLTMSKGETARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVSD	222
Stickleback	RQTKFLSFRVGLGRVIRGWDEALLTMSKGETARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVSD	221
Tetraodon	RQTKFLSFRVGLGRVIRGWDEALLTMSKGETARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVSD	221
Zebrafish	RQTKFLSFRVGLGRVIRGWDEALLTMSKGETARLEIEPENAYGKKGGQDAKIPPNAKLIFEVELVSD	221
Consensus	kp k v rg s e w p ippn l fevelv	

FIGURE 1: Sequence and structural alignment of FKBP25.

the presence of sites with $\omega > 1$ is suggested. On the contrary, the results of P value > 0.05 proved the absence of sites with $\omega > 1$. The twice log likelihood difference between the two compared models ($2\Delta l$) is compared against χ^2 with

FIGURE 5: Sequence logos (MEME LOGOs) of conserved motifs identified in vertebrate FKBP25.

critical values 5.99 and 9.21 at 0.05 and 0.01 significance levels, respectively [15].

2.4. Protein Domain and Motif Analyses. Protein domain analyses of FKBP25 were shown at Pfam domains database (<http://pfam.sanger.ac.uk>) [16]. SMART (<http://smart.embl-heidelberg.de/>) was used to make sure the presence of FKBP25 domains [17]. The motifs of FKBP25 were analyzed by the MEME software (<http://meme.sdsc.edu/meme/web-site/intro.html>) with a maximum of 10 motifs to find [18].

2.5. Exon-Intron Conservation Analyses. We collected elaborate information about FKBP25 exon and intron from ENSEMBL (<http://www.ensembl.org/index.html>) [19]. The number and length of FKBP25 exon and intron in 28 sequences were investigated for exon-intron conservation analyses.

3. Results

3.1. Phylogenetic Analyses of FKBP25. All the FKBP25 gene and protein sequences were collected from the ENSEMBL and checked by BLAST at NCBI. The sequence and structural alignment of FKBP25 was shown in Figure 1. The phylogenetic tree was constructed according to the protein coding sequences of FKBP25 using the maximum likelihood method (Figure 2, left panel). The FKBP25 genes from the primate lineage and teleost lineage form a species-specific cluster, respectively. Four FKBP25 isoforms of *dog* exhibited a close relationship and clustered together, according to the phylogenetic tree. There were similar phenomena in *rabbit* and *elephant*.

3.2. Selection Pressure Analyses. The nonsynonymous to synonymous rate ratio (dN/dS) may demonstrate the selective pressures of involved protein. We calculated the pairwise distance of FKBP25 sequences using MEGA 5.05. There was a significantly lower dN than dS in the pairwise comparisons of these sequences. Most values of dN/dS in these sequences were distributed below the diagonal, showing that the presence of a purifying selection existed in the FKBP25 (Figure 3). The comparisons of average dN and dS in various vertebrate groups were shown in Figure 4, respectively. Furthermore, site-specific tests were performed for searching the positive selection sites in vertebrate, mammalian, primate, and mammalian excluding primate, rodent and teleost lineages. Although some positive selection sites were computed, each $2\Delta l$ of M7 and M8 < 5.99 indicated that the M8 model was not significantly better than the M7 model to fit the data. Consequently, we concluded that the site-specific analyses also compute no positive selection sites acting on FKBP25 using PAML4.7 (Table 1).

3.3. Protein Domain and Motif Analyses. Early studies reported that mammalian FKBP25 have two portions: one is a putative helix-loop-helix motif within N-terminal unique sequence (Figure 5(a)) and the other is the PPIase domain at its C-terminus (Figure 5(b)) [20].

The domain distribution of FKBP25 was investigated using FKBP25 to search amino acid sequences at the Pfam database firstly. Only one domain (PPIase domain) was found in the Pfam database. The PPIase domain within FKBP25 sequences generally started at position 122 and ended at position 221. Similarly, we further make sure that the FKBP25 domain is at SMART, resulting in the single PPIase domain at position 119 to 221.

FIGURE 6: Exon-intron conservation among FKBP25 genes.

We then performed a detailed domain and motif analyses using the MEME software. Except two *dog* isoforms, *dog2* and *dog3*, the FKBP25 sequences used in this study contain a conserved PPIase domain within motif 1 (shown in Figure 2) at its C-terminus. In addition, the result implied that motif 2 located in the N-terminal contained an HLH motif [6], which was associated with DNA binding and dimerization [21]. However, HLH motif was not found in *dog3*, *anole lizard*, and *teleost* lineage, implying that these FKBP25 proteins may function on gene expression in another pathway.

3.4. Exon-Intron Conservation Analyses. The exon-intron information collected from the ENSEMBL database was shown in Table 2 and Figure 6. Most of the FKBP25 genes have 7 exons with similar length in different species (Table 2). Mammalian FKBP25 shows exon-intron conservation with 6 introns and similar sizes of each intron. Intron deletions existed in several isoforms of species. The *rabbit2* isoform had 2 exons, and *elephant2* isoform had only one exon. The exon numbers of *dog2*, *dog3*, and *dog4* isoforms were less

than seven. Except mammalian FKBP25 genes, *anole lizard* reduced one exon compared with mammalian and birds, but the *xenopus* and teleost maintained 7 exons. The intron deletions of FKBP25 genes may happen in the evolutionary process from amphibian to reptile. Then, a subsequent intron insertion occurred in the evolution from reptile to more advanced animals. The FKBP25 genes also had intron insertion in *zebra finch* and *zebra fish*.

4. Discussion

FKBP25 is a nuclear member of the FKBP family that is associated with transcription and chromatin structure [2]. The interactions of FKBP25 with nuclear proteins are closely associated with HLH motif at the N-terminal of FKBP25. However, whether the PPIase domain at C-terminus is important for these interactions remains uncertain. The selection pressure analyses revealed that the purifying selection triggered a whole evolutionary history of FKBP25 in vertebrates, even in each lineage of vertebrates. Purifying selection is one

of the natural selections that resist deleterious mutations with negative selective coefficients [22]. The mutations that disrupt the correct folding of the FKBP25 domain can weaken PPIase activity and may be the deleterious mutations [5]. It was hypothesized that the mutations of PPIase domain were one of explanations behind the purifying selection throughout FKBP25 evolution. Therefore, although the PPIase domain of FKBP25 was not found to be involved in the protein interactions previously, the PPIase domain might have some associations with the YY1 DNA-binding, MDM2 autoubiquitination and degradation, and HDACs complex formation. These inferences will become a potent direction for exploring the relationship between nuclear proteins and PPIase domain in the future.

The protein-coding sequence length of vertebrate FKBP25 is highly conserved that almost all the taxa are 224 bp; nevertheless the original gene length and exon-intron status are tremendously various among vertebrate species. However, mammalian FKBP25 exhibit exon-intron conservation with 6 introns and similar sizes of each intron. *Chicken* FKBP25 maintains 6 introns, but *zebra finch* has one more intron that inserts in the gene. Similarly, a large variability of intron number and sizes among all the taxa shown in Figure 6 revealed that intron insertion and deletion events happened frequently during the FKBP25 evolutionary history from teleost to birds. In particular, *zebrafish* demonstrated the maximum number of introns in this study, and the size of exon is much smaller than other teleost species (Figure 6(g)). The intron loss of FKBP25 gene from species more advanced than *zebrafish* is likely to induce alterations of gene expression due to the absence of specific intron splicing. Under the purifying selection, the FKBP25 gene expression event continuously removes the pernicious mutations that may associate with intron splicing regulation [23].

FKBP25 gene knockdown declined the expression levels of p53 and p21, which emphasized the significance of FKBP25 in regulating p53 and subsequently p21 expression through controlling the ubiquitination of MDM2. Both the FKBP25 PPIase domain and its N-terminal portion were critical for the ubiquitination and degradation of MDM2 [2]. Moreover, Jin et al. reported that FKBP25 prefers to bind to rapamycin rather than FK506, implying that FKBP25 may be an important target molecule for immunosuppression by rapamycin [8]. All the evolution analyses indicated the conservation of FKBP25 gene in vertebrates. Therefore, FKBP25 possesses some basic functions in vertebrate species, like regulating p53 and p21 expression and binding to rapamycin for immunosuppression, reinforcing the suggestion that the purifying selection triggered the evolution of vertebrate FKBP25.

In conclusion, FKBP25 as a nuclear FKBP subjects to the purifying selection throughout the whole evolution, which implied the complete role of the PPIase domain involved in the interaction between FKBP25 and the nuclear proteins that are needed to be discovered continually. Additionally, incomplete exon-intron conservation of FKBP25 meets the vertebrate lineage. The intron gain or loss among the taxa is likely to be involved in the purifying selection.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Authors' Contribution

Fei Liu and Xiao-Long Wei contributed to this paper equally.

Acknowledgments

This project was sponsored by the Grants from the National Natural Science Foundation of China (81273593, 81273274, and 81302331), the Priority Academic Program Development of Jiangsu Higher Education Institutions, National Major Scientific, Technological Special Project for "Significant New Drugs Development" (2011ZX09302-003-02), Jiangsu Province Major Scientific and Technological Special Project (BM2011017), Jiangsu Province's Key Provincial Talents Program (RC201170 and H201108), and the Foundation of the Nanjing Pharmaceutical Association, China (Nanjing, China) (Grant no. H2011YX001).

References

- [1] E. A. Blackburn and M. D. Walkinshaw, "Targeting FKBP isoforms with small-molecule ligands," *Current Opinion in Pharmacology*, vol. 11, no. 4, pp. 365–371, 2011.
- [2] Y.-L. Yao, Y.-C. Liang, H.-H. Huang, and W.-M. Yang, "FKBPs in chromatin modification and cancer," *Current Opinion in Pharmacology*, vol. 11, no. 4, pp. 301–307, 2011.
- [3] S. Riviere, A. Menez, and A. Galat, "On the localization of FKBP25 in T-lymphocytes," *FEBS Letters*, vol. 315, no. 3, pp. 247–251, 1993.
- [4] J. Liang, D. T. Hung, S. L. Schreiber, and J. Clardy, "Structure of the human 25 kDa FK506 binding protein complexed with rapamycin," *Journal of the American Chemical Society*, vol. 118, no. 5, pp. 1231–1232, 1996.
- [5] G. Gudavicius, H. Soufari, S. Upadhyay et al., "Resolving the functions of peptidylprolyl isomerases: insights from the mutagenesis of the nuclear FKBP25 enzyme," *Biochemical Society Transactions*, vol. 41, no. 3, pp. 761–768, 2013.
- [6] W.-M. Yang, Y.-L. Yao, and E. Seto, "The Fk506-binding protein 25 functionally associates with histone deacetylases and with transcription factor YY1," *EMBO Journal*, vol. 20, no. 17, pp. 4814–4825, 2001.
- [7] A. M. Ochocka, P. Kampanis, S. Nicol et al., "FKBP25, a novel regulator of the p53 pathway, induces the degradation of MDM2 and activation of p53," *FEBS Letters*, vol. 583, no. 4, pp. 621–626, 2009.
- [8] Y.-J. Jin, S. J. Burakoff, and B. E. Bierer, "Molecular cloning of a 25-kDa high affinity rapamycin binding protein, FKBP25," *Journal of Biological Chemistry*, vol. 267, no. 16, pp. 10942–10945, 1992.
- [9] C. Mas, I. Guimiot-Maloum, F. Guimiot et al., "Molecular cloning and expression pattern of the Fkbp25 gene during cerebral cortical neurogenesis," *Gene Expression Patterns*, vol. 5, no. 5, pp. 577–585, 2005.
- [10] P. Flicek, I. Ahmed, M. R. Amode et al., "Ensembl 2013," *Nucleic Acids Research*, vol. 41, no. D1, pp. D48–D55, 2013.

- [11] M. Johnson, I. Zaretskaya, Y. Raytselis, Y. Merezuk, S. McGinnis, and T. L. Madden, "NCBI BLAST: a better web interface," *Nucleic Acids Research*, vol. 36, supplement 2, pp. W5–W9, 2008.
- [12] S. Kumar, M. Nei, J. Dudley, and K. Tamura, "MEGA: a biologist-centric software for evolutionary analysis of DNA and protein sequences," *Briefings in Bioinformatics*, vol. 9, no. 4, pp. 299–306, 2008.
- [13] X. Xia and Z. Xie, "DAMBE: software package for data analysis in molecular biology and evolution," *Journal of Heredity*, vol. 92, no. 4, pp. 371–373, 2001.
- [14] K. Tamura, D. Peterson, N. Peterson, G. Stecher, M. Nei, and S. Kumar, "MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods," *Molecular Biology and Evolution*, vol. 28, no. 10, pp. 2731–2739, 2011.
- [15] Z. Yang, "PAML 4: phylogenetic analysis by maximum likelihood," *Molecular Biology and Evolution*, vol. 24, no. 8, pp. 1586–1591, 2007.
- [16] R. D. Finn, J. Tate, J. Mistry et al., "The Pfam protein families database," *Nucleic Acids Research*, vol. 36, no. 1, pp. D281–D288, 2008.
- [17] I. Letunic, T. Doerks, and P. Bork, "SMART 7: recent updates to the protein domain annotation resource," *Nucleic Acids Research*, vol. 40, no. D1, pp. D302–D305, 2012.
- [18] T. L. Bailey and C. Elkan, "Fitting a mixture model by expectation maximization to discover motifs in biopolymers," in *Proceedings of the International Conference on Intelligent Systems for Molecular Biology*, vol. 2, pp. 28–36, 1994.
- [19] T. Hubbard, D. Barker, E. Birney et al., "The Ensembl genome database project," *Nucleic Acids Research*, vol. 30, no. 1, pp. 38–41, 2002.
- [20] M. Leclercq, F. Vinci, and A. Galat, "Mammalian FKBP-25 and its associated proteins," *Archives of Biochemistry and Biophysics*, vol. 380, no. 1, pp. 20–28, 2000.
- [21] W. D. Kohn, C. T. Mant, and R. S. Hodges, " α -helical protein assembly motifs," *Journal of Biological Chemistry*, vol. 272, no. 5, pp. 2583–2586, 1997.
- [22] Z. Yang and J. R. Bielawski, "Statistical methods for detecting molecular adaptation," *Trends in Ecology and Evolution*, vol. 15, no. 12, pp. 496–503, 2000.
- [23] A. Resch, Y. Xing, A. Alekseyenko, B. Modrek, and C. Lee, "Evidence for a subpopulation of conserved alternative splicing events under selection pressure for protein reading frame preservation," *Nucleic Acids Research*, vol. 32, no. 4, pp. 1261–1269, 2004.