Research Article

Effect of Platelet-Rich Plasma on CCl₄-Induced Chronic Liver Injury in Male Rats

Zahra Hesami,¹ Akram Jamshidzadeh,² Maryam Ayatollahi,³ Bita Geramizadeh,³ Omid Farshad,⁴ and Akbar Vahdati¹

¹ Department of Biology, Science and Research Branch, Islamic Azad University, Fars, Iran

² Pharmaceutical Sciences Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

³ Transplant Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

⁴ International Branch, Shiraz University of Medical Sciences, Shiraz, Iran

Correspondence should be addressed to Zahra Hesami; z_hesami59@yahoo.com

Received 11 November 2013; Accepted 2 January 2014; Published 23 February 2014

Academic Editor: Dirk Uhlmann

Copyright © 2014 Zahra Hesami et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Platelet-rich plasma (PRP) has been of great concern to the scientists and doctors who are involved in wound healing and regenerative medicine which focuses on repairing and replacing damaged cells and tissues. Growth factors of platelet-rich plasma are cost-effective, available, and is more stable than recombinant human growth factors. Given these valuable properties, we decided to assess the effect of PRP on CCl₄-induced hepatotoxicity on rats. The rats received CCl₄ (1 mL/kg, i.p. 1:1 in olive oil) twice per week for 8 weeks. Five weeks after CCl₄ injection, the rats also received PRP (0.5 mL/kg, s.c.) two days a week for three weeks. Twenty-four hours after last CCl₄ injection, the animals bled and their livers dissected for biochemical and histopathological studies. Blood analysis was performed to evaluate enzyme activity. The results showed that PRP itself was not toxic for liver and could protect the liver from CCl₄-induced histological damages and attenuated oxidative stress by increase in glutathione content and decrease in lipid peroxidative marker of liver tissue. The results of the present study lend support to our beliefs in hepatoprotective effects of PRP.

1. Introduction

Liver is considered the key organ in the metabolism, detoxification, and secretory functions in the body, and its disorders are numerous with no effective remedies; however, the search for new medicines is still ongoing. It is a hematopoietic organ in the fetal period, and mature hepatocytes produce thrombopoietin, which can stimulate platelet production in bone marrow. However, few studies have investigated the relationship between hematic components, that is, platelets and liver regeneration [1–4].

The platelet-rich plasma (PRP) used in tissue regeneration serves as a developing area for clinicians and researchers. It is well known that platelets have a thrombotic effect. Platelets contain not only proteins needed for hemostasis but also many growth factors such as transforming growth factor (TGF), platelet-derived growth factor (PDGF), vascular

endothelial growth factor (VEGF), epidermal growth factor (EGF), and insulin-like growth factor (IGF) [5].

Some study has been reported; platelets accumulate in the liver under some kinds of pathologic conditions, like ischemia/reperfusion injury [6–8], liver cirrhosis [9], cholestatic liver [10], and viral hepatitis [11].

Many in vitro studies demonstrated that platelets contain several growth factors which may theoretically contribute to the process of liver regeneration [12, 13]. However, there are few studies on the role of platelets in liver regeneration in rats that failed to identify a correlation between platelets and liver regeneration [14, 15].

Carbon tetrachloride (CCl₄) is a widely used chemical for experimental induction of fatty liver and liver fibrosis in animals [16]. Its biotransformation produces hepatotoxic metabolites, the highly reactive trichloromethyl-free radical, subsequently converted to the peroxytrichloromethyl radical

[17]. Based on the findings of different studies in this field, the present study was undertaken to investigate hepatoprotective activity of PRP on the liver injury induced by CCl_4 in experimental animal model.

2. Materials and Methods

2.1. Chemicals. CCl_4 , calcium gluconate, sodium dodecyl sulfate, ethylenediaminetetraacetic acid (EDTA), 5, 5'-dithiobis-(2-nitrobenzoic acid) or DTNB, tris, thiobarbituric acid (TBA), and trichloroacetic acid were from Sigma Chemical Company, Germany. All other chemicals were of highest quality available in the market.

2.2. Preparing Platelet-Rich Plasma. Fourteen female rats (170–200 g) were selected from the Laboratory Animals Research Center in Shiraz University of Medical Sciences. The rats were maintained under controlled temperature and 12 hours light/12 hours dark conditions for one week before the start of the experiments. They were allowed to feed on standard laboratory chaw and tap water ad libitum. The research protocol complied with the guidelines for animal care of our institution.

The rats were anesthetized with ether, followed by blood collection from the rats via open chest cardiac puncture. About 100 mL of blood was collected from them after killing. The blood was then mixed with sodium citrate (3.8%) (9 parts of blood to 1 part of sodium citrate) anticoagulant solution. Then, the blood was centrifuged at 1000 rpm for 15 min at 20°C for separation of platelet rich plasma. Also, the plasma was centrifuged at 3000 rpm for 10 min at 20°C to obtain platelet pellet. The platelet concentrate dissolved in phosphate buffer saline (PBS), pooled and incubated at room temperature for 30 min on a rotating platform to eliminate platelet agglomerates. Platelets were counted using Sysmex KX-21 (Japan), resulting in a platelet number of $679 \times 10^3/\mu\text{L}$ [18]. Afterwards, autologous thrombin was prepared as per Lucarelli et al.'s method [19]. At this step, 330 μL of calcium gluconate (100 mg/mL) was added to 10 mL of plasma and 1 mL of thrombin preparation to 4 mL of platelet concentrate and incubated for 1 h at room temperature to facilitate growth factors release. The platelet secretion was centrifuged at 4000 rpm for 5 min to reduce the presence of platelet membrane fragments. The supernatant was filtered with a 0.22 μm pore filter, divided into aliquots, and frozen at -80°C for subsequent use [20]. The protein concentration in the filtrates was determined by Bradford method (150 mg/mL) [21].

2.3. Study Design. As designed, the 24 male Wistar rats (250–300 g) were randomly divided into 4 groups, each consisting of 6 animals. Group I received olive oil (0.5 mL/kg, i.p., $n = 6$) as normal control; group II received CCl_4 (1 mL/kg body weight as a 1:1 mixture with olive oil i.p., $n = 6$) twice per week for 8 weeks; group III received PRP (0.5 mL/kg 1:1 in PBS, s.c. $n = 6$) two days a week for three weeks; and group IV received the CCl_4 (1 mL/kg body weight as a 1:1 mixture with olive oil i.p.) twice per week for 8 weeks. Five weeks after

CCl_4 injection, the rats received PRP (0.5 mL/kg 1:1 in PBS s.c. $n = 6$) two days a week for three weeks. Twenty-four hours after CCl_4 injection, the animals were anaesthetized by sodium thiopental injection (50 mg/kg) and their blood samples collected from the vena cava. Then, the respective sera were separated for subsequent use of different enzyme measurements. The rats were then decapitated and their livers carefully dissected and cleaned of extraneous tissues, and parts of the liver tissue were immediately transferred to 10% formalin for histopathological assessments.

2.4. Histopathological Studies. Part of the liver was removed from the animals and the tissue fixed in 10% formalin for at least 24 hours. Then, the paraffin sections were prepared (by Automatic tissue processor, Autotechnique) and cut into 5 μm thick sections by a rotary microtome. The sections then were stained with Haematoxylin-Eosin dye and studied for histopathological changes, that is, necrosis, fatty changes, ballooning degeneration, and Inflammation. Histological damage is scored as follows: 0: absent; +: mild; ++: moderate; and +++: severe.

2.5. Measurement of ALT, AST, and Albumin in Serum. Biocon standard kits and DAX-48 autoanalyzer were used to measure alanine aminotransferase (ALT), aspartate aminotransferase (AST), and albumin (ALB) activities in serum, according to Wilkinson et al.'s and Bessay et al.'s method [22, 23].

2.6. Determination of Lipid Peroxidation. The lipid peroxidation extent was assessed by measuring the amount of thiobarbituric acid-reactive substances (TBARs). In Brief, 500 mg of liver tissue gently minced in 4.5 mL of 0.25 M sucrose. The minced tissues gently homogenized and then centrifuged at 2000 rpm for 30 min. Afterwards, 0.1 mL of the supernatant was treated with a buffer containing 0.75 mL of thiobarbituric acid (0.8%, w/v), 0.75 mL of 20% acetic acid (pH = 3.5), and 0.1 mL of sodium dodecyl sulfate (8.1%, w/v). The solution was mixed up with 2 mL of distilled water and heated in a boiling water bath for 60 min. The absorbance then was measured at 532 nm by a Beckman DU-7 spectrophotometer [24].

2.7. GSH Determination. Glutathione reductase 5,5-dithiobis-2 nitrobenzoic acid (DTNB) recycling procedure [25] was used to determine the reduced glutathione. In brief, 100 mg of liver tissues was homogenized in a buffer containing EDTA (0.2 M) to obtain 4% (w/v) whole homogenate. Then, 1.5 mL of the suspension was taken and mixed with a buffer containing 2.5 mL distilled water and 0.5 mL of 50% TCA. The mixture then was centrifuged at 3000 rpm for 15 min and 1 mL of the supernatant mixed with 1 mL of Tris buffer (0.4 M, pH = 8.9) and 0.1 mL of DTNB (0.01 M). The absorbance was measured after 5 min at 412 nm using a Beckman DU-7 spectrophotometer [26].

2.8. Statistical Analysis. The data were analyzed by student's Tukey test and one-way ANOVA, followed by Graph pad

FIGURE 1: Effect of the platelet-rich plasma (PRP) on histopathological changes induced by CCl₄ in rats. (a) (H&E ×100) liver section of normal rats showing normal hepatocytes with prominent nucleus, cytoplasm, and central vein; (b) (H&E ×250) liver sections of CCl₄-treated (0.5 mL/kg i.p.) rats showing fatty chain, necrosis, and infiltration of inflammatory cells; (c) (H&E ×250) liver sections of the rats treated with PRP (0.5 mL/kg s.c.) showing well-brought out central vein hepatocytes with well-preserved cytoplasm and normal hepatocytes with prominent nucleus; (d) (H&E ×250) liver sections of the rats treated with CCl₄ + PRP (0.5 mL/kg + 0.5 mL/kg s.c.) showing normal architecture of hepatocytes and mild infiltration of inflammatory cells. H: hepatocyte, CV: central vein, N: nucleus, F: foamy macrophage cells, FC: fatty chain, NC: necrosis, I: infiltration of inflammatory cells, and BD: ballooning degeneration.

TABLE 1: Effect of the platelet-rich plasma (PRP) on histopathological liver damages induced by CCl₄ in rats.

Groups	Ballooning Degeneration	Fatty change	Hepatocyte necrosis	Inflammation	Others
Control	0	0	0	0	—
CCl ₄	++	+	+++	+	Many foamy macrophages and old necrosis
PRP	0	0	0	0	—
PRP + CCl ₄	0	0	+	+	—

0: absent; +: mild; ++: moderate; +++: severe.
Rats were injected (i.p.) CCl₄ with doses (0.5 mL/kg 1:1 in olive oil) twice per week for 8 weeks. Five weeks after CCl₄ injection, rats received PRP (0.5 mL/kg 1:1 in PBS, s.c.). The PRP (0.5 mL/kg 1:1 in PBS, s.c.) alone did not increase the levels of the enzymes. Values are mean ± SD of 6 rats per group.

Prism 5. The difference between the control and experimental groups was considered significant at $P \leq 0.05$.

3. Results

Histopathological studies revealed that CCl₄ imposed focal necrosis, fatty changes, ballooning degeneration, and infiltration of lymphocytes around the central veins (Figure 1(b); Table 1). Necrosis, which is a more severe form of injury, was markedly prevented by treatment with PRP (Figure 1(d))

and demonstrated a normal appearance, except for mild inflammation (+ in Table 1) in pericentral hepatocytes to the vein and mild necrosis (+ in Table 1). In the PRP groups that received only PRP there was no significant toxicity, which shows that PRP did not induce hepatotoxicity (Figure 1(c)).
Results of enzyme activity analysis are presented in Figure 2. Administration of CCl₄ to rats caused a significant elevation in serum ALT and AST activities after 8 weeks. Albumin (Figure 2(c)) did not show significant changes in CCl₄-treated group, compared to the control group.

FIGURE 2: Effect of platelet rich plasma (PRP) on rat hepatic enzymes and albumin levels changed by CCl₄. (a) ALT, (b) AST, (c) albumin. Rats were injected CCl₄ with doses (0.5 mL/kg i.p. 1:1 in olive oil) twice per week for 8 weeks. Five weeks after CCl₄ injection rats received PRP (0.5 mL/kg s.c.) 2 days a week for 3 weeks. Values are mean $\pm$ SD of 6 rats per group. *Significantly different from CCl₄-treated group ($P \leq 0.05$). **Significantly different from CCl₄-treated group ($P \leq 0.01$).

TABLE 2: Effects of platelet-rich plasma (PRP) on GSH and TBARs levels of the liver damaged by CCl₄ in rats.

Groups	GSH (nmol/g liver)	TBARs (nmol/g liver)
Control	0.35454 $\pm$ 0.035*	0.7 $\pm$ 0.057**
CCl ₄ (0.5 mL/kg)	0.18727 $\pm$ 0.016	4.96 $\pm$ 0.84
PRP (0.5 mL/kg)	0.33187 $\pm$ 0.058*	0.962 $\pm$ 0.23**
CCl ₄ + PRP	0.28774 $\pm$ 0.034*	0.981 $\pm$ 0.035**

GSH: reduced glutathione; TBARs: thiobarbituric acid-reactive substances. Values are mean $\pm$ SD, ($n = 6$).

* $P \leq 0.05$ mean difference, compared to CCl₄-treated rats.

** $P \leq 0.001$ mean difference, compared to CCl₄-treated rats.

Treatment of rats with 0.5 mg/kg i.p. of the PRP markedly prevented CCl₄-induced elevation of serum ALT and AST (Figures 2(a) and 2(b)).

As shown in Table 2, the liver's lipid peroxidation was significantly increased in the CCl₄ group when compared with the controls ($P \leq 0.01$) and PRP ameliorated CCl₄-induced increases of MDA concentration ($P \leq 0.01$). These findings indicate that the oxidative stress in the liver was effectively decreased when treated with PRP.

Glutathione (GSH) is measured as an index of antioxidant status of liver. There was a significant increase of GSH content in the PRP groups, compared to the group that received CCl₄ alone (Table 2).

4. Discussion

Carbon tetrachloride-induced hepatic injury is commonly used as an experimental method for the study of hepatoprotective effects of drugs or medicinal plants extracts, by in vivo and in vitro techniques [27, 28].

CCl₄ is believed to be metabolized by microsomal CYP450 in the liver to a highly reactive trichloromethyl-free radical (*CCl₃) which can start a chain of reactive free radical formation resulting in peroxidation of lipids and damage to the proteins and components of the cells leading to cell lyses [29, 30].

Effect of platelets on liver regeneration was not addressed till the beginning of the 21st century. There are some reported studies in which platelets were shown to promote liver regeneration [27]. The following study was conducted to determine the role of platelets in liver regeneration using a thrombocytosis model in mice after 90% partial hepatectomy [28]. The entire disrupted platelets and the soluble fraction were with significant proliferative effects, whereas the membrane fraction had no significant effect. Studies indicate that the direct contact between platelets and hepatocytes could spark the release of soluble factors from the platelets such as IGF-1 and HGF; IGF-1 is contained in human platelet as the most important mediator for liver regeneration, which had a proliferative effect on them [29]. The findings in the other experiment demonstrate that exogenous platelets also

enhance liver regeneration [30]. Meanwhile, growth factors, like vascular insulin-like growth factor I (IGF-I), endothelial growth factor (VEGF), and hepatocyte growth factor (HGF), contribute to hepatocyte proliferation that induced by platelet [31]. The growth factors stimulate onset of hepatocyte mitosis, which ultimately promote liver regeneration, especially in humans, since it was reported that human platelets do not contain a significant amount of HGF [32]. Most reports lend support to a decrease in platelet count associated with the severity of liver injury [33–35]. Carbon tetrachloride (CCl_4) is a chemical agent used for experimental promotion of fatty liver and liver fibrosis in animals [16]. The present study was undertaken to investigate of the hepatoprotective activity of PRP against CCl_4 -induced damage in rat.

Lipid peroxidation is among the actual causes of CCl_4 -induced liver injury [36, 37] and is mediated by the free radical derivatives of CCl_4 . CCl_3 radicals produced in reactions by animals microsomes liver exposed to CCl_4 were assumed to attack the membrane lipid in endoplasmic reticulum of hepatocyte. When they attacked the membrane lipid in hepatocyte endoplasmic reticulum, malondialdehyde (MDA) emerged promptly [38]. The rise of MDA levels in the liver is indicative of an enhanced peroxidation that causes tissue damage and breakdown of the antioxidant defense mechanisms and thus inhibits the formation of superabundant free radicals [39]. In the present study, PRP administration caused a significant decrease in MDA levels, compared to the CCl_4 -treated rat, suggesting that PRP could protect against CCl_4 -induced lipid peroxidation in rats.

Unlike the toxic consequences of CCl_4 metabolism through the $\text{CYP}_{2\text{E}1}$ pathway, the detoxification pathway involves GSH conjugation of trichloromethyl-free radicals [40]. A reduced level of GSH is crucial in the detoxification of the reactive toxic metabolites of CCl_4 ; liver necrosis is initiated when reserves of GSH are remarkably depleted [41]. In the present study, the hepatic content of GSH was found to be decreased significantly in CCl_4 intoxicated rats, compared to the controls. Table 2 shows that PRP treatment significantly inhibited the CCl_4 -induced decrease of hepatic GSH content. When CCl_4 is administered to rats, the actions of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) in rat plasma rise remarkably with necrosis and lipid accumulation of hepatocyte [42]. Both enzymes are indicators of liver injury. ALT is more sensitive to acute liver injury test, whereas AST is more sensitive to chronic injury [43]. The present study showed that PRP treatment significantly improved levels of ALT, AST, and albumin after CCl_4 administration (Figures 2(a), 2(b), and 2(c)). The most important protein synthesized by the liver is serum albumin whose main function is to regulate the colloidal osmotic pressure of blood and it reflects the extent of functioning liver cell mass. However, measuring of albumin can provide information to identify chronic injury among the experimental rats [44].

The biochemical observations are supported by the histopathological examination of rats' livers. As a result of hepatotoxicity of CCl_4 , significant regenerative cellular proliferation occurs to compensate for the necrotic or damaged tissue. The Histopathological result (Figure 1) shows

severe necrosis in central vein of the rats treated with CCl_4 compared to the control group that showed +++ degree (Table 1). This result is in accordance with other results [16] and shows fatty changes (+ grade in Table 1), compared to the control group.

We showed that fatty changes in PRP + CCl_4 groups decrease (0 degree, Table 1), compared to CCl_4 rat models (Figure 1(b)). The platelets are stimulated by a lot of motivation like infection, inflammation, and injury. Platelets have modulatory effects on inflammatory cell responses [45]. It seems that the histological changes in CCl_4 group, in addition to necrosis, have a foci apoptotic lesion in their livers (Figure 1(b)). The grade of necrotic foci in the liver reduced from +++ in CCl_4 to + in PRP treatment groups (Table 1).

There is evidence showing platelets to be effective in antifibrosis [46, 47], antiapoptosis [48], and liver regeneration [29]. Platelet therapy can open a new horizon to develop novel strategies for the treatments of liver diseases.

The results presented in this study indicated that the treatment of rats with the PRP 5 weeks after CCl_4 -induced toxicity leads to the reduction of hepatotoxicity. The hepatoprotective effects of PRP may be due to inhibited lipid peroxidation and effective recovery of the antioxidative defense system and has a remarkable effect on signal transduction. Overall, platelet-rich plasma can be used as a complementary procedure to decrease the destructive effects of hepatotoxics.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] S. Murata, N. Ohkohchi, R. Matsuo, O. Ikeda, A. Myronovych, and R. Hoshi, "Platelets promote liver regeneration in early period after hepatectomy in mice," *World Journal of Surgery*, vol. 31, no. 4, pp. 808–816, 2007.
- [2] R. Matsuo, N. Ohkohchi, S. Murata et al., "Platelets strongly induce hepatocyte proliferation with IGF-1 and HGF in vitro," *Journal of Surgical Research*, vol. 145, no. 2, pp. 279–286, 2008.
- [3] R. Hoshi, S. Murata, R. Matsuo et al., "Freeze-dried platelets promote hepatocyte proliferation in mice," *Cryobiology*, vol. 55, no. 3, pp. 255–260, 2007.
- [4] M. Lesurtel, R. Graf, B. Aleil et al., "Platelet-derived serotonin mediates liver regeneration," *Science*, vol. 312, no. 5770, pp. 104–107, 2006.
- [5] Y. Hiyama, I. Mahmud, and F. Karimi-Tari, "Platelet-derived growth factor and thromboxane are necessary for liver regeneration," *Cellular and Molecular Biology*, vol. 27, no. 6, pp. 593–599, 1981.
- [6] A. Khandoga, P. Biberthaler, K. Messmer, and F. Krombach, "Platelet-endothelial cell interactions during hepatic ischemia-reperfusion in vivo: a systematic analysis," *Microvascular Research*, vol. 65, no. 2, pp. 71–77, 2003.
- [7] A. Khandoga, M. Hanschen, J. S. Kessler, and F. Krombach, "CD4+ T cells contribute to postischemic liver injury in mice by interacting with sinusoidal endothelium and platelets," *Hepatology*, vol. 43, no. 2, pp. 306–315, 2006.

- [8] S. Pak, T. Kondo, Y. Nakano et al., "Platelet adhesion in the sinusoid caused hepatic injury by neutrophils after hepatic ischemia reperfusion," *Platelets*, vol. 21, no. 4, pp. 282–288, 2010.
- [9] M. M. Zaldivar, K. Pauels, P. Von Hundelshausen et al., "CXC chemokine ligand 4 (CXCL4) is a platelet-derived mediator of experimental liver fibrosis," *Hepatology*, vol. 51, no. 4, pp. 1345–1353, 2010.
- [10] M. W. Laschke, S. Dold, M. D. Menger, B. Jeppsson, and H. Thorlacius, "Platelet-dependent accumulation of leukocytes in sinusoids mediates hepatocellular damage in bile duct ligation-induced cholestasis," *British Journal of Pharmacology*, vol. 153, no. 1, pp. 148–156, 2008.
- [11] P. A. Lang, C. Contaldo, P. Georgiev et al., "Aggravation of viral hepatitis by platelet-derived serotonin," *Nature Medicine*, vol. 14, no. 7, pp. 756–761, 2008.
- [12] K. Matsumoto and T. Nakamura, "Hepatocyte growth factor: molecular structure, roles in liver regeneration, and other biological functions," *Critical Reviews in Oncogenesis*, vol. 3, no. 1-2, pp. 27–54, 1992.
- [13] A. J. Strain, J. A. McGowan, and N. L. R. Bucher, "Stimulation of DNA synthesis in primary cultures of adult rat hepatocytes by rat platelet-associated substance(s)," *In Vitro*, vol. 18, no. 2, pp. 108–116, 1982.
- [14] Y. Kuwashima, K. Aoki, K. Kohyama, and T. Ishikawa, "Hepatocyte regeneration after partial hepatectomy occurs even under severely thrombocytopenic conditions in the rat," *Japanese Journal of Cancer Research*, vol. 81, no. 6-7, pp. 607–612, 1990.
- [15] M. Tomikawa, M. Hashizume, H. Higashi, M. Ohta, and K. Sugimachi, "The role of the spleen, platelets, and plasma hepatocyte growth factor activity on hepatic regeneration in rats," *Journal of the American College of Surgeons*, vol. 182, no. 1, pp. 12–16, 1996.
- [16] R. O. Recknagel, "Carbon tetrachloride hepatotoxicity," *Pharmacological Reviews*, vol. 19, no. 2, pp. 145–208, 1967.
- [17] A. T. Williams and R. F. Burk, "Carbon tetrachloride hepatotoxicity: an example of free radical-mediated injury," *Seminars in Liver Disease*, vol. 10, no. 4, pp. 279–284, 1990.
- [18] X. Xie, S. Zhao, H. Wu et al., "Platelet-rich plasma enhances autograft revascularization and reinnervation in a dog model of anterior cruciate ligament reconstruction," *Journal of Surgical Research*, vol. 183, no. 1, pp. 214–222, 2013.
- [19] E. Lucarelli, A. Beccheroni, D. Donati et al., "Platelet-derived growth factors enhance proliferation of human stromal stem cells," *Biomaterials*, vol. 24, no. 18, pp. 3095–3100, 2003.
- [20] S. Kazemnejad, A. Allameh, A. Gharehbaghian, M. Soleimani, N. Amirzadeh, and M. Jazayeri, "Efficient replacing of fetal bovine serum with human platelet releasate during propagation and differentiation of human bone marrow-derived mesenchymal stem cells to functional hepatocytes-like cells," *Vox Sanguinis*, vol. 95, no. 2, pp. 149–158, 2008.
- [21] N. J. Kruger, "The Bradford method for protein quantitation," in *The Protein Protocols Handbook*, pp. 17–24, Springer, 2009.
- [22] J. H. Wilkinson, J. H. Boutwell, and S. Winsten, "Evaluation of a new system for the kinetic measurement of serum alkaline phosphatase," *Clinical Chemistry*, vol. 15, no. 6, pp. 487–495, 1969.
- [23] O. A. Bessay, O. H. Lowry, and M. J. Brock, "A method for rapid determination of alkaline phosphatase with five cubic milliliters of serum," *The Journal of Biological Chemistry*, vol. 164, pp. 321–329, 1946.
- [24] I. S. Jamall and J. C. Smith, "Effects of cadmium treatment on selenium-dependent and selenium-independent glutathione peroxidase activities and lipid peroxidation in the kidney and liver of rats maintained on various levels of dietary selenium," *Archives of Toxicology*, vol. 58, no. 2, pp. 102–105, 1985.
- [25] F. Tietze, "Enzymic method for quantitative determination of nanogram amounts of total and oxidized glutathione: applications to mammalian blood and other tissues," *Analytical Biochemistry*, vol. 27, no. 3, pp. 502–522, 1969.
- [26] J. Sedlak and R. H. Lindsay, "Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent," *Analytical Biochemistry*, vol. 25, pp. 192–205, 1968.
- [27] S. Murata, N. Ohkohchi, R. Matsuo, O. Ikeda, A. Myronovych, and R. Hoshi, "Platelets promote liver regeneration in early period after hepatectomy in mice," *World Journal of Surgery*, vol. 31, no. 4, pp. 808–816, 2007.
- [28] A. Myronovych, S. Murata, M. Chiba et al., "Role of platelets on liver regeneration after 90% hepatectomy in mice," *Journal of Hepatology*, vol. 49, no. 3, pp. 363–372, 2008.
- [29] N. Ohkohchi, S. Murata, and K. Takahashi, *Platelet and Liver Regeneration, Tissue Regeneration—from Basic Biology to Clinical Application*, 2012.
- [30] R. Matsuo, Y. Nakano, and N. Ohkohchi, "Platelet administration via the portal vein promotes liver regeneration in rats after 70% hepatectomy," *Annals of Surgery*, vol. 253, no. 4, pp. 759–763, 2011.
- [31] R. Matsuo, N. Ohkohchi, S. Murata et al., "Platelets strongly induce hepatocyte proliferation with IGF-1 and HGF in vitro," *Journal of Surgical Research*, vol. 145, no. 2, pp. 279–286, 2008.
- [32] T. Nakamura, T. Nishizawa, M. Hagiya et al., "Molecular cloning and expression of human hepatocyte growth factor," *Nature*, vol. 342, no. 6248, pp. 440–443, 1989.
- [33] S. Panzer, E. Seel, M. Brunner et al., "Platelet autoantibodies are common in hepatitis C infection, irrespective of the presence of thrombocytopenia," *European Journal of Haematology*, vol. 77, no. 6, pp. 513–517, 2006.
- [34] P. Schöffski, F. Tacke, C. Trautwein et al., "Thrombopoietin serum levels are elevated in patients with hepatitis B/C infection compared to other causes of chronic liver disease," *Liver*, vol. 22, no. 2, pp. 114–120, 2002.
- [35] A. Panasiuk, D. Prokopowicz, J. Zak, J. Matowicka-Karna, J. Osada, and J. Wysocka, "Activation of blood platelets in chronic hepatitis and liver cirrhosis P-selectin expression on blood platelets and secretory activity of β -thromboglobulin and platelet factor-4," *Hepato-Gastroenterology*, vol. 48, no. 39, pp. 818–822, 2001.
- [36] S. Basu, "Carbon tetrachloride-induced lipid peroxidation: eicosanoid formation and their regulation by antioxidant nutrients," *Toxicology*, vol. 189, no. 1-2, pp. 113–127, 2003.
- [37] M. K. Manibusan, M. Odin, and D. A. Eastmond, "Postulated carbon tetrachloride mode of action: a review," *Journal of Environmental Science and Health C*, vol. 25, no. 3, pp. 185–209, 2007.
- [38] Y.-W. Hsu, C.-F. Tsai, W.-K. Chen, and F.-J. Lu, "Protective effects of seabuckthorn (*Hippophae rhamnoides* L.) seed oil against carbon tetrachloride-induced hepatotoxicity in mice," *Food and Chemical Toxicology*, vol. 47, no. 9, pp. 2281–2288, 2009.
- [39] K. J. Lee, J. H. Choi, T. Khanal, Y. P. Hwang, Y. C. Chung, and H. G. Jeong, "Protective effect of caffeic acid phenethyl ester against carbon tetrachloride-induced hepatotoxicity in mice," *Toxicology*, vol. 248, no. 1, pp. 18–24, 2008.

- [40] Y. Ou, S. Zheng, L. Lin, Q. Jiang, and X. Yang, "Protective effect of C-phycoerythrin against carbon tetrachloride-induced hepatocyte damage in vitro and in vivo," *Chemico-Biological Interactions*, vol. 185, no. 2, pp. 94–100, 2010.
- [41] R. O. Recknagel, E. A. Glende Jr., J. A. Dolak, and R. L. Waller, "Mechanisms of carbon tetrachloride toxicity," *Pharmacology & Therapeutics*, vol. 43, no. 1, pp. 139–154, 1989.
- [42] R. Yachi, O. Igarashi, and C. Kiyose, "Protective effects of vitamin E analogs against carbon tetrachloride-induced fatty liver in rats," *Journal of Clinical Biochemistry and Nutrition*, vol. 47, no. 2, pp. 148–154, 2010.
- [43] P. L. Wolf, "Biochemical diagnosis of liver disease," *Indian Journal of Clinical Biochemistry*, vol. 14, no. 1, pp. 59–90, 1999.
- [44] D. Vasudevan and S. Sreekumari, "Mineral metabolism," in *Textbook of Biochemistry for Medical Students*, p. 305, Jaypee Brothers, 4th edition, 2005.
- [45] M. R. Yeaman, "The role of platelets in antimicrobial host defense," *Clinical Infectious Diseases*, vol. 25, no. 5, pp. 951–970, 1997.
- [46] M. Watanabe, S. Murata, I. Hashimoto et al., "Platelets contribute to the reduction of liver fibrosis in mice," *Journal of Gastroenterology and Hepatology*, vol. 24, no. 1, pp. 78–89, 2009.
- [47] T. Kodama, T. Takehara, H. Hikita et al., "Thrombocytopenia exacerbates cholestasis-induced liver fibrosis in mice," *Gastroenterology*, vol. 138, no. 7, pp. 2487.e7–2498.e7, 2010.
- [48] K. Hisakura, S. Murata, K. Fukunaga et al., "Platelets prevent acute liver damage after extended hepatectomy in pigs," *Journal of Hepato-Biliary-Pancreatic Sciences*, vol. 17, no. 6, pp. 855–864, 2010.