Research Article

Differential Effect of the Dopamine D₃ Agonist ($\pm$)-7-Hydroxy-2-(N,N-di-*n*-propylamino) Tetralin (7-OH-DPAT) on Motor Activity between Adult Wistar and Sprague-Dawley Rats after a Neonatal Ventral Hippocampus Lesion

Sonia Guzmán-Velázquez,¹ Linda Garcés-Ramírez,¹ Gonzalo Flores,²
Fidel De La Cruz,¹ and Sergio R. Zamudio¹

¹ Departamento de Fisiología, Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional, Prolongación de Carpio y Plan de Ayala, 11340 México, DF, Mexico

² Laboratorio de Neuropsiquiatría, Instituto de Fisiología, Universidad Autónoma de Puebla, 14 Sur 6301, 72570 Puebla, PUE, Mexico

Correspondence should be addressed to Sergio R. Zamudio, zrzamudio@hotmail.com

Received 15 February 2011; Accepted 24 March 2011

Academic Editor: Maarten Reith

Copyright © 2011 Sonia Guzmán-Velázquez et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The neonatal ventral hippocampal lesion (nVHL) has been widely used as an animal model for schizophrenia. Rats with an nVHL show several delayed behavioral alterations that mimic some symptoms of schizophrenia. Sprague-Dawley (SD) rats with an nVHL have a decrease in D₃ receptors in limbic areas, but the expression of D₃ receptors in Wistar (W) rats with an nVHL is unknown. The 7-Hydroxy-2-(N,N-di-*n*-propylamino) tetralin (7-OH-DPAT) has been reported as a D₃-preferring agonist. Thus, we investigated the effect of ($\pm$)-7-OH-DPAT (0.25 mg/kg) on the motor activity in male adult W and SD rats after an nVHL. The 7-OH-DPAT caused a decrease in locomotion of W rats with an nVHL, but it did not change the locomotion of SD rats with this lesion. Our results suggest that the differential effect of 7-OH-DPAT between W and SD rats with an nVHL could be caused by a different expression of the D₃ receptors. These results may have implications for modeling interactions of genetic and environmental factors involved in schizophrenia.

1. Introduction

Dopamine (DA) receptors are classified into two broad families, namely, the D₁-like (D₁ and D₅) and D₂-like (D₂, D₃, and D₄) receptors [1]. The D₃ receptor was first cloned and characterized by Sokoloff et al. in 1990 [2]. It is negatively coupled to adenylate cyclase. In rats, the D₃ receptor is mostly distributed in projection areas of the mesocorticolimbic dopaminergic system, for example, the nucleus accumbens, olfactory tubercle, islands of Calleja, and prefrontal cortex [3–6]. Although the role of the D₃ receptor in the brain function has not been completely established, it has been related to behavioral aspects such as locomotion, emotion, and cognition [6–11]. The D₃ receptor has also been implicated in disorders, such as schizophrenia and drug abuse,

because its pharmacology and pattern of location in the brain is consistent with defective neural circuits seen in such disorders [12, 13]. For example, postmortem studies suggest a D₃ receptor dysfunctionality in some cortical regions of brains obtained from schizophrenic patients [14, 15].

The locomotor responses to novelty and psychostimulants seem to be regulated by D₃ receptors [6, 10, 16, 17]. 7-Hydroxy-2-(N,N-di-*n*-propylamino) tetralin (7-OH-DPAT) has been described as a D₃-preferring agonist [4, 18]. In rats, administration of low doses of 7-OH-DPAT decreases locomotion, with such reduction in locomotion attributed to a D₃ autoreceptor stimulation [7], but some findings suggest that the inhibitory action of the D₃ receptors on locomotion can also occur via postsynaptic mechanisms [8, 19].

To date, the main theory of the origin of schizophrenia points to the neurodevelopmental model, in which developmental abnormalities early in life lead to the activation of pathologic neural circuits during adolescence or young adulthood leading to the appearance of the schizophrenic symptoms [20]. The most thoroughly characterized neurodevelopmental model of schizophrenia is the neonatal ventral hippocampal lesion (nVHL) model [21]. Rats with a bilateral neonatal ventral hippocampus lesion show a delayed onset of motor, social, and cognitive behaviors comparable to many symptoms of schizophrenia [3, 22–28]. For example, animals with an nVHL show increased locomotor responses to stress and psychostimulants after puberty [3, 22, 29, 30]. Changes in the D_3 expression in the limbic areas have also been observed in this model [3].

Although Sprague-Dawley (SD) and Wistar (W) rats are two of the most used strains of rats in research, the vast majority of studies of the nVHL model have been made in SD rats, though, there are some reports using W rats [31, 32]. However, although W and SD rats show differences in some neurobiological functions [33–39], to the best of our knowledge, a comparative study of the nVHL model between W and SD rats has not yet been reported. Therefore, to assess possible strain differences between adult Wistar and Sprague-Dawley rats in the behavior related to the dopaminergic dysregulation caused by the nVHL, we investigated the effect of the dopamine D_3 -preferring agonist 7-OH-DPAT on open-field activity in W and SD rats after an nVHL.

2. Methods

2.1. Animals. Pregnant Sprague-Dawley and Wistar rats were obtained at gestational day 14 to 17 from our facilities (Harlan México was the original source). The animals were individually housed in a Plexiglas cage with a stainless steel cover in a light-(0700 to 1900 lights on) and temperature-(20–22°C) controlled room. Food and water were always available. All surgical and behavioral procedures described in this study were in accordance with *The Guide for the Care and Use of Laboratory Animals* of the Mexican Council for Animal Care (NOM-062-ZOO-1999). Every effort was made to alleviate any pain or distress that might be experienced by the animals during this experiment. Behavioral testing was done between 1000 and 1400 and was recorded on videotape using a VHS video camera (NV-N3000PN, Panasonic) for later examination.

2.2. Neonatal Lesions. The method followed was in essence as previously described [3, 22]. The day following birth, eight litters of 8 male pups each were culled to maintain same litter size across dams. On postnatal day 7 (PD7), pups within each litter (weighing 15 to 17 g) were randomly assigned to either the sham or the lesion group [40, 41]. The pups were anesthetized by hypothermia (placed on wet ice for 12–15 min, with a latex cover placed on the ice to protect the pups' skin), until they were immobile [40]. They were then placed on a modified platform [42] that was positioned in the ear bars of the stereotaxic device (David Kopf Instruments, Tujunga, CA, USA). To achieve a flat-

skull position, the platform was adjusted until the heights of lambda and bregma skull points were equal. Then, 0.3 μ L of 10 μ g/ μ L ibotenic acid (Sigma-Aldrich, México) or an equal volume of 0.1 M phosphate-buffered saline (PBS), pH 7.4, was injected into the ventral hippocampus over 90 s through a 31-gauge stainless steel cannula aimed at the following coordinates: AP -3.0 mm, ML ± 3.5 mm to bregma, and DV -5.0 mm from dura. The cannula was left inserted 60 s more to avoid backflow of the drug up the cannula tract. The injection cannula was connected to a 1- μ L syringe (Hamilton Co., Reno, NV, USA) with TYGON microbore tubing (ID: 0.25 mm; OD: 0.76 mm) filled with sterile water. After the procedure, the pups were placed on a heat pad for recovery and then returned to their dams. On PD21, animals were weaned and segregated into the sham or lesion group (3 or 4 animals per cage) [40, 43].

2.2.1. Brain Histology. After behavioral testing, all animals were overdosed with sodium pentobarbital (Pfizer, México; 150 mg/kg, i.p.) and perfused intracardially with 0.9% saline solution followed by 4% formalin. The brains were removed manually and stored for at least 48 h in 10% formalin. Coronal sections 100- μ m thick were obtained using a vibroslicer (752 M, Cambden Instrument, Lafayette, IN, USA), the sections stained with 0.5% cresyl violet, mounted with resinous medium and examined under a microscope (SMZ-10A, Nikon Instruments Inc., Melville, NY, USA) using visible light. The lesion sites were located with reference to the stereotaxic atlas of Paxinos and Watson [44].

2.3. Motor Activity. The animal was placed on the middle of the open field (black-painted wooden box 60 cm $\times$ 60 cm $\times$ 30 cm; $w \times l \times h$). The light conditions were comparable to the light intensity in the housing room; two 32-W overhead fluorescent bulbs were suspended 208 cm above the center of the field and provided 195 Lx at the floor of the box. The spontaneous motor activity in an unfamiliar environment was measured using a video image analyzer (Videomex-V, Columbus Instruments, Columbus, OH, USA), which keeps track of the distance the animal travels (DT), the amount of time spent travelling (TA), the amount of time spent in a nonambulatory activity (TNA), and the amount of time resting (TR). It also displays the tracings of the path during the session. The floor of the open field was wiped with a detergent solution between each rat.

2.4. Experimental Procedure. To test the effect of the 7-OH-DPAT in male rats with a bilateral, neonatal excitotoxic lesion of the ventral hippocampus, four groups of animals were formed; (1) Wistar rats with a sham neonatal excitotoxic lesion of the ventral hippocampus (W-sham; $n = 14$), (2) Wistar rats with a neonatal excitotoxic lesion of the ventral hippocampus (W-lesion; $n = 14$), (3) Sprague-Dawley rats with a sham neonatal excitotoxic lesion of the ventral hippocampus (SD-sham; $n = 15$), and (4) Sprague-Dawley rats with a neonatal excitotoxic lesion of the ventral hippocampus (SD-lesion; $n = 12$). At PD60, rats were brought, in individual polysulfone cages, to the testing area, which was in the same building and floor as the

FIGURE 1: Schematic drawing of coronal sections from Paxinos and Watson [44] illustrating lesion boundaries and the areas of neural loss and gliosis obtained from cresyl violet-stained coronal sections from male adult W and SD rats with a neonatal ibotenic acid lesion of the ventral hippocampus. The gray area shows the largest, and black area shows the smallest lesions.

colony room. The animals from each group were randomly assigned to receive either a sc injection of ($\pm$)-7-Hydroxy-2-(N,N-di-*n*-propylamino) tetralin hydrobromide (7-OH-DPAT; 0.25 mg/kg; Sigma-Aldrich, México) or vehicle (saline solution; 0.9% NaCl), in a volume of 1 mL/kg body weight. Approximately in each group half of the rats received the 7-OH-DPAT and half saline. Thirty minutes after injection, rats were individually placed in the testing box and 3 min of motor activity was recorded. It was previously reported, under these experimental conditions, that the motor activity of the animals declined during the initial 9 min, with the highest levels of activity in the first 3 min of the test [35, 45]. The 30-min postinjection time for the motor activity recording was chosen because it allows observation of changes in rat behavior produced by low doses of 7-OH-DPAT (motor activity [8, 46], amphetamine-induced stereotypy [47], and latent inhibition [9]). The test time during the day was balanced across strains.

2.5. Statistical Analysis. The behavioral data were compared by a three-way analysis of variance (ANOVA) with strain, lesion, and drug treatment as independent factors. A $P < .05$ was considered statistically significant. For multiple comparisons, in absence of a significant interaction between factors, independent analyses (separated from the ANOVA) of Student-Newman-Keuls (S-N-K) were made. For all analysis, the software SigmaStat version 3.5 (Systat Software Inc., San Jose, CA) was used.

3. Results

3.1. Verification of the Lesion. Bilateral reduction in the size of the ventral hippocampus (VH) was seen in both adult W and SD rats after making the neonatal ventral hippocampus lesions. Cresyl violet-stained sections obtained from brains of adult animals with nVHL showed important bilateral damage of the VH, with neural loss, atrophy, cavities, and apparent retraction of the VH (Figure 1), as previously reported [3]. Only animals that showed evidence of bilateral ventral hippocampus damage were included in the study.

3.2. Motor Activity. Figure 2 shows the locomotor activity as the distance traveled (DT) by Wistar and Sprague-Dawley rats. The three-way ANOVA showed significant differences for strain ($F_{1,47} = 10.26, P < .01$) and drug treatment ($F_{1,47} = 22.06, P < .01$) factors, though there were no significant differences of the main effect of the lesion ($F_{1,47} = 0.22, P > .05$) or strain by lesion by treatment interaction ($F_{1,47} = 0.42, P > .05$). In W rats, the Student-Newman-Keuls (S-N-K) tests, independent of the ANOVA, show that 7-OH-DPAT reduced the distance traveled of sham rats, as well as of rats with a lesion. In SD rats, the S-N-K tests, independent of the ANOVA, show that the 7-OH-DPAT reduced the distance traveled in SD rats but only in the sham group. In addition the sham W rats were more active than the sham SD rats (Figure 2).

FIGURE 2: Locomotor activity in a novel environment measured as the distance traveled (DT) in male adult Wistar (W) and Sprague-Dawley (SD) rats with a neonatal ventral hippocampus lesion. Data are the mean $\pm$ SE ($n = 6-8$ animals per group) of the DT in 3 min. The administration of 7-OH-DPAT (0.25 mg/kg sc) decreased the DT in all groups except in SD rats with a lesion. Sham SD rats showed a smaller amount of DT compared to sham W rats. * $P < .05$, Student-Newman-Keuls test.

Similar results were observed for the time spent in ambulatory movements (TA) (Figure 3). The three-way ANOVA showed for the time in locomotor activity significant effects of strain ($F_{1,47} = 6.90$, $P < .05$) and drug treatment ($F_{1,47} = 13.76$, $P < .01$). Nevertheless, for the main effect of the lesion and interaction of strain by treatment by lesion factors ($F_{1,47} = 0.42$, $P > .05$ and $F_{1,47} = 0.13$, $P > .05$) there were no significant differences. When the independent S-N-K test was used in W rats, it is showed that the 7-OH-DPAT causes a reduction in the time spent in ambulation in both sham and the rats with lesions. As was seen for the distance traveled, in SD rats, the independent S-N-K test showed a different effect of the 7-OH-DPAT on the time spent in locomotion. The neonatal ventral hippocampus lesions prevented the decrease in locomotion caused by the 7-OH-DPAT, though such reduction was measured in sham animals. The sham W rats also spent more time in ambulation compared to the sham SD rats (Figure 3).

In Figure 4, the amount of time spent by rats in nonambulatory activity (TNA) is shown. The TNA includes postural adjustments and stereotypical behavior (grooming, rearing, sniffing, etc.). The three-way ANOVA did not show significant differences caused by the strain ($F_{1,47} = 1.04$, $P > .05$). Both the W and SD rats showed a nonsignificant effect in the TNA caused by the 7-OH-DPAT ($F_{1,47} = 0.14$, $P > .05$). In addition, there were no statistical differences in the TNA caused by the neonatal HV lesion in both W and SD rats ($F_{1,47} = 2.35$, $P > .05$). The ANOVA also did not show a statistically significant interaction for strain by treatment by lesion factors ($F_{1,47} = 0.12$, $P > .05$).

Figure 5 shows the time spent in resting (TR) of W and SD rats. The three-way ANOVA did not show a significant

FIGURE 3: Locomotor activity in a novel environment measured as the time spent in ambulation (TA) in male adult Wistar (W) and Sprague-Dawley (SD) rats with a neonatal ventral hippocampus lesion. Data are the mean $\pm$ SE ($n = 6-8$ animals per group) of the TA in 3 min. The administration of 7-OH-DPAT (0.25 mg/kg sc) decreased the TA in all groups except in SD rats with a lesion. In addition, the sham W rats were more active than the sham SD rats. * $P < .05$, Student-Newman-Keuls test.

FIGURE 4: Motor activity in the novel environment measured as the amount of time spent in nonambulatory activity (TNA) in male adult Wistar (W) and Sprague-Dawley (SD) rats with a neonatal ventral hippocampus lesion. Data are the mean $\pm$ SE ($n = 6-8$ animals per group) of the TNA in 3 min. The administration of 7-OH-DPAT (0.25 mg/kg sc) did not change the TNA in all groups tested.

difference for strain ($F_{1,47} = 3.14$, $P > .05$). The 7-OH-DPAT produced a nonsignificant effect in the TR in both the sham and rats with lesions ($F_{1,47} = 3.12$, $P > .05$). Furthermore, there was no statistical differences in the lesion factor ($F_{1,47} = 1.74$, $P > .05$) or strain by lesion by treatment interaction ($F_{1,47} = 0.15$, $P > .05$).

FIGURE 5: Time spent in resting (TR) in the novel environment in male adult Wistar (W) and Sprague-Dawley (SD) rats with a neonatal ventral hippocampus lesion. Data are the mean $\pm$ SE ($n = 6-8$ animals per group) of the TR in 3 min. The administration of 7-OH-DPAT (0.25 mg/kg sc) did not change the TR in all groups tested.

Finally, there were no pattern differences in the tracings of the path of rats in the open field during the tests therefore these data are not shown.

4. Discussion

Our results confirm that the administration of 7-OH-DPAT (at a D_3 -preferring agonist dose) reduces the locomotion in the open-field test of both adult W and SD rats. In addition, we found a different effect of the 7-OH-DPAT between adult W and SD rats with neurodevelopmental excitotoxic hippocampal damage. After administering the 7-OH-DPAT, the W rats with an nVHL had a decrease in locomotion similar to the W sham rats. In contrast, the nVHL prevented the reduction in locomotion caused by the 7-OH-DPAT in SD rats. These results may have implications for modeling interactions of genetic and environmental factors involved in schizophrenia.

The mesolimbic and nigrostriatal dopamine pathways in the mammalian brain play a role in mediating motor behaviors, including locomotion and stereotypical behavior. Accumulating evidence suggests that stimulation of the D_3 receptors inhibits spontaneous and psychostimulant-caused locomotion, different than the synergistic D_1 and D_2 receptors mediating behavioral sensitivity [7, 17, 19, 48–52]. In agreement with previous work [8, 46, 53], the D_3 -preferring receptor agonist 7-OH-DPAT administered peripherally in our work reduced the spontaneous locomotion in the sham animals. The 7-OH-DPAT has been reported as a D_3 -preferring agonist [4, 18]. The 7-OH-DPAT produces a biphasic effect on behavioral activity in which locomotion is inhibited at lower doses and stimulated at higher doses [7, 54], suggesting a D_3 -receptor activation at low doses and increasing D_2 -receptor occupancy at higher

doses. In our study, a low dose (0.25 mg/kg) of 7-OH-DPAT was used to cause hypolocomotion, suggesting primarily a D_3 -receptor stimulation. The estimates of the D_2 -receptor occupancy in vivo suggest that 7-OH-DPAT doses below 0.3 mg/kg are devoid of significant D_2 -receptor occupancy [55]. The decrease in locomotion has been attributed to a D_3 -autoreceptor stimulation [7, 56], which in turn inhibits dopamine release [57]. Studies in D_3 -receptor knockout mice support the idea of a D_3 -receptor regulation of dopamine levels. In a novel environment, the D_3 mutant mice are transiently more active than wild-type mice [52], and microdialysis studies in vivo have shown that basal extracellular dopamine levels are increased in the nucleus accumbens [58] and dorsal striatum [59] of the D_3 -receptor knockout mice. However, some findings suggest that the inhibitory action of the D_3 receptors on locomotion can also occur via postsynaptic mechanisms [8, 19].

The predominate expression of the D_3 receptors in the limbic areas, for example, nucleus accumbens, has led the field to consider this receptor as a potential target for the treatment of schizophrenia, which is associated with a dysregulation of dopamine neurotransmission [12]. Rats with an nVHL show an alteration of the mesolimbic and mesostriatal dopaminergic transmission [60]. After puberty, rats with lesions develop hypersensitivity to stress and dopamine agonists [3, 22, 43]. In postpubertal SD rats, a decrease in D_3 receptors was measured in the limbic areas of the rats with lesions compared to sham controls, particularly in the nucleus accumbens [3]. This decrease in the D_3 receptors could explain the hyperdopaminergic behaviors observed in SD rats after an nVHL. This explanation could also explain our results with SD rats with lesions treated with the D_3 -preferring agonist 7-OH-DPAT. Here, SD sham rats (with normal D_3 receptor levels in the limbic areas) showed a reduction in locomotion after 7-OH-DPAT administration. In contrast, SD rats with an nVHL (with low D_3 receptor levels in the limbic areas) were not susceptible to the 7-OH-DPAT administration. A reduction in D_3 receptor levels has also been suggested in brains of schizophrenic patients [14, 15], in these postmortem studies, the expression of D_3 mRNA was found selectively lost in the motor, parietal, and anterior cingulate cortices of brains obtained from patients with chronic schizophrenia. In contrast, a truncated D_3 -like mRNA (named D_{3nf}) was abundantly expressed in the same brain areas, this D_{3nf} mRNA could encode a dysfunctional D_3 protein receptor [14, 61]. Thus, another possible explanation about the lack effect of 7-OH-DPAT on locomotion in SD rats with nVHL arises from the above described works, where the nVHL in these rats could produce a dysfunctional expression of D_3 receptors in some brain areas. Future studies will show if truncated D_3 -like mRNAs are also abundantly expressed in brains of rats with nVHL.

The Wistar rats have also been used in the nVHL model [31, 32, 62], but to the best of our knowledge, a direct comparative study of the effect of an nVHL between SD and W rats has not been reported. We found that W rats with an nVHL were susceptible to the 7-OH-DPAT administration. There was a comparable reduction in the locomotor activity, after 7-OH-DPAT treatment, between the sham W rats and

those with lesions. We hypothesized that there is a different effect of an nVHL on the expression of the D₃ receptors in the limbic areas between the W and the SD animals. In the W rats, the reduction in the D₃ receptors seen in the SD rats with lesions [3] could be absent or attenuated. Data from studies with Lewis and Fisher 344 rats are consistent with the above hypothesis. Lipska and Weinberger [43] found that Fisher 344 rats with an nVHL have an increase in spontaneous and amphetamine-caused locomotion. In contrast, Lewis rats with an nVHL were not affected by having a lesion. Interestingly in the Lewis rats without a lesion, the D₃ receptors in the nucleus accumbens and olfactory tubercle are lower than levels found in Fisher 344 rats [63]. This suggests that because Lewis rats already have a reduced amount of D₃ receptors, a further decrease in D₃ receptors caused by an nVHL may not occur. Further investigations of dopamine receptor levels in the brain in the nVHL model with different strains will be of interest to extend these findings.

Another interpretation of our results arises from the use of a single dose of 7-OH-DPAT. The SD rats exhibited less locomotor activity than W rats and therefore strain differences in the sensitivity for detecting 7-OH-DPAT-caused decreases in locomotion could explain our results. Certainly, the lower locomotor activity in SD rats could account for the preventive effect of the nVHL on 7-OH-DPAT-caused decreases in locomotion in SD rats. However, it seems less likely because the single dose of 7-OH-DPAT used here was able to reduce the locomotor activity in sham SD rats in a percentage similar to the observed in sham W rats (the 7-OH-DPAT-caused DT reduction was 59% for sham SD and 44% for sham W rats). Both rat strains showed similar sensitivity for the drug in sham animals, thus, it seems likely that, in rats with a lesion, the differences observed in the 7-OH-DPAT-caused decreases in locomotion between strains can be attributed to the nVHL.

Because the D₃ receptor may be an important target for antipsychotic drugs, our results with the nVHL model suggest a strain-dependent different effect on the expression and/or functionality of D₃ receptors in the limbic areas. It could help explain individual differences in the response to antipsychotic drugs in schizophrenic patients.

Acknowledgments

This work was supported by funds from the Instituto Politécnico Nacional (SIP-IPN, 20100852). S. Guzmán-Velázquez, L. Garcés-Ramírez, G. Flores, F. De La Cruz, and S. Zamudio are fellows of Consejo Nacional de Ciencia y Tecnología. The authors thank Ellis Glazier for editing the English text.

References

- [1] C. Missale, S. R. Nash, S. W. Robinson, M. Jaber, and M. G. Caron, "Dopamine receptors: from structure to function," *Physiological Reviews*, vol. 78, no. 1, pp. 189–225, 1998.
- [2] P. Sokoloff, B. Giros, M. P. Martres, M. L. Bouthenet, and J. C. Schwartz, "Molecular cloning and characterization of a novel dopamine receptor (D₃) as a target for neuroleptics," *Nature*, vol. 347, no. 6289, pp. 146–151, 1990.
- [3] G. Flores, D. Barbeau, R. Quirion, and L. K. Srivastava, "Decreased binding of dopamine D₃ receptors in limbic subregions after neonatal bilateral lesion of rat hippocampus," *Journal of Neuroscience*, vol. 16, no. 6, pp. 2020–2026, 1996.
- [4] D. Levesque, J. Diaz, C. Pilon et al., "Identification, characterization, and localization of the dopamine 3D receptor in rat brain using 7-[3H]hydroxy-N,N-di-n-propyl-2-aminotetralin," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 89, no. 17, pp. 8155–8159, 1992.
- [5] J. Diaz, C. Pilon, B. Le Foll et al., "Dopamine D₃ receptors expressed by all mesencephalic dopamine neurons," *Journal of Neuroscience*, vol. 20, no. 23, pp. 8677–8684, 2000.
- [6] J. Flores-Tochihuitl, G. Vargas, J. C. Morales-Medina et al., "Enhanced apomorphine sensitivity and increased binding of dopamine D₂ receptors in nucleus accumbens in prepubertal rats after neonatal blockade of the dopamine D₃ receptors by (+)-S14297," *Synapse*, vol. 62, no. 1, pp. 40–49, 2008.
- [7] S. A. Daly and J. L. Waddington, "Behavioural effects of the putative D-3 dopamine receptor agonist 7-OH-DPAT in relation to other "D-2-like" agonists," *Neuropharmacology*, vol. 32, no. 5, pp. 509–510, 1993.
- [8] K. Svensson, A. Carlsson, and N. Waters, "Locomotor inhibition by the D₃ ligand R-(+)-7-OH-DPAT is independent of changes in dopamine release," *Journal of Neural Transmission-General Section*, vol. 95, no. 1, pp. 71–74, 1994.
- [9] L. Chagas-Martinich, R. J. Carey, and M. P. Carrera, "7-OH-DPAT effects on latent inhibition: low dose facilitation but high dose blockade: implications for dopamine receptor involvement in attentional processes," *Pharmacology Biochemistry and Behavior*, vol. 86, no. 3, pp. 441–448, 2007.
- [10] L. M. Pritchard, A. D. Logue, B. C. Taylor et al., "Relative expression of D₃ dopamine receptor and alternative splice variant D₃nf mRNA in high and low responders to novelty," *Brain Research Bulletin*, vol. 70, no. 4–6, pp. 296–303, 2006.
- [11] C. Wilson and M. Pulido, "Effects of the dopamine antagonist PD 152255 on juvenile rats' responses to dorsal stimulation, the transport response, and related behaviors," *Behavioral Neuroscience*, vol. 116, no. 6, pp. 1098–1102, 2002.
- [12] J. C. Schwartz, J. Diaz, C. Pilon, and P. Sokoloff, "Possible implications of the dopamine D(3) receptor in schizophrenia and in antipsychotic drug actions," *Brain Research Reviews*, vol. 31, no. 2-3, pp. 277–287, 2000.
- [13] M. Pilla, S. Perachon, F. Sautel et al., "Selective inhibition of cocaine-seeking behaviour by a partial dopamine D₃ receptor agonist," *Nature*, vol. 400, no. 6742, pp. 371–375, 1999.
- [14] C. Schmauss, "Enhanced cleavage of an atypical intron of dopamine D₃-receptor pre-mRNA in chronic schizophrenia," *Journal of Neuroscience*, vol. 16, no. 24, pp. 7902–7909, 1996.
- [15] C. Schmauss, V. Haroutunian, K. L. Davis, and M. Davidson, "Selective loss of dopamine D₃-type receptor mRNA expression in parietal and motor cortices of patients with chronic schizophrenia," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 90, no. 19, pp. 8942–8946, 1993.
- [16] R. K. McNamara, A. Logue, K. Stanford, M. Xu, J. Zhang, and N. M. Richtand, "Dose-response analysis of locomotor activity and stereotypy in dopamine D₃ receptor mutant mice following acute amphetamine," *Synapse*, vol. 60, no. 5, pp. 399–405, 2006.
- [17] L. M. Pritchard, A. D. Logue, S. Hayes et al., "7-OH-DPAT and PD 128907 selectively activate the D₃ dopamine receptor in a novel environment," *Neuropsychopharmacology*, vol. 28, no. 1, pp. 100–107, 2003.

- [18] M. Hillefors-Berglund and G. Von Euler, "Pharmacology of dopamine D3 receptors in the islands of Calleja of the rat using quantitative receptor autoradiography," *European Journal of Pharmacology*, vol. 261, no. 1-2, pp. 179–183, 1994.
- [19] N. Waters, K. Svensson, S. R. Haadsma-Svensson, M. W. Smith, and A. Carlsson, "The dopamine D3-receptor: a post-synaptic receptor inhibitory on rat locomotor activity," *Journal of Neural Transmission-General Section*, vol. 94, no. 1, pp. 11–19, 1993.
- [20] S. H. Fatemi and T. D. Folsom, "The neurodevelopmental hypothesis of schizophrenia, revisited," *Schizophrenia Bulletin*, vol. 35, no. 3, pp. 528–548, 2009.
- [21] K. Y. Tseng, R. A. Chambers, and B. K. Lipska, "The neonatal ventral hippocampal lesion as a heuristic neurodevelopmental model of schizophrenia," *Behavioural Brain Research*, vol. 204, no. 2, pp. 295–305, 2009.
- [22] B. K. Lipska, G. E. Jaskiw, and D. R. Weinberger, "Post-pubertal emergence of hyperresponsiveness to stress and to amphetamine after neonatal excitotoxic hippocampal damage: a potential animal model of schizophrenia," *Neuropsychopharmacology*, vol. 9, no. 1, pp. 67–75, 1993.
- [23] F. Sams-Dodd, B. K. Lipska, and D. R. Weinberger, "Neonatal lesions in the rat ventral hippocampus result in hyperlocomotion and deficits in social behaviour in adulthood," *Psychopharmacology*, vol. 132, no. 3, pp. 303–310, 1997.
- [24] A. Becker, G. Grecksch, H. G. Bernstein, V. Höllt, and B. Bogerts, "Social behaviour in rats lesioned with ibotenic acid in the hippocampus: quantitative and qualitative analysis," *Psychopharmacology*, vol. 144, no. 4, pp. 333–338, 1999.
- [25] J. P. Marquis, S. Goulet, and F. Y. Doré, "Neonatal lesions of the ventral hippocampus in rats lead to prefrontal cognitive deficits at two maturational stages," *Neuroscience*, vol. 140, no. 3, pp. 759–767, 2006.
- [26] R. A. Chambers, J. Moore, J. P. McEvoy, and E. D. Levin, "Cognitive effects of neonatal hippocampal lesions in a rat model of schizophrenia," *Neuropsychopharmacology*, vol. 15, no. 6, pp. 587–594, 1996.
- [27] B. K. Lipska, J. M. Aultman, A. Verma, D. R. Weinberger, and B. Moghaddam, "Neonatal damage of the ventral hippocampus impairs working memory in the rat," *Neuropsychopharmacology*, vol. 27, no. 1, pp. 47–54, 2002.
- [28] G. K. Wood, R. Quirion, and L. K. Srivastava, "Early environment contributes to developmental disruption of MPFC after neonatal ventral hippocampal lesions in rats," *Synapse*, vol. 50, no. 3, pp. 223–232, 2003.
- [29] H. A. Al-Amin, D. R. Weinberger, and B. K. Lipska, "Exaggerated MK-801-induced motor hyperactivity in rats with the neonatal lesion of the ventral hippocampus," *Behavioural Pharmacology*, vol. 11, no. 3-4, pp. 269–278, 2000.
- [30] T. Hori, S. Subramaniam, L. K. Srivastava, and R. Quirion, "Behavioral and neurochemical alterations following repeated phencyclidine administration in rats with neonatal ventral hippocampal lesions," *Neuropharmacology*, vol. 39, no. 12, pp. 2478–2491, 2000.
- [31] V. Blas-Valdivia, E. Cano-Europa, A. Hernández-García, and R. Ortiz-Butrón, "Neonatal bilateral lidocaine administration into the ventral hippocampus caused postpubertal behavioral changes: an animal model of neurodevelopmental psychopathological disorders," *Neuropsychiatric Disease and Treatment*, vol. 5, no. 1, pp. 15–22, 2009.
- [32] K. S. Alexander, J. M. Brooks, M. Sarter, and J. P. Bruno, "Disruption of mesolimbic regulation of prefrontal cholinergic transmission in an animal model of schizophrenia and normalization by chronic clozapine treatment," *Neuropsychopharmacology*, vol. 34, no. 13, pp. 2710–2720, 2009.
- [33] S. Bekris, K. Antoniou, S. Daskas, and Z. Papadopoulou-Daifoti, "Behavioural and neurochemical effects induced by chronic mild stress applied to two different rat strains," *Behavioural Brain Research*, vol. 161, no. 1, pp. 45–59, 2005.
- [34] E. L. Abel, "Response to alarm substance in different rat strains," *Physiology and Behavior*, vol. 51, no. 2, pp. 345–347, 1992.
- [35] S. Zamudio, T. Fregoso, A. Miranda, F. De La Cruz, and G. Flores, "Strain differences of dopamine receptor levels and dopamine related behaviors in rats," *Brain Research Bulletin*, vol. 65, no. 4, pp. 339–347, 2005.
- [36] G. G. Kinney, L. O. Wilkinson, K. L. Saywell, and M. D. Tricklebank, "Rat strain differences in the ability to disrupt sensorimotor gating are limited to the dopaminergic system, specific to prepulse inhibition, and unrelated to changes in startle amplitude or nucleus accumbens dopamine receptor sensitivity," *Journal of Neuroscience*, vol. 19, no. 13, pp. 5644–5653, 1999.
- [37] P. K. Pilz, R. Linke, D. M. Yilmazer-Hanke, and H. Schwegler, "Comparison of two sensitization paradigms of the acoustic startle response in Wistar and Sprague-Dawley rats," *Behavior Genetics*, vol. 29, no. 1, pp. 59–63, 1999.
- [38] I. C. Weiss, L. Di Iorio, J. Feldon, and A. M. Domeney, "Strain differences in the isolation-induced effects on prepulse inhibition of the acoustic startle response and on locomotor activity," *Behavioral Neuroscience*, vol. 114, no. 2, pp. 364–373, 2000.
- [39] A. Rex, J. P. Voigt, C. Gustedt, S. Beckett, and H. Fink, "Anxiolytic-like profile in Wistar, but not Sprague-Dawley rats in the social interaction test," *Psychopharmacology*, vol. 177, no. 1-2, pp. 23–34, 2004.
- [40] K. Y. Tseng, B. L. Lewis, T. Hashimoto et al., "A neonatal ventral hippocampal lesion causes functional deficits in adult prefrontal cortical interneurons," *Journal of Neuroscience*, vol. 28, no. 48, pp. 12691–12699, 2008.
- [41] H. A. Al Amin, S. F. Atweh, S. J. Jabbur, and N. E. Saadé, "Effects of ventral hippocampal lesion on thermal and mechanical nociception in neonates and adult rats," *European Journal of Neuroscience*, vol. 20, no. 11, pp. 3027–3034, 2004.
- [42] A. Sierra, I. Camacho-Abrego, J. V. Negrete-Díaz, L. Rodríguez-Sosa, C. Escamilla, and G. Flores, "Economic body platform for neonatal rats stereotaxic surgery," *Revista de Neurologia*, vol. 48, no. 3, pp. 141–146, 2009.
- [43] B. K. Lipska and D. R. Weinberger, "Genetic variation in vulnerability to the behavioral effects of neonatal hippocampal damage in rats," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 92, no. 19, pp. 8906–8910, 1995.
- [44] G. Paxinos and C. Watson, *The Rat Brain in Stereotaxic Coordinates*, Academic Press, New York, NY, USA, 1986.
- [45] M. López-Martínez, H. Salgado-Zamora, E. R. San-Juan et al., "Anti-anxiety and sedative profile evaluation of imidazo[1,2-a]pyridine derivatives," *Drug Development Research*, vol. 71, no. 6, pp. 371–381, 2010.
- [46] K. Frantz, D. Babcock, and C. Van Hartesveldt, "The locomotor effects of a putative dopamine D3 receptor agonist in developing rats," *European Journal of Pharmacology*, vol. 302, no. 1-3, pp. 1–6, 1996.

- [47] T. V. Khroyan, D. A. Baker, R. A. Fuchs, N. Manders, and J. L. Neisewander, "Differential effects of 7-OH-DPAT on amphetamine-induced stereotypy and conditioned place preference," *Psychopharmacology*, vol. 139, no. 4, pp. 332–341, 1998.
- [48] P. De Boer, P. Enrico, J. Wright et al., "Characterization of the effect of dopamine D3 receptor stimulation on locomotion and striatal dopamine levels," *Brain Research*, vol. 758, no. 1-2, pp. 83–91, 1997.
- [49] T. Kling-Petersen, E. Ljung, and K. Svensson, "Effects on locomotor activity after local application of D3 preferring compounds in discrete areas of the rat brain," *Journal of Neural Transmission-General Section*, vol. 102, no. 3, pp. 209–220, 1995.
- [50] A. M. Ouagazzal and I. Creese, "Intra-accumbens infusion of D(3) receptor agonists reduces spontaneous and dopamine-induced locomotion," *Pharmacology Biochemistry and Behavior*, vol. 67, no. 3, pp. 637–645, 2000.
- [51] L. Zhang, D. Lou, H. Jiao et al., "Cocaine-induced intracellular signaling and gene expression are oppositely regulated by the dopamine D1 and D3 receptors," *Journal of Neuroscience*, vol. 24, no. 13, pp. 3344–3354, 2004.
- [52] M. Xu, T. E. Koeltzow, G. T. Santiago et al., "Dopamine D3 receptor mutant mice exhibit increased behavioral sensitivity to concurrent stimulation of D1 and D2 receptors," *Neuron*, vol. 19, no. 4, pp. 837–848, 1997.
- [53] S. Ahlenius and P. Salmi, "Behavioral and biochemical effects of the dopamine D3 receptor-selective ligand, 7-OH-DPAT, in the normal and the reserpine-treated rat," *European Journal of Pharmacology*, vol. 260, no. 2-3, pp. 177–181, 1994.
- [54] R. Depoortere, G. Perrault, and D. J. Sanger, "Behavioural effects in the rat of the putative dopamine D3 receptor agonist 7-OH-DPAT: comparison with quinpirole and apomorphine," *Psychopharmacology*, vol. 124, no. 3, pp. 231–240, 1996.
- [55] B. Levant, G. N. Bancroft, and C. M. Selkirk, "In vivo occupancy of D2 dopamine receptors by 7-OH-DPAT," *Synapse*, vol. 24, no. 1, pp. 60–64, 1996.
- [56] D. B. Gilbert and S. J. Cooper, "7-OH-DPAT injected into the accumbens reduces locomotion and sucrose ingestion: D3 autoreceptor-mediated effects?" *Pharmacology Biochemistry and Behavior*, vol. 52, no. 2, pp. 275–280, 1995.
- [57] R. R. Gainetdinov, T. D. Sotnikova, T. V. Grekhova, and K. S. Rayevsky, "In vivo evidence for preferential role of dopamine D3 receptor in the presynaptic regulation of dopamine release but not synthesis," *European Journal of Pharmacology*, vol. 308, no. 3, pp. 261–269, 1996.
- [58] T. E. Koeltzow, M. Xu, D. C. Cooper et al., "Alterations in dopamine release but not dopamine autoreceptor function in dopamine D3 receptor mutant mice," *Journal of Neuroscience*, vol. 18, no. 6, pp. 2231–2238, 1998.
- [59] J. D. Joseph, Y. M. Wang, P. R. Miles et al., "Dopamine autoreceptor regulation of release and uptake in mouse brain slices in the absence of D(3) receptors," *Neuroscience*, vol. 112, no. 1, pp. 39–49, 2002.
- [60] Y. Goto and P. O'Donnell, "Delayed mesolimbic system alteration in a developmental animal model of schizophrenia," *Journal of Neuroscience*, vol. 22, no. 20, pp. 9070–9077, 2002.
- [61] K. Liu, C. Bergson, R. Levenson, and C. Schmauss, "On the origin of mRNA encoding the truncated dopamine D3-type receptor D3nf and detection of D3nf-like immunoreactivity in human brain," *Journal of Biological Chemistry*, vol. 269, no. 46, pp. 29220–29226, 1994.
- [62] E. W. Daenen, G. Wolterink, J. A. Van Der Heyden, C. G. Kruse, and J. M. Van Ree, "Neonatal lesions in the amygdala or ventral hippocampus disrupt prepulse inhibition of the acoustic startle response; implications for an animal model of neurodevelopmental disorders like schizophrenia," *Neuropsychopharmacology*, vol. 13, no. 3, pp. 187–197, 2003.
- [63] G. Flores, G. K. Wood, D. Barbeau, R. Quirion, and L. K. Srivastava, "Lewis and Fischer rats: a comparison of dopamine transporter and receptors levels," *Brain Research*, vol. 814, no. 1-2, pp. 34–40, 1998.