Research Article

Boric Acid Disturbs Cell Wall Synthesis in *Saccharomyces cerevisiae*

Martin Schmidt, Jaron Z. Schaumberg, Courtney M. Steen, and Michael P. Boyer

Biochemistry and Nutrition, Des Moines University, 3200 Grand Avenue, Des Moines, IA 50312, USA

Correspondence should be addressed to Martin Schmidt, mschmidt@dmu.edu

Received 6 August 2010; Revised 20 October 2010; Accepted 16 November 2010

Academic Editor: Robert P. Gunsalus

Copyright © 2010 Martin Schmidt et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Boric acid (BA) has broad antimicrobial activity that makes it a popular treatment for yeast vaginitis in complementary and alternative medicine. In the model yeast *S. cerevisiae*, BA disturbs the cytoskeleton at the bud neck and impairs the assembly of the septation apparatus. BA treatment causes cells to form irregular septa and leads to the synthesis of irregular cell wall protuberances that extend far into the cytoplasm. The thick, chitin-rich septa that are formed during BA exposure prevent separation of cells after abscission and cause the formation of cell chains and clumps. As a response to the BA insult, cells signal cell wall stress through the Slt2p pathway and increase chitin synthesis, presumably to repair cell wall damage.

1. Introduction

In the right amounts, boron is an essential nutrient for animals, plants, and fungi [1–3]. However, at high concentrations boric acid (BA) becomes an effective poison that is widely used for the killing of diverse organisms ranging from bacteria to rodents [4]. In medicine, BA is used as an alternative treatment for vaginal yeast infections [5]. While the molecular details of BA action on yeast remain unclear, it was recently shown that BA interferes with morphogenesis, to the effect that it inhibits the transition from the yeast to the hyphal form of the pathogenic yeast *C. albicans* [6]. Because the ability to switch to hyphal growth is an important virulence factor in *C. albicans* [7], suppression of such elongated growth by BA may in part explain its therapeutic effect. The present study was undertaken to assess the effect of BA on morphogenesis and cell wall synthesis in yeast, using the well-established model organism *Saccharomyces cerevisiae* as a study subject.

In *S. cerevisiae*, morphogenesis and cell wall synthesis depend on the correct assembly of cytoskeletal proteins. To guide cell wall synthesis during cytokinesis, a ring of septin filaments forms during the G1 phase of the cell cycle and is subsequently completed into a contractile actomyosin

ring (CAR) by the addition of myosin and actin, among other proteins [8, 9]. To complete abscission, the last phase of cytokinesis, the cells first separate mother and daughter cells with a chitin primary septum. The deposition of glucan and mannoprotein-rich cell wall material on the mother and daughter side of the primary septum later completes the trilaminar septum that can be observed under normal culture conditions. A disturbance in the assembly of the septation apparatus—the cohesive functional unit that constructs the primary septum—leads to the formation of highly aberrant septa [10, 11]. The septa formed under these conditions do not allow for the separation of cells after cytokinesis, leading to the formation of chains and clumps of misshaped cells. Based on the observation that BA causes such clumping and chain formation in *S. cerevisiae*, the present study was initiated to assess the influence of BA on the function of the septation apparatus.

2. Methods

2.1. Strains and Culture Conditions. Growth media and culture conditions were as described in [12]. Strains are listed in Table 1.

TABLE 1: Strains.

Strain	Genotype	Source
BY4742	<i>MATα his3Δ1 leu2Δ 0 lys2Δ0 ura3Δ0</i>	Invitrogen
BY4742 <i>slt2::kanMX6</i>	<i>MATα his3Δ1 leu2Δ0 lys2Δ 0 ura3Δ 0 <i>slt2::kanMX6</i></i>	Invitrogen
ECY46-1-8D	<i>MATα ura3-52 lys2-801 ade2-101 <i>trp1-Δ63 his3-Δ200 leu2-Δ 1chs1::HIS3</i></i>	[13]
YMS415	<i>MATα ura3-52 lys2-801 ade2-101 <i>trp1-Δ63 his3-Δ200 leu2-Δ1 CHS3::3HA-HIS3</i></i>	This study
YPH499	<i>MATα ura3-52 lys2-801 ade2-101 <i>trp1-Δ63 his3-Δ200 leu2-Δ1</i></i>	[14]

Strain YMS415 (*CHS3::HA-HIS3*) was constructed by the short flanking homology method [15]. The *CHS3::HA-HIS3* fragment was amplified by PCR from plasmid pFA6a-3HA-His3MX6 with primers 5'-TATTCTCAATCGGAAGGA-GGAAAGTGACTCCTTCGTTGCAGGTGGAGGTGG-AGGTGGAGGTGGACGGATCCCCGGGTTAATTAA-3' and 5'-TCAACTTGTAAGTATCACAGTAAAAAT-ATTTTCATACTGTGAATTTCGAGCTCGTTTAAAC-3'. Integration of the fragment was verified by PCR and western blotting. Transformants showed a wild-type like distribution of chitin in calcofluor white stained preparations, demonstrating full functionality of the Chs3p-HA fusion protein.

2.2. Determination of BA Sensitivity. A visual representation of BA sensitivity was obtained by serially diluting an overnight culture of yeast grown in YPD and spotting 5 μ L of cell dilutions on YPD plates with the indicated concentrations of BA. Growth was analyzed after 3-day incubation at 30°C. Minimal inhibitory concentrations were determined according to CLSI standards with the broth microdilution assay [16].

2.3. Staining Procedures and Fluorescence Microscopy. Viability staining of yeast cultures was performed by incubating cells in 0.2 mg/mL methylene blue in 50 mM KH₂PO₄ for 5 minutes at RT.

In order to visualize the distribution of chitin and glucan in the cell wall, cells were washed and incubated for 5 minutes in 0.01% calcofluor white or 0.5% aniline blue, respectively [17]. Fluorescence was observed with a standard diamidino-2-phenylindol (DAPI) filter set (Zeiss).

Filamentous actin was visualized by Alexa 568-phalloidin staining of yeast cells. Fifty mL yeast cultures at a titer of 1–5 $\times 10^6$ cells/mL in YPD were fixed by the addition of formaldehyde to a final concentration of 4% and incubated for 10 minutes at room temperature. Cells were pelleted and incubated for 1 hour in phosphate buffered saline (PBS; 137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, 1.76 mM KH₂PO₄, pH 7.4) with 4% formaldehyde. After 2 washes in PBS, cells were suspended in 100 μ L PBS with 10 μ L of an Alexa 568-phalloidin solution (6.6 μ M in methanol; Invitrogen) and 1 μ L of 1 mg/mL calcofluor white. Cells were incubated for 1 hour in the dark and washed twice

with PBS. Cells were then pelleted and taken up in 50 μ L ProLong Antifade reagent (Invitrogen) before mounting on slides.

GFP-tagged Cdc3p and Myo1p were observed with the GFP-filter set (Zeiss) in cells transformed with pRS316CDC 3GFP [12] and pMS55 [11].

2.4. Electron Microscopy. Electron microscopic examination of yeast cell walls followed a previously published protocol [18].

2.5. Western Blotting. Analysis of Slt2p phosphorylation and total Slt2p by western blotting followed established protocols [19, 20]. As a loading control, an antibody directed against Tub4p (goat χ -tubulin γ K18, Santa Cruz Biotechnology) was used at 1/1,000 dilution. For the determination of Chs3p-HA, cell membranes were isolated according to Orlean [21]. One volume of mercaptoethanol-free 2x sample buffer with 2% SDS (Bio-Rad) was added to the protein extracts and samples were loaded on 6% SDS polyacrylamide gels without boiling. Incubations with antibodies were performed as described above [18]. Antibodies used were a 1/5,000 dilution of anti-HA/1/5,000 dilution of goat antimouse HRP (Santa Cruz Biotechnology). Four independent experiments were performed and representative results are shown. Membranes were stained in 0.02% Coomassie blue R250 in 40% Methanol, 5% acetic acid to serve as loading controls.

2.6. Enzyme Measurements. For the determination of enzyme activities, cell membranes were isolated as previously described [21].

Determination of glucan synthase activity followed a protocol by Mol and coworkers [22]. A 40 μ L glucan synthase assay contained 20 μ L of membrane suspension at a protein concentration of 1 mg mL⁻¹, 5 mM ¹⁴C-UDP-glucose (activity 1 $\times 10^9$ cpm mmol⁻¹), 75 mM Tris-Cl pH 7.5, 25 mM KF, and—when indicated—with 20 μ M GTP- γ -S for the determination of maximal GS activity. The reaction was incubated for 1 hour at 30°C and then stopped by adding 1 mL 10% trichloroacetic acid (TCA). Reaction mixtures were filtered through a type A/E glass fiber filter (Pall). Filters were washed twice with 1 mL 10% TCA and four times with 70% ethanol. Filters were dried and taken up in 10 mL cytoscent ES scintillation fluid (ICN), and activity was recorded in a scintillation counter. Activities were calculated as c.p.m. incorporated h⁻¹ μ g protein⁻¹.

Chitin synthesis was determined as described by Choi and Cabib [23]. To measure chitin synthase 2 and 3 activities, experiments were performed in the *chs1* deletion strain ECY46-1-8D. The *chs1* deletion was found to have no influence on BA sensitivity (Figure 1(a)). The chitin synthase assay mixture contained 20 μ L of membrane suspension at 1 mg protein mL⁻¹, 5 μ L of 0.5 M Tris/Cl pH 7.8, 5 μ L of 20 mM cobalt acetate, 5 μ L of 10 mM ¹⁴C-UDP-GlcNAc (5000 c.p.m. μ L⁻¹), and 2 μ L of trypsin solutions (Sigma) at concentrations from 0.25 to 2.0 mg mL⁻¹ in a total volume of 46 μ L. For determination of chitin synthase 3 activity, 5 μ L of water were substituted with 5 μ L of 50 mM

nickel acetate. Proteolysis was stopped after incubating for 15 minutes at 30°C by adding 2 μ L of a soybean trypsin inhibitor solution at 1.5x the concentration of the trypsin solution. Chitin synthesis was initiated by adding 2 μ L of 0.8 M GlcNAc. After incubating for 60 minutes at 30°C, chitin synthesis was stopped by adding 1 mL 10% TCA. Reaction mixtures were filtered, washed, and assayed in cytosol fluid as described above. Assays containing cobalt only show activities of chitin synthases 2 and 3 while cobalt/nickel assays show the activity of chitin synthase 3. Chitin synthase 2 activity was calculated from the difference of both assays. Two independent experiments were conducted for chitin and glucan synthesis.

3. Results

3.1. BA Acts as a Fungistatic Agent That Inhibits Cytokinesis.

Vital staining of strains YPH499 and ECY46-1-8D with 0.05% methylene blue showed that BA concentrations between 0.1 and 0.4% do not severely reduce cell viability (Figure 1(a)). In the examined range around the minimal inhibitory concentration of 0.31%, BA thus functions as a fungistatic agent that slows down proliferation but does not kill cells. Note that the decline in viability in *chs1* deletion strain ECY46-1-8D parallels the decline in wildtype viability. Due to the previously reported lysis of daughter cells in *chs1* mutants [24], strain ECY46-1-8D shows a higher fraction of dead cells in all of the examined samples.

Clumping and chain formation of cells occurs at BA concentrations above 0.2%, with the most striking effect observed at 0.4% (Figure 1(b)). This clumping is the hallmark of a cytokinesis defect that causes daughter cells to remain attached to mother cells. A spot test of BY4742 cells on YPD plates showed that no growth occurs at BA concentrations above 0.5%—although viable cells can be retrieved even after 10 days of incubation under these conditions (data not shown).

3.2. BA Exposure Leads to Abnormal Deposition of Chitin and Glucan.

Staining of chitin and glucan in walls of cells grown with 0.4% BA revealed a buildup of cell wall material at bud necks, particularly in cell chains (Figure 2).

3.3. BA Exposure Interferes with the Localization of the Septation Apparatus.

In order to characterize the cytokinesis defect in BA-treated cells, the localization of key morphogenetic proteins at the bud neck was examined by fluorescence microscopy. It was found that BA influences the localization of the septin Cdc3, the cytokinetic myosin Myo1p, and the ring of filamentous actin that form sequentially in preparation for cytokinesis. Increasing concentrations of BA leads to the formation of Cdc3GFP rings that are disorganized, uneven, and not centered at the bud neck. Moreover, BA causes the formation of Cdc3GFP patches at sites other than the bud neck (Figure 3).

Imaging of Myo1GFP and actin shows that the formation of the CAR is impaired in a manner similar to the disturbance in septin ring organization. Increasing BA concentrations

FIGURE 1: (a) Viability of strains YPH499 and ECY46-1-8D in the concentration range of 0–0.4% BA as determined by methylene blue staining. (b) Changes in morphology of YPH499 during exposure to BA. Cultures grown with BA show clumps and chains of incompletely separated cells. The scale bar represents 5 μ m.

FIGURE 2: Analysis of chitin and glucan distribution in BA-treated cells: fluorescence of calcofluor white (top) and aniline blue (bottom) stained cells. At BA concentrations of 0.4%, chitin-rich—and to a lesser extent, glucan-rich—material accumulates at the bud neck. The thickening of septa is evident particularly between cells in a chain.

FIGURE 3: Appearance of septin (Cdc3-GFP) rings in BA-treated cells. Starting at a BA concentration of 0.2% and worsening at 0.4%, septin rings show irregular thickness and wide spacing between the mother and the daughter-side rings. BA also induces ectopic localization of septin patches (arrowheads).

lowers the fraction of cells with Myo1GFP rings at the bud neck from $49 \pm 5\%$ at 0% to $25 \pm 11\%$ at 0.2% to $11 \pm 11\%$ at 0.4%. In addition, Myo1GFP rings formed under the influence of BA are often irregular in shape and at high concentrations (0.4%) ectopic localization of Myo1GFP patches at sites other than the bud neck is common (Figure 4(a)).

BA also impairs the assembly of the actin ring at the bud neck, which is the last component to be incorporated into the CAR before contraction starts (Figure 4(b)). Actin rings formed under the influence of BA become blurry, faint, and irregular in thickness. Under these conditions, it can be observed that some actin rings fail to localize to the narrowest point of the bud neck. In some cases, BA-treated cells even attempt cytokinesis without a detectable actin ring. Due to the low abundance of cells with actin rings in culture ($< 2\%$ in controls), quantification of rings yielded no statistically significant data.

3.4. BA Causes Ultrastructural Abnormalities in the Cell Wall. Cultures of cells grown with 0.3% and 0.4% BA were examined by electron microscopy using a protocol that allowed for the visualization of chitin (Figure 5). The analysis of BA-treated cells showed massive abnormalities in the cell wall, particularly in the area of the septum, worse at 0.4% than at 0.3%. At these concentrations, BA prevented the construction of a single, straight chitin septum to separate mother and daughter cells. Instead, the cells synthesized protuberances extending far into the cytosol and large irregular septa, often at ectopic locations at the cell periphery.

3.5. Function of Slt2 in BA-Stressed Cells. In order to assess BA-induced cell wall integrity signaling, the amounts and the phosphorylation status of the signaling kinase Slt2p were determined by western blotting (Figure 6(a)). Phosphorylation of Slt2p increases with escalating concentrations of BA, suggesting that BA induces a cell wall integrity stress response. The dramatic increase in phosphorylation of Slt2p is accompanied by a weak increase in protein expression. Furthermore, it was determined that a *slt2* deletion mutant is sensitive to BA (Figure 6(b)), suggesting that Slt2p-signaling improves BA resistance by causing a transcriptional response to BA stress.

3.6. BA Stimulates Chitin Synthesis. Western blotting showed a nonlinear correlation between BA concentration and abundance of Chs3p-HA (Figure 7(a)). While at BA concentrations of 0.1 and 0.2% the amount of Chs3p-HA decreases, there is an increase of Chs3p-HA at 0.4% BA. The increase of Chs3p-HA parallels the increase of chitin-rich septa visible in calcofluor white-stained cultures.

The multiple posttranslational and protein targeting processes involved in Chs3p and Chs2p activation [9] necessitate enzymatic determination of chitin synthase activities in addition to detection of protein amounts. Figure 7(b) shows that BA-induced changes in chitin synthase activities mirror the changes in Chs3p-HA amount (Figure 7(a)) and

FIGURE 4: Appearance of the actomyosin rings in BA-treated cells. (a) Localization of Myo1GFP. Increasing concentrations of BA leads to irregular localization of Myo1GFP. Images show samples of normal, uneven rings not centered at the bud neck (u) and ectopic localization (e) of Myo1GFP patches. Note that at 0.4% BA Myo1GFP forms rings at locations other than the bud site that appear as bands in the selected images (arrows). (b) Localization of actin. The selection of images shows that increasing concentrations of BA causes actin rings to appear blurred (b) and uneven/not centered at the bud neck (u).

bud neck thickness (Figure 2). While at 0.1% and 0.2% BA decreases chitin synthase activities, presence of 0.3% and 0.4% BA increases chitin synthesis. At 0.4% BA, activities of both chitin synthase 2 and 3 exceed activities in control cultures.

A measurement of glucan synthase activity in the membranes of strain YPH499 showed a constant decline with increasing BA concentrations (Figure 7(b)). The addition of GTP- γ S activates the GTP binding protein Rho1p and maximizes Fks1p activity. Neither maximal nor physiological (without addition of GTP- γ S) glucan synthase activity showed an increase at high concentrations of BA.

4. Discussion

Living organisms are constantly exposed to boron, a mineral that is abundant in soil and water. Boron is a weak acid and, at physiological pH, is present mostly as boric acid

FIGURE 5: Electron micrographs of YPH499 cells grown in the presence of 0.3% and 0.4% BA. Cell wall chitin appears electron translucent. (a) No BA. Left: a mother-daughter pair; Middle: a close-up of the bud neck; Right: a completed trilaminar septum. Note the thin layer of chitin (primary septum; arrow) in the center of the structure. (b) Cells grown with 0.3% BA. Left: a mother-daughter pair; Middle: a close-up of the bud neck; Right: a completed septum. (c) Examples of other cell wall abnormalities in cells grown with 0.3% BA. Left: cell wall thickening and lacunae at the cell periphery; Middle: massive cell wall protuberances; Right: a septum forming at an ectopic location. (d) Cells grown with 0.4% BA. Left: a mother-daughter pair; Middle: a close-up of the bud neck; Right: a completed default septum. (e) Examples of other cell wall abnormalities in cells grown with 0.4% BA. Left: a chain of cells connected by default septa; Middle: a large cell wall structure spanning the entire length of the cell; Right: chitin-rich cell wall material deposited at the bud neck with two embedded aberrant septa (arrows).

FIGURE 6: Slt2p-signaling in BA-treated cells. (a) Detection of phosphorylated Slt2p by western blotting in strain YPH499. Phosphorylation of Slt2p increases with escalating concentrations of BA. The total amount of Slt2p increases only slightly under the same condition. Gamma tubulin (Tub4p) detection by western blotting and the Coomassie blue-stained membrane are shown as loading controls. (b) BA sensitivity of an *slt2* mutant strain compared to its WT BY4742 as determined by a spot assay.

(H₃BO₃; BA). For each organism, there is an optimal BA concentration. Too little BA causes symptoms of deficiency while too much BA has a poorly defined cytotoxic effect [25]. Particularly the toxic effects of boron overload have led to a variety of applications for BA and related compounds, ranging from pest control to the treatment of vaginal yeast infections.

Boron toxicity is not yet understood. Molecular and *in vitro* studies suggest that boron participates in both enzymatic and nonenzymatic processes. The biological function of BA might be due to its reactivity with *cis*-hydroxyl groups on carbohydrate molecules [26]. Examples for boron-dependent reactions include plant hormone-sensitive NADH oxidase activity [27], the crosslinking of plant cell wall

FIGURE 7: Glucan and chitin synthase activities in BA-treated cells. (a) Abundance of HA-tagged Chs3p in strain YMS415 treated with BA. Note the increase of Chs3p-HA at 0.4% BA. (b) Chitin synthase 2 and 3 activities in strain ECY46-1-8D. After a decline at lower BA concentrations, chitin synthase activities increase at 0.4% (c) Glucan synthase activities. Solid bars indicate activity after activation by GTP- χ S. Both maximal and physiological glucan synthase activities decline with increasing BA concentrations.

carbohydrates [28], and the mineralization of bone [29]. The molecular effects of boron lead to developmental defects in animals, particularly in the formation of the skeleton [30, 31]. These teratogenic effects of boron might be caused by direct or indirect inhibition of histone deacetylase activity [32] and a shift in Hox gene expression [31].

However, none of the above observations about BA toxicity serve to explain its effect on yeast—an organism that does not undergo genomic imprinting or Hox-dependent development. The only published data about the BA effect on yeast come from a recent study of the yeast *C. albicans* that suggested that BA impairs oxidative metabolism [6]. While this observation in itself is interesting and deserves further study, the authors also show that BA directly or indirectly influences the morphology of *C. albicans*. The present study expands on this observation by showing that BA disturbs morphogenesis in the model yeast *Saccharomyces cerevisiae*, particularly during cytokinesis.

The data presented here show that BA impairs the formation of the primary septum in *S. cerevisiae*—a defect that can be explained by incorrect assembly of the cytoskeleton at the bud neck. The molecular machinery that constructs the chitin cell wall between mother and daughter cells, the septation apparatus, consists of chitin synthase 2, and a contractile actomyosin ring (CAR). The septation apparatus is assembled sequentially from myosin, chitin synthase 2 and actin on a scaffold of septins at the bud neck [8, 9]. Should the assembly of a functional septation apparatus fail, the cell is unable to construct an orderly primary septum by chitin synthase 2 and is forced to divide by depositing large amounts of chitin-rich cell wall material at the bud neck. The chitin in these so-called default septa is provided by chitin synthase 3 [10, 33, 34]. Since these irregular septum structures are resistant to degradation by chitinase, cells remain connected after cytokinesis and form cell chains and clumps [35]. In *S. cerevisiae*, the localization of the septum is determined by the position of the septins. A septin assembly that erroneously localizes to a site other than the bud neck will direct the formation of a septum-like structure at the respective location [34].

Our data show that in *S. cerevisiae* increasing concentrations of BA leads to an irregular assembly of the septin scaffold, an inability to position the Myo1 ring, and a failure to correctly assemble an actin ring at the bud neck. The aberrant localization and irregular appearance of the septin Cdc3GFP in BA-treated cells is an important key to understanding the septation defect. We propose that BA causes problems with the assembly of the septin scaffold which later impair the localization and function of the CAR, resulting in the formation of highly irregular cell wall structures.

Like other threats to the integrity of the cell wall, septation defects trigger cell wall integrity signaling through the protein kinase C (PKC) pathway. Under these conditions, cell wall integrity signaling leads to hyperphosphorylation of the PKC downstream effector Slt2p [36–38], accompanied with a much weaker increase in Slt2p amount [39]. This will ultimately lead to the activation of cell wall repair enzymes, most notably chitin synthase 3 [36, 40]. The

present study shows that BA activates yeast cell wall integrity signaling pathways as evidenced by BA concentration-dependent phosphorylation of Slt2p. Presumably in response to cell wall integrity signaling, cells increase the activity of the cell wall repair enzyme chitin synthase 3 in a BA concentration-dependent manner. The somewhat unusual pattern of chitin synthase induction—a decline at low concentrations followed by an increase at higher doses—should be interpreted based on the impact boric acid on growth. We propose that at concentrations below the MIC where the impact on growth is measurable but weak, boric acid stress reduces cell wall synthesis activity along with other metabolic activities. Once boric acid stress exceeds a tolerable limit—at concentrations above the MIC—the cell responds forcefully by induction of stress survival mechanisms such as chitin synthase 3-mediated cell wall reinforcement.

It is worth noting that the activity of the chitin septum-forming chitin synthase 2 also increases during BA exposure. It is evident that the increase in chitin synthase 2 activity correlates well with the increased number and size of chitin septa in BA-treated cells. However, since the regulation of chitin synthase 2 activity is not well understood, we dare not hazard an explanation for this phenomenon.

The data presented in this study show that in *S. cerevisiae* BA disturbs the localization of the contractile actomyosin ring secondary to causing irregularities in the septin scaffold at the bud neck. In agreement with the reviewed literature we propose that the aberrant localization of the septins ultimately impedes the formation of the primary septum, which leads to the synthesis of thick, chitin-rich default septa. In addition, the localization of septins at sites other than the bud neck explains the synthesis of cell wall protuberances that should be interpreted as incomplete ectopic septa [34]. Furthermore, our data show that a BA-induced septation defect, just like other septation defects, triggers cell wall integrity signaling through the Pkc1-Slt2 pathway and results in increased chitin synthase 3 activity.

Acknowledgment

This work was supported by a grant from the Iowa Osteopathic Education and Research Foundation (IOER).

References

- [1] M. Benderdour, T. Bui-Van, A. Dicko, and F. Belleville, “In vivo and in vitro effects of boron and boronated compounds,” *Journal of Trace Elements in Medicine and Biology*, vol. 12, no. 1, pp. 2–7, 1998.
- [2] A. Fleischer, C. Titel, and R. Ehwald, “The boron requirement and cell wall properties of growing and stationary suspension-cultured *Chenopodium album* L cells,” *Plant Physiology*, vol. 117, no. 4, pp. 1401–1410, 1998.
- [3] A. Bennett, R. I. Rowe, N. Soch, and C. D. Eckhart, “Boron stimulates yeast (*Saccharomyces cerevisiae*) growth,” *Journal of Nutrition*, vol. 129, no. 12, pp. 2236–2238, 1999.
- [4] E. Siegel and S. Wason, “Boric acid toxicity,” *Pediatric Clinics of North America*, vol. 33, no. 2, pp. 363–367, 1986.
- [5] S. M. Prutting and J. D. Cervený, “Boric acid vaginal

- suppositories: a brief review," *Infectious Diseases in Obstetrics and Gynecology*, vol. 6, no. 4, pp. 191–194, 1998.
- [6] F. De Seta, M. Schmidt, B. Vu, M. Essmann, and B. Larsen, "Antifungal mechanisms supporting boric acid therapy of *Candida* vaginitis," *Journal of Antimicrobial Chemotherapy*, vol. 63, no. 2, pp. 325–336, 2009.
 - [7] J. D. Sobel, G. Muller, and H. R. Buckley, "Critical role of germ tube formation in the pathogenesis of candidal vaginitis," *Infection and Immunity*, vol. 44, no. 3, pp. 576–580, 1984.
 - [8] J. B. Moseley and B. L. Goode, "The yeast actin cytoskeleton: from cellular function to biochemical mechanism," *Microbiology and Molecular Biology Reviews*, vol. 70, no. 3, pp. 605–645, 2006.
 - [9] G. Lesage and H. Bussey, "Cell wall assembly in *Saccharomyces cerevisiae*," *Microbiology and Molecular Biology Reviews*, vol. 70, no. 2, pp. 317–343, 2006.
 - [10] M. Schmidt, B. Bowers, A. Varma, D. H. Roh, and E. Cabib, "In budding yeast, contraction of the actomyosin ring and formation of the primary septum at cytokinesis depend on each other," *Journal of Cell Science*, vol. 115, no. 2, pp. 293–302, 2002.
 - [11] M. Schmidt, A. Varma, T. Drgon, B. Bowers, and E. Cabib, "Septins, under Cla4p regulation, and the chitin ring are required for neck integrity in budding yeast," *Molecular Biology of the Cell*, vol. 14, no. 5, pp. 2128–2141, 2003.
 - [12] D. Burke, D. Dawson, and T. Stearns, *Methods in Yeast Genetics*, Cold Spring Harbor Press, Cold Spring Harbor, NY, USA, 2000.
 - [13] L. B. Crotti, T. Drgon, and E. Cabib, "Yeast cell permeabilization by osmotic shock allows determination of enzymatic activities *in situ*," *Analytical Biochemistry*, vol. 292, no. 1, pp. 8–16, 2001.
 - [14] R. S. Sikorski and P. Hieter, "A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*," *Genetics*, vol. 122, no. 1, pp. 19–27, 1989.
 - [15] M. S. Longtine, A. McKenzie, D. J. Demarini et al., "Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*," *Yeast*, vol. 14, no. 10, pp. 953–961, 1998.
 - [16] P. A. Wayne, *National Committee for Clinical Laboratory Standards. Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts; Approved Standard*, NNCLS Document M27-A3. Clinical and Laboratory Standards Institute, 3rd edition, 2008.
 - [17] M. Schmidt, T. Drgon, B. Bowers, and E. Cabib, "Hyperpolarized growth of *Saccharomyces cerevisiae* *clb1*^{P212S} and *cla4* mutants weakens cell walls and renders cells dependent on chitin synthase 3," *FEMS Yeast Research*, vol. 8, no. 3, pp. 362–373, 2008.
 - [18] M. Schmidt, "Survival and cytokinesis of *Saccharomyces cerevisiae* in the absence of chitin," *Microbiology*, vol. 150, no. 10, pp. 3253–3260, 2004.
 - [19] H. Martín, J. M. Rodríguez-Pachón, C. Ruiz, C. Nombela, and M. Molina, "Regulatory mechanisms for modulation of signaling through the cell integrity Slt2-mediated pathway in *Saccharomyces cerevisiae*," *Journal of Biological Chemistry*, vol. 275, no. 2, pp. 1511–1519, 2000.
 - [20] G. Griffioen, S. Swinnen, and J. M. Thevelein, "Feedback inhibition on cell wall integrity signaling by Zds1 involves Gsk3 phosphorylation of a cAMP-dependent protein kinase regulatory subunit," *Journal of Biological Chemistry*, vol. 278, no. 26, pp. 23460–23471, 2003.
 - [21] P. Orlean, "Two chitin synthases in *Saccharomyces cerevisiae*," *Journal of Biological Chemistry*, vol. 262, no. 12, pp. 5732–5739, 1987.
 - [22] P. C. Mol, H. M. Park, J. T. Mullins, and E. Cabib, "A GTP-binding protein regulates the activity of (1–3)- β -glucan synthase, an enzyme directly involved in yeast cell wall morphogenesis," *Journal of Biological Chemistry*, vol. 269, no. 49, pp. 31267–31274, 1994.
 - [23] W. J. Choi and E. Cabib, "The use of divalent cations and pH for the determination of specific yeast chitin synthetases," *Analytical Biochemistry*, vol. 219, no. 2, pp. 368–372, 1994.
 - [24] E. Cabib, A. Sburlati, B. Bowers, and S. J. Silverman, "Chitin synthase 1, an auxiliary enzyme for chitin synthesis in *Saccharomyces cerevisiae*," *Journal of Cell Biology*, vol. 108, no. 5, pp. 1665–1672, 1989.
 - [25] M. Park, Q. Li, N. Shcheynikov, S. Muallem, and W. Zeng, "Borate transport and cell growth and proliferation: not only in plants," *Cell Cycle*, vol. 4, no. 1, pp. 24–26, 2005.
 - [26] D. H. Kim, K. F. Faull, A. J. Norris, and C. D. Eckhart, "Borate-nucleotide complex formation depends on charge and phosphorylation state," *Journal of Mass Spectrometry*, vol. 39, no. 7, pp. 743–751, 2004.
 - [27] R. Barr, M. Bottger, and F. L. Crane, "The effect of boron on plasma membrane electron transport and associated proton secretion by cultured carrot cells," *Biochemistry and Molecular Biology International*, vol. 31, no. 1, pp. 31–39, 1993.
 - [28] M. A. O'Neill, T. Ishii, P. Albersheim, and A. G. Darvill, "Rhamnogalacturonan II: structure and function of a borate cross-linked cell wall pectic polysaccharide," *Annual Review of Plant Biology*, vol. 55, pp. 109–139, 2004.
 - [29] R. E. Chapin, W. W. Ku, M. A. Kenney et al., "The effects of dietary boron on bone strength in rats," *Fundamental and Applied Toxicology*, vol. 35, no. 2, pp. 205–215, 1997.
 - [30] J. W. Cherrington and N. Chernoff, "Periods of vertebral column sensitivity to boric acid treatment in CD-1 mice in utero," *Reproductive Toxicology*, vol. 16, no. 3, pp. 237–243, 2002.
 - [31] N. Wéry, M. G. Narotsky, N. Pacico, R. J. Kavlock, J. J. Picard, and F. Gofflot, "Defects in cervical vertebrae in boric acid-exposed rat embryos are associated with anterior shifts of box gene expression domains," *Birth Defects Research Part A—Clinical and Molecular Teratology*, vol. 67, no. 1, pp. 59–67, 2003.
 - [32] F. Di Renzo, G. Cappelletti, M. L. Broccia, E. Giavini, and E. Menegola, "Boric acid inhibits embryonic histone deacetylases: a suggested mechanism to explain boric acid-related teratogenicity," *Toxicology and Applied Pharmacology*, vol. 220, no. 2, pp. 178–185, 2007.
 - [33] J. A. Shaw, P. C. Mol, B. Bowers et al., "The function of chitin synthases 2 and 3 in the *Saccharomyces cerevisiae* cell cycle," *Journal of Cell Biology*, vol. 114, no. 1, pp. 111–123, 1991.
 - [34] D. H. Roh, B. Bowers, M. Schmidt, and E. Cabib, "The septation apparatus, an autonomous system in budding yeast," *Molecular Biology of the Cell*, vol. 13, no. 8, pp. 2747–2759, 2002.
 - [35] M. J. Kuranda and P. W. Robbins, "Chitinase is required for cell separation during growth of *Saccharomyces cerevisiae*," *Journal of Biological Chemistry*, vol. 266, no. 29, pp. 19758–19767, 1991.
 - [36] H. Martín, M. C. Castellanos, R. Cenamor, M. Sánchez, M. Molina, and C. Nombela, "Molecular and functional characterization of a mutant allele of the mitogen-activated protein-kinase gene SLT2(MPK1) rescued from yeast autolytic mutants," *Current Genetics*, vol. 29, no. 6, pp. 516–522, 1996.
 - [37] D. E. Levin, "Cell wall integrity signaling in *Saccharomyces*

- cerevisiae*,” *Microbiology and Molecular Biology Reviews*, vol. 69, no. 2, pp. 262–291, 2005.
- [38] J. F. Rodríguez-Quinones and J. R. Rodríguez-Medina, “Differential gene expression signatures for cell wall integrity found in chitin synthase II (*chs2Δ*) and myosin II (*myo1Δ*) deficient cytokinesis mutants of *Saccharomyces cerevisiae*,” *BMC Research Notes*, vol. 2, article no. 87, 2009.
- [39] H. de Nobel, C. Ruiz, H. Martin et al., “Cell wall perturbation in yeast results in dual phosphorylation of the Slt2/Mpk1 MAP kinase and in an Slt2-mediated increase in *FKS2-lacZ* expression, glucanase resistance and thermotolerance,” *Microbiology*, vol. 146, no. 9, pp. 2121–2132, 2000.
- [40] R. H. Valdivia and R. Schekman, “The yeasts Rho1p and Pkc1p regulate the transport of chitin synthase III (Chs3p) from internal stores to the plasma membrane,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 100, no. 18, pp. 10287–10292, 2003.