ON NONAMENABLE GROUPS

SU-SHING CHEN

School of Mathematics
Georgia Institute of Technology
Atlanta, Georgia 30332
U.S.A.

(Received April 10, 1978 and in revised form October 3, 1978)

ABSTRACT. A sufficient condition is given for a countable discrete group G to contain a free subgroup of two generators.

KEY WORDS AND PHRASES. *Nonamenable group, free group.*

AMS (MOS) SUBJECT CLASSIFICATION (1970) CODES. 22D05.

Given a topological group G , we denote by L the Banach algebra of all real valued bounded left uniformly continuous functions on G with the supremum norm. A mean m on L is a continuous, positive, linear functional such that $m(1) = 1$. A mean is called invariant if $m(f^g) = m(f)$ for every $f \in L$ and $g \in G$, where f^g is the translate of f by g .

G is called amenable if there exists an invariant mean on L . G has the fixed point property if whenever G acts on a compact convex set Q affinely in a locally convex topological vector space E , then G has a fixed point in Q [2].

It is well known that G is amenable if and only if G has the fixed point property for any topological group G .

In [4], von Neumann proved that if G has a free subgroup of two generators then G is not amenable and conjectured that the converse is true. In this paper, we shall give a sufficient condition for a discrete group G to contain a free subgroup of two generators. This result may be interesting to the investigation of von Neumann's conjecture.

Let ϕ be an affine transformation of a compact convex set Q in a locally convex topological vector space E . Then ϕ has a fixed point in Q by the famous Tychonoff fixed point theorem. Furthermore, one can prove easily that the fixed point set F_ϕ of an affine transformation ϕ of Q is a compact convex subset of Q .

Let us consider a discrete group G acting affinely on Q . The fixed point set F_ϕ of each element ϕ of G coincides with the fixed point set $F_{\phi^{-1}}$ of the inverse ϕ^{-1} . An element ϕ of G is said to be attractive if for each weak neighborhood U_ϕ of the fixed point set F_ϕ of ϕ , the orbit $\{\phi^n(S) \mid n \in \mathbb{Z}\}$ of any compact convex subset S in $Q - U_\phi$ converges to the fixed point set F_ϕ of ϕ , that is, there is a positive integer N such that for all $|n| > N$, $\phi^n(S) \subset U_\phi$. An element ϕ of G is said to be weakly attractive if, for each weak neighborhood U_ϕ of the fixed point set F_ϕ of ϕ , there is a positive integer N' such that for all $n \in \mathbb{Z}^{*(1)}$, $\phi^{nN'}(S) \subset U_\phi$. It is obvious that an attractive element ϕ of G is weakly attractive. [Note: (1) $\mathbb{Z}^* = \mathbb{Z} - \{0\}$]

THEOREM. If a discrete group G acts on a compact convex set of Q of a locally convex topological vector space E affinely such that G contains at least two weakly attractive elements without common fixed points, then G contains a free subgroup of two generators.

PROOF. Let ϕ and ψ be two weakly attractive elements of G . Then the fixed point sets F_ϕ and F_ψ are disjoint. By the separation theorem [6], there exist

a linear functional L on E and real numbers c_1 and c_2 such that $Lx < c_1 < c_2 < Ly$ for every x in F_ϕ and every y in F_ψ . Without loss of generality, we may assume that $c_1 < 0 < c_2$.

Thus $K_1 = \{x \in Q \mid Lx < 0\}$ is a weak convex neighborhood of F_ϕ and $K_2 = \{x \in Q \mid Lx > 0\}$ is a weak convex neighborhood of F_ψ . The complements K_1^c and K_2^c of K_1 and K_2 respectively are compact and convex sets in $Q - K_1$ and $Q - K_2$. By the definition of weak attractiveness, there exist positive integers N' and N'' such that $\phi^{nN'}(K_1^c) \subset K_1$ and $\psi^{nN''}(K_2^c) \subset K_2$ for all $n \in \mathbb{Z}^*$. Let $s = \phi^{N'}$ and $t = \psi^{N''}$. Then the group F generated by s and t is a free group. In fact, for any relation $s^p t^q \dots = \text{id}$, we have $s^p t^q \dots(z) = z$ for each z in the hyperplane section $K_1^c \cap K_2^c = \{z \in Q \mid Lz = 0\}$ of Q . But clearly $Ls^p t^q \dots(z) \neq 0$, while $Lz = 0$. We have a contradiction.

COROLLARY. If a nonamenable discrete group G acts on a compact convex set Q of a locally convex topological vector space E affinely such that G contains all weakly attractive elements then G contains a free subgroup of two generators.

PROOF. This follows from the theorem and the non-fixed point property of nonamenable groups.

ACKNOWLEDGEMENT. The author is indebted to the referee for his comments.

REFERENCES

1. Furstenberg, H., A Poisson Formula for Semisimple Lie Groups, Ann. of Math. 77 (1963) 335-383.
2. Glasner, S., Proximal Flows, Lecture Notes in Math., Springer-Verlag, Vol. 517, 1976.
3. Greenleaf, F. P., Invariant Means on Topological Groups, Van Nostrand Math. Studies, 1969.
4. Klee, V., Some Topological Properties of Convex Sets, Trans. Amer. Math. Soc., 78 (1955) 30-45.
5. Von Neumann, J., Zur allgemeinen theorie des masses, Fund. Math. 13 (1929) 73-116.
6. Rudin, W., Functional Analysis, McGraw-Hill 1973.
7. Tits, J., Free Subgroups in Linear Groups, Jour. of Algebra 20 (1972) 250-270.