RESEARCH NOTES

PIS FOR n -COUPLED NONLINEAR SYSTEMS

S. K. DEY

Department of Mathematics
Eastern Illinois University
Charleston, Illinois 61920 U.S.A.

(Received October 15, 1979 and in revised form February 13, 1980)

ABSTRACT. A numerical algorithm dealing with solutions of equations with one variable may not be extended to solve nonlinear systems with n unknowns. Even when such extensions are possible, properties of these two similar algorithms are, in general, different. In [2] a perturbed iterative scheme (PIS) has been developed to solve nonlinear equations with one variable. Its properties with regard to nonlinear systems were analyzed in [1]. Here these properties were extended to n -coupled nonlinear systems.

KEY WORDS AND PHRASES. *Perturbed numerical iterations, nonlinear equations.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODES: 65H10, 65D15.

1. THE ALGORITHM.

In [2] a simple functional iterative scheme has been developed to solve nonlinear equations with one variable by adding a unique perturbation parameter to Picard's iterations. The algorithm was directly extended to nonlinear systems

[1] and convergence properties were analyzed using a special mapping called D-mapping [1]. We will study here that for n-coupled nonlinear systems analysis of convergence may be done in a similar way.

Let us consider a nonlinear system having n-coupled equations:

$$x^i = F^i(x^1, x^2, \dots, x^n) \quad (1.1)$$

where, $x^i = (x_1^i \ x_2^i \ \dots \ x_n^i)^T \in D^i \subset R^n$

$$F^i = (F_1^i \ F_2^i \ \dots \ F_n^i)^T \in D^i \subset R^n,$$

$i = 1, 2, \dots, n$ and $R^n =$ real n-dimensional space. Thus, each $F^i: D^1 \times D^2 \times \dots \times D^n \subset R^n \times R^n \times \dots \times R^n \rightarrow D^i$. Let $R = R^n \times R^n \times \dots \times R^n$ and $D = D^1 \times D^2 \times \dots \times D^n$. As before [1] we assume that (1.1) has a solution in D given by $x^* = (x^1,^* \ x^2,^* \ \dots \ x^n,^*)^T \in D$ where $x^{i,*} = (x_1^{i,*} \ x_2^{i,*} \ \dots \ x_n^{i,*})^T \in D^i$. Hence,

$$x^{i,*} = F^i(x^{1,*}, x^{2,*}, \dots, x^{n,*}) \quad (1.2)$$

In the element form, PIS is:

$$w_j^{i,k} = w_j^{i,k} + F_j^i(x^{1,k}, \dots, x^{i-1,k}, x_1^{i,k}, \dots, x_{j-1}^{i,k}, x_j^{i,k-1}, \dots, x_n^{i,k-1}, x_{j+1}^{i,k-1}, \dots, x^{n,k-1}) \quad (1.3)$$

$i = 1, 2, \dots, n$ and $j = 1, 2, \dots, n$.

To compute the perturbation parameters $w_j^{i,k}$ we assume that they are small and their squares may be neglected; also the functionals F_j^i are $\partial F_j^i / \partial x_j^i \neq 1$, and $\partial^2 F_j^i / \partial x_j^{i2}$ are bounded $\forall x^i \in D^i$. Then assuming convergence after (k-1) iterations we have:

$$w_j^{i,k} + F_j^{i,k} = F_j^i(x^{1,k}, \dots, x^{i-1,k}, x_1^{1,k}, \dots, x_{j-1}^{i,k}, w_j^{i,k} + F_j^{i,k}, x_{j+1}^{i,k-1}, \dots, x_n^{i,k-1}, x_{j+1}^{i,k-1}, \dots, x^{n,k-1}) \quad (1.4)$$

where $F_j^{i,k}$ is given by the second term of the right side of (1.3). Expanding the right hand side of (1.4) by Taylor's theorem and using the above assumptions we have:

$$w_j^{i,k} = (\bar{F}_j^{i,k} - F_j^{i,k}) / (1 - \partial_i F_j^{i,k}) \quad (1.5)$$

where $\bar{F}_j^{i,k} = F_j^i(x^{1,k} \dots x^{i-1,k}, x_1^i, \dots x_{j-1}^i, F_j^i, x_{j+1}^{i,k-1} \dots x_n^{i,k-1}, x^{i+1,k-1}, \dots, x^{n,k-1})$

and $\partial_j F_j^{i,k} = \left[\frac{\partial F_j^i}{\partial x_j^i} \right] x^{1,k} \dots x^{i-1,k}, x_1^i, \dots x_{j-1}^i, F_j^i, x_{j+1}^{i,k-1} \dots x_n^{i,k-1}, x^{i+1,k-1}, \dots x^{n,k-1}$

Once $w_j^{i,k}$ is known from (1.5) we use PIS to get:

$$x_j^{i,k} = w_j^{i,k} + F_j^{i,k} \quad (1.6)$$

for $i = 1, 2, \dots, n$, $j = 1, 2, \dots, n$. Comparing this equation with (1.2) it is clear that a necessary condition for convergence is:

$$\lim_{k \rightarrow \infty} |w_j^{i,k}| = 0, \forall i, j \quad (1.7)$$

If we write, $x^k = (x^{1,k} \dots x^{n,k})^T \in D$ and $w^k = (w^{1,k} \dots w^{n,k})^T \in R$ then

(1.6) may be expressed as:

$$x^k = w^k + F(x^k, x^{k-1}) \quad (1.8)$$

where $F : D \times D \subset R \times R \rightarrow D$

Now it is clear that $X^* = F(X^*, X^*)$ giving X^* as the fixed image of F on $D \times D$.

Also, if F is a D -mapping [1] on $D \times D$, (1.7) will be both necessary and sufficient condition for convergence of PIS.

Hence, after studying the convergence analysis of PIS in [1], it is now easy to see the same concept being extended for n -coupled nonlinear systems.

REFERENCES

1. Dey, S. K. Convergence Analysis of a Perturbed Iterative Scheme (PIS) for Solution of Nonlinear Systems, (to appear) International Journal of Math and Math Sciences.
2. Dey, S. K. Numerical Solution of Nonlinear Equations with Real and Complex Variables, Journal of the Franklin Inst., Vol. 307, No. 1, 1979.