ON RANK 4 PROJECTIVE PLANES

O. BACHMANN

Département de mathématiques
Ecole polytechnique fédérale
CH-1007 Lausanne, Suisse

(Received October 4, 1979)

ABSTRACT. Let a finite projective plane be called rank m plane if it admits a collineation group G of rank m , let it be called strong rank m plane if moreover $G_P = G_l$ for some point-line pair (P, l) . It is well known that every rank 2 plane is desarguesian (Theorem of Ostrom and Wagner). It is conjectured that the only rank 3 plane is the plane of order 2. By [1] and [7] the only strong rank 3 plane is the plane of order 2. In this paper it is proved that no strong rank 4 plane exists.

KEY WORDS AND PHRASES. Projective planes, rank 4 groups.

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE: 50 Geometry.

1. INTRODUCTION.

In [6] Kallaher gives restrictions for the order n of a finite rank 3 pro-

jective plane and conjectures that no such plane exists if $n \neq 2$. Let a finite projective plane be called a strong rank m projective plane if it admits a rank m collineation group G such that $G_P = G_l$ for some point-line pair (P, l) . By Bachmann [1] and Kantor [7] no strong rank 3 projective plane of order $n \neq 2$ exists. If the conjecture is true that for projective designs the representations on the points and on the blocks of an arbitrary transitive collineation group are similar (see Dembowski [2], p. 78), then every rank m projective plane is a strong rank m plane.

We shall prove in this article the following

THEOREM: No strong rank 4 projective plane exists.

To prove the Theorem we first divide the strong rank 4 planes into 3 classes (see Lemma 2 and 3). Then we associate with each such plane $(0,1)$ -matrices A and C of trace 0 (see [3]). Finally we show that for each class the trace condition contradicts the integrality of the multiplicities of the eigenvalues of A or C .

We shall use the following notations, definitions and basic results (see Dembowski [2]):

A collineation group of a projective plane has equally many point orbits and line orbits. The rank of a transitive permutation group is the number of orbits of the stabilizer of one of the permuted elements. If G is a (point or line) transitive collineation group of a projective plane, then the point and line ranks are equal (Kantor [8]). A rank m projective plane is a projective plane which admits a transitive collineation group whose (point or line) rank is m ($m \geq 2$). The lines (points) are identified with the set of points (lines) on them. We write $P \in l^G$ if and only if $P \in l^\gamma$ for all $\gamma \in G$.

2. PROOF OF THE THEOREM.

Let $\mathbb{P} = (P, L, \epsilon)$ be a projective plane of finite order n and let G be a rank 4 collineation group of $\mathbb{P}$ such that $G_{P_0} = G_{l_0}$ for some point-line pair (P_0, l_0) . It is easily seen that $n \geq 3$. A bijective map $\sigma : P \rightarrow L$ is defined by $P^\sigma = l$ if and only if $P = P_0^\gamma$ and $l = l_0^\gamma$ for some $\gamma \in G$. If $i \in \mathbb{N}$ we write l_i for P_i^σ . Clearly $P_0^\sigma = l_0$ and

$$P^{\sigma\gamma} = P^{\gamma\sigma}, \quad l^{\sigma^{-1}\gamma} = l^{\gamma\sigma^{-1}} \quad \text{for all } P \in \mathcal{P}, l \in L, \gamma \in G. \quad (1)$$

For $P \in \mathcal{P}$ G_P has exactly 4 orbits $\{P\}$, $\Delta(P)$, $\Gamma(P)$, $\Pi(P)$. We choose the notation in such a way that

$$(\Delta(P))^\gamma = \Delta(P^\gamma), \quad (\Gamma(P))^\gamma = \Gamma(P^\gamma), \quad (\Pi(P))^\gamma = \Pi(P^\gamma) \quad \text{for all } P \in \mathcal{P}, \gamma \in G \quad (2)$$

LEMMA 2.1: If $\Delta_1, \Delta_2, \Delta_3 \in \{\Delta, \Gamma, \Pi\}$, then $|\Delta_1(A) \cap \Delta_2(B)| = |\Delta_1(A') \cap \Delta_2(B')|$ if $A \in \Delta_3(B)$ and $A' \in \Delta_3(B')$.

PROOF: If $A \in \Delta_3(B)$, $A' \in \Delta_3(B')$, then for some $\gamma \in G$, $\gamma_0 \in G_B$

$$B' = B^\gamma = B^{\gamma_0\gamma}, \quad A' = A^{\gamma_0\gamma}, \quad \text{whence by (2)}$$

$$|\Delta_1(A') \cap \Delta_2(B')| = |\Delta_1(A)^{\gamma_0\gamma} \cap \Delta_2(B)^{\gamma_0\gamma}| = |(\Delta_1(A) \cap \Delta_2(B))^{\gamma_0\gamma}| = |\Delta_1(A) \cap \Delta_2(B)|.$$

LEMMA 2.2: Suppose that $P_0 \in l_0$. Then $l_0 - \{P_0\}$ and $P_0 - \{l_0\}$ are G_{P_0} -orbits, say $\Delta(P_0) = l_0 - \{P_0\}$ and $l_2^{P_0} = P_0 - \{l_0\}$ with $P_2^{P_0} = \Gamma(P_0)$. $P_1 \in \Delta(P_0)$ and $P_3 \in \Pi(P_0)$ can be chosen such that $P_1 \in l_0$; $P_0, P_2, P_3 \in l_2$; $P_2 \in l_1$; $P_1 \notin l_3$ (Fig. 1).

The case described by Lemma 2 will be called case I.

PROOF: If $l_0 - \{P_0\}$ is not a G_{P_0} -orbit, then it is the union of 2 orbits, say $l_0 - \{P_0\} = \Delta(P_0) \cup \Gamma(P_0)$. Then $P_0 - \{l_0\}$ is a line orbit l^{P_0} and $\Pi(P_0) = P^{P_0}$ with $l = P^\sigma$. This leads to the contradiction

$$n = |1_{P_0}^{G_{P_0}}| = |P^\sigma G_{P_0}| = |P^{G_{P_0} \sigma}| = |P^{G_{P_0}}| = |P - 1_0| = n^2.$$

Hence we may assume that $\Delta(P_0) = 1_0 - \{P_0\}$.

Dually: $P_0 - \{1_0\}$ is a G_{P_0} -orbit, say $1_2^{G_{P_0}} = P_0 - \{1_0\}$ where $P_2^{G_{P_0}} = \Gamma(P_0)$ (note that $P_0 \notin 1_1$).

$$|P_2^{G_{P_0}}| = |1_2^{G_{P_0}}| = |P_0 - \{1_0\}| = n \text{ implies } \Gamma(P_0) \cap 1_2 = \{P_2\}.$$

For any point $Q \neq P_0, P_2$ on 1_2 holds $Q^{G_{P_0}} = \Gamma(P_0)$.

Let $P_1' \in \Delta(P_0)$. If $P_2 \in 1_1'$ put $P_1 = P_1'$. If $P_2 \notin 1_1'$ then

$$\begin{aligned} |P_1^{G_{P_0}, P_2}| &= |1_1^{G_{P_0}, P_2}| \geq |(1_1' \cap 1_2)^{G_{P_0}, P_2}| = n-1 = |(1_1' \cap 1_2)^\sigma G_{P_0}, P_2| \\ &> |(1_1' \cap 1_2)^\sigma \cap 1_0)^{G_{P_0}, P_2}|; \text{ this implies } P_1^{G_{P_0}, P_2} = P_1 \text{ for some point} \end{aligned}$$

$P_1 \in \Delta(P_0)$ and hence $P_2 \in 1_1$.

It remains to prove that $P_3 \in \Gamma(P_0) \cap 1_2$ exists such that $P_1 \notin 1_3$. If no

such P_3 exists then $P_1 \in Q^\sigma$ for all $Q \in 1_2 - \{P_0, P_2\}$ and hence $G_{P_0, P_2} \leq G_{P_0, P_1}$.

Let $\gamma \in G$ be such that $P_2^\gamma = P_0$. Then $1_2^\gamma = 1_0$ and therefore $P_0^\gamma \in 1_0, P_0^\gamma \neq P_0$ and $P_0^{\gamma\gamma_0} = P_1$ for some $\gamma_0 \in G_{P_0}$. It follows that $G_{P_0, P_1} = (\bar{\gamma}\gamma_0)^{-1} G_{P_0, P_2} \bar{\gamma}\gamma_0$.

Hence

$$G_{P_0, P_1} = G_{P_0, P_2}. \quad (3)$$

Further

$$P_2 \notin 1_1^{\gamma_0} \text{ for some } \gamma_0 \in G_{P_0}, \quad (4)$$

for otherwise $P_2^{\gamma_0'} \in 1_1^{\gamma_0''}$ for all $\gamma_0', \gamma_0'' \in G_{P_0}$ which cannot occur.

$$P_2^{\gamma_0} \in 1_1 \text{ for some } \gamma_0 \in G_{P_0} \text{ if and only if } \gamma_0 \in G_{P_2}. \quad (5)$$

To prove (5) note that by (4) through any point of $1_2 - \{P_0\}$ goes at least one and hence exactly one line of $1_1^{G_{P_0}}$. (3) and $P_2 \in 1_1$ then imply (5).

Let's apply (5) to G_{P_1} in place of G_{P_0} :

$$\Delta(P_1) = 1_1 - \{P_1\}; \Gamma(P_1) = \Gamma(P_0^\gamma) = (\Gamma(P_0))^\gamma = P_2^{G_{P_0}^\gamma} = P_0^{\gamma^{-1} G_{P_0} \gamma} = P_0^{G_{P_1}}$$

where $\gamma \in G$ such that $P_0^\gamma = P_1$, $P_2^\gamma = P_0$; $\mathbb{I}(P_1) = S^{G_{P_1}}$ for some $S \in l_0 - \{P_0, P_1\}$; hence $P_0^{\gamma_1} \in l_2$ for some $\gamma_1 \in G_{P_1}$ if and only if $\gamma_1 \in G_{P_0}$.

It follows that $R \notin P_0^{G_{P_1}}$ for any $R \in l_2 - \{P_0, P_2\}$. Let $r = R^\delta$ for some such R .

Of the 3 orbits $(P_0, l_1)^G$, $(P_0, l_2)^G$, $(P_0, r)^G$ induced by G on $P \times L - (P_0, l_0)^G$ only one consists of flags. Thus (P_1, l_0) and (P_1, r) and then also (P_0, l_1) and (R, l_1) belong to the same G -orbit. This contradicts $R \notin P_0^{G_{P_1}}$. Hence there exists $P_3 \in \mathbb{I}(P_0) \cap l_2$ such that $P_1 \notin l_3$.

LEMMA 2.3: Suppose that $P_0 \notin l_0$. Then l_0 and dually P_0 are G_{P_0} -orbits, say $\Delta(P_0) = l_0$. $P_1 \in \Delta(P_0)$, $P_2 \in \Gamma(P_0)$, $P_3 \in \mathbb{I}(P_0)$ can be chosen such that either $P_0, P_2, P_3 \in l_1$; $P_1 \in l_2, l_3$; $P_2 \notin l_3$; $P_3 \notin l_2$ or $P_0, P_1, P_3 \in l_2$; $P_2 \in l_0$; $\Gamma(P_0) \cap l_2 = \{P_2^{\gamma_0}\}$ for some $\gamma_0 \in G_{P_0}$; $P_2^{\gamma_0} \in l_1$; $P_1, P_2, P_3 \notin l_1, l_3$. In both cases $n \geq 4$.

Figure 1

The 2 cases described by Lemma 3

will be called case III1 resp. case III2 (Fig. 2).

PROOF: It is easily seen that

l_0 and P_0 are G_{P_0} -orbits; say

$\Delta(P_0) = l_0$. Let $P_1 \in \Delta(P_0)$. We

have to distinguish 2 cases:

Case III1: $P_0 \in l_1$

Case III2: $P_0 \notin l_1$.

CASE III1: Clearly $P_0 = l_1^{G_{P_0}}$ and $\Gamma(P_0) = P_2^{G_{P_0}}$, $\mathbb{I}(P_0) = P_3^{G_{P_0}}$ for some

$P_2, P_3 \in l_1 - \{P_0, l_0 \cap l_1\}$. If $P_2 \in l_3$ then $(P_2, l_3) \in (P_0, l_1)^G$, hence

$(P_3, l_2) \in (P_1, l_0)^G$, so $P_3 \in l_2$.

Figure 2

Analogously $P_2 \in l_3$ if $P_3 \in l_2$. Thus

$$P_2 \in l_3 \quad \text{if and only if} \quad P_3 \in l_2. \quad (6)$$

Similarly one proves

$$P_1 \in l_2, l_3. \quad (7)$$

If $n > 3$ then, by (6), we can choose P_2, P_3 such that $P_2 \notin l_3, P_3 \notin l_2$.

Let's show that $n > 3$ (Fig. 3). Suppose that $n = 3$. Put $P_4 = l_0 \cap l_1$.

Figure 3

Then, since $P_0 \in l_1$ and $P_1 \in l_0$,

$l_4 = P_0 P_1$. Let $P_5 \in l_0 - \{P_1, P_4\}$.

Then $P_0 \in l_5$ and then $l_5 \cap l_2 =$

$(P_0 P_5 \cap l_3) P_4 \cap l_2$. Denote this

point by T . Clearly $P_2 P_5 \cap l_2 =$

T . Since $(P_2 P_5)^{\sigma^{-1}} \in l_2 \cap l_5$ we

obtain the contradiction

$$(P_2 P_5)^{\sigma^{-1}} \in P_2 P_5.$$

CASE II2: We may assume that $P_0, P_1, P_3 \in l_2$ where $P_2 \in \Gamma(P_0)$. Then

$G_{P_0, P_1} = G_{P_0, P_2}$. We first assume that $n > 3$. $|P_2^o| = |l_2^o| = n+1$, hence $|l_2 \cap \Gamma(P_0)| = 1$; let $l_2 \cap \Gamma(P_0) = \{P_2^{\gamma_0}\}$ with some $\gamma_0 \in G_{P_0}$. Then $P_3^{G_{P_0, P_1}} = l_2 - \{P_0, P_1, P_2^{\gamma_0}\}$ and hence, since $P_2^{\gamma_0}$ is invariant under G_{P_0, P_1} and since $n > 3$, $l_1 \cap l_2 = P_2^{\gamma_0}$.

The only G -orbit of $\mathcal{P} \times L$ consisting of flags is $(P_0, l_2)^G$. Hence $(P_1, l_2), (P_3, l_2), (P_1, l_0) \in (P_0, l_2)^G$. $P_0 \notin l_1$ then implies that $(P_2, l_1), (P_2, l_3), (P_0, l_1), (P_2, l_0) \notin (P_0, l_2)^G$, in particular $P_2 \notin l_0, l_1, l_3$.

If $P_1 \in l_3$ then $(P_1, l_3) \in (P_0, l_2)^G$ and hence $(P_3, l_1) \in (P_2, l_0)^G$.

Since also $(P_0, l_1) \in (P_2, l_0)^G$ we have $P_0^{\gamma_1} = P_3$ for some $\gamma_1 \in G_{l_1} = G_{P_1}$.

This implies that G_{P_1} is transitive on $l_2 - \{P_1, P_2^{\gamma_0}\}$ which is impossible.

Hence $P_1 \notin l_3$.

If $n=3$ then $l_2 = \{P_0, P_1, P_2^{\gamma_0}, P_3\}$. γ_0 is of order 4, for if $\gamma_0^2 = 1$ then $(P_2, l_2^{\gamma_0}) \in (P_2^{\gamma_0}, l_2)^{G_{P_0}}$ which is impossible. Moreover $P_2^{\gamma_0^2} \neq P_2$ since otherwise $\gamma_0^2 \in G_{P_0, P_2} = 1$. It follows that $|(P_2 P_2^{\gamma_0^2})^{G_{P_0}}| = 4$ which contradicts

$(P_2 P_2^{\gamma_0^2})^{G_{P_0}} = \{P_2 P_2^{\gamma_0^2}, P_2^{\gamma_0} P_2^{\gamma_0^3}\}$. This completes the proof of the Lemma.

Let us now associate with $(G, \mathbb{P})$ 3 $(0,1)$ -matrices.

If $\rho(P)$ is a G -orbit then let $\rho'(P)$ denote the paired orbit (see Wielandt

[9]). If $Q \in \rho(P)$ then $Q = P^{\gamma}$ for some $\gamma \in G$ and $Q^{\gamma} \in (\rho(P))^{\gamma} = \rho(P^{\gamma}) = \rho(Q)$.

Hence $Q^{\gamma^{-1}} = P \in \rho'(Q)$, i.e.

$Q \in \rho(P)$ implies that $P \in \rho'(Q)$. (8)

This implies that in

Case I:

$$\Delta'(P) = \Gamma(P)$$

$$\Gamma'(P) = \Delta(P)$$

$$\Pi'(P) = \Pi(P)$$

Case II1:

$$\Delta'(P) = \Delta(P)$$

$$\Gamma'(P) = \Pi(P) \text{ resp. } \Gamma(P)$$

$$\Pi'(P) = \Gamma(P) \text{ resp. } \Pi(P)$$

Case II2:

$$\Delta'(P) = \Gamma(P)$$

$$\Gamma'(P) = \Delta(P)$$

$$\Pi'(P) = \Pi(P).$$

Now let $P = \{P_1, P_2, \dots, P_v\}$, $L = \{l_1, l_2, \dots, l_v\}$, $l_k = P_k^\sigma$ ($k = 1, 2, \dots, v$). Let A be the $(0,1)$ -matrix with rows enumerated by the points P_k and columns by $\Delta(P_k)$ and such that $(P_k, \Delta(P_i)) = 1$ if and only if $P_k \in \Delta(P_i)$. Let B, C be the analogous matrices with $\Gamma(P_k)$ resp. $\Pi(P_k)$ in place of $\Delta(P_k)$.

We have in

case I:

$$A^t = B, C^t = C$$

case II1:

$$A^t = A, B^t = C$$

$$\text{if } \Gamma'(P) = \Pi(P)$$

$$A^t = A, B^t = B, C^t = C \text{ if } \Gamma'(P) = \Gamma(P)$$

case II2:

$$A^t = B, C^t = C$$

Let $k = |\Delta(P)|$, $l = |\Gamma(P)|$, $m = |\Pi(P)|$,

$$|\Delta(P) \cap \Delta(Q)| = \begin{cases} \lambda \\ \mu \\ \nu \end{cases} \text{ if } Q \in \begin{cases} \Delta(P) \\ \Gamma(P) \\ \Pi(P) \end{cases}$$

$$|\Pi(P) \cap \Pi(Q)| = \begin{cases} \lambda' \\ \mu' \\ \nu' \end{cases} \text{ if } Q \in \begin{cases} \Pi(P) \\ \Delta(P) \\ \Gamma(P) \end{cases}.$$

A straightforward calculation shows that

$I + A + B + C = J$, the $v \times v$ -matrix with 1's in every entry

$$A^t A = k I + \lambda A + \mu B + \nu C$$

$$C^t C = m I + \mu' A + \nu' B + \lambda' C$$

$$A J = k J$$

$$B J = l J$$

$$C J = m J.$$

Now we determine the eigenvalues of A in case II1 and of C in the cases I and II2.

$$\begin{aligned} \text{CASE II1: } \quad k &= n + 1 \\ l &= n_2(n + 1) \quad \text{where } n_2 = |P_2^{G_{P_0}, P_1}| \\ m &= n_3(n + 1) \quad \text{where } n_3 = |P_3^{G_{P_0}, P_1}| \\ k + l + m + 1 &= v = n^2 + n + 1, \quad n_2 + n_3 = n - 1, \quad \lambda = \mu = v = 1. \end{aligned}$$

It follows that $A^2 = A^t A = (n + 1) I + A + B + C = n I + J$; hence $(A - (n + 1) I)(A^2 - n I) = 0$. This gives the eigenvalues of A:

$$\lambda_1 = n + 1, \quad \lambda_{2,3} = \pm \sqrt{n}.$$

$$\text{CASE I:} \quad k = l = n, \quad m = n(n - 1), \quad k + l + m + 1 = v = n^2 + n + 1.$$

$$\begin{aligned} \text{We have} \quad \lambda' &= |\Pi(P_0) \cap \Pi(P_3)| \\ \mu' &= |\Pi(P_0) \cap \Pi(P_1)| \\ v' &= |\Pi(P_0) \cap \Pi(P_2)|. \end{aligned}$$

Let's calculate λ' :

$$n(n - 1) = |\Pi(P_3)| = |\Pi(P_3) \cap \Delta(P_0)| + |\Pi(P_3) \cap \Gamma(P_0)| + |\Pi(P_3) \cap \Pi(P_0)| + 1 \quad (9)$$

(note that $\Gamma(P_3) = P_2^{G_{P_3}}$ and hence $P_0 \in \Pi(P_3)$).

$$n = |\Delta(P_0)| = |\Delta(P_0) \cap \Delta(P_3)| + |\Delta(P_0) \cap \Gamma(P_3)| + |\Delta(P_0) \cap \Pi(P_3)|. \quad (10)$$

Clearly

$$|\Delta(P_0) \cap \Delta(P_3)| = 1 \quad (11)$$

$$|\Delta(P_0) \cap \Gamma(P_3)| = |\Delta(P_3) \cap \Gamma(P_0)| = 2. \quad (12)$$

PROOF of (12): $P_o \in \Pi(P_3)$ and $P_3 \in \Pi(P_o)$, hence, by Lemma 1, $|\Delta(P_o) \cap \Gamma(P_3)|$
 $= |\Delta(P_3) \cap \Gamma(P_o)|$. $P_2 \notin 1_1^{\gamma_o}$ for some $\gamma_o \in G_P$. Thus $|P_3^{G_{P_o, P_2}}| = n - 1$ implies
 that $|1_1^{\gamma_o G_{P_o, P_2}}| \geq n - 1$. Hence $|1_1^{\gamma_o G_{P_o, P_2}}| = n - 1$. Since $P_1 \notin P_1^{\gamma_o G_{P_o, P_2}}$ we
 then have $P_1^{G_{P_o, P_2}} = P_1$, i.e. $G_{P_o, P_2} \leq G_{P_o, P_1}$. Since both groups are conjugate
 (see the proof of Lemma 2) this gives $G_{P_o, P_1} = G_{P_o, P_2}$. $\gamma'_o, \gamma''_o \in 1_3^o, P_2 \in 1_3^o, P_2 \neq$
 P_2 for some $\gamma'_o, \gamma''_o \in G_P$. Since, by the above, G_{P_o, P_2} is transitive on $1_o -$
 $\{P_o, P_1\}$, $|P_2^{G_{P_o, P_2}}| = n - 1$; hence $|1_3^o \cap P_2^{G_{P_o, P_2}}| = |1_3 \cap P_2^{G_{P_o, P_2}}| = 2$. This
 proves (12).

Equations (10), (11), (12) imply

$$|\Pi(P_3) \cap \Delta(P_o)| = n - 3. \quad (13)$$

To determine $|\Pi(P_3) \cap \Gamma(P_o)|$ we use

$$n = |\Gamma(P_o)| = |\Gamma(P_o) \cap \Delta(P_3)| + |\Gamma(P_o) \cap \Gamma(P_3)| + |\Gamma(P_o) \cap \Pi(P_3)|. \quad (14)$$

By (12) $|\Gamma(P_o) \cap \Delta(P_3)| = 2$. Since $\Gamma(P_3) = P_2^{G_{P_3}}$, $|\Gamma(P_o) \cap \Gamma(P_3)| =$
 $|P_2^{G_{P_o}} \cap P_2^{G_{P_3}}| = |1_2^{G_{P_o}} \cap 1_2^{G_{P_3}}| = 1$. It follows that

$$|\Pi(P_3) \cap \Gamma(P_o)| = n - 3. \quad (15)$$

Equations (9), (13) and (15) imply that

$$\lambda' = n^2 - 3n + 5. \quad (16)$$

Analogously we calculate μ' and ν' :

$$n(n - 1) = |\Pi(P_1)| = |\Pi(P_1) \cap \Delta(P_o)| + |\Pi(P_1) \cap \Gamma(P_o)| + |\Pi(P_1) \cap \Pi(P_o)|$$

with $|\Pi(P_1) \cap \Delta(P_o)| = n - 1$.

$$\text{In } n = |\Gamma(P_o)| = |\Gamma(P_o) \cap \Delta(P_1)| + |\Gamma(P_o) \cap \Gamma(P_1)| + |\Gamma(P_o) \cap \Pi(P_1)|$$

$|\Gamma(P_0) \cap \Delta(P_1)| = 1$ by the proof of (12) and $|\Gamma(P_0) \cap \Gamma(P_1)| = |P_2^{G_{P_0}} \cap P_0^{G_{P_1}}| = |1_2^{G_{P_0}} \cap 1_0^{G_{P_1}}| = 0$. Hence $|\Pi(P_1) \cap \Gamma(P_0)| = n - 1$ and thus

$$\mu' = (n - 1)(n - 2). \quad (17)$$

$$n(n - 1) = |\Pi(P_2)| = |\Pi(P_2) \cap \Delta(P_0)| + |\Pi(P_2) \cap \Gamma(P_0)| + |\Pi(P_2) \cap \Pi(P_0)|.$$

In $n = |\Delta(P_0)| = |\Delta(P_0) \cap \Delta(P_2)| + |\Delta(P_0) \cap \Gamma(P_2)| + |\Delta(P_0) \cap \Pi(P_2)|$
 $|\Delta(P_0) \cap \Delta(P_2)| = 0$ and $|\Delta(P_0) \cap \Gamma(P_2)| = |P_1^{G_{P_0}} \cap P_1^{G_{P_2}}| = |1_1^{G_{P_0}} \cap 1_1^{G_{P_2}}| = 1$
 (note that $|1_1^{G_{P_2}}| = |P_1^{G_{P_2}}| = n$ and hence $\Gamma(P_2) = P_1^{G_{P_2}}$). Hence $|\Pi(P_2) \cap \Delta(P_0)| = n - 1$.

Further $|\Pi(P_2) \cap \Gamma(P_0)| = |\Gamma(P_0)| - |\Gamma(P_0) \cap \Delta(P_2)| - |\Gamma(P_0) \cap \Gamma(P_2)| - 1$
 where $|\Gamma(P_0)| = n$, $|\Gamma(P_0) \cap \Delta(P_2)| = 0$ and $|\Gamma(P_0) \cap \Gamma(P_2)| = |P_2^{G_{P_0}} \cap P_1^{G_{P_2}}| = |1_2^{G_{P_0}} \cap 1_1^{G_{P_2}}| = 0$. Hence $|\Pi(P_2) \cap \Gamma(P_0)| = n - 1$. It follows that

$$\nu' = (n - 1)(n - 2). \quad (18)$$

Equations (16), (17) and (18) imply that $C^2 = C^t C = n(n - 1) I + (n - 1)(n - 2)(A + B) + (n^2 - 3n + 5)C = n(n - 1)I + (n - 1)(n - 2)(J - I) + 3C$
 and then $(C - n(n - 1)I)(C^2 - 3C - 2(n - 1)I) = 0$.

The eigenvalues of C are $\lambda_1 = n(n - 1)$; $\lambda_{2,3} = (3 \pm \sqrt{8n + 1})/2$.

CASE II2: $k = 1 = n + 1$, $m = (n - 2)(n + 1)$, $k + 1 + m + 1 = v = n^2 + n + 1$.

By the proof of Lemma 3 $n \geq 4$. Let's determine λ' , μ' , ν' :

$(n + 1)(n - 2) = |\mathbb{I}(P_3)| = |\mathbb{I}(P_3) \cap \Delta(P_0)| + |\mathbb{I}(P_3) \cap \Gamma(P_0)| + |\mathbb{I}(P_3) \cap \Pi(P_0)| - 1$. In $n + 1 = |\Delta(P_0)| = |\Delta(P_0) \cap \Delta(P_3)| + |\Delta(P_0) \cap \Gamma(P_3)| + |\Delta(P_0) \cap \Pi(P_3)|$ clearly $|\Delta(P_0) \cap \Delta(P_3)| = 1$. Let's show that

$$|\Delta(P_0) \cap \Gamma(P_3)| = |\Delta(P_3) \cap \Gamma(P_0)| = 2 \quad (19)$$

$$|\Delta(P_1) \cap \Gamma(P_0)| = 2. \quad (20)$$

PROOF of (19) and (20): By Lemma 1 $|\Delta(P_o) \cap \Gamma(P_3)| = |\Delta(P_3) \cap \Gamma(P_o)|$.

For $\gamma_o' \in G_{P_o}$

$$P_2^{\gamma_o'} \in l_1^{\gamma_o'}, l_1^{\gamma_o'} \neq P_2^{\gamma_o'} P_2 \text{ if and only if } \gamma_o' \in G_{P_o, P_1}, \quad (21)$$

for otherwise $P_2^{\gamma_o'} \in l_1^{\gamma_o' G_{P_o, P_1}}, l_1^{\gamma_o' G_{P_o, P_1}} \neq l_1^{\gamma_o' G_{P_o, P_1}}, |l_1^{\gamma_o' G_{P_o, P_1}}| = n - 2, P_2 \in$

$l_1^{\gamma_o' G_{P_o, P_1}} \gamma_o'^{-1}$, which leads to the contradiction $n + 1 = |l_1^{\gamma_o'}| \geq (|P_2^{\gamma_o'}| - 2) +$

$$(|P_2| - 2) = 2(n - 1).$$

Further

$$P_2^{\bar{\gamma}_o} \notin l_1 \cup P_2^{\gamma_o} P_2 \text{ for some } \bar{\gamma}_o \in G_{P_o}. \quad (22)$$

otherwise, since $|P_2^{\gamma_o}| \geq 5$, every line of $l_1^{\gamma_o}$ would contain at least 3 points of $P_2^{\gamma_o}$ and this would imply that a point of $P_2^{\gamma_o} - (l_1 \cup P_2^{\gamma_o} P_2)$ exists. By (21)

and (22) $P_2^{\gamma_o}, P_2^{\bar{\gamma}_o} \in l_3^{\gamma_o^*}, P_2^{\gamma_o} \neq P_2^{\bar{\gamma}_o}$ for some $\gamma_o^* \in G_{P_o}$. Since $|l_3^{\gamma_o^*}| = n - 2, |P_2^{\gamma_o} \cap l_3| = |\Gamma(P_o) \cap \Delta(P_3)| = 2$. This proves (19).

Each of the $n - 2$ lines of $l_3^{\gamma_o} - \{P_2^{\gamma_o} P_2\}$ through $P_2^{\gamma_o}$ contains exactly one point of $P_2^{\gamma_o} - \{P_2^{\gamma_o}\}$. Together with $P_2^{\gamma_o}, P_2$ this gives n points of $P_2^{\gamma_o}$. It follows that exactly one point of $P_2^{\gamma_o} - \{P_2^{\gamma_o}\}$ lies on l_1 . This proves (20).

By means of (19) we obtain $|\Pi(P_3) \cap \Delta(P_o)| = n - 2$.

In $n + 1 = |\Gamma(P_o)| = |\Gamma(P_o) \cap \Delta(P_3)| + |\Gamma(P_o) \cap \Gamma(P_3)| + |\Gamma(P_o) \cap \Pi(P_3)|$
 $|\Gamma(P_o) \cap \Delta(P_3)| = 2$ by (19) and $|\Gamma(P_o) \cap \Gamma(P_3)| = |P_2^{\gamma_o} \cap P_2^{\gamma_o}| = |l_2^{\gamma_o} \cap l_2^{\gamma_o}|$
 $= 1$. Hence $|\Pi(P_3) \cap \Gamma(P_o)| = n - 2$. It follows that $\lambda' = n^2 - 3n + 1$.

$\mu' = |\Pi(P_o) \cap \Pi(P_1)| = |\Pi(P_1)| - |\Pi(P_1) \cap \Delta(P_o)| - |\Pi(P_1) \cap \Gamma(P_o)|$ where
 $|\Pi(P_1)| = (n + 1)(n - 2), |\Pi(P_1) \cap \Delta(P_o)| = n - 2$ and $|\Pi(P_1) \cap \Gamma(P_o)| = |\Gamma(P_o)| -$
 $|\Gamma(P_o) \cap \Delta(P_1)| - |\Gamma(P_o) \cap \Gamma(P_1)|$. $|\Gamma(P_o) \cap \Delta(P_1)| = 2$ by (20) and $|\Gamma(P_o) \cap \Gamma(P_1)| =$
 $|P_2^{\gamma_o} \cap P_2^{\gamma_o}| = |l_2^{\gamma_o} \cap l_2^{\gamma_o}| = 1$. Hence $|\Pi(P_1) \cap \Gamma(P_o)| = n - 2$ and $\mu' =$

$$(n-2)(n-1).$$

$$v' = |\Pi(P_0) \cap \Pi(P_2)| = |\Pi(P_2)| - |\Pi(P_2) \cap \Delta(P_0)| - |\Pi(P_2) \cap \Gamma(P_0)| \text{ where} \\ |\Pi(P_2)| = (n+1)(n-2),$$

$$|\Pi(P_2) \cap \Delta(P_0)| = |\Pi(P_0) \cap \Delta(P_1)| \text{ by Lemma 1} \\ = |\Delta(P_1)| - |\Delta(P_1) \cap \Delta(P_0)| - |\Delta(P_1) \cap \Gamma(P_0)| \\ = (n+1) - 1 - 2 = n-2 \text{ by (20),}$$

$$|\Pi(P_2) \cap \Gamma(P_0)| = |\Pi(P_0) \cap \Gamma(P_1)| \text{ by Lemma 1} \\ = |P_3^{G_P} \cap P_0^{G_P}| = |\{l_3^{\gamma_0} : \gamma_0 \in G_P, P_1 \in l_3^{\gamma_0}\}| = n-2 \text{ since} \\ \text{through any point on } l_0 \text{ goes exactly one line of } l_1^{G_P} \text{ and one of } l_2^{G_P}. \text{ Hence} \\ v' = (n-2)(n-1).$$

$$\text{It follows that } C^2 = C^t C = (n+1)(n-2)I + (n-1)(n-2)(A+B) + \\ (n^2 - 3n + 1)C = (n+1)(n-2)I + (n^2 - 3n + 2)(A+B+C) - C \text{ and } C^2 + C - \\ 2(n-2)I = (n-1)(n-2)J \text{ whence } (C - (n+1)(n-2)I)(C^2 + C - 2(n-2)I) \\ = 0. \text{ The eigenvalues of } C \text{ are } \lambda_1 = (n+1)(n-2), \lambda_{2,3} = (-1 \pm \sqrt{8n-15})/2.$$

REMARK: Let $\phi: G \rightarrow GL_V(\mathbb{C})$ be the matrix representation of G obtained by associating with each $\gamma \in G$ the corresponding permutation matrix $\phi(\gamma)$ (the ordering of P is the same as used in constructing the matrices A, B, C). By (2) $\phi(\gamma)$ commutes with A, B, C for all $\gamma \in G$. Hence, by [9] Theorem 28.4, $\{I, A, B, C\}$ is the basis of the commuting algebra $\mathcal{V}(G)$ of ϕ . By [9] Theorem 29.5 $\mathcal{V}(G)$ is commutative and hence, by [9] Theorem 29.4, the representation ϕ has 4 irreducible constituents $D_1 = 1, D_2, D_3, D_4$, each with multiplicity 1. If f_i is the degree of D_i then $f_1 = 1$ and $\sum_{i=1}^4 f_i = v$.

Let us finally show how the fact that A and C have trace 0 contradicts the integrality of the multiplicities of $\lambda_1, \lambda_2, \lambda_3$.

In the 3 cases λ_1 appears with multiplicity 1. Let f denote the multiplicity of λ_2 ; then $v - f - 1$ is the multiplicity of λ_3 . This leads to

$$0 = n(n-1) + f(3 + \sqrt{8n+1})/2 + (n(n+1) - f)(3 - \sqrt{8n+1})/2 \quad \text{in case I,}$$

$$0 = (n+1) + f\sqrt{n} + (-\sqrt{n})(n(n+1) - f) \quad \text{in case II1,}$$

$$0 = (n+1)(n-2) + f(-1 + \sqrt{8n-15})/2 + (n(n+1) - f)(-1 - \sqrt{8n-15})/2$$

in case II2.

In any case this contradicts the fact that $n \geq 2$ and $f \geq 1$ are integers:

In case II1 this is clear.

In case I suppose that a prime p divides $\sqrt{8n+1}$. Then $p \nmid n$, hence $p \mid 5n+1$ and then $p \mid 3n$, i.e. $p = 3$. This implies that $8n+1 = 3^{2i}$ for some $i \geq 2$ and that $n(5n+1)/\sqrt{8n+1} = (3^{2i}-1)(5 \cdot 3^{2i-1}+1)/8^2 \cdot 3^{i-1} \notin \mathbb{N}$.

In case II2 suppose that a prime p divides $\sqrt{8n-15}$. Then $p \in \{17, 23\}$ and $p^2 \nmid n+1$, $p^2 \nmid n-4$. Hence $8n-15 \in \{17^2, 23^2, 17^2 \cdot 23^2\}$, i.e. $n \in \{38, 68, 19112\}$. Suppose that $n = 38$. Since G_{P_0, P_2} is transitive on $1_2 - \{P_0, P_1, P_2^{\gamma_0}\}$ $|G|$ is even. This contradicts the fact that if $n \equiv 2 \pmod{4}$, then the full collineation group is of odd order (Hughes [5]).

Suppose that $n \in \{68, 19112\}$. Then n is not a square and $n^2 + n + 1$ not a prime. Hence, since G is flag-transitive, n is a prime power (Higman and Mc Laughlin [4]) which is absurd.

REFERENCES

1. Bachmann, O., On rank 3 projective designs, Mh. Math. **89** (1980), 175-183.
2. Dembowski, P., Finite Geometries, New York, Springer, 1968.
3. Higman, D.G., Finite permutation groups of rank 3, Math. Z. **86** (1964), 145-156.
4. Higman, D.G. and J.E. Mc Laughlin, Geometric ABA-groups, Ill. J. Math. **5** (1961), 382-397.

5. Hughes, D.R., Generalized incidence matrices over group algebras, Ill. J. Math. 1 (1957), 545-551.
6. Kallaher, M.J., On rank 3 projective planes, Pacif. J. Math. 39 (1971), 207-214.
7. Kantor, W.M., Moore geometries and rank 3 groups having $\mu=1$, Quart. J. Math. Oxford (2) 28 (1977), 309-328.
8. Kantor, W.M., Automorphism groups of designs, Math. Z. 109 (1969), 246-252.
9. Wielandt, H., Finite permutation groups, New York, Academic Press, 1964.