A RESULT ON CO-CHROMATIC GRAPHS

E.J. FARRELL

Department of Mathematics
The University of the West Indies
St. Augustine, Trinidad.

(Received December 17, 1979 and in revised
form April 18, 1980)

ABSTRACT. A sufficient condition for two graphs with the same number of nodes to have the same chromatic polynomial is given.

KEY WORDS AND PHRASES. Graph, Co-chromatic, Chromatic polynomial.

1980 MATHEMATICS SUBJECT CLASSIFICATION CODES: Primary 05 C15, Secondary 26 C10.

1. INTRODUCTION.

We prove a theorem which gives a sufficient condition for two graphs to be co-chromatic i.e. to have the same chromatic polynomial.

The chromatic polynomial $\chi(G;\lambda)$ of a graph G with p nodes has degree p and constant term equal to 0. Hence the chromatic polynomial has p coefficients. If the graph has at least one edge, then the sum of these coefficients is equal to 0. Hence the chromatic polynomial is uniquely determined if $p-1$ of the coefficients are known. Our result is a generalization of this.

2. MAIN RESULTS.

THEOREM 2. If two graphs with p nodes have chromatic numbers $\geq n$ and have at least $p+1-n$ corresponding coefficients of their chromatic polynomials equal, then they are co-chromatic.

In the proof we shall use a special case of the following Lemma.

LEMMA 2.1

Let $P(x) = c_1 x^{n_1} + c_2 x^{n_2} + \dots + c_s x^{n_s}$, where c_i and n_i are real numbers for $i = 1, 2, \dots, s$. We assume that $c_i \neq 0$ for all i and that $n_i \neq n_j$ for $i \neq j$. Then the equation $P(x) = 0$ has at most $s-1$ real positive solutions.

PROOF OF THE LEMMA 2.1 By induction over s . For $s = 1$ the statement is obvious. Suppose that it is true for $s - 1$ and let $P(x)$ be the above expression.

The expression

$$Q(x) = x^{-n_1} \cdot P(x) = c_1 + c_2 x^{n_2 - n_1} + \dots + c_s x^{n_s - n_1}$$

has the derivative

$$Q'(x) = c_2(n_2 - n_1) \cdot x^{n_2 - n_1 - 1} + \dots + c_s(n_s - n_1) \cdot x^{n_s - n_1 - 1}.$$

By induction, $Q'(x) = 0$ for at most $s-2$ positive x . But between any two positive solutions of $P(x) = x^{n_1} \cdot Q(x) = 0$, there exists a solution of $Q'(x) = 0$. Hence $P(x) = 0$ for at most $s-1$ positive x .

Q.E.D.

PROOF OF THE THEOREM 2. Let G and H be the two graphs. Let us assume that m of the coefficients of $\chi(G; \lambda)$ and $\chi(H; \lambda)$ are equal. Then $m \geq p + 1 - n$, by our assumption. Let us assume that $m < p$. We can therefore write

$$\chi(G; \lambda) = f(\lambda) + g(\lambda)$$

and

$$\chi(H; \lambda) = f(\lambda) + h(\lambda),$$

where $f(\lambda)$ contains m terms and $g(\lambda)$ and $h(\lambda)$ are the remaining terms of $\chi(G; \lambda)$ and $\chi(H; \lambda)$ respectively.

Since G and H have chromatic numbers $\geq n$, all integers $1, 2, \dots, n-1$ are roots of $\chi(G; \lambda)$ and $\chi(H; \lambda)$.

Let

$$g(\lambda) = a_1 \lambda^{n_1} + a_2 \lambda^{n_2} + \dots + a_{p-m} \lambda^{n_{p-m}}$$

$$\text{and } h(\lambda) = b_1 \lambda^{n_1} + b_2 \lambda^{n_2} + \dots + b_{p-m} \lambda^{n_{p-m}}$$

If r is an integer such that $1 \leq r \leq n-1$, then $g(r) = h(r)$, i.e.

$$(a_1 - b_1)r^{n_1} + (a_2 - b_2)r^{n_2} + \dots + (a_{p-m} - b_{p-m})r^{n_{p-m}} = 0.$$

Since $n - 1 \geq p - m$, this is a contradiction by the Lemma.

Q.E.D.

3. ILLUSTRATIONS

We will now illustrate the theorem. We will assume that the chromatic polynomial of a graph G with p nodes is written in descending powers of λ .

$$\text{i.e. } \chi(G; \lambda) = \sum_{k=0}^p a_{p-k} \lambda^{p-k}.$$

It is well known that if G contains p nodes and q edges, then a_p , a_{p-1} and a_{p-2} are 1, $-q$ and $\binom{q}{2} - A$ respectively, where A is the number of triangles in G . It was also shown in Farrell [1] (Theorem 1) that

$$a_{p-2} = -\binom{q}{3} + (q-2)A + B - 2C,$$

where B and C are the numbers of subgraphs of G which are quadrilaterals (without diagonals) and complete graphs with four nodes.

Let G_1 and G_2 be the graphs shown below

Let $\chi(G_1; \lambda) = \sum_{k=0}^6 a_{6-k} \lambda^{6-k}$ and $\chi(G_2; \lambda) = \sum_{k=0}^6 b_{6-k} \lambda^{6-k}$. Then $a_6 = b_6 = 1$ and $a_5 = b_5 = 11$. Since G_1 and G_2 contain 7 triangles, $a_4 = b_4 = \binom{11}{2} - 7 = 48$. Therefore G_1 and G_2 have 6 nodes, their chromatic number is $\geq 4 = n$ and $(p+1-n)=3$ of their corresponding coefficients are equal. It follows from the above theorem that G_1 and G_2 are co-chromatic.

The chromatic polynomial of G_1 and G_2 has been computed. It is

$$\chi(G_1; \lambda) = \chi(G_2; \lambda) = \lambda^6 - 11\lambda^5 + 48\lambda^4 - 103\lambda^3 + 107\lambda^2 - 42\lambda.$$

Consider the graphs H_1 , H_2 and H_3 shown below.

H_1 H_2 H_3

All three graphs contain 8 nodes and 18 edges. Each contains 17 triangles.

Therefore, the third coefficient of their chromatic polynomial is $\binom{18}{2} - 17 = 136$.

Finally, each contains 7 subgraphs which are complete graphs with 4 nodes and 0 quadrilaterals without diagonals. Therefore the fourth coefficients are equal.

Hence from the above theorem H_1 , H_2 and H_3 are co-chromatic.

The chromatic polynomial of H_1 , H_2 and H_3 has been computed. It is

$$\chi(H_1; \lambda) = \chi(H_2; \lambda) = \chi(H_3; \lambda) = \lambda^8 - 18\lambda^7 + 136\lambda^6 - 558\lambda^5 + 1339\lambda^4 - 1872\lambda^3 + 1404\lambda^2 - 432\lambda.$$

REFERENCES

- [1] Farrell, E.J., On Chromatic Coefficients, Discrete Mathematics, 29(1980), 257-264.