ON ELATIONS IN SEMI-TRANSITIVE PLANES

N.L. JOHNSON

Department of Mathematics
The University of Iowa
Iowa City, Iowa 52242
U.S.A.

(Received January 2, 1980)

ABSTRACT. Let π be a semi-transitive translation plane of even order with reference to the subplane π_0 . If π admits an affine elation fixing π_0 for each axis in π_0 and the order of π_0 is not 2 or 8, then π is a Hall plane.

KEY WORDS AND PHRASES. Elations, Semi-transitive planes.

1980 MATHEMATICS SUBJECT CLASSIFICATION CODES. 50D05, 05B25.

1. INTRODUCTION.

Kirkpatrick [9] and Rahilly [10] have characterized the Hall planes as those generalized Hall planes of order q^2 that admit $q+1$ central involutions.

In [7] the author has shown that the derived semifield planes of characteristic $\neq 3$ and order q^2 are Hall planes precisely when they admit $q+1$ central involutions. This extends Kirkpatrick and Rahilly's work as generalized Hall planes are certain derived semifield planes.

If a translation plane π of order q^2 admits $q+1$ affine elations with distinct axes then the generated group $\mathcal{G}$ contains $SL(2, q)$, $S_2(q)$ or contains a normal subgroup N of odd order and index 2 (Hering [5]). In the latter case, little is known about $\mathcal{G}$ except that it is usually dihedral.

In this article, we study semi-transitive translation planes of order q^2 that admit $q+1$ affine elations.

In [8], the author introduces the concept of the generalized Hall planes of type 1. These are derivable translation planes that admit a particular collineation group which is transitive on the components outside the derivable net. In this situation the group is generated by Baer collineations.

More generally, Jha [6] has considered the "semi-transitive" translation planes.

(1.1) Let π be a translation plane with subplane π_0 . If there is a collineation group $\mathcal{L}$ such that

- 1) $\mathcal{L}$ fixes $\pi_0 \cap \ell_\infty$ pointwise,
- 2) leaves π_0 invariant, and
- 3) acts transitively on $\ell_\infty - \pi_0 \cap \ell_\infty$,

then π is said to be a semi-transitive translation plane with reference to π_0 and with respect to $\mathcal{L}$.

Our main result is that semi-transitive planes of order not 16 or 64 that admit elations with axis $\mathcal{L}$ fixing π_0 for every component $\mathcal{L}$ of π_0 are Hall planes. We also give a necessary and sufficient condition that a translation plane of order $q^2 \neq 64$ admitting $q+1$ elations with distinct axes is derivable.

2. TRANSLATION PLANES OF EVEN ORDER q^2 ADMITTING $q+1$ ELATIONS.

(2.1) THEOREM. Let π be a translation plane of even order $q^2 \neq 64$ that admits $q+1$ affine elations with distinct axes. Let $\mathcal{N}$ denote the net of degree $q+1$ that is defined by the elation axes and assume the group D generated by these elations leaves $\mathcal{N}$ invariant. Then $\mathcal{N}$ is derivable if and only if D is either isomorphic to $SL(2, q)$ or is dihedral of order $2(q+1)$ where the cyclic stem fixes at least two components.

PROOF. If D is isomorphic to $SL(2, q)$ then $\mathcal{N}$ is derivable and actually π is Desarguesian by Foulser-Johnson-Ostrom [3].

Let $D = \langle \sigma, \chi \mid \sigma^2 = \chi^{q+1} = 1, \sigma\chi = \chi^{-1}\sigma \rangle$. If $\langle \chi \rangle$ fixes the components $X = \mathcal{O}$, $Y = \mathcal{O}$ then we may choose coordinates so that σ is $(x, y) \rightarrow (y, x)$ and χ is $(x, y) \rightarrow (xT, yT^{-1})$ for some matrix T of order $q+1$.

By Ostrom [11], Theorem 3, there is a Desarguesian plane Σ containing the two χ -fixed components and η . Clearly η is an André net in Σ and thus derivable in π .

Conversely, suppose η is derivable. Since each elation fixes η , D must fix each Baer subplane of η incident with $\mathcal{O}$. By Foulser [2], Theorem 3, $D \leq \text{GL}(2, q)$ in its action on π so that $D \leq \text{SL}(2, q)$ (each elation is then in $\text{SL}(2, q)$). By Gleason [4], D is transitive on the elation axes so $q+1 \mid |D|$. Thus, D is clearly $\text{SL}(2, q)$ or is dihedral of order $2(q+1)$. Moreover, if η is derivable then χ fixes at least two infinite points of $\pi - \eta$. Let $\bar{\eta}$ replace η so $\mathcal{L}$ fixes $\bar{\eta}$ componentwise in the derived plane $\bar{\pi}$. Let $\langle \bar{\chi} \rangle \triangleleft \langle \chi \rangle$ such that $|\bar{\chi}|$ is a prime 2-primitive divisor of $q^2 - 1$ (one exists since $q^2 \neq 64$). Then $\bar{\chi}$ fixes at least two infinite points of $\bar{\pi} - \bar{\eta}$ so there is a unique Desarguesian plane Σ containing the $\bar{\chi}$ -fixed components of $\bar{\pi}$ (see Ostrom [11], Cor. to Theorem 1—uniqueness comes from the fact that the degree of $\Sigma \cap \bar{\pi}$ is greater than $q+1$). Since $\mathcal{L}$ permutes the components of $\Sigma \cap \bar{\pi}$ (i.e., $\langle \bar{\chi} \rangle$ is characteristic in $\langle \chi \rangle$), $\mathcal{L}$ is a collineation group of Σ . The collineation χ has the form $(x, y) \rightarrow (x^\phi a, y^\phi a)$ where ϕ is an automorphism of $\text{GF}(q^2)$ and $a \in \text{GF}(q^2)$. (Note χ fixes $\bar{\eta}$ componentwise.) Since $q+1$ is odd, $\langle \chi^2 \rangle = \langle \chi \rangle$. Choosing coordinates so that the components of $\bar{\eta}$ are $X = \mathcal{O}$, $Y = \mathcal{O}$, $y = x\alpha$, $\alpha \in \text{GF}(q^2)$ then χ fixes $y = x\alpha$ for all $\alpha \in \text{GF}(q^2)$ if and only if $\alpha^\phi = \alpha$. Since $\langle \chi^2 \rangle = \langle \chi \rangle$, we may assume $\phi = 1$. Thus, χ fixes ℓ_∞ of Σ pointwise. Since Σ and π share at least two components (those fixed by $\bar{\chi}$), χ must fix at least two components of π .

3. SEMI-TRANSITIVE TRANSLATION PLANES OF EVEN ORDER.

Let π be a translation plane of even order q^2 that admits $q+1$ elations as in section 2. Then, π is a derivable plane provided the generated group D is dihedral and the cyclic stem fixes at least 2 points or $\text{SL}(2, q)$. In any case let η denote the net defined by the elation axes. Let $\mathcal{L}$ be a collineation group that commutes with D . Then clearly, $\mathcal{L}$ must fix $\eta \cap \ell_\infty$ pointwise.

(3.1) THEOREM. Let π be a translation plane of even order $q^2 \neq 64$ that admits $q+1$ elations with distinct axes. Assume the group D generated by these

$q+1$ elations leaves the net $\mathcal{N}$ of the elation axes invariant. Let $\mathcal{L}$ be a collineation group which commutes with D and is transitive on $\ell_\infty - \mathcal{N} \cap \ell_\infty$. Then π is a Hall plane.

PROOF. Since $q^2 \neq 64$, there is a prime 2-primitive divisor m of q^2-1 . By Gleason [4], $q+1 \mid |D|$. Clearly, $m \mid q+1$. Let χ be an element of D of order m . χ acts on the $q(q-1)$ points of $\ell_\infty - \mathcal{N} \cap \ell_\infty$ so must fix at least two points of $\ell_\infty - \mathcal{N} \cap \ell_\infty$. Since $\mathcal{L}$ commutes with χ and $\mathcal{L}$ is transitive on $\ell_\infty - \mathcal{N} \cap \ell_\infty$, χ must fix $\ell_\infty - \mathcal{N} \cap \ell_\infty$ pointwise.

By the corollary to Theorem 1, Ostrom [11], there is a Desarguesian plane Σ such that the components fixed by χ in π are exactly the common components of Σ and π . Let $\pi = \mathcal{N} \cup \mathcal{M}$ where $\mathcal{M}$ is the net complementary to $\mathcal{N}$ in π . Then $\Sigma = \overline{\mathcal{N}} \cup \mathcal{M}$ for some net $\overline{\mathcal{N}}$ of degree $q+1$. So Σ and π are two extensions of a net $\mathcal{M}$ of critical deficiency (see Ostrom [12]). Then π must be Hall since Σ and π must be related by derivation (i.e., π cannot be itself Desarguesian) by Ostrom [12].

The conditions of (3.1) are close to giving the definition of a "semi-transitive" translation plane (see (1.1)). In (3.1), it is possible that $\mathcal{L}$ may not satisfy condition 2. Also, it is not clear that a semi-transitive translation plane is derivable. However, Jha [6] shows if π has order not 16 and there is a nontrivial kern homology in π then π is derivable and π_0 is a Baer subplane.

We may overcome this restriction on the kern in our situation:

(3.2) THEOREM. Let π be a semi-transitive translation plane of even order with respect to a collineation group $\mathcal{L}$ and with reference to a subplane π_0 . Let π admit an affine elation for each axis in π_0 .

- 1) If the order of π_0 is not 8 then π is derivable.
- 2) If the order of π_0 is not 2 or 8 then π is a Hall plane.

PROOF. Following Jha's [6] ideas, let π_1 be a minimal subplane of π properly containing π_0 . Clearly, the stabilizer $\mathcal{L}_{\pi_1}$ of π_1 is a semi-transitive collineation group of π_1 with reference to π_0 . Moreover, a Sylow 2-subgroup of $\mathcal{L}_{\pi_1}$ must leave π_0 pointwise fixed since $\mathcal{L}$ fixes π_0 and fixes $\pi_0 \cap \ell_\infty$ pointwise. (Note $|\mathcal{L}_{\pi_1}|$ is divisible by $(2^r+1) - (2^s+1)$ for some r, s .) Clearly, π_0 is a Baer subplane of π_1 .

Every elation which leaves π_0 invariant must also leave any superplane invariant. So the group D generated by the elations leaves π_1 invariant and, clearly, $\mathcal{E}$ commutes with D since $\mathcal{E}$ fixes $\pi_0 \cap \ell_\infty$ pointwise ($\mathcal{E}$ must commute with each central collineation fixing π_0).

By (3.1), if the order of π_0 is not 8 then π_1 is a Hall plane and π_1 is derivable. We may now directly use Jha [6] to show that if the order of π_0 is not 2 then $\pi_1 = \pi$ (that is, Jha uses the hypothesis that there is a kern homology to show that π_1 is derivable).

Actually, our proof of (3.2) proves the following more general theorem for arbitrary order.

(3.3) THEOREM. Let π be a semi-transitive translation plane with reference to π_0 and with respect to $\mathcal{E}$ and order p^r . Let χ be a collineation generated by central collineations leaving π_0 invariant such that $|\chi|$ is a prime p -primitive divisor of $(\text{order } \pi_0)^2 - 1$ (where the order of π_0 is not 2). Then π is a Hall plane.

Note that a semi-transitive plane of odd order p^{2r} must admit Baer p -elements (see Jha [6]). By Foulser [1], we could then not have both Baer p -elements and elations so we could restate our Theorem (3.2) without reference to order.

(3.2)2) is also valid if the order π_0 is 8. The arguments supporting this will appear in a related article.

REFERENCES

1. Foulser, D. A. Baer p -elements in translation planes, J. Algebra 31(1974) 354-366.
2. Foulser, D. A. Subplanes of partial spreads in translation planes, Bull. London Math. Soc. 4(1972) 1-7.
3. Foulser, D. A., N. L. Johnson and T. G. Ostrom. Characterization of the Desarguesian and Hall planes of order q^2 by $SL(q, 2)$, submitted.
4. Gleason, A. M. Finite Fano planes, Amer. J. Math. 78(1956) 797-806.
5. Hering, Ch. On shears of translation planes, Abh. Math. Sem. Hamburg 27(1972) 258-268.
6. Jha, V. Finite semi-transitive translation planes, to appear.
7. Johnson, N. L. On central collineations of derived semifield planes, J. Geom. 11/2(1978) 139-149.

8. Johnson, N. L. Distortion and generalized Hall planes, Geom. Ded. 4(1975) 1-20.
9. Kirkpatrick, P. B. A characterization of the Hall planes of odd order, Bull. Austral. Math. Soc. 6(1972) 407-415.
10. Rahilly, A. Finite collineation groups and their collineation groups, Thesis, Univ. of Sydney, 1973.
11. Ostrom, T. G. Linear transformations and collineations of translation planes, J. Algebra (3) 14(1970) 405-416.
12. Ostrom, T. G. Nets with critical deficiency, Pacific J. Math. 14(1964), 1381-1387.