SOME GENERATING FUNCTIONS OF MODIFIED BESSEL POLYNOMIALS FROM THE VIEW POINT OF LIE GROUP

ASIT KUMAR CHONGDAR

Department of Mathematics
Bangabasi Evening College
19 Scott Lane, Calcutta - 700 009.
India

(Received May 31, 1984)

ABSTRACT. In this paper we have derived a class of bilateral generating relation for modified Bessel polynomials from the view point of Lie group. An application of our theorem is also pointed out.

KEY WORDS AND PHRASES. *Bessel Polynomials, Generating functions.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE. 33A70

1. INTRODUCTION.

In [1], the modified Bessel polynomials are defined by

$$y_n(x; \alpha-n, \beta) = {}_2F_0(-n, \alpha-1; -; -x/\beta), \quad (1.1)$$

where $y_n(x; \alpha, \beta)$ denotes the Bessel polynomials introduced by H. L. Krall and O. Frink [2].

The object of the present paper is to derive the following theorem on bilateral generating relation for modified Bessel polynomials from the view point of Lie-group.

THEOREM. If there exists a generating relation of the form,

$$G(x, w) = \sum_{n=0}^{\infty} a_n w^n y_n(x; \alpha-n, \beta) \quad (1.2)$$

then

$$(1-nw)^{1-\alpha} e^{\beta w} \left(G\left(\frac{x}{1-xw}, wz\right) \right) = \sum_{n=0}^{\infty} w^n g_n(z) y_n(x; \alpha-n, \beta) \quad (1.3)$$

where

$$g_n(z) = \sum_{m=0}^n \frac{a_m}{m!} (\beta z)^m.$$

The importance of our result lies in the fact that if one knows a generating relation of the type (1.2) for a particular value of a_n , then the corresponding bilateral generating relation follows atonce from (1.3).

2. PROOF OF THE THEOREM.

From [1] we observe

$$\exp(wR) f(x, y) = (1-wxy)^{1-\alpha} e^{\beta wy} f\left(\frac{x}{1-wxy}, y\right) \quad (2.1)$$

where

$$R = x^2 y \frac{\partial}{\partial x} + \beta y + (\alpha-1)xy$$

and

$$R(F_n(x, y, \alpha-n, \beta)) = \beta F_{n+1}(x, y, \alpha-n-1, \beta) \quad (2.2)$$

where

$$F_n(x, y, \alpha-n, \beta) = y^n y_n(x; \alpha-n, \beta).$$

In the formula:

$$G(x, w) = \sum_{n=0}^{\infty} a_n w^n y_n(x; \alpha-n, \beta)$$

replacing w by wyz and then operating both sides by $\exp(wR)$, we get

$$\exp(wR) G(x, wyz) = \exp(wR) \sum_{n=0}^{\infty} a_n (wz)^n F_n(x, y, \alpha-n, \beta).$$

The first member of (2.3) is equal to

$$(1-wxy)^{1-\alpha} e^{\beta wy} G\left(\frac{x}{1-wxy}, wyz\right),$$

and the second member of (2.3) is equal to

$$\begin{aligned} & \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} a_n \frac{w^{m+n}}{m!} z^n R^m (F_n(x, y, \alpha-n, \beta)) \\ &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} a_n \frac{w^{m+n}}{m!} z^n \beta^m F_{n+m}(x, y, \alpha-n-m, \beta) \\ &= \sum_{n=0}^{\infty} (wy)^n g_n(z) y_n(x; \alpha-n, \beta) \end{aligned}$$

where

$$g_n(z) = \sum_{m=0}^{\infty} a_m \frac{(\beta z)^m}{m!}.$$

Equating the above two results and then putting $y=1$, we get

$$(1-wx)^{1-\alpha} e^{\beta w} G\left(\frac{x}{1-wx}, wz\right) = \sum_{n=0}^{\infty} w^n g_n(z) y_n(x; \alpha-n, \beta)$$

where

$$g_n(x) = \sum_{m=0}^n a_m \frac{(\beta z)^m}{m!},$$

this completes the proof of our theorem.

3. APPLICATION.

As an application of our theorem we consider the following generating relation [3].

$$\sum_{n=0}^{\infty} \frac{w^n}{n!} y_n(x; \alpha-n, \beta) = e^w (1 - \frac{wx}{\beta})^{1-\alpha} \quad (3.1)$$

If, in our theorem, we put $a_n = 1/n!$ we obtain

$$e^{(\beta+z)w} (1 - \frac{(\beta+z)w}{\beta} x)^{1-\alpha} = \sum_{n=0}^{\infty} w^n g_n(z) y_n(x; \alpha-n, \beta) \quad (3.2)$$

where

$$g_n(z) = \sum_{m=0}^{\infty} \frac{(\beta z)^m}{(m!)^n}.$$

REFERENCES

1. GUHATHAKURTA, B.K., and CHONGDAR, A. K., Some generating functions of modified Bessel polynomials by Lie algebraic method, communicated, 1983.
2. KRALL, H. L., and FRINK, O., A new class of orthogonal polynomials: The Bessel polynomials, Trans. Amer. Math. Soc. 65 (1949), 100-115.
3. GROSSWALD, E., Bessel polynomials, Lecture notes in Mathematics: Springer-Verlag, Berlin, Heidelberg, New York 1978, 44.