RESEARCH NOTES

AN INTEGRAL EQUATION ASSOCIATED WITH LINEAR HOMOGENEOUS DIFFERENTIAL EQUATIONS

A.K. BOSE

Department of Mathematical Sciences
Clemson University
Clemson, South Carolina 29631 U.S.A.

(Received March 7, 1985)

ABSTRACT. Associated with each linear homogeneous differential equation

$y^{(n)} = \sum_{i=0}^{n-1} a_i(x)y^{(i)}$ of order n on the real line, there is an equivalent integral equation

$$f(x) = f(x_0) + \int_{x_0}^x h(u)du + \int_{x_0}^x \left[\int_{x_0}^u G_{n-1}(u,v)a_0(v)f(v)dv \right] du$$

which is satisfied by each solution $f(x)$ of the differential equation.

KEY WORDS AND PHRASES. *Linear homogeneous differential equations, Integral equations, Initial value problems, Variation of parameters formula, Uniform convergence.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE. 34A

1. INTRODUCTION.

Let n be a positive integer, I be an interval on the real line R and $C(I)$ be the class of all functions continuous on I . Let

$$y^{(n)} = a_{n-1}(x)y^{(n-1)} + \dots + a_0(x)y, \quad x \in I \quad (1.1)$$

be any n -th order (normalized) ordinary linear homogeneous differential equation, where $a_i(x) \in C(I)$, $i=0,1,2,\dots,(n-1)$.

The purpose of this article is to derive an equivalent integral equation satisfied by the solutions of the linear homogeneous differential equation (1.1).

2. MAIN RESULTS.

THEOREM. Let $f(x)$ be a solution of (1.1) defined on I and $x_0 \in I$. Then $f(x)$ is also a solution of the integral equation

$$f(x) = f(x_0) + \int_{x_0}^x h(u)du + \int_{x_0}^x \left[\int_{x_0}^u G_{n-1}(u,v)a_0(v)f(v)dv \right] du, \quad (2.1)$$

where $h(x)$ is the unique solution of the $(n-1)$ -th order linear homogeneous differential equation

$$y^{(n-1)} = a_{n-1}(x)y^{(n-2)} + \dots + a_1(x)y, \quad x \in I \quad (2.2)$$

$x_0 \in [a, b]$.

Let $M = \max |G_{n-1}(u, v)a_0(v)|$, $s = \max |f_1(v)|$ for each $u, v \in [a, b]$. One can now see very easily that for each $x \in [a, b]$,

$$|f_{k+1}(x) - f_k(x)| \leq M^k s^k (b-a)^{2k}/2k!$$

$$|f'_{k+1}(x) - f'_k(x)| \leq M^k s^k (b-a)^k/k!,$$

using recursively the bounds for $|f_{i+1}(x) - f_i(x)|$, $i = 1, 2, 3, \dots$. Since each of the series $\sum M^k s^k (b-a)^{2k}/2k!$, $\sum M^k s^k (b-a)^k/k!$ converges, we conclude by Weierstrass' M-test that each of the series of functions

$$f_1(x) + \sum_{k=1}^{\infty} (f_{k+1}(x) - f_k(x)), \quad f'_1(x) + \sum_{k=1}^{\infty} (f'_{k+1}(x) - f'_k(x))$$

converges uniformly on $[a, b]$ and hence on B . Therefore there is a function $g(x) \in C^1(I)$ such that

$$f_1(x) + \sum_{k=1}^{\infty} (f_{k+1}(x) - f_k(x)) = \lim_{k \rightarrow \infty} f_k(x) = g(x)$$

$$f'_1(x) + \sum_{k=1}^{\infty} (f'_{k+1}(x) - f'_k(x)) = \lim_{k \rightarrow \infty} f'_k(x) = g'(x)$$

for all $x \in I$. In particular $g(x_0) = f(x_0)$.

Also from (2.5) we get by taking limit as $k \rightarrow \infty$

$$g'(x) = h(x) + \int_{x_0}^x G_{n-1}(x, u)a_0(u)g(u)du, \quad x \in I. \quad (2.7)$$

Hence

$$g(x) = f(x_0) + \int_{x_0}^x h(u)du + \int_{x_0}^x \left[\int_{x_0}^u G_{n-1}(u, v)a_0(v)g(v)dv \right] du \quad (2.8)$$

Again, relation (2.7) implies by (2.3) that $g'(x)$ is the unique solution of the initial value problem

$$y^{(n-1)} = a_{n-1}(x)y^{(n-2)} + \dots + a_1(x)y + a_0(x)g(x)$$

$$y'(x_0) = f'(x_0), \dots, y^{(n-1)}(x_0) = f^{(n-1)}(x_0).$$

Therefore $g(x) \in C^n(I)$ and

$$g^{(n)}(x) = a_{n-1}(x)g^{(n-1)}(x) + \dots + a_0(x)g(x), \quad x \in I.$$

In other words, $g(x)$ is the unique solution of the homogeneous initial value problem

$$y^{(n)} = a_{n-1}(x)y^{(n-1)}(x) + \dots + a_0(x)y, \quad x \in I.$$

$$y(x_0) = f(x_0), \quad y'(x_0) = f'(x_0), \dots, y^{(n-1)}(x_0) = f^{(n-1)}(x_0).$$

Hence $f(x) = g(x)$ for all $x \in I$. Therefore, by (2.8)

$$f(x) = f(x_0) + \int_{x_0}^x h(u)du + \int_{x_0}^x \left[\int_{x_0}^u G_{n-1}(u, v)a_0(v)f(v)dv \right] du.$$

This completes the proof.

REMARK. The above proof clearly shows how a solution of a linear homogeneous equation with prescribed initial values can be constructed out of a solution $h(x)$ and the Green's Function $G_{n-1}(x,u)$ of a lower order homogeneous linear equation. This is specially significant in case of second order homogeneous equations, as solutions $\{ce^{A(x)}\}$ and the Green's function $G_1(x,u) = e^{A(x)-A(u)}$,

$[A(x) = \int_{x_0}^x a_1(u)du]$, of first order homogeneous equation $y' = a_1(x)y$ are readily

available.

REFERENCES

1. CODDINGTON, E.A. An Introduction to Ordinary Differential Equations, Englewood Cliffs, New Jersey: Prentice-Hall, Inc.
2. WIDDER, D.V. Advanced Calculus, Englewood Cliffs, New Jersey: Prentice-Hall, Inc.