ON ALPHA-CLOSE-TO-CONVEX FUNCTIONS OF ORDER BETA

M.A. NASR

Faculty of Science
University of Mansoura
Mansoura, Egypt

(Received January 9, 1985)

ABSTRACT Let $M_\beta(\alpha)$ [$\alpha \geq 0$ and $\beta \geq 0$] denote the class of all functions

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

analytic in the unit disc U with $f'(z)f(z)/z \neq 0$ and which satisfy for $z=re^{i\theta} \in U$ the condition

$$\int_{\theta_1}^{\theta_2} \operatorname{Re} \left\{ (1-\alpha) \frac{zf'(z)}{f(z)} + \alpha \left(1 + \frac{zf''(z)}{f'(z)} \right) \right\} d\theta > -\beta\pi$$

for all $\theta_2 > \theta_1$. In this note we show that each $f \in M_\beta(\alpha)$ is close-to-star of order β when $0 < \beta \leq \alpha$.

KEY WORDS AND PHRASES. Close-to-star functions, close-to-convex functions, α -convex functions.

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE. 30 C45

I. INTRODUCTION.

A shall denote the class of all functions

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

analytic in the unit disc $U = \{z: |z| < 1\}$ and S shall denote the subclass of functions in A which are univalent in U .

Let $\alpha \geq 0$, $\beta \geq 0$, and let $f \in A$ with $f'(z)f(z)/z \neq 0$ in U , and let

$$J(\alpha, f) = (1-\alpha) \frac{zf'(z)}{f(z)} + \alpha \left(1 + \frac{zf''(z)}{f'(z)} \right). \quad (1.1)$$

If for $z=re^{i\theta} \in U$,

$$\int_{\theta_1}^{\theta_2} \left\{ \operatorname{Re} J(\alpha, f) \right\} d\theta > -\beta\pi \quad (1.2)$$

whenever $0 \leq \theta_1 < \theta_2 < 2\pi$, then f is said to be an α -close-to-convex function of order β or $f \in M_\beta(\alpha)$. The class $M_\beta(\alpha)$ was introduced by Nasr[1].

It was shown [1] that $M_\beta(\alpha) \subset S$ if and only if $0 \leq \beta \leq \alpha$.

Note that $M_\beta(1) = K_\beta$, is the class of close-to-convex functions of order β introduced by Reade [2] and studied by Goodman [3] for $\beta \geq 1$ and by Pommerenke [4] for $0 \leq \beta \leq 1$, and $M_\beta(0) = R_\beta$ is the class of close-to-star functions of order β introduced by Goodman [3]. Moreover $M_0(\alpha) = M(\alpha)$ is the class of α -convex functions introduced by Mocanu [5], and $M_{\gamma/\alpha}(\frac{1}{\alpha}) = B_\gamma(\alpha)$, $\alpha > 0$, is the class of Bazilevič functions of order γ introduced by Nasr [6].

In this note we continue the investigation of α -close-to-convex functions of order β studied in [1].

2. RESULTS

In this section we show that each $f \in M_\beta(\alpha)$ is close-to-star of order β when $0 < \beta \leq \alpha$. For $\alpha \geq 1$ we show each $f \in M_\beta(\alpha)$ is close-to-convex of order β when $0 < \beta \leq \alpha$ and if $f \in M_\beta(\alpha)$, then $f \in M_\beta(\gamma)$ when $0 < \beta \leq \gamma \leq \alpha$.

We assume, unless otherwise stated, that θ is a real number, that $0 < r < 1$ and that $z = re^{i\theta}$. Also that $0 < \beta \leq \alpha$.

We shall need the following result.

LEMMA 1: If $f \in M_\beta(\gamma)$, then the function h given by

$$h(z) = f(z) \cdot (zf'(z)/f(z))^\alpha \quad (2.1)$$

belongs to R_β . (The powers taken are the principal values).

PROOF: Let $f \in M_\beta(\alpha)$. If we choose the branch of $(zf'(z)/f(z))^\alpha$ which is equal to 1 when $z = 0$, a simple calculation shows that the function h defined by (2.1) belongs to R_β .

THEOREM 1: If $f \in M_\beta(\alpha)$ then $f \in R_\beta$.

PROOF: Since $f \in M_\beta(\alpha)$, it follows from Lemma 1 that

$$\int_{\theta_1}^{\theta_2} \operatorname{Re} \left\{ \frac{\frac{d}{dz} h(z)^{1/\alpha}}{\frac{d}{dz} f(z)^{1/\alpha}} \right\} d\theta = \int_{\theta_1}^{\theta_2} \operatorname{Re} \left\{ \frac{zh'(z)}{h(z)} \right\} d\theta > -\pi\beta. \quad (2.2)$$

In (2.2) we choose the branches for $f(z)^{1/\alpha}$ and $h(z)^{1/\alpha}$ for which

$$h(z)^{1/\alpha}/f(z)^{1/\alpha} = (h(z)/f(z))^{1/\alpha} \quad (2.3)$$

with value 1 for $z=0$. If we use (2.1), (2.2) and (2.3) it is easy to prove that

$$\int_{\theta_1}^{\theta_2} \operatorname{Re} \left\{ h(z)/f(z)^{1/\alpha} \right\} d\theta > -\pi\beta. \quad (2.4)$$

In fact, since f is univalent, we can let $w = f(z)$, $z = z(w) = f^{-1}(w)$ and $w = \rho e^{i\phi}$ to obtain

$$\begin{aligned}
\frac{h(z)^{1/\alpha}}{f(z)^{1/\alpha}} &= \frac{1}{w^{1/\alpha}} \int_0^w \frac{d}{dw} \left[h(z(w)) \right]^{1/\alpha} dw \\
&= \frac{1}{\rho^{1/\alpha}} \int_0^\rho \frac{d \left[\frac{h(z(\rho e^{i\phi}))}{(\rho e^{i\phi})^{1/\alpha}} \right]^{1/\alpha}}{d \left(\frac{\rho e^{i\phi}}{\rho e^{i\phi}} \right)^{1/\alpha}} \frac{1/\alpha}{\rho^{1/\alpha - 1}} d\rho \\
&= \frac{1}{\rho^{1/\alpha}} \int_0^\rho \frac{d \left[\frac{h(z)}{[f(z)]^{1/\alpha}} \right]^{1/\alpha}}{d \left[\frac{h(z)}{[f(z)]^{1/\alpha}} \right]^{1/\alpha}} \frac{1/\alpha}{\rho^{1/\alpha - 1}} d\rho.
\end{aligned}$$

Hence

$$\begin{aligned}
&\int_{\theta_1}^{\theta_2} \operatorname{Re} \left\{ \frac{h(z)^{1/\alpha}}{f(z)^{1/\alpha}} \right\} d\theta = \\
&\frac{1}{\rho^{1/\alpha}} \int_0^\rho \int_{\theta_1}^{\theta_2} \left[\operatorname{Re} \left\{ \frac{d[h(z)]^{1/\alpha}}{d[f(z)]^{1/\alpha}} \right\} d\theta \right] \frac{1/\alpha}{\rho^{1/\alpha - 1}} d\rho.
\end{aligned}$$

The result now follows from (2.1) and (2.4).

COROLLARY 1: If $f \in M_\beta(\alpha)$, $\alpha \geq 1$, then $f \in K_\beta$.

PROOF: Let $f \in M_\beta(\alpha)$, $\alpha \geq 1$, then

$$\alpha \int_{\theta_1}^{\theta_2} \operatorname{Re} \left\{ \left(1 + \frac{zf''(z)}{f'(z)} \right) \right\} d\theta > (\alpha-1) \int_{\theta_1}^{\theta_2} \operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} d\theta - \pi\beta.$$

Now from THEOREM 1, we have $\int_{\theta_1}^{\theta_2} \operatorname{Re} \left\{ zf'(z)/f(z) \right\} d\theta > -\pi\beta$,

and therefore

$$\alpha \int_{\theta_1}^{\theta_2} \operatorname{Re} \left\{ \left(1 + \frac{zf''(z)}{f'(z)} \right) \right\} d\theta > -(\alpha-1)\pi\beta - \pi\beta = -\alpha\pi\beta$$

and the proof of Corollary 1 is complete.

COROLLARY 2: If $f \in M_\beta(\alpha)$, $0 < \beta \leq \gamma \leq \alpha$, then $f \in M_\beta(\gamma)$.

PROOF:

By THEOREM 1, $f \in R_\beta$. Suppose there exists a γ , $0 < \beta \leq \gamma \leq \alpha$, such that $f \in M_\beta(\gamma)$. Then there is $\tau \in U$ for which

$$\begin{aligned}
&\int_{\theta_1}^{\theta_2} \left\{ \operatorname{Re} \left[\frac{\tau f''(\tau)}{f'(\tau)} + 1 - \frac{\tau f'(\tau)}{f(\tau)} \right] \right\} d\theta \\
&< -\frac{\pi\beta}{\gamma} - \frac{1}{\gamma} \int_{\theta_1}^{\theta_2} \left\{ \operatorname{Re} \left[\frac{\tau f'(\tau)}{f(\tau)} \right] \right\} d\theta. \tag{2.5}
\end{aligned}$$

However, for $f \in M_\beta(\alpha)$,

$$0 < \pi\beta + \int_{\theta_1}^{\theta_2} \left\{ \operatorname{Re} \left[\frac{\tau f'(\tau)}{f(\tau)} \right] \right\} d\theta + \alpha \int_{\theta_1}^{\theta_2} \left\{ \operatorname{Re} \left[\frac{\tau f''(\tau)}{f'(\tau)} + 1 - \frac{\tau f'(\tau)}{f(\tau)} \right] \right\} d\theta. \tag{2.6}$$

Substituting (2.5) into (2.6), we obtain

$$0 < (1 - \frac{\alpha}{\gamma}) [\beta\pi + \int_{\theta_1}^{\theta_2} \left\{ \operatorname{Re} \left[\frac{\tau f'(\tau)}{f(\tau)} \right] \right\} d\theta].$$

But $(1 - \frac{\alpha}{\gamma}) < 0$ implies $\int_{\theta_1}^{\theta_2} \left\{ \operatorname{Re} \left[\frac{\tau f'(\tau)}{f(\tau)} \right] \right\} d\theta < -\pi\beta$, which contradicts the assumption that $f \in R_\beta$. Thus $f \in M_\beta(\gamma)$.

REMARK:

For $\beta = 0$ we obtain results due to Miller, Mocanu and Reade [7].

ACKNOWLEDGMENT

The author would like to thank Professor M. O. Reade for reading the manuscript and giving some helpful suggestions.

REFERENCES

1. NASR, M. A., Alpha-Close-to-Convex Functions of Order Beta, Mathematica 24 (1982) 79-83.
2. READE, M. O., The Coefficients of Close-to-Convex Functions, Duke Math. Journal 23 (1950), 459-462.
3. GOODMAN, A. W., On Close-to-Convex Functions of Higher Order, Ann. Univ. Sci. Budapest Eotvos sect. Math. 25 (1972), 17-30.
4. POMMERENKE, C., On the Coefficients of Close-to-Convex Functions, Michigan Math Journal, 9 (1962), 259-269.
5. MOCANU, P. T., Une Propriete de Convexite Generalisee dans la Theorie de la Representation Conform, Mathematica 11 (1969) 127-133.
6. NASR, M. A., Bazilevici Functions of Higher Order, Rev. Roum. Math. Pures et Appl. 28 (1983) 709-714.
7. MILLER, S. S., MOCANU, P. T., AND READE, M. A., All α -Convex Functions are Univalent and Starlike, Proc. Amer. Math. Soc. 37 (1973) 552-554.