ON THE DUAL SPACE OF A WEIGHTED BERGMAN SPACE ON THE UNIT BALL OF C^n

J.S. CHOA and H.O. KIM

Department of Applied Mathematics
KAIST
P.O. Box 150, Cheongryang
Seoul, KOREA

(Received January 30, 1987 and in revised form March 24, 1987)

ABSTRACT. The weighted Bergman space $A_\alpha^p(B_n)$ ($0 < p < \infty$), of the holomorphic functions on the unit ball B_n of C^n forms an F-space. We find the dual space of $A_\alpha^p(B_n)$ by determining its Mackey topology.

KEY WORDS AND PHRASES. Hardy space, Bergman space, Mackey topology.

1980 AMS SUBJECT CLASSIFICATION CODE. 30D55, 32A35.

1. INTRODUCTION.

Let B_n be the unit ball of C^n , ν be the normalized Lebesgue measure and σ be the rotation invariant positive Borel measure on S , the boundary of B_n , with $\sigma(S) = 1$. The weighted Bergman space $A_\alpha^p(B_n)$ ($0 < p < \infty, \alpha \geq -1$) consists of all functions holomorphic in B_n for which

$$\|f\|_{p,\alpha}^p = \begin{cases} \int_0^1 M_p^p(r;f) (1-r)^\alpha d\sigma(\zeta) 2nr^{2n-1} dr < \infty, & \text{if } \alpha > -1, \\ \sup_{0 \leq r < 1} M_p^p(r;f) < \infty, & \text{if } \alpha = -1, \end{cases}$$

where $M_p^p(r;f) = \int_S |f(r\zeta)|^p d\sigma(\zeta)$. Note that the weighted Bergman space $A_\alpha^p(B_n)$ is, in fact, the Hardy space $H^p(B_n)$ if $\alpha = -1$ (See [1]).

The purpose of this paper is to compute the dual space $(A_\alpha^p(B_n))^*$ for $0 < p \leq 1$ by determining the Mackey topology of $A_\alpha^p(B_n)$. The corresponding problems for the case $n = 1$ are settled by Duren, Romberg and Shields [2], Shapiro [3] and Ahern [4]. Our computations are very similar to those of them.

Throughout this work, $C_{\alpha,\beta,\dots}$ denotes a positive constant depending only on $\alpha, \beta, \dots$ which may vary in the various places, and the notation $a(z) \sim b(z)$ means that the ratio $a(z)/b(z)$ has a positive finite limit as $|z| \rightarrow 1$.

2. SOME PRELIMINARY RESULTS.

LEMMA 2.1. If $f \in A_\alpha^p(B_n)$ ($0 < p < \infty, \alpha \geq -1$), then

$$|f(z)| \leq C_{n,p,\alpha} \|f\|_{p,\alpha} (1 - |z|)^{\frac{n+\alpha+1}{p}}.$$

PROOF. The case $\alpha = -1$ is proved in [1, Thm 7.2.5]. For the proof of the case $\alpha > -1$, it is enough to prove the result for $\frac{1}{2} \leq |z| < 1$ since $|f(z)|$ is bounded for $|z| \leq \frac{1}{2}$. For this range of ρ we have:

$$\begin{aligned} \|f\|_{p,\alpha}^p &= \int_0^1 \int_S |f(r\zeta)|^p (1-r)^{\alpha} 2nr^{2n-1} dr d\sigma(\zeta) \\ &\geq C_{n,p} \int_\rho^1 \int_S |f(r\zeta)|^p (1-r)^{\alpha} d\sigma(\zeta) dr \\ &\geq C_{n,p} M_p^p(\rho; f) \int_\rho^1 (1-r)^{\alpha} dr \\ &= C_{n,p,\alpha} M_p^p(\rho; f) (1-\rho)^{\alpha+1}. \end{aligned}$$

By the result of the case $\alpha = -1$ and the above result, we get

$$\begin{aligned} |f(\rho z)| &\leq C_{n,p} M_p(\rho; f) (1-|z|)^{-\frac{n}{p}} \\ &\leq C_{n,p,\alpha} \|f\|_{p,\alpha} (1-\rho)^{-\frac{\alpha+1}{p}} (1-|z|)^{-\frac{n}{p}}. \end{aligned}$$

Consequently we have

$$\begin{aligned} |f(z)| &= |f(\sqrt{r}\sqrt{r}\zeta)| \quad (z = r\zeta) \\ &\leq C_{n,p,\alpha} \|f\|_{p,\alpha} (1-\sqrt{r})^{-\frac{\alpha+1}{p}} (1-\sqrt{r})^{-\frac{n}{p}} \\ &\leq C_{n,p,\alpha} \|f\|_{p,\alpha} (1-r)^{-\frac{n+\alpha+1}{p}}. \end{aligned}$$

COROLLARY 2.2. (a) The convergence of $A_\alpha^p(B_n)$ with its invariant metric

$$d(f,g) = \begin{cases} \|f-g\|_{p,\alpha}^p, & (0 < p < 1), \\ \|f-g\|_{p,\alpha}, & (p \geq 1) \end{cases}$$

implies the uniform convergence on any compact subset of B_n .

(b) $A_\alpha^p(B_n)$ is an F-space if $0 < p < 1$ and a Banach space if $p \geq 1$.

PROOF. (a) follows immediately from Lemma 2.1. The proof of (b) is routine and is omitted.

COROLLARY 2.3. $A_\alpha^p(B_n) \subset A_\beta^q(B_n)$ if $0 < p < q$ and $\frac{n+\alpha+1}{p} = \frac{n+\beta+1}{q}$.

In particular,

$$A_\alpha^p(B_n) \subset A_\sigma^1(B_n), \quad \text{where } \sigma = \frac{n+\alpha+1}{p} - (n+1).$$

PROOF. First, we prove the case $\alpha > -1$. We use Lemma 2.1 in the first inequality of the following.

$$\begin{aligned}
\|f\|_{q,\beta}^q &= \int_0^1 \int_S |f(r\zeta)|^q (1-r)^{\beta} 2nr^{2n-1} dr d\sigma(\zeta) \\
&\leq \int_0^1 \int_S |f(r\zeta)|^p C_{n,p,\alpha,q} \left[(1-r)^{-\frac{n+\alpha+1}{p}} \right]^{q-p} \|f\|_{p,\alpha}^{q-p} \\
&\quad \times (1-r)^{\beta} 2nr^{2n-1} dr d\sigma(\zeta) \\
&\leq C_{n,p,\alpha,q} \|f\|_{p,\alpha}^{q-p} \int_0^1 \int_S |f(r\zeta)|^p (1-r)^{\alpha} 2nr^{2n-1} dr d\sigma(\zeta) \\
&\leq C_{n,p,\alpha,q} \|f\|_{p,\alpha}^q.
\end{aligned}$$

This completes the proof of the case $\alpha > -1$. The remaining case is essentially a result of Hardy and Littlewood, but we give a proof using Ahern's technique in [5]. Let $f \in H^p(B_n)$. By Lemma 2.1, we have

$$|f(z)| \leq K_{n,p} (1 - |z|)^{-\frac{n}{p}} \|f\|_p.$$

Set

$$Mf(\zeta) = \sup_{0 \leq r < 1} |f(r\zeta)|.$$

Then we have

$$\int_0^1 |f(r\zeta)| (1-r)^{\left(\frac{1}{p}-1\right)n-1} 2nr^{2n-1} dr \quad (2.1)$$

$$\begin{aligned}
&\leq K_{n,p} \|f\|_p \int_0^\lambda (1-r)^{-n-1} dr + C_{n,p} Mf(\zeta) \int_\lambda^1 (1-r)^{\left(\frac{1}{p}-1\right)n-1} dr \\
&\leq K_{n,p} \|f\|_p \frac{(1-\lambda)^{-n}}{n} + C_{n,p} Mf(\zeta) \frac{(1-\lambda)^{\left(\frac{1}{p}-1\right)n}}{\left(\frac{1}{p}-1\right)n}.
\end{aligned} \quad (2.2)$$

If $Mf(\zeta) \leq K_{n,p} \|f\|_p$, by setting $\lambda = 0$ in (2.2), (2.1) is dominated by $C_{n,p} \|f\|_p$.

If $Mf(\zeta) \geq K_{n,p} \|f\|_p$, by setting

$$\lambda = 1 - \left[\frac{K_{n,p} \|f\|_p}{Mf(\zeta)} \right]^{\frac{p}{n}}$$

in (2.2), (2.1) is dominated by

$$C_{n,p} \|f\|_p^{1-p} Mf(\zeta)^p.$$

Hence, for any $\zeta \in S$,

$$(2.1) \leq C_{n,p} \|f\|_p + C_{n,p} \|f\|_p^{1-p} Mf(\zeta)^p. \quad (2.3)$$

Integrating (2.3) with respect to $d\sigma(\zeta)$ over S and using the complex maximal theorem [1, Thm. 5.6.5], we obtain

$$\begin{aligned}
& \int_S \int_0^1 |f(r\zeta)| (1-r)^{\left(\frac{1}{p}-1\right)n-1} 2nr^{2n-1} dr d\sigma(\zeta) \\
& \leq C_{n,p} \|f\|_p + C_{n,p} \|f\|_p^{1-p} \|f\|_p^p \\
& \leq C_{n,p} \|f\|_p.
\end{aligned}$$

3. THE MACKEY TOPOLOGY OF $A_\alpha^p(B_n)$.

In this section, we will show that the Mackey topology of $A_\alpha^p(B_n)$ is the restriction of the topology of $A_\sigma^1(B_n)$, where $\sigma = \frac{n+\alpha+1}{p} - (n+1)$.

First we give necessary definitions.

DEFINITION 3.1. The Mackey topology of a non-locally convex topological vector space (X, τ) is the unique locally convex topology m on X satisfying the following conditions:

- (1) m is weaker than τ ,
- (2) the τ -closure of the absolutely convex hull of each τ -neighborhood of the origin contains an m -neighborhood of the origin (See [6, Thm 1]).

DEFINITION 3.2. For $\beta > -n$ and $z, w \in B_n$, we define

$$K_\beta(z, w) = \binom{n+\beta}{n} \frac{(1-|w|^2)^\beta}{(1-\langle z, w \rangle)^{\beta+n+1}}$$

and

$$J_{\beta, \sigma}(w)(z) = (1-|w|^2)^{-\sigma} K_\beta(z, w).$$

The following proposition is useful in the sequel:

PROPOSITION 3.3. [1, p. 120] If $\beta > -n$, then $K_\beta(z, w)$ is a reproducing kernel for the holomorphic functions in $L^1((1-|w|^2)^\beta dv(w))$. In other words, if f is holomorphic on B_n and integrable with respect to the measure $(1-|w|^2)^\beta dv(w)$, then

$$f(z) = \int_{B_n} K_\beta(z, w) f(w) dv(w).$$

LEMMA 3.4. [1, Prop. 1.4.10] For $z \in B_n$ and c real, we define

$$I_c(z) = \int_S \frac{d\sigma(\zeta)}{|1-\langle z, \zeta \rangle|^{n+c}}.$$

If $c > 0$, then $I_c(z) \sim (1-|z|^2)^{-c}$.

LEMMA 3.5. [7, Lemma 6] If $0 < r, \rho < 1$ and $\alpha - \beta + 1 < 0$, then

$$\int_0^1 (1-r)^\alpha (1-\rho r)^\beta dr \leq C_{\alpha, \beta} (1-\rho)^{\alpha-\beta+1}$$

for some positive constant $C_{\alpha, \beta}$.

The next lemma is an easy application of the above two lemmas.

LEMMA 3.6. Let $0 < p < 1$ and fix $\beta > \frac{n+\alpha+1}{p} - (n+1) = \sigma$. Then

$$\sup\{\|J_{\beta, \sigma}(w)\|_{p, \alpha}^p : w \in B_n\} < \infty.$$

PROOF. We only prove the case $\alpha > -1$. Let $w \in B_n$, and $0 < r < 1$.

Then we have, by Lemma 3.4 and 3.5,

$$\begin{aligned}
 & \|J_{\beta, \sigma}(w)\|_{p, \alpha}^p \\
 &= \int_0^1 \int_S |J_{\beta, \sigma}(w)(r\zeta)|^p (1-r)^\alpha 2nr^{2n-1} dr d\sigma(\zeta) \\
 &\leq C_{n, p, \beta} (1-|w|^2)^{(\beta-\sigma)p} \int_0^1 (1-r)^\alpha \int_S \frac{d\sigma(\zeta)}{|1-\langle r\zeta, w \rangle|^{(\beta+n+1)p}} dr \\
 &\leq C_{n, p, \beta} (1-|w|^2)^{(\beta-\sigma)p} \int_0^1 (1-r)^\alpha (1-r|w|^2)^{n-(\beta+n+1)p} dr \\
 &\leq C_{n, p, \alpha, \beta} (1-|w|^2)^{(\beta-\sigma)p} (1-|w|^2)^{\alpha+n+1-(\beta+n+1)p} \\
 &\leq C_{n, p, \alpha, \beta}.
 \end{aligned}$$

Thus we have

$$\sup\{\|J_{\beta, \sigma}(w)\|_{p, \alpha}^p : w \in B_n\} < \infty.$$

The proofs of the following theorems are essentially the same as those of [3] (Prop. 4.4 and Prof. 4.5) and are omitted.

THEOREM 3.7. Let $0 < p < 1$ and $\beta > \frac{n+\alpha+1}{p} - (n+1) \equiv \sigma$. Then there exists $C_{n, p, \alpha, \beta} < \infty$ such that for each $f \in A_{\sigma}^1(B_n)$ there exist a sequence (w_j) of the points in B_n and a sequence (λ_j) of the complex numbers such that

$$\sum_j |\lambda_j| \leq C_{n, p, \alpha, \beta} \|f\|_{1, \sigma} \quad (3.1)$$

and

$$f = \sum_j \lambda_j J_{\beta, \sigma}(w_j), \quad (3.2)$$

where the last series converges in $A_{\sigma}^1(B_n)$.

THEOREM 3.8. The Mackey topology of $A_{\alpha}^p(B_n)$ is the restriction of the topology of $A_{\sigma}^1(B_n)$ where $\sigma = \frac{n+\alpha+1}{p} - (n+1)$.

4. THE DUAL SPACE OF $A_{\alpha}^p(B_n)$.

Finally, we will find the dual space of $A_{\alpha}^p(B_n)$. For the proof of this main result, the following definition is needed:

DEFINITION 4.1. (Radial fractional derivatives of holomorphic functions in B_n) Let $g(z) = \sum_{k=0}^{\infty} G_k(z)$ be the homogeneous expansion of g . For any real number q , the radial fractional derivative of g of order q is defined by

$$R^q g(z) = \sum_{k=0}^{\infty} (k+1)^q G_k(z).$$

Let

$$f(z) = \sum_{k=0}^{\infty} F_k(z) = \sum_{k=0}^{\infty} \sum_{|\gamma|=k} c(\gamma) z^{\gamma},$$

and

$$g(z) = \sum_{\ell=0}^{\infty} G_{\ell}(z) = \sum_{\ell=0}^{\infty} \sum_{|\delta|=\ell} d(\delta) z^{\delta}$$

be the homogeneous expansions of f and g , respectively. We note that for $q > 0$, $0 \leq \rho < 1$, we have

$$\begin{aligned} & \sum_{k=0}^{\infty} \sum_{|\gamma|=k} c(\gamma) \overline{d(\gamma)} \frac{(n-1)!k!}{(n-1+k)!} \frac{2n(k+1)^q}{(k+n)^q} \rho^k \\ &= \frac{2^q}{\Gamma(q)} \int_0^1 \left(\log \frac{1}{r}\right)^{q-1} \int_S R^{-a} f(r\zeta) \overline{R^{q+a} g(r\rho\zeta)} 2nr^{2n-1} dr d\sigma(\zeta). \end{aligned} \quad (4.1)$$

We can now prove the duality relation. We use the idea of Ahern [4] in the proof of the following.

THEOREM 4.2. Let $0 < p < 1$ and $\sigma = \frac{n+\alpha+1}{p} - (n+1)$. Then

$$(A_{\alpha}^p(B_n))^* = \{f \in H(B_n) : \sup(1 - |z|) |R^{\sigma+2} f(z)| = \|f\|_{\Lambda_1} < \infty\}.$$

PROOF. By Theorem 3.8, $(A_{\alpha}^p)^* = (A_{\sigma}^1)^*$. It suffices to compute $(A_0^1)^*$. For simplicity we assume $\sigma = 0$. Take g such that

$$\sup_{z \in B_n} (1 - |z|) |R^2 g(z)| < \infty$$

and let f be a polynomial. Then by (4.1)

$$\begin{aligned} & \sum_{k=0}^{\infty} \sum_{|\gamma|=k} c(\gamma) \overline{d(\gamma)} \frac{(n-1)!k!}{(n-1+k)!} \frac{2n(k+1)^q}{(k+n)^q} \rho^k \\ &= 2 \int_0^1 \int_S R^{-1} f(r\zeta) \overline{R^2 g(r\rho\zeta)} 2nr^{2n-1} dr d\sigma(\zeta) \\ &= 4 \int_0^1 \int_S \left(\log \frac{1}{r}\right) f(r\zeta) \overline{R^2 g(r\rho\zeta)} 2nr^{2n-1} dr d\sigma(\zeta). \end{aligned}$$

Hence

$$\begin{aligned} & \left| \lim_{\rho \rightarrow 1} \left(\sum_{k=0}^{\infty} \sum_{|\gamma|=k} c(\gamma) \overline{d(\gamma)} \frac{(n-1)!k!}{(n-1+k)!} \frac{2n(k+1)^q}{(k+n)^q} \rho^k \right) \right| \\ & \leq 4 \int_0^1 \int_S \frac{\log \frac{1}{r}}{1-r} |f(r\zeta)| \sup_{z \in B_n} (1-r) |R^2 g(r\zeta)| 2nr^{2n-1} dr d\sigma(\zeta). \end{aligned} \quad (4.2)$$

Since $\log \frac{1}{r} \sim 1-r$ as $r \rightarrow 1$, (4.2) is dominated by

$$C_{k,n} \|f\|_{A_0^1} \|g\|_{\Lambda_1}.$$

Since polynomials are dense in A_0^1 , the mapping

$$\psi(f) = \lim_{\rho \rightarrow 1} \left(\sum_{k=0}^{\infty} \sum_{|\gamma|=k} c(\gamma) \overline{d(\gamma)} \frac{(n-1)!k!}{(n-1+k)!} \frac{2n(k+1)^q}{(k+n)^q} \rho^k \right)$$

extends to be a bounded linear functional on A_0^1 . Conversely, let $\psi \in (A_0^1)^*$. Since $A_0^1 \subset L^1(2nr^{2n-1}drd\sigma(\zeta))$, by the Hahn-Banach theorem ψ extends to be a bounded linear functional ψ on the space $L^1(2nr^{2n-1}drd\sigma(\zeta))$. But since $(L^1)^* = L^\infty$, there exists G in $L^\infty(2nr^{2n-1}drd\sigma(\zeta))$ such that

$$\psi(f) = \int_0^1 \int_S f(r\zeta) \overline{G(r\zeta)} 2nr^{2n-1} drd\sigma(\zeta)$$

for each f in A_0^1 . Let

$$H(z) = \int_0^1 \int_S \frac{G(w)}{(1-\langle z, w \rangle)^{n+1}} 2np^{2n-1} dpd\sigma(\eta) \quad (w = \rho\eta)$$

be the holomorphic projection of G . If f is a holomorphic polynomial, then

$$\begin{aligned} \psi(f) &= \int_0^1 \int_S f(r\zeta) \overline{G(r\zeta)} 2nr^{2n-1} drd\sigma(\zeta) \\ &= \int_0^1 \int_S f(r\zeta) \overline{H(r\zeta)} 2nr^{2n-1} drd\sigma(\zeta) \\ &= \int_0^1 \int_S R^{-1} f(r\zeta) \overline{RH(r\zeta)} 2nr^{2n-1} drd\sigma(\zeta) \\ &= \int_0^1 \int_S R^{-1} f(r\zeta) \overline{R^2 g(r\zeta)} 2nr^{2n-1} drd\sigma(\zeta), \end{aligned}$$

where g is defined to be $R^{-1}H$. The proof will be complete if we can show that

$$\sup_{z \in B_n} (1 - |z|) |R^1 H(z)| < \infty.$$

Since

$$\frac{\partial H(r\zeta)}{\partial r} = \int_0^1 \int_S \frac{(n+1)}{r} \frac{\langle r\zeta, \rho\eta \rangle G(\rho\eta)}{(1-\langle r\zeta, \rho\eta \rangle)^{n+2}} 2np^{2n-1} dpd\sigma(\zeta), \quad (z = r\zeta)$$

we have

$$\begin{aligned} r \left| \frac{\partial H(r\zeta)}{\partial r} \right| &\leq C_n \|G\|_\infty \int_0^1 \int_S \frac{d\sigma(\eta)}{|1-\langle r\zeta, \rho\eta \rangle|^{n+2}} dp \\ &\leq C_n \|G\|_\infty \int_0^1 \frac{1}{(1-\rho r)^2} dp \\ &= C_n \|G\|_\infty \frac{1}{1-r}. \end{aligned} \quad (4.3)$$

By (4.3) and $R^1 H(r\zeta) = r \frac{\partial H(r\zeta)}{\partial r} + H(r\zeta)$, we have

$$\sup_{z \in B_n} (1 - |z|) |R^1 H(z)| < \infty.$$

REFERENCES

1. RUDIN, W. Function Theory in the Unit Ball of C^n , Springer-Verlag, New York, 1980.
2. DUREN, P.L. ROMBERG, W. and SHIELDS, A.L. Linear Functionals on H^p space with $0 < p < 1$, J. Reine Angew. Math. 38 (1969), 32-60.
3. SHAPIRO, J.H. Some F -spaces of Harmonic Functions for which Orlicz-Pettis Theorem Fails, Proc. London Math. Soc. 50 (1985), 299-313.
4. AHERN, P.R. and JEVTIĆ, M. Duality and Multipliers for Mixed Normed Spaces, Michigan Math. J. 30 (1983), 53-64.
5. KIM, H.O. Derivatives of Blaschke Products, Pacific J. of Math. 114 (1984), 175-190.
6. SHAPIRO, J.H. Mackey Topologies, reproducing Kernels and Diagonal Maps on the Hardy and Bergman Spaces, Duke Math. J. 43 (1976), 187-202.
7. SHIELDS, A.L. and WILLIAMS, D.L. Bounded Projections, Duality, and Multipliers in Space of Analytic Functions, Trans. Amer. Math. Soc. 162 (1971), 287-302.