SMOOTH STRUCTURES ON SPHERE BUNDLES OVER SPHERES

SAMUEL OMOLOYE AJALA

Institute for Advanced Study
School of Mathematics
Princeton, New Jersey 08543
USA

and

Department of Mathematics
University of Lagos
Akoka - Yaba
Lagos - Nigeria
West Africa

(Received May 30, 1986)

ABSTRACT. In [1] R. De Sapia gave a classification of smooth structures of a p -sphere bundle over a q -sphere with one cross-section and $p < q$. In [2] J. Munkres also gave a classification up to concordance of differential structures in the case where the bundle has at least two cross-sections. In [3] R. Schultz gave a classification in the case $p \geq q$. Here we will give a classification of the p -sphere bundle over a q -sphere without any cross-section and $p < q$.

KEY WORDS AND PHRASES. Smooth structures, differential classification, internal groups.

1980 AMS SUBJECT CLASSIFICATION CODES: 57R55.

1. INTRODUCTION

Let E represent p -sphere bundle over a q -sphere with $\beta \in \pi_{q-1}SO(p+1)$ the characteristic class of the corresponding $p+1$ -disc bundle over the q -sphere. In [4] R. De Sapia gave a complete classification of the special case where $\beta = 0$. In [5] and [6] Kawakubo and Schultz respectively also gave a classification of E for this special case. This author in [7] gave a generalization of this special case to product of three ordinary spheres. In [1] a classification of E was given for $p < q - 1$ and where E has a cross-section and $\beta \neq 0$. In [3] Schultz gave a classification of E for $p \geq q$ and E is without cross-section. We shall here remove the fact that E has a cross-section so that not every element of $\pi_{q-1}SO(p+1)$ can be pulled back to the element $\pi_{q-1}SO(p)$ in the homomorphism $S_* : \pi_{q-1}SO(p) \rightarrow \pi_{q-1}SO(p+1)$ induced by the inclusion $s : SO(p) \rightarrow SO(p+1)$. S^n denotes the unit n -sphere with the usual differential structure in the Euclidean

$(n+1)$ -space $R^{n+1} \cdot \Sigma^n$ denotes an homotopy n -sphere and θ^n denotes the group of homotopy n -spheres. $H(p,k)$ denotes the subset of θ^p which consists of those homotopy p -sphere Σ^p such that $\Sigma^p \times S^k$ is diffeomorphic to $S^p \times S^k$. By [4, Lemma 4], $H(p,k)$ is a subgroup of θ^p and it is not always zero and in fact in [7] we showed that if $k \geq p-3$, $H(p,k) = \theta^p$. We shall adopt the notation $E(\Sigma^q)$ to represent the total space of a p -sphere bundle over a homotopy q -sphere Σ^q . We will then prove the following:

THEOREM. If M is a smooth, n -manifold homeomorphic to a p -sphere bundle over a q -sphere with total space E where $n = p+q \geq 6$ and $p < q$ then there exists homotopy spheres Σ^q and Σ^n such that M is diffeomorphic to $E(\Sigma^q) \# \Sigma^n$. We shall define a pairing

$$G : \pi_p SO(q) \times \pi_{q-1} SO(p+1) \rightarrow \theta^{p+q}$$

and show that if $\beta \in \pi_{q-1} SO(p+1)$ is the characteristic class of a p -sphere bundle over an homotopy q -sphere Σ^q , then $G(\pi_p SO(q), \beta)$ equals the inertial group of $E(\Sigma^q)$. The above theorem together with the latter will give us the following.

THEOREM. Let E be the total space of a p -sphere bundle over a q -sphere then the diffeomorphism classes of $(p+q)$ -manifolds that are homeomorphic to E are in one-to-one correspondence with the group

$$\frac{\theta^q}{H(p,q)} \times \frac{\theta^n}{\text{Image } G_\beta} \quad \text{where } n = p+q \geq 6 \quad \text{and } p < q.$$

2. CLASSIFICATION THEOREM

In this section, we will prove the classification theorem for any manifold M^n homeomorphic to E . We will apply the obstruction theory to smoothing of manifolds developed by Munkres in [8]. Since $p+q \geq 6$ and $2 \leq p < q$ then E is simply-connected and the homology of E has no 2-torsion, hence the "Hauptvermutung" of D. Sullivan [9] applies and this means that piecewise linear homeomorphism can be replaced by homeomorphism, we shall not distinguish the two.

DEFINITION. Let M and N be smooth closed n -manifolds and L a closed subset of M of dimension less than n . Let $f : M \rightarrow N$ be a homeomorphism such that for each simplex γ of L , $\bar{\gamma}$ and $f(\bar{\gamma})$ are contained in coordinate systems under which they are flat. f is said to be a diffeomorphism modulo L if $f|(M-L)$ is a diffeomorphism and each simplex γ of L has a neighborhood V such that f is smooth on $V-L$ near γ . By [8, Theorem 2.8], if M and N are homeomorphic then there is a diffeomorphism modulo $(n-1)$ -skeleton of M . If $f : M \rightarrow N$ is a diffeomorphism modulo m -skeleton $m < n$ then the obstruction to deforming

f to a diffeomorphism modulo $(m-1)$ -skeleton $g : M \rightarrow N$ is an element $\lambda(f) \in H_m(M, \Gamma^{n-m})$ where Γ^{n-m} is a group of diffeomorphism of S^{n-m-1} modulo those that extend to diffeomorphisms of D^{n-m} . g is called the smoothing of f . If $\lambda(f) = 0$ then by [8, §4] smoothing g exist.

THEOREM 2.1. If M is a smooth n -manifold homeomorphic to E where E denotes the total space of a p -sphere bundle over a q -sphere, $2 \leq p < q$ and $n = p + q$ then there exist homotopy spheres Σ^q and Σ^n such that M is diffeomorphic to $E(\Sigma^q) \# \Sigma^n$ where $E(\Sigma^q)$ denotes the total space of a p -sphere bundle over the homotopy q -sphere Σ^q .

PROOF. E is the total space of a p -sphere bundle over a q -sphere with characteristic class $[b] \in \pi_{q-1}SO(p+1)$ then $E = D^q \times S^p \bigcup_{f_b} D^q \times S^p$ where $f_b : S^{q-1} \times S^p \rightarrow S^{q-1} \times S^p$ is a diffeomorphism defined by $f_b(x, y) = (x, b(x) \cdot y)$, $(x, y) \in S^{q-1} \times S^p$

$$H_i(E) = \begin{cases} \mathbb{Z} & \text{for } i = 0, p, q, p+q \\ 0 & \text{elsewhere} \end{cases}$$

Since M^n is homeomorphic to E where $n = p+q \geq 6$ $2 \leq p < q$, then M^n is simply connected and since $H_3(M, \mathbb{Z})$ has no 2-torsion, then "Hauptvermutung" of D. Sullivan [9] implies that there is a piecewise linear homeomorphism $h : M^n \rightarrow E$ which by [8, §5] is a diffeomorphism modulo $(n-1)$ -skeleton. Since $H_i(M, \mathbb{Z}) = 0$ for $n-p+1 \leq i \leq n-1$ then we can assume that h is a diffeomorphism modulo $n-p = q$ skeleton. The obstruction to a diffeomorphism modulo $q-1$ skeleton is $\lambda(h) \in H_q(M, \Gamma^p) = \Gamma^p$. If $[\phi] = \lambda(h) \in \Gamma^p$ where $\phi : S^{p-1} \rightarrow S^{p-1}$ is a diffeomorphism that represents $\lambda(h)$ and let Σ^p denote the homotopy p -sphere where $\Sigma^p = D_1^p \bigcup_{\phi} D_2^p$. We define a map

$$j : S^p \rightarrow \Sigma^p \quad \text{where} \quad S^p = D_1^p \bigcup_{\text{id.}} D_2^p$$

such that

$$j(x) = \begin{cases} x & \text{if } x \in D_1^p \\ x \phi^{-1}(\frac{x}{|x|}) & \text{if } x \in D_2^p. \end{cases}$$

So j is an homeomorphism which is identity on D_1^p and the radial extension of ϕ^{-1} on D_2^p and so the first obstruction $\lambda(j)$ to deforming j to a diffeomorphism is $[\phi^{-1}] = -\lambda(h)$. We then define $\text{id} \times j : D^q \times S^p \rightarrow D^q \times \Sigma^p$ where id is the identity, then $\text{id} \times j$ is a homeomorphism and it follows from [8, Def. 3.4] that the first obstruction $\lambda(\text{id} \times j)$ to

deforming $\text{id} \times j$ to a diffeomorphism is also $-\lambda(h)$. We can form a manifold E' by identifying two copies of $D^q \times \Sigma^p$ along their common boundaries $S^{q-1} \times \Sigma^p$ by the diffeomorphism $f_b : S^{q-1} \times \Sigma^p \rightarrow S^{q-1} \times \Sigma^p$ where $f_b(x, y) = (x, b(x) \cdot y)$ and

$[b] \in \pi_{q-1} SO(p+1)$. So $E' = D^q \times \Sigma^p \bigcup_{f_b} D^q \times \Sigma^p$. We define a map

$g : E = (D^q \times S^p)_1 \bigcup_{f_b} (D^q \times S^p)_2 \rightarrow (D^q \times \Sigma^p)_1 \bigcup_{f_b} (D^q \times \Sigma^p)_2 = E'$ by $g(x, y) = \text{id} \times j(x, y)$

on both $(D^q \times \Sigma^p)_1$, and $(D^q \times \Sigma^p)_2$, the map looks like

$$\begin{array}{ccccccc} E = (D^q \times S^p)_1 & \bigcup_{f_b} & (D^q \times S^p)_2 & = & (D^q \times S^p)_1 & \bigcup_{f_b} & S^{q-1} \times S^p \bigcup_{\text{id}} (D^q \times S^p)_2 \\ \downarrow g & & & = & \downarrow \text{id} \times j & & \downarrow \text{id} \times j & \downarrow \text{id} \times j \\ E' = (D^q \times \Sigma^p)_1 & \bigcup_{f_b} & (D^q \times \Sigma^p)_2 & = & (D^q \times \Sigma^p)_1 & \bigcup_{f_b} & S^{q-1} \times S^p \bigcup_{\text{id}} (D^q \times \Sigma^p)_2 \end{array}$$

g is an homeomorphism and the first obstruction to a diffeomorphism is $\lambda(\text{id} \times j) = -\lambda(h)$.

It follows that the obstructions to smoothing the composition $g \cdot h : M \rightarrow E'$ is

$\lambda(g \cdot h) = \lambda(g) + \lambda(h) = -\lambda(h) + \lambda(h) = 0$. It follows that $g \cdot h : M \rightarrow E'$ is a diffeomorphism modulo $(q-1)$ -skeleton. However in [7, Remark 1] we showed that $D^q \times \Sigma^p$ is

diffeomorphic to $D^q \times S^p$ if $p \leq q + 2$ and so by our hypothesis $p < q$ then it follows that $D^q \times \Sigma^p$ is diffeomorphic to $D^q \times S^p$. This implies that E and E' are diffeomorphic

hence $g' : M \rightarrow E$ is a diffeomorphism modulo $(q-1)$ -skeleton. Since $H_i(M, \mathbb{Z}) = 0$ for $p+1 < i < q-1$, there is no more obstruction to deforming g' to a diffeomorphism until

we get to $(p-1)$ skeleton. We can then assume that g' is a diffeomorphism modulo p -skeleton. The first obstruction to deforming g' to a diffeomorphism modulo $(p-1)$ -

skeleton is $\lambda(g';) \in H_p(M, \mathbb{R}^q) = \mathbb{R}^p$. Let $[\phi] = \lambda(g') \in \mathbb{R}^q$ where $\phi : S^{q-1} \rightarrow S^{q-1}$ is a

diffeomorphism which represents $\lambda(g') \in \mathbb{R}^q$. We define $(\phi \times \text{id}) : S^{q-1} \times S^p \rightarrow S^{q-1} \times S^p$

where $(\phi \times \text{id})(x, y) = (\phi(x), y)$ and if $\beta = [b] \in \pi_{q-1} SO(p+1)$ we also define $f_b : S^{q-1} \times S^p$

$\rightarrow S^{q-1} \times S^p$ where $f_b(x, y) = (x, b(x) \cdot y)$. We then have two orientation preserving diffeomorphisms of $S^{q-1} \times S^p$ unto itself which we can compose to get $(\phi \times \text{id}) \cdot f_b : S^{q-1} \times S^p \rightarrow$

$S^{q-1} \times S^p$ where $(\phi \times \text{id}) \cdot f_b(x, y) = (\phi(x), b(x) \cdot y)$. We then construct a manifold by

attaching two copies of $D^q \times S^p$ along their common boundary $S^{q-1} \times S^p$ using the diffeomorphism

$(\phi \times \text{id}) \cdot f_b$ to have $D_1^q \times S^p \bigcup_{(\phi \times \text{id}) \cdot f_b} D_2^q \times S^p$. Notice that this manifold is

a p -sphere bundle over a homotopy q -sphere $\Sigma^q = D_1^q \bigcup_{\phi} D_2^q$ whose characteristic map is

$\beta = [b] \in \pi_{q-1}SO(p+1)$. We define a map

$$h : D_1^q \times S^p \cup_{f_b} D_2^q \times S^p \rightarrow D_1^q \times S^p \cup_{(\phi \times id) \cdot f_b} D_2^q \times S^p$$

by

$$h(x, y) = \begin{cases} (x, y) & \text{if } (x, y) \in D_1^q \times S^p \\ (x \cdot \phi^{-1}(\frac{x}{|x|}), y) & \text{if } (x, y) \in D_2^q \times S^p \end{cases}$$

Hence h is identity on $D_1^q \times S^p$ and a radial extension of ϕ^{-1} on D_2^q . It then follows that h is an homeomorphism with the first obstruction to a diffeomorphism being $[\phi^{-1}] = -\lambda(g')$. Then by [8, 3.8] the first obstruction to deforming the composition $g' \circ h =$

$g : M \rightarrow D_1^q \times S^p \cup_{(\phi \times id) \cdot f_b} D_2^q \times S^p$ into a diffeomorphism is $\lambda(g) = \lambda(g' \circ h) = \lambda(g') + \lambda(h) = -\lambda(h) + \lambda(h) = 0$ and hence g is a diffeomorphism modulo $(p-1)$ -skeleton. Since $H_i(M, \mathbb{Z}) = 0$ for $0 < i < p$ then we can assume that g is a diffeomorphism modulo one point.

Since $D_1^q \times S^p \cup_{(\phi \times id) \cdot f_b} D_2^q \times S^p$ is a p -sphere bundle over a homotopy q -sphere

$\Sigma_1^q = D_2^q \cup_{\phi} D_1^q$ with characteristic map $[b] \in \pi_{q-1}SO(p+1)$, we shall denote it by $E(\Sigma^q)$.

Since g is a diffeomorphism modulo one point then it is known that there is an homotopy n -sphere Σ^n such that M is diffeomorphic to $E(\Sigma^q) \# \Sigma^n$. Hence the proof.

3. INERTIAL GROUPS

Since by Theorem 2.1, every manifold homeomorphic to E is diffeomorphic to $E(\Sigma^q) \# \Sigma^n$ for some homotopy spheres Σ^q, Σ^n , classification of such manifolds reduces to classification of manifolds of the form $E(\Sigma^q) \# \Sigma^n$. To complete this classification, we then need to investigate what happens when we vary the homotopy spheres and in particular we need to investigate the Inertial group of $E(\Sigma^q)$. We will investigate these in this section.

LEMMA 3.1. Let Σ_1^q and Σ_2^q be homotopy q -spheres such that $\Sigma_i^q = D_1^q \cup_{\phi_i} D_2^q$ $i = 1, 2$

then $E(\Sigma_1^q)$ is diffeomorphic to $E(\Sigma_2^q)$ if and only if $\Sigma_1^q \pm \Sigma_2^q \in H(q, p)$.

PROOF. Suppose $E(\Sigma_1^q)$ is diffeomorphic to $E(\Sigma_2^q)$. This means that

$D_1^q \times S^p \cup_{(\phi_1 \times id) \cdot f_b} D_2^q \times S^p$ is diffeomorphic to $D_1^q \times S^p \cup_{(\phi_2 \times id) \cdot f_b} D_2^q \times S^p$ where

$\phi_i \times id : S^{q-1} \times S^p \rightarrow S^{q-1} \times S^p$ is the diffeomorphism defined by $\phi_i(x, y) = (\phi_i(x), y)$ and

$f_b : S^{q-1} \times S^p \rightarrow S^{q-1} \times S^p$ is defined by $f_b(x, y) = (x, b(x) \cdot y)$ where

$[b] = \beta \in \pi_{q-1}SO(p+1)$ is the characteristic map of the bundle. The manifold $E(\Sigma_2^q)$ can

be regarded as the boundary of the $(p+1)$ -disc bundle over Σ_2 which is denoted by

$D_1^q \times D^{p+1} \bigcup_{(\phi_2 \times \text{id}) \cdot f_b} D_2^q \times D^{p+1} = D(\Sigma_2^q)$. So if $E(\Sigma_1^q)$ is diffeomorphic to $E(\Sigma_2^q)$ then since Σ_1^q can be embedded in $E(\Sigma_1^q)$ it follows that Σ_1^q embeds in $E(\Sigma_2^q)$. But Σ_2^q naturally embeds in $E(\Sigma_2^q)$ and so we have Σ_1^q and Σ_2^q sitting in $E(\Sigma_2^q)$, if we translate Σ_1^q away from Σ_2^q we can run a tube between them to obtain an embedding $\Sigma_1^q \# (-\Sigma_2^q) \rightarrow E(\Sigma_2^q)$ so that the embedding is homotopically trivial and so by the engulfing result of [10, chapter 7] it means that $\Sigma_1^q \# (-\Sigma_2^q)$ can be embedded in the interior of a $(p+q+1)$ -disc in $E(\Sigma_2^q)$ and by [11, 3.5] the embedding is isotopic to a nuclear embedding into the interior of $S^q \times D^{p+1}$. However the embedding $\Sigma_1^q \# (-\Sigma_2^q) \rightarrow S^q \times D^{p+1}$ is a homotopy equivalence, it then follows by Smale's theorem [12, Theorem 4.1] that $\Sigma_1^q \# (-\Sigma_2^q) \times D^{p+1}$ is diffeomorphic to $S^q \times D^{p+1}$ and so it follows that $\Sigma_1^q \# (-\Sigma_2^q) \times S^p$ is diffeomorphic to $S^q \times S^p$ hence $\Sigma_1^q \# (-\Sigma_2^q) \in H(q, p)$. Conversely suppose $\Sigma_1^q \# (-\Sigma_2^q) \in H(q, p)$ then this implies $(\Sigma_1^q \# (-\Sigma_2^q)) \times S^p$ is diffeomorphic to $S^q \times S^p$. Since $S^q \times S^p$ embeds in R^{p+q+1} with trivial normal bundle then it follows that $\Sigma_1^q \# (-\Sigma_2^q)$ embeds in R^{p+q+1} with trivial normal bundle. This shows that each Σ_i^q for $i = 1, 2$ embeds in R^{p+q+1} with trivial normal bundle and by [11, §3.5] the embedding is isotopic to an embedding of Σ_i^q into the interior of $S^q \times D^{p+1}$. However for $i = 1, 2$ the embedding $\Sigma_i^q \rightarrow S^q \times D^{p+1}$ is a homotopy equivalence hence it follows from [12, Theorem 4.1] that $\Sigma_i^q \times D^{p+1}$ is diffeomorphic to $S^q \times D^{p+1}$ which implies that $\Sigma_1^q \times D^{p+1}$ is diffeomorphic to $\Sigma_2^q \times D^{p+1}$. Now since $\Sigma_i^q = D_1^q \bigcup D_2^q$ where $\phi_i : S^{q-1} \rightarrow S^{q-1}$ represents $\Sigma_i^q \in \pi^q$ $i = 1, 2$, then we can write $\Sigma_i^q \times D^{p+1} \stackrel{\phi_i}{=} D_1^q \times D^{p+1} \bigcup_{\phi_i \times \text{id}} D_2^q \times D^{p+1}$ where we identify two copies of $D^q \times D^{p+1}$ along $S^{q-1} \times D^{p+1}$ by the diffeomorphism $\phi_i \times \text{id} : S^{q-1} \times D^{p+1} \rightarrow S^{q-1} \times D^{p+1}$ defined by $(\phi_i \times \text{id})(x, y) = (\phi_i(x), y)$ where $(x, y) \in S^{q-1} \times D^{p+1}$. So $\Sigma_1^q \times D^{p+1}$ is diffeomorphic to $\Sigma_2^q \times D^{p+1}$ implies $D_1^q \times D^{p+1} \bigcup_{\phi_1 \times \text{id}} D_2^q \times D^{p+1}$ is diffeomorphic to $D_1^q \times D^{p+1} \bigcup_{\phi_2 \times \text{id}} D_2^q \times D^{p+1}$. Now consider the manifold $D(S^q) = D_+^q \times D^{p+1} \bigcup_{f_b} D_-^q \times D^{p+1}$ which is a $(p+1)$ -disc bundle over a q -sphere with characteristic map $[b] \in \pi_{q-1} SO(p+1)$. We then form the quotient space

$$D(S^q) \bigcup \Sigma_1^q \times D^{p+1} = (D_+^q \times D^{p+1} \bigcup_{f_b} D_-^q \times D^{p+1}) \bigcup (D_1^q \times D^{p+1} \bigcup_{\phi_1 \times \text{id}} D_2^q \times D^{p+1})$$

by identifying $D_-^q \times D^{p+1} \subset D(S^q)$ and $D_1^q \times D^{p+1} \subset \Sigma_1^q \times D^{p+1}$ by the relation $(x, y) = (x, y)(x \in D_-^q = D_1^q, y \in D^{p+1})$. The manifold $D(S^q) \bigcup \Sigma_2^q \times D^{p+1}$ is similarly constructed. Since $\Sigma_1^q \times D^{p+1}$ is diffeomorphic to $\Sigma_2^q \times D^{p+1}$. Let $d : \Sigma_1^q \times D^{p+1} \rightarrow \Sigma_2^q \times D^{p+1}$ be the

diffeomorphism and since any diffeomorphism fixes a disc, we can assume that d is identity on the disc $D^{p+q+1} = D_1^q \times D^{p+1}$, then we can define a diffeomorphism.

$$g : D(S^q) \cup \Sigma_1^q \times D^{p+1} \rightarrow D(S^q) \cup \Sigma_2^q \times D^{p+1}$$

where

$$g(x) = \begin{cases} d(x) & \text{for } x \in \Sigma_1^q \times D^{p+1} \\ x & \text{for } x \in D(S^q). \end{cases}$$

This means that $g = d$ on $\Sigma_1^q \times D^{p+1}$ and identity on $D(S^q)$. g is well defined because d is identity on the disc connecting $D(S^q)$ and $\Sigma_1^q \times D^{p+1}$ and g is a diffeomorphism. The manifold $D(S^q) \cup \Sigma_1^q \times D^{p+1}$ can be clearly seen as follows. Let $(\phi_i \times id) \cdot f_b : S^{q-1} \times D^{p+1} \rightarrow S^{q-1} \times D^{p+1}$ be the diffeomorphism defined by $((\phi_i \times id) \cdot f_b)(x, y) = (\phi_i(x), b(x) \cdot y)$, $(x, y) \in S^{q-1} \times D^{p+1}$ then attaching two manifolds $D_+^q \times D^{p+1}$ and $D_-^q \times D^{p+1}$ by the diffeomorphism $(\phi_i \times id) \cdot f_b$ we have $D_+^q \times D^{p+1} \cup_{(\phi_i \times id) \cdot f_b} D_-^q \times D^{p+1}$ we get a $(p+1)$ -disc bundle over the homotopy q -sphere $\Sigma_i^q = D_+^q \cup_{\phi_i} D_-^q$ $i = 1, 2$. However, from the way

$D(S^q) \cup \Sigma_1^q \times D^{p+1}$ is constructed it is easily seen that $D(S^q) \cup \Sigma_1^q \times D^{p+1} = D_+^q \times D^{p+1} \cup_{(\phi_1 \times id) \cdot f_b} D_-^q \times D^{p+1} = D(\Sigma_1^q)$ hence g is the diffeomorphism of $D(\Sigma_1^q)$ onto $D(\Sigma_2^q)$ then it follows that $\partial(D(\Sigma_1^q)) = E(\Sigma_1^q)$ is diffeomorphic to $\partial(D(\Sigma_2^q)) = E(\Sigma_2^q)$.

Hence the theorem is proved.

REMARK 1. This theorem implies that $E(\Sigma_1^q)$ is diffeomorphic to $E(\Sigma_2^q)$ if and only if Σ_1^q and Σ_2^q are equivalent in the quotient group $\theta^q/H(q, p)$.

To complete this classification, we need to determine the inertial group of $E(\Sigma^q)$. The inertial group ${}_*(M)$ of an oriented closed smooth n -dimensional manifold M is defined to be the subgroup of θ^n consisting of those homotopy n -spheres Σ^n such that $M \# \Sigma^n$ diffeomorphic to M .

Let E_β represent the total space of a p -sphere bundle over a real q -sphere with characteristic class $\beta \in \pi_{q-1}SO(p+1)$. In [13] we defined a map $G_\beta : \pi_p SO(q) \rightarrow \theta^{p+n}$ and showed that the image of this map equals the inertial group of E_β where $p < q$ and E_β has no cross-section. We shall similarly define a map $G_{\phi, \beta} : \pi_p SO(q) \rightarrow \theta^{p+q}$ and show that the image of this map equals the inertial group of $E(\Sigma^q)$ where $E(\Sigma^q)$ is the total space of p -sphere bundle over a homotopy sphere $\Sigma^q = D_1^q \cup_\phi D_2^q$. Let $\alpha \in \pi_p SO(q)$ we define $G_{\phi, \beta}(\alpha) = S^{q-1} \times D^{p+1} \cup_{f_{-1}(\phi \times id) \cdot f_b} D^q \times S^p$ where $[a] = \alpha$ and $[b] = \beta \in \pi_{q-1}SO(p+1)$ and

$f_{a-1}(\phi \times id) \cdot f_b : S^{q-1} \times S^p \rightarrow S^{q-1} \times S^p$ is a diffeomorphism defined by $f_{a-1}(\phi \times id) \cdot f_b(x, y) = (a^{-1}(b(x) \cdot y) \cdot (x), b(x) \cdot y)$. One can easily show that $G_{\phi \cdot b}$ is well-defined and that its image is an homotopy $(p+q)$ -sphere as similarly shown in [13].

LEMMA 3.2. Let $E(\Sigma^q)$ denote the total space of a p -sphere bundle over an homotopy q -sphere $\Sigma^q = D_1^q = D_1^q \cup D_2^q$ with characteristic class $\beta \in \pi_{q-1}SO(p+1)$ then $G_{\phi \cdot b} \pi_p(SO(q)) = I(E(\Sigma^q))$.

PROOF. If $\Sigma^{p+q} \in I(E(\Sigma^q))$ then this means there is a diffeomorphism $d : E(\Sigma^q) \# \Sigma^{p+q} \rightarrow E(\Sigma^q)$, that is,

$$d : (D_1^q \times S^p \cup_{(\phi \times id) \cdot f_b} D_2^q \times S^p) \# \Sigma^{p+q} \rightarrow D_1^q \times S^p \cup_{(\phi \times id) \cdot f_b} D_2^q \times S^p$$

since $p < q$ then $\pi_p(E(\Sigma^q))$ is infinitely cyclic and $d(o \times S^q)$ represents a generator and so is homotopic to the inclusion $0 \times S^p \rightarrow E(\Sigma^q)$. By Haefliger's theorem [14], $d|_{0 \times S^p}$ and the inclusion $0 \times S^p \rightarrow E(\Sigma^q)$ are isotopic and by isotopy extension theorem and tubular neighborhood theorem, d is isotopic to a map which we shall again denote by d such that $d|_{D^q \times S^p} = D^q \times S^p$ where $d(x, y) = (a(y) \cdot x, y)$ for $[a] \in \pi_p SO(q)$ and $(x, y) \in D^q \times S^p$. We now remove $D^q \times S^p$ from $E(\Sigma^q) \# \Sigma^{p+q} = (D^q \times S^p \cup_{(\phi \times id) \cdot f_b} D^q \times S^p) \# \Sigma^{p+q}$ by

surgery away from the connected sum and replace it with $S^{q-1} \times D^{p+1}$. After this operation on the summand $E(\Sigma^q)$ of the connected sum, we have the manifold $S^{q-1} \times D^{p+1} \cup_{(\phi \times id) \cdot f_b} D^q \times S^p$. Since the diffeomorphism $(\phi \times id) \cdot f_b : S^{q-1} \times S^p \rightarrow S^{q-1} \times S^p$ extend to the diffeomorphism of $S^{q-1} \times D^{p+1}$ onto itself then $S^{q-1} \times D^{p+1} \cup_{(\phi \times id) \cdot f_b} D^q \times S^p$ is diffeomorphic to $S^{q-1} \times D^{p+1} \cup_{id} D^q \times S^p$, the diffeomorphism g is defined thus

$$\begin{array}{ccc} S^{q-1} \times D^{p+1} & \cup & D^q \times S^p \\ (\phi \times id) \cdot f_b \downarrow & id & \downarrow \\ S^{q-1} \times D^{p+1} & \cup & D^q \times S^p \\ & (\phi \times id) \cdot f_b & \end{array}$$

where

$$g(x, y) = \begin{cases} (x, y) & \text{if } (x, y) \in D^q \times S^p \\ ((\phi \times id) \cdot f_b)(x, y) & \text{if } (x, y) \in S^{q-1} \times D^{p+1}. \end{cases}$$

However, by [7, Lemma 2.1.2], $S^{q-1} \times D^{p+1} \cup_{id} D^q \times S^p$ is diffeomorphic to the standard $(p+q)$ -sphere S^{p+q} , hence after this surgery $E(\Sigma^q)$ is reduced to S^{p+q} and so $E(\Sigma^q) \# \Sigma^{p+q}$ is reduced to $S^{p+q} \neq \Sigma^{p+q} = \Sigma^{p+q} = \Sigma^{p+q}$.

We perform the corresponding modification (under d) on $E(\Sigma^q)$ to remove the p -sphere $0 \times S^p$ with product structure $d(D_1^q \times S^p)$ in $E(\Sigma^q)$. From this modification we obtain a manifold $S^{q-1} \times D^{p+1} \cup_{\psi} D^q \times S^p$ where $\psi = (d^{-1}|_{S^{q-1} \times S^p}) \cdot (\phi \times \text{id}) \cdot f_b$ and this is

diffeomorphic to Σ^{p+q} because of the way we performed the surgery using d . However, this manifold $S^{q-1} \times D^{p+1} \cup_{\psi} D^q \times S^p = G_{\phi \cdot \beta}(\alpha)$ by the definition of $G_{\phi \cdot \beta}$, thus there

exists an element $\alpha \in \pi_p SO(q)$ (namely $d|(D_1^q \times S^p)$ which gives $\alpha \in \pi_p SO(q)$ such that $\Sigma^{p+q} = G_{\phi \cdot \beta}(\alpha)$ and so $\Sigma^{p+q} \in G_{\phi \cdot \beta}(\pi_p SO(q))$, hence $I(E(\Sigma^q)) \subset G_{\phi \cdot \beta}(\pi_p SO(q))$. Conversely suppose $\Sigma^{p+q} \in G_{\phi \cdot \beta}(\pi_p SO(q))$ then for some $\alpha \in \pi_p SO(q)$, $\Sigma^{p+q} = S^{q-1} \times D^{p+1} \cup_{f_a^{-1} \cdot (\phi \times \text{id}) \cdot f_b}$

$D^q \times S^p$ where ϕ is a diffeomorphism of S^{q-1} onto itself representing $\Sigma^q = D_1^q \cup_{\phi} D_2^q$ and f_a^{-1} and f_b are as defined earlier. Notice that $G_{\phi \cdot \beta}(\alpha)$ is thus the obstruction to the construction of a diffeomorphism $S^{p+q} \rightarrow \Sigma^{p+q}$. To construct a diffeomorphism from $S^{p+q} \rightarrow \Sigma^{p+q}$, we map $S^{q-1} \times D^{p+1} \subset S^{p+q}$ to itself using $(\phi \times \text{id}) \cdot f_b$ to have

$$\begin{array}{ccc} S^{p+q} = S^{q-1} \times D^{p+1} & \cup & D^q \times S^p \\ & \downarrow (\phi \times \text{id}) \cdot f_b & \\ \Sigma^{p+q} = S^{q-1} \times D^{p+1} & \cup & D^q \times S^p \\ & \downarrow f_a^{-1} \cdot (\phi \times \text{id}) \cdot f_b & \end{array}$$

and try to extend it to $D^q \times S^p$. On the boundary $S^{q-1} \times S^p$ of $D^q \times S^p$, the map is $f_b^{-1} \cdot (\phi^{-1} \times \text{id}) \cdot f_a \cdot (\phi \times \text{id}) \cdot f_b$. So this means that $\Sigma^{p+q} = G_{\phi \cdot \beta}(\alpha)$ is the obstruction to extending the diffeomorphism $f_b^{-1} \cdot (\phi^{-1} \times \text{id}) \cdot f_a \cdot (\phi \times \text{id}) \cdot f_b : S^{q-1} \times S^p \rightarrow S^{q-1} \times S^p$ to a

diffeomorphism of $D^q \times S^p$ onto itself. We can then define a map $E(\Sigma^p) \rightarrow E(\Sigma^q)$ using the diffeomorphism $f_a : D_1^q \times S^p \rightarrow D_1^q \times S^p$ where $f_a(x, y) = (a(y) \cdot x, y)$ ($x, y \in D_1^q \times S^p$) we then have

$$\begin{array}{ccc} E(\Sigma^q) = D_1^q \times S^p & \cup & D_2^q \times S^p \\ & \downarrow f_a & \\ E(\Sigma^q) = D_1^q \times S^p & \cup & D_2^q \times S^p \\ & \downarrow (\phi \times \text{id}) \cdot f_b & \end{array}$$

On the boundary $S^{q-1} \times S^p$ of $D_1^q \times S^p$, this map is $f_b^{-1} \cdot (\phi^{-1} \times \text{id}) \cdot f_a \cdot (\phi \times \text{id}) \cdot f_b$ and the obstruction to extending this to a diffeomorphism of $E(\Sigma^q)$ onto itself is the

obstruction to extending the map $f_{b-1} \cdot (\phi^{-1} \times \text{id}) \cdot f_a \cdot (\phi \times \text{id}) \cdot f_b$ to the diffeomorphism of $D_2^q \times S^p$ onto itself which is Σ^{p+q} . It then follows that $E(\Sigma^q) \rightarrow E(\Sigma^q) \# \Sigma^{p+q}$ is a diffeomorphism and so $\Sigma^{p+q} \in I(E(\Sigma^q))$ hence

$$E(E(\Sigma^q)) = G_{\phi \cdot \beta} \pi_p(SO(q))$$

REMARK 2. We note that if $p = 2, 4, 5, 6 \pmod{8}$ and $p < q-1$ then $\pi_p SO(q) = 0$ and so the image of G is trivial and hence in this particular case, the inertial group of $E(\Sigma^q)$ is trivial and this coincides with the result of [4, Proposition 1].

REMARK 3. By [15], inertial group $I(M)$ of a smooth manifold M is a diffeotopy invariant of M . So if $2p \geq q+1$ then we can deduce that the inertial group $I(E(\Sigma^q))$ of a p -sphere bundle over an homotopy q -sphere Σ^q is equal to the inertial group $I(E_\beta)$ of a p -sphere bundle over the standard q -sphere, where $\beta \in \pi_{q-1} SO(p+1)$ classifies the associated disc bundle. Let $D(\Sigma^q)$ be the associated $(p+1)$ -disc bundle over the homotopy q -sphere where $E(\Sigma^q)$ is the boundary of $D(\Sigma^q)$. Σ^q has the homotopy type of $D(\Sigma^q)$ and Σ^q has the homotopy type of S^q , it follows that S^q has the homotopy type of $D(\Sigma^q)$. Since $2p \geq q+1$ then it follows that $2(p+q+1) \geq 3q+3$ and since $p+q > 5$ and $p \geq 3$ then $D(\Sigma^q)$ and $E(\Sigma^q)$ are simply connected and from [12: Theorem 4.4], it follows that $D(\Sigma^q)$ is diffeomorphic to a $(p+1)$ -disc bundle $D(S^q)$ over the q -sphere S^q hence the boundary $\partial D(\Sigma^q) = E(\Sigma^q)$ of $D(\Sigma^q)$ is diffeomorphic to the boundary $\partial D(S^q) = E_\beta$ of $D(S^q)$. It then follows by [15] that $I(E(\Sigma^q)) = I(E_\beta)$. This means that the inertial group of S_β in [13] coincides with Lemma 3.2.

Combination of Lemmas 3.1 and 3.2 give the following.

THEOREM 3.3. Let E be the total space of a p -sphere bundle over a q -sphere with characteristic map $\beta \in \pi_{q-1} SO(p+1)$ then the diffeomorphism classes of $p+q$ -manifolds that are homeomorphic to E are in one-to-one correspondence with the group

$$\frac{\theta^q}{H(q,p)} \times \frac{\theta^n}{\text{Image } G_\beta}$$

where $p+q = n \geq 6$ and $p < q$.

REFERENCES

1. De Sapio, R., Manifolds Homeomorphic to Sphere Bundles Over Spheres. Bull. American Math. Soc. 75 (1969), 59-63.
2. Munkres, J., Concordance of Differentiable Structures - Two Approaches. Michigan Math. J. 14 (1967), 183-191.

3. Schultz, R., Smoothing of Sphere Bundles of Spheres in Stable Range. Inventiones Math 9 (1969), 81-88.
4. De Sapio, R., Differential Structures on a Product of Spheres. Comm. Math. Helv. Vol. 44, 1 (1969), 61-69.
5. Kawakubo, K., Smooth Structures on $S^p \times S^q$. Proc. Japan Acad. 45 (1969), 215-218.
6. Schultz, R., Smooth Structures on $S^p \times S^q$. Annals of Math. 90 (1969), 187-198.
7. Ajala, S. O., Differentiable Structures on Product of Spheres. Houston Journal of Math., Vol. 10, 1 (1984), 1-14.
8. Munkres, J., Obstruction to Smoothing of Piecewise-Differentiable Homeomorphisms. Annals of Math. Vol. 72, 3 (1960), 521-554.
9. Sullivan, D., On Hauptvanmutung for Manifolds. Bull. Amer. Math. Soc. 73 (1967), 598-600.
10. Zeeman, E., Seminar on Combinatorial Topology (mimeographed notes), Inst. Hautes Etudes, Sci. Publ. Math. (1965).
11. Levine, J., Classification of Differentiable Knots. Annals of Math. 82 (1965), 15-50.
12. Smale, S., On the Structure of Manifolds. Amer. Math. J. 84 (1962), 387-399.
13. Ajala, S. O., Inertial Group of p-sphere Bundle over a q-sphere without Cross-section. Nigerian Journal of Science Vol. 18, (1984), to appear.
14. Haefliger, A., Plongements Differentiables de Varietes dans Varietes, Comm. Math. Helv. 36 (1961), 47-81.
15. Kawakubo, K., Inertial Group of Homology Tori. J. Math. Soc. Japan Vol. 21, 1 (1969).
16. Hsiang, W. C., Leving, J., and Szczarba, R. H., On the Normal Bundle of a Homotopy Sphere Embedded in Euclidean Space. Topology 3 (1965), 173-181.