ON UNIFORM CONNECTEDNESS

D. BABOOLAL

Department of Mathematics
University of Durban-Westville
Private Bag X54001, Durban 4000
South Africa

(Received July 6, 1987 and in revised form August 29, 1988)

ABSTRACT. The concept of uniform connectedness, which generalizes the concept of well-chainedness for metric spaces, is used to prove the following: (a) If two points a and b of a compact Hausdorff uniform space $(X, \mathcal{U})$ can be joined by a $\mathcal{U}$ -chain for every $U \in \mathcal{U}$, then they lie together in the same component of X ; (b) Let $(X, \mathcal{U})$ be a compact Hausdorff uniform space, A and B non-empty disjoint closed subsets of X such that no component of X intersects both A and B . Then there exists a separation $X = X_A \cup X_B$, where X_A and X_B are disjoint compact sets containing A and B respectively. These generalize the corresponding results for metric spaces.

KEY WORDS AND PHRASES. Uniform spaces, uniform connectedness and well-chainedness.
1980 AMS SUBJECT CLASSIFICATION CODES. Primary 54E15, Secondary 54D05.

1. INTRODUCTION.

All topological spaces and all uniform spaces considered are Hausdorff, $(X, \mathcal{U})$ being used to denote a uniform space with $\mathcal{U}$ the family of entourages of X . In [1] Mrowka and Pervin introduced into the theory of uniform spaces the concept of uniform connectedness: $(X, \mathcal{U})$ is uniformly connected iff every uniformly continuous function on X to a discrete uniform space is constant. As is evident from the definition, connectedness is a stronger property than uniform connectedness. However, for the class of compact uniform spaces these two properties are equivalent [1].

In [2] a criterion for a completely regular Hausdorff space to have a locally connected compactification was obtained. This criterion, which involved the concept of uniform connectedness, is that the space must possess a compatible uniformity with respect to which it has weak property S . A uniform space $(X, \mathcal{U})$ is said to have weak property S if given $U \in \mathcal{U}$, X can be covered by a finite collection of uniformly connected sets each of which is U -small [2]. In view of the fact that the concept of uniform connectedness bears on the problem of existence of locally connected compactifications a further study of this concept was felt to be justified. Our specific aim in this paper is to show how we can use this concept to generalize to compact Hausdorff uniform spaces the following two propositions appearing in [3], pp. 15.

PROPOSITION 1.1. If two points a and b of a compact metric space X can be joined by an ε -chain for every $\varepsilon > 0$, they lie together in the same component of X .

PROPOSITION 1.2. If A and B are disjoint closed sets of a compact metric space X such that no component of X intersects both A and B , there exists a separation

$X = X_A \cup X_B$ where X_A and X_B are disjoint compact sets containing A and B respectively. Along the way a problem in [4], p. 170 is strengthened.

2. PRELIMINARIES.

We need some basic facts on Efremovič proximities, (EF-proximities, for short). Proofs of these can be found in [5]. If (X, δ) is an EF-proximity space, its associated topology defined by

$$T(\delta) = \{G \subset X \mid \delta(X-G) \neq 0 \text{ for each } x \in G\}$$

is completely regular and is Hausdorff iff (X, δ) is separated. Conversely every completely regular space (X, T) admits an EF-proximity δ such that $T(\delta) = T$. Each uniform space (X, U) has an associated EF-proximity $\delta = \delta(U)$ given by

$$A \delta B \iff (A \times B) \cap U \neq \emptyset \text{ for every } U \in U. \text{ Furthermore the uniform topology } T(U) = T(\delta).$$

As pointed out in [1], (topological) connectedness implies uniform connectedness but the converse is not true in general, the rational numbers with the usual uniformity providing a counterexample.

We now define the concept of a well-chained uniform space which Mrowka and Pervin have called 'chain-connected' in [1]. This is a natural generalization of the concept of well-chained metric spaces (see eg [3]).

DEFINITION 2.1. A U -chain from a point $x \in X$ to a point $y \in X$ in a uniform space (X, U) , where $U \in U$, is a finite sequence of points $x = x_1, x_2, \dots, x_n = y$ such that $(x_i, x_{i+1}) \in U$ for each $i = 1, 2, \dots, n-1$. (X, U) is said to be well-chained iff any two points x and y in X can be joined by a U -chain for every $U \in U$.

We require the following two results due to Mrowka and Pervin [1].

THEOREM 2.2. Let (X, U) be a uniform space and δ the induced proximity relation. The following conditions are equivalent:

- (a) (X, U) is uniformly connected
- (b) For every $A \subset X$, $\phi \neq A \neq X$, we have $A \delta (X-A)$
- (c) (X, U) is well-chained.

THEOREM 2.3. A uniform space (X, U) is connected iff it is uniformly connected with respect to every uniformity compatible with its topology. Thus a compact uniform space which is uniformly connected must be connected, as it has a unique compatible uniformity.

In [6], Sieber and Pervin discuss connectedness in syntopogeneous spaces (which generalize the notions of topological, proximity and uniform spaces; see [5] eg.). The following results which we have stated in the context of uniform spaces, have been obtained in the general setting of syntopogenous spaces in [6], and are stated here for subsequent reference. First a definition.

DEFINITION 2.4. Two sets A and B in a uniform spaces $(X, \mathcal{U})$ are said to be uniformly separated iff $A \delta B$, where $\delta = \delta(\mathcal{U})$ is the induced EF-proximity.

THEOREM 2.5 [6]. (a) The uniformly continuous image of a uniformly connected space is again uniformly connected.

(b) If A is a uniformly connected subset of $(X, \mathcal{U})$ and $A \subset (M \cup N)$ where M and N are uniformly separated subsets of X , then either $A \subset M$ or $A \subset N$.

(c) If A is a uniformly connected subset of $(X, \mathcal{U})$ and $B \subset X$ is such that $A \subset B \subset \bar{A}$, then B is uniformly connected.

(d) If $\{A_\lambda \mid \lambda \in \Lambda\}$ is a family of uniformly connected subsets of $(X, \mathcal{U})$ and A_{λ_0} ($\lambda_0 \in \Lambda$) is not uniformly separated from any A_λ , ($\lambda \neq \lambda_0$), then $A = \bigcup_{\lambda} A_\lambda$ is uniformly connected.

From (d) above it follows that the union of all uniform connected subsets of X which contain a fixed point $p \in X$ is a uniformly connected set, which is called the uniform component of P denoted by $C^{\mathcal{U}}(p)$. It is clearly the maximal uniformly connected set containing p .

If $C^{\mathcal{U}}(p)$ and $C^{\mathcal{U}}(q)$ are not uniformly separated, then by (d), $C^{\mathcal{U}}(p) \cup C^{\mathcal{U}}(q)$ is uniformly connected, and by maximality we have $C^{\mathcal{U}}(p) = C^{\mathcal{U}}(q)$. Thus a uniform space is partitioned into uniformly separated uniform components. Clearly by (c) above these uniform components are closed sets. Since each connected set is uniformly connected, the component C_p containing $p \in X$ is clearly contained in $C^{\mathcal{U}}(p)$. An example where $C^{\mathcal{U}}(p)$ is strictly larger than C_p is easily found: Let Q be the rational numbers with the usual uniformity. Then $C_p = \{p\}$ for each $p \in Q$, but $C^{\mathcal{U}}(p) = Q$ since Q is uniformly connected as was remarked in [1].

3. THE RESULTS.

Let $(X, \mathcal{U})$ be a uniform space. For each subset A of X and $U \in \mathcal{U}$ let

$$C_U(A) = \{x \in X \mid x \text{ can be joined to some point of } A \text{ by a } U\text{-chain}\}$$

and let $C(A) = \bigcap_{U \in \mathcal{U}} C_U(A)$

We thus have the following result which is a generalization of problem T(a) in [4] p. 170.

PROPOSITION 3.1. For each $A \subset X$, $U \in \mathcal{U}$, $C_U(A)$ is clopen.

PROOF. To show $C_U(A)$ open, let $x \in C_U(A)$. Find $a \in A$ and a U -chain

$x = x_1, x_2, \dots, x_n = a$ joining x to a . Find open symmetric $V \in \mathcal{U}$ such that $V \subset U$. Then $x \in V[x] \subset C_U(A)$: for if $y \in V[x]$ then $(x, y) \in V$ and hence $(y, x) \in V \subset U$. Thus $y, x_1, x_2, \dots, x_n = a$ is a U -chain joining y to a and hence $y \in C_U(A)$, thus showing $C_U(A)$ open.

To show $C_U(A)$ closed, let $y \in \overline{C_U(A)}$. Find $x \in C_U(A)$ such that $x \in U[y]$. There exists a U -chain $x = x_1, x_2, \dots, x_n = a$ joining x to some point $a \in A$, and hence $y, x_1, x_2, \dots, x_n = a$ is a U -chain joining y to $a \in A$, ie $y \in C_U(A)$, thus showing $C_U(A)$ is closed.

The next proposition proved in the setting of uniform spaces, when interpreted in the setting of metric spaces, strengthens Problem T(b) of [4], p. 170 viz, that for a compact metric space X , if $A \subset X$ is connected, then $C(A)$ is connected, $C(A)$ being defined as $\bigcap_{\epsilon > 0} C_\epsilon(A)$ for metric spaces. It is of course a strengthening since A may be taken to be uniformly connected instead of (the stronger) connected in the hypothesis. Use is made of this result in the generalizations of Proposition 1.1 and 1.2. First we need the following lemma which can be easily proved or can be found in [4], p. 163, Problem F(a).

LEMMA 3.2. Let F be a family of closed compact sets in a topological space such that $\bigcap \{A \mid A \in F\} \subset U$ where U is open. Then there exists a finite subfamily H of F such that $\bigcap \{A \mid A \in H\} \subset U$.

PROPOSITION 3.3. Let (X, U) be a compact uniform space, $A \subset X$ uniformly connected. Then $C(A)$ is connected.

PROOF. Suppose $C(A)$ is disconnected. Now since $C(A) = \bigcap_{U \in \mathcal{U}} C_U(A)$ and $C_U(A)$ is closed for each $U \in \mathcal{U}$ by Proposition 3.1, $C(A)$ is closed in X and hence compact. This implies by Theorem 2.3 that $C(A)$ is not uniformly connected. By Theorem 2.2 $C(A) = B \cup D$ where $B \neq \emptyset$, $D \neq \emptyset$ and $B \not\delta D$, where $\delta = \delta(U)$ is the EF-proximity induced by U . Hence there exists $V_1 \in \mathcal{U}$ such that $(B \times D) \cap V_1 = \emptyset$. Now find open symmetric $V \in \mathcal{U}$ such that $V \circ V \circ V \subset V_1$. Noting that each $C_U(A)$ is closed and compact and that $C(A) = \bigcap_{U \in \mathcal{U}} C_U(A) \subset V[B] \cup V[D]$ we have, by the above lemma, a finite collection $\{C_{U_i}(A)\}_{i=1}^n$, $U_i \in \mathcal{U}$, such that

$$\bigcap_{i=1}^n C_{U_i}(A) \subset V[B] \cup V[D]$$

Let $U = \bigcap_{i=1}^n U_i$. Then $U \in \mathcal{U}$ and clearly we have

$$C_U(A) = \bigcap_{i=1}^n C_{U_i}(A) \subset V[B] \cup V[D].$$

We may assume without loss of generality that $U \subset V$. Now since A is uniformly connected and $A \subset C(A) = B \cup D$ with $B \not\delta D$ we have from Theorem 2.5(b) that either $A \subset B$ or $A \subset D$. Let us assume $A \subset D$.

Since $B \neq \emptyset$, find $x \in B$. Then $x \in C(A) \subset C_U(A)$ and hence there exists a U -chain

$x = x_1, x_2, \dots, x_n = a$ joining x to some point $a \in A$. Clearly each x_i , for $1 \leq i \leq n$, belongs to $C_U(A)$, and hence $\{x_1, x_2, \dots, x_n\} \subset V[B] \cup V[D]$. As is easily verified $V[B] \cap V[D] = \emptyset$.

Now let x_k be the last point in this chain such that $x_k \in V[B]$. Then $x_k \neq x_n = a$ otherwise $V[B] \cap V[D] \neq \emptyset$. Thus x_{k+1} belongs to this chain and $x_{k+1} \in V[D]$. Thus we can find $b \in B$, $d \in D$ such that $x_k \in V[b]$ and $x_{k+1} \in V[d]$. Since $(x_k, x_{k+1}) \in U \subset V$, we have $(b, d) \in V \circ V \circ V \subset V_1$ contradicting the fact that $(B \times D) \cap V_1 = \emptyset$. Hence $C(A)$ is connected.

Noting that every connected set is well chained we have:

COROLLARY 3.4. If $(X, \mathcal{U})$ is a compact uniform space and $x \in X$, then $C(\{x\}) = C_x$, the component of x in X .

We now generalize Proposition 1.1.

COROLLARY 3.5. If two points a and b of a compact uniform space $(X, \mathcal{U})$ can be joined by a U -chain for every $U \in \mathcal{U}$, then they lie together in the same component of X .

PROOF. From Corollary 3.4, $C(\{a\}) = C_a$, $C(\{b\}) = C_b$. By hypothesis $a \in C(\{b\})$. Thus $C_a \cap C_b \neq \emptyset$ and hence $C_a = C_b$.

We remark that the condition that X is compact cannot be dropped, the rationals with the usual uniformity providing an easy counterexample.

The next result generalizes Proposition 1.2.

THEOREM 3.6. Let $(X, \mathcal{U})$ be a compact uniform space, A and B non-empty disjoint closed subsets of X such that no component of X intersects both A and B . Then there exists a separation $X = X_A \cup X_B$, where X_A and X_B are disjoint compact sets containing A and B respectively.

PROOF. We claim there exists $U \in \mathcal{U}$ such that no U -chain in X joins a point of A to a point of B . For suppose not. Then for each $U \in \mathcal{U}$, there exists $a_U \in A$ and $b_U \in B$ such that there is a U -chain in X joining a_U to b_U . Now $(U, \leq)$ is a directed set according to the definition $U_1 \leq U_2$ iff $U_1 \supset U_2$. Consequently (a_U) and (b_U) are nets in A and B respectively. Now (a_U) has a cluster point $a \in A$ since A is compact. Let (a_{U_λ}) be a subnet of (a_U) such that $a_{U_\lambda} \rightarrow a$. Then (b_{U_λ}) is a subnet of (b_U) in the compact set B , and hence has a cluster point $b \in B$. Now let $U \in \mathcal{U}$ be arbitrary. Find symmetric $U' \in \mathcal{U}$ such that $U' \subset U$. Find λ_0 such that $U' \leq U_{\lambda_0}$.

Since $a_{U_\lambda} \rightarrow a$, we can find $\lambda_1 \geq \lambda_0$ such that $\lambda \geq \lambda_1 \Rightarrow a_{U_\lambda} \in U'[a]$. Since b is a cluster point of (b_{U_λ}) we can find $\lambda_2 \geq \lambda_1$ such that $b_{U_{\lambda_2}} \in U'[b]$. Now we have $a_{U_{\lambda_2}} \in U'[a]$, $b_{U_{\lambda_2}} \in U'[b]$, $U_{\lambda_2} \geq U_{\lambda_1} \geq U_{\lambda_0} \geq U'$. Now there exists a U_{λ_2} -chain in X joining $a_{U_{\lambda_2}}$ to $b_{U_{\lambda_2}}$. Since $U \subset U'$, this is also a U' -chain, and since $(a, a_{U_{\lambda_2}}) \in U'$ and $(b, b_{U_{\lambda_2}}) \in U'$, there is thus a U' -chain joining a to b , and hence a U -chain

joining a to b . By Corollary 3.5, a and b lie together in the same component of X , contradicting the fact that no component of X intersects both A and B . Thus a $U \in \mathcal{U}$ as claimed above exists.

Let $X_A = C_U(A)$ and $X_B = X - X_A$. By Proposition 3.1 X_A is clopen and hence X_A and X_B are compact. Furthermore $A \subset X_A$ is clear, $B \subset X_B$ follows by the above claim, and $X_A \cup X_B$ is a separation of X .

REFERENCES

1. MROWKA, S.G. and PERVIN, W.J., On uniform connectedness, Proc. Amer. Math. Soc. 15., (1964), 446-449.
2. BABOOLAL, D., On Some Uniform Connection Properties related to Local Connectedness, Quaestiones Mathematicae 7 (1984), 155-160.
3. WHYBURN, G.T., Analytic Topology, Amer. Math. Soc. Colloq. Publications, 28 (1942).
4. KELLEY, J.L., General Topology, Springer-Verlag, New York, Heidelberg, Berlin, (1955).
5. NAIMPALLY, S.A. and WARRACK, B.D., Proximity Spaces, Cambridge Tracts in math. and Math. Phys., Cambridge (1970).
6. SIEBER, J.L. and PERVIN, W.J., Connectedness in syntopogenous spaces, Proc. Amer. Math. Soc. 15 (1964), 590-595.