BOUNDED SPIRAL-LIKE FUNCTIONS WITH FIXED SECOND COEFFICIENT

M.K. AOUF

Department of Mathematics
Faculty of Science
University of Mansoura
Mansoura, Egypt

(Received July 10, 1986)

ABSTRACT. Let $F_p(\alpha, \beta, M)$ ($0 < p < 1$, $|\alpha| < \frac{\pi}{2}$, $0 < \beta < 1$ and $M > \frac{1}{2}$), denote the class of functions $f(z)$ which are regular in $U = \{z: |z| < 1\}$ and of the form $f(z) = z + |a_2| e^{-i\alpha} z^2 + \dots$, where $|a_2| = p(1 + \sigma)(1 - \beta) \cos \alpha$, which satisfy for

fixed M , $z = re^{i\theta} \in U$ and

$$\left| \frac{e^{i\alpha} \frac{zf'(z)}{f(z)} - \beta \cos \alpha - i \sin \alpha}{(1 - \beta) \cos \alpha} - M \right| < M.$$

In this paper we have found the sharp radius of γ -spiralness of the functions belonging to the class $F_p(\alpha, \beta, M)$.

KEY WORDS AND PHRASES. Spirallike, bounded functions, radius of γ -spiralness. .

1980 AMS CLASSIFICATION CODES. 30A32, 30A36.

1. INTRODUCTION. Let A denote the class of functions which are regular and univalent in the unit disc $U = \{z: |z| < 1\}$ and satisfy the conditions $f(0) = 0 = f'(0) - 1$.

Let $F(\alpha, \beta, M)$ ($|\alpha| < \frac{\pi}{2}$, $0 < \beta < 1$ and $M > \frac{1}{2}$) denote the class of bounded α -spirallike functions of order β , that is $f \in F(\alpha, \beta, M)$ if and only if for fixed M ,

$$\left| \frac{e^{i\alpha} \frac{zf'(z)}{f(z)} - \beta \cos \alpha - i \sin \alpha}{(1 - \beta) \cos \alpha} - M \right| < M, \quad z \in U. \quad (1.1)$$

The class $F(\alpha, \beta, M)$ introduced by Aouf [1], he proved that if

$f(z) = z + a_2 z^2 + \dots \in F(\alpha, \beta, M)$ then,

$$|a_2| < (1 + \sigma)(1 - \beta) \cos \alpha, \quad \sigma = 1 - \frac{1}{M}. \quad (1.2)$$

If $\epsilon = \exp(-i \arg a_2 - i\alpha)$, then $\frac{f(\epsilon z)}{\epsilon} = z + |a_2| e^{-i\alpha} z^2 + \dots \in F(\alpha, \beta, M)$, whenever $f(z) \in F(\alpha, \beta, M)$. Thus without loss of generality we can replace the second coefficient a_2 of $f(z) \in F(\alpha, \beta, M)$ by $|a_2| e^{-i\alpha}$.

Let $F_p(\alpha, \beta, M)$ denote the class of functions $f(z) = z + |a_2| e^{-i\alpha} z^2 + \dots$,

which satisfy (1.1), where $|a_2| = p(1+\sigma)(1-\beta) \cos \alpha$. In view of (1.2) it follows that $0 < p < 1$.

Let $G_p(\alpha, \beta, M)$ denote the class of functions $g(z) = z + |b_2| e^{-i\alpha} z^2 + \dots$, regular in U and satisfy the condition

$$\left| \frac{e^{i\alpha} (1 + \frac{zg''(z)}{g'(z)}) - \beta \cos \alpha - i \sin \alpha}{(1-\beta) \cos \alpha} - M \right| < M, \quad z \in U, \quad (1.3)$$

where $|b_2| = \frac{1}{2} p(1+\sigma)(1-\beta) \cos \alpha$.

It follows from (1.1) and (1.3) that

$$g(z) \in G_p(\alpha, \beta, M), \text{ if and only if } zg'(z) \in F_p(\alpha, \beta, M). \quad (1.4)$$

We note that by giving specific values to p, α, β and M , we obtain the following important subclasses studied by various authors in earlier papers:

(i) $F_1(\alpha, \beta, M) = F_M(\alpha, \beta)$ and $G_1(\alpha, \beta, M) = G_M(\alpha, \beta)$, are respectively the class of bounded spirallike functions of order β and the class of bounded Robertson functions of order β investigated by Aouf [1] and $F_1(\alpha, 0, M) = F_{\alpha, M}$ and $G_1(\alpha, 0, M) = G_{\alpha, M}$, are respectively the class of bounded spirallike functions and the class of bounded Robertson functions investigated by Kulshrestha [2].

(ii) $F_p(\alpha, \beta, \infty) = F_p(\alpha, \beta)$ and $G_p(\alpha, \beta, \infty) = G_p(\alpha, \beta, \cdot)$, are considered by Umarani [3].

In this paper we determine the sharp radius of γ -spirallness of the functions belonging to the class $F_p(\alpha, \beta, M)$, generalizing an earlier result due to Kulshrestha [2], Libera [4], Umarani [5,3].

The technique employed to obtain this result is similar to that used by McCarty [6] and Umarani [3].

2. THE SHARP RADIUS OF γ -SPIRALNESS OF THE CLASS $F_p(\alpha, \beta, M)$, $M > 1$.

LEMMA 1. If $f(z) \in F_p(\alpha, \beta, M)$, $M > 1$, then $\left| \frac{zf'(z)}{f(z)} - w_0 \right| < \rho_0$, (2.1)

where

$$w_0 = \frac{(1+pr)^2 + \{[(1-\beta)(\frac{1+\sigma}{\sigma}) - 1] \cos \alpha - i \sin \alpha\} e^{-i\alpha} r^2 (r+p)^2}{(1-r^2)(1+2pr+r^2)} \quad (2.2)$$

and

$$\rho_0 = \frac{(1+\sigma)(1-\beta)\cos \alpha \ r(1+pr)(r+p)}{(1-r^2)(1+2pr+r^2)}. \quad (2.3)$$

This result is sharp.

PROOF. Let $f(z) \in F_p(\alpha, \beta, M)$, $M > 1$, then there exists a function $w(z)$ analytic in U and $|w(z)| < 1$ in U such that

$$e^{i\alpha} \frac{zf'(z)}{f(z)} = \cos \alpha \left\{ \frac{1 + [(1-\beta)(\frac{1+\sigma}{\sigma}) - 1] \sigma w(z)}{1 - \sigma w(z)} \right\} + i \sin \alpha, \quad \sigma = 1 - \frac{1}{M}$$

or

$$\frac{zf'(z)}{f(z)} = \frac{1 + \{[(1-\beta)(\frac{1+\sigma}{\sigma}) - 1] \cos \alpha - i \sin \alpha\} e^{-i\alpha} \sigma w(z)}{1 - \sigma w(z)}.$$

Solving for $w(z)$,

$$w(z) = \frac{\frac{zf'(z)}{f(z)} - 1}{\sigma \left[\frac{zf'(z)}{f(z)} + \{[(1-\beta)(\frac{1+\sigma}{\sigma}) - 1] \cos \alpha - i \sin \alpha\} e^{-i\alpha} \right]}.$$

Since $f(z) = z + |a_2| e^{-i\alpha} z^2 + \dots$, we obtain $w(z) = pz + \dots = z\phi(z)$, where $\phi(z)$

is analytic in U , $\phi(0) = p$ and $|\phi(z)| < 1$ in U . Now $\frac{\phi(z)-p}{1-p\phi(z)} z$. Therefore

$$\phi(z) = \frac{z+p}{1+pz}. \quad \text{Also } |w(z)| = |z\phi(z)| < \frac{|z|+p}{1+|z|p} |z|. \quad \text{Let } g(z) = \frac{|z|+p}{1+p|z|} z$$

and

$$h(z) = \frac{1 + \{[(1-\beta)(\frac{1+\sigma}{\sigma}) - 1] \cos \alpha - i \sin \alpha\} e^{-i\alpha} \sigma z}{1 - \sigma z}.$$

Since the image of $|z| < r$ under $g(z)$ is a disc and $h(z)$ is a bilinear transformation, then $\frac{zf'(z)}{f(z)}$ is subordinate to $(hog)(z)$. That is, the image of $|z| < r$ under $\frac{zf'(z)}{f(z)}$ is contained in the image of $|z| < r$ under $(hog)(z)$.

Equality in (2.1) can be attained by a function

$$f(z) = z(1-2p\sigma z + \sigma z^2) e^{-\frac{(1+\sigma)}{2\sigma}(1-\beta)\cos \alpha e^{-i\alpha} z} \quad (2.4)$$

$$= z + p(1+\sigma)(1-\beta)\cos\alpha e^{-i\alpha} z^2 + \dots ;$$

hence

$$\begin{aligned} \frac{zf'(z)}{f(z)} &= \frac{1-2p\sigma z + \sigma z^2 - (1+\sigma)(1-\beta)\cos\alpha e^{-i\alpha} z(z-p)}{1-2p\sigma z + \sigma z^2} \\ &= \frac{1 + \sigma \psi - (1+\sigma)(1-\beta)\cos\alpha e^{-i\alpha} \psi}{1 + \sigma \psi}, \end{aligned} \quad (2.5)$$

where $\psi = \frac{z(z-p)}{1-p\sigma z}$.

Since $p < 1$, $0 < \sigma < 1$, $|\psi| < 1$ for $z \in U$.

This shows that

$$e^{i\alpha} \frac{zf'(z)}{f(z)} = \cos\alpha \left\{ \frac{1 + [1 - (\frac{1+\sigma}{\sigma})(1-\beta)] \sigma \psi(z)}{1 + \sigma \psi(z)} \right\} + i \sin\alpha$$

and

$$\frac{e^{i\alpha} \frac{zf'(z)}{f(z)} - i \sin\alpha - \beta \cos\alpha}{(1-\beta)\cos\alpha} = \frac{1 - \psi(z)}{1 + \sigma \psi(z)}.$$

Then it is easy to show that $\left| \frac{1 - \psi(z)}{1 + \sigma \psi(z)} - M \right| < M$, $\sigma = 1 - \frac{1}{M}$. Thus $f \in F_p(\alpha, \beta, M)$.

Substituting $\psi = -\frac{\delta(\delta - \sigma e^{i\alpha})}{\sigma(1 - \sigma \delta e^{i\alpha})}$, where $\delta = \frac{r(r+p)}{1+rp}$ in (2.5), we find that

$$\left| \frac{zf'(z)}{f(z)} - w_0 \right| = \rho_0, \text{ where } w_0 \text{ and } \rho_0 \text{ are given by (2.2) and (2.3).}$$

This completes the proof of the lemma.

REMARK 1.

- (i) If $p=1$ and $\beta=0$ in Lemma 1, we obtain a result of Kulshrestha [2].
- (ii) If $M = \infty (\sigma=1)$ in Lemma 1, we obtain a result of Umarani [3].
- (iii) If $\alpha=0$ and $M=\infty (\sigma=1)$ in Lemma 1, we obtain a result of McCarty [6].

THEOREM 1. If $f(z) \in F_p(\alpha, \beta, M)$, > 1 , then $f(z)$ is γ -spiral
 $|z| < r_\gamma$, where r_γ is the smallest positive root of the equation

$$\begin{aligned} &\cos \gamma + p [2 \cos \gamma - (1+\sigma)(1-\beta)\cos\alpha]r + \\ &[p^2 \cos \gamma + cp^2 - (1+\sigma)(1-\beta)\cos\alpha(1+p^2)] r^2 \\ &+ p [2c - (1+\sigma)(1-\beta)\cos\alpha]r^3 + cr^4 = 0, \end{aligned} \quad (2.6)$$

where $c = \cos(\gamma - 2\alpha) + [(1-\beta)(\frac{1+\sigma}{\sigma}) - 2] \cos \alpha \cos(\gamma - \alpha)$. The result is sharp.

PROOF. Let $f(z) \in F_p(\alpha, \beta, M)$, $M > 1$, then by the above Lemma, we have

$$\left| \frac{zf'(z)}{f(z)} - w_0 \right| < \rho_0.$$

$$\text{Hence } \operatorname{Re} e^{i\gamma} \frac{zf'(z)}{f(z)} > \operatorname{Re} e^{i\gamma} \cdot w_0 - \rho_0$$

$$\begin{aligned} & \cos \gamma (1+pr)^2 + \operatorname{Re} \{ [(1-\beta)(\frac{1+\sigma}{\sigma}) - 1] \cos \alpha - i \sin \alpha \} e^{i(\gamma-\alpha)} r^2 (r+p)^2 \\ &= \left[\frac{-(1+\sigma)(1-\beta) \cos \alpha r(1+pr)(r+p)}{(1-r^2)(1+2pr+r^2)} \right] \\ & \cos \gamma (1+pr)^2 + \{ \cos(\gamma-2\alpha) + [(1-\beta)(\frac{1+\sigma}{\sigma}) - 2] \cos \alpha \cos(\gamma-\alpha) \} r^2 (r+p)^2 \\ &= \left[\frac{-(1+\sigma)(1-\beta) \cos \alpha r(1+pr)(r+p)}{(1-r^2)(1+2pr+r^2)} \right]. \end{aligned} \quad (2.7)$$

$f(z)$ is γ -spiral if the R.H.S. of (2.7) is positive. Hence $f(z)$ is γ -spiral for $|z| < r_\gamma$ where r_γ is the smallest positive root of the equation

$$\begin{aligned} & \cos \gamma (1+pr)^2 + \{ \cos(\gamma-2\alpha) + [(1-\beta)(\frac{1+\sigma}{\sigma}) - 2] \cos \alpha \cos(\gamma-\alpha) \} r^2 (r+p)^2 \\ & - (1+\sigma)(1-\beta) \cos \alpha r(1+pr)(r+p) = 0. \end{aligned}$$

Simplifying the above equation, we obtain (2.6).

If $\gamma=0$ in the above theorem, we obtain the radius of starlikeness of the class $F_p(\alpha, \beta, M)$.

COROLLARY 1. $f(z) \in F_p(\alpha, \beta, M)$, $M > 1$, is starlike for $|z| < r_0$, where r_0 is the least positive root of the equation

$$\begin{aligned} & 1+p [2-(1+\sigma)(1-\beta) \cos \alpha] r + \\ & \left(\frac{1+\sigma}{\sigma} \right) (1-\beta) \cos \alpha [\cos \alpha p^2 - \sigma (1+p^2)] r^2 + \\ & p [2c - (1+\sigma)(1-\beta) \cos \alpha] r^3 + cr^4 = 0, \end{aligned} \quad (2.8)$$

where $c = (\frac{1+\sigma}{\sigma})(1-\beta) \cos^2 \alpha - 1$.

If $p=1$, $\gamma=0$ and $\beta=0$ in Theorem 1, we obtain a result of Kulshrestha [2].

COROLLARY 2. $f(z) \in F_{\alpha, M}$, $M > 1$, is starlike for $|z| < r_0$, where r_0 is the least positive root of the equation

$$1 - (1+\sigma) \cos \alpha r + \left(\frac{1+\sigma}{\sigma} \right) \cos^2 \alpha - 1 \quad r^2 = 0.$$

REMARK 2.

(i) If $M=\infty$ ($\sigma=1$) in Theorem 1, we obtain a result of Umarani [3].

(ii) If $p=1$ and $M=\infty$ ($\sigma=1$) in Theorem 1, we obtain a result of Libera [4] and Umarani [5].

(iii) If $p=1$, $\beta=0$, $\gamma=0$ and $M=\infty$ ($\sigma=1$) in Theorem 1, we obtain a result of Robertson [7].

Since $g(z) \in G_p(\alpha, \beta, M)$ if and only if $zg'(z) \in F_p(\alpha, \beta, M)$ we obtain from Theorem 1,

THEOREM 2. If $g(z) \in G_p(\alpha, \beta, M)$, $M > 1$, then $\operatorname{Re} e^{i\gamma} \left(1 + \frac{zg''(z)}{g'(z)}\right) > 0$ for

$|z| < r_\gamma$, where r_γ is the least positive root of equation (2.6).

The result is sharp.

If $\gamma=0$ in Theorem 2, we obtain the radius of convexity of the class $G_p(\alpha, \beta, M)$.

COROLLARY 3. If $g(z) \in G_p(\alpha, \beta, M)$, $M > 1$, then the radius of convexity of $g(z)$ is the least positive root of equation (2.8).

REMARK 3.

(i) For $M=\infty$ ($\sigma=1$) in Theorem 2, and Corollary 3, we obtain a results of Umarani [3].

(ii) If $p=1$ and $\beta=0$ in Corollary 3, we obtain a result of Kulshrestha [2].

(iii) For $p=1$ and $M=\infty$ ($\sigma=1$), Theorem 2, generalizes the result of Umarani [5].

REFERENCES

1. AOUF, M.K. Bounded p -valent Robertson functions of order α , Indian J. of Pure and Appl. Maths., **16** (1985), 775-790.
2. KULSHRESTHA, P.K. Bounded Robertson functions, Rend. di Matematica, **(6)9**, (1976), 137-150.
3. UMARANI, P.G. Spiral-like functions with fixed second coefficient, Indian J. Pure appl. Math. **13(3)** (1982), 370-374.
4. LIBERA, R.J. Univalent α -spiral functions, Canad. J. Math. **19** (1967), 449-456.
5. UMARANI, P.G. Some studies in univalent functions, Ph.D. Thesis, Karnataka University, Dharwad (1976).
6. McCARTY, C.P. Two radius of convexity problems, Proc. Amer. Math. Soc. **42** (1972), 153-160.
7. ROBERTSON, M.S. Radii of starlikeness and close-to-convexity, Proc. Amer. Math. Soc. **16** (1965), 847-852.