RING HOMOMORPHISMS ON $H(G)$

N.R. NANDAKUMAR

Department of Mathematical Sciences
University of Delaware
Newark, Delaware 19716, U.S.A.

and

Department of Mathematics and Computer Science
University of Nebraska at Omaha
Omaha, Nebraska 68182, U.S.A.

(Received January 18, 1989)

ABSTRACT. It is shown that a ring homomorphism on $H(G)$, the algebra of analytic functions on a regular region G in the complex plane, is either linear or conjugate linear provided that the ring homomorphism takes the identity function into a nonconstant function.

KEY WORDS AND PHRASES. Ring homomorphism, Algebra of analytic function, Linear, Conjugate linear.

1980 AMS SUBJECT CLASSIFICATION CODE. 30H05.

1. INTRODUCTION.

An operator M on a commutative algebra A is called a **ring homomorphism** if for all $x, y \in A$, $M(x+y) = M(x) + M(y)$ and $M(xy) = M(x)M(y)$. Throughout this paper G denotes a region, i.e., a connected open set in the complex plane, $H(G)$ denotes the algebra of analytic functions on a region G in the complex plane equipped with the topology of uniform convergence on compact subsets of G , M denotes a nontrivial ring homomorphism on $H(G)$, and I denotes the identity function on G . A region G in C is called **regular** if $G = \text{interior}(\text{closure } G)$. The rationals, reals and complex numbers are denoted by Q , R , and C respectively.

If N is a maximal ideal in $H(G)$ then the quotient algebra $H(G)/N$ is isomorphic (as an algebra) to C if and only if N is the kernel of a linear homomorphism. Henriksen [1] has shown that $H(G)/N$ is isomorphic (as a ring) to C where $G=C$ and the maximal ideal M is not closed. This implies that there exist discontinuous homomorphisms from the ring of entire functions onto C .

In this paper we show that if G is a regular region in C and a ring homomorphism M on $H(G)$ takes the identity function I to a non-constant function, then M is necessarily continuous. Essentially we prove that homomorphisms under consideration preserve constants. However, the results of this fact can be obtained by the techniques used in [2] and [3] except in the case $G = C$.

If M is a ring homomorphism on $H(G)$ then the following assertions are equivalent:

- 1) M is continuous,
- 2) either $M(k) = k$ for all $k \in C$ or $M(k) = \bar{k}$ for all $k \in C$,
- 3) M is either linear or conjugate linear,
- 4) there exists $h \in H(G)$ with $h(G) \subset G$ such that $M(f) = foh$ for all $f \in H(G)$
or there exist $h \in H(G)$ with $\overline{h(G)} \subset G$ such that $M(f) = \overline{foh}$ for all $f \in H(G)$.

The implications $4 \Rightarrow 1) \Rightarrow 2) \Rightarrow 3)$ are trivial or easy to prove; $3) \Rightarrow 4)$ is the content of Lemma 2.1.

We show that a ring homomorphism M on $H(G)$ which takes the identity function to a non-constant function is necessarily linear or conjugate linear using Nienhuys-Thiemann's theorem [4] which states that given any two countable dense subsets A and B of R there exists an entire function which is real valued and increasing on the real line R such that $f(A) = B$. In Section 2 we give some lemmas and a theorem of Nienhuys and Thiemann. In Section 3 we prove the following main theorem.

THEOREM 1.1. Let G be a regular region in C and let M be a ring homomorphism on $H(G)$ such that $M(I)$ is not a constant function where I is the identity function. Then $M(i) = \pm i$. Further

- a) if $M(i) = i$ then M is linear,
- b) if $M(i) = -i$ then M is conjugate linear,

2. LEMMAS.

The following lemma is well known and we give the proof for the sake of completeness.

LEMMA 2.1. Let M be a ring homomorphism on $H(G)$. If M is linear then there exists a $h \in H(G)$ with $h(G) \subset G$ such that $M(f) = foh$ for all $f \in H(G)$.

PROOF. Let $M(I) = h$ and $z_0 \in G$. We claim that $h(z_0) \in G$. Suppose not, then

$$(I - h(z_0)) \left(\frac{1}{I - h(z_0)} \right) = 1.$$

Applying M on both sides and evaluating at z_0 with the observation that $M(h(z_0)) = h(z_0)$ we obtain

$$\begin{aligned} 0 &= (M(I)(z_0) - h(z_0)) M\left(\frac{1}{I - h(z_0)}\right)(z_0) \\ &= M(I - h(z_0))(z_0) M\left(\frac{1}{I - h(z_0)}\right)(z_0) \\ &= M(1)(z_0) \\ &= 1. \end{aligned}$$

which is a contradiction. Since z_0 is arbitrary we have $h(G) \subset G$.

$$\begin{aligned} \text{Since } h(z_0) \in G \text{ we have } \frac{f - f(h(z_0))}{I - h(z_0)} \in H(G) \text{ and} \\ f - f(h(z_0))0 = (I - h(z_0)) \left(\frac{f - f(h(z_0))}{I - h(z_0)} \right). \end{aligned}$$

Applying M on both sides and evaluating at z_0 we obtain

$$M(f)(z_0) = M(f(h(z_0)))(z_0) = f(h(z_0)) .$$

Since z_0 is arbitrary the result follows.

LEMMA 2.2. Let G be a regular region in C and M be a ring homomorphism on $H(G)$ with $M(i) = i$. If $M(I) = h$ is not a constant function then $h(G) \subset G$.

PROOF. Since M is a nontrivial ring homomorphism it is easy to show that $M(\alpha) = \alpha$ for all $\alpha \in Q$. Since $M(i) = i$ we have $M(\alpha + i\beta) = \alpha + i\beta$ where $\alpha, \beta \in Q$. Let $z_0 \in G$ such that $h(z_0) \in Q + iQ$. Just as in the above lemma it is easy to show that $h(z_0) \in G$. Since h is not a constant function we have $h(z_0) \in G$ for a dense set of z_0 in G and since $h(G)$ is open we have $h(G) \subset \text{interior}(\text{closure } G) = G$.

Let $K \in Q$. Denote by H_K the set of all entire functions which map $Q + iK$ into Q except possibly for one point of $Q + iK$ and also denote by EM the class of entire functions whose restriction to R is a real monotonically increasing function. The proof of Lemma 2.3 follows the proof of the following theorem of Nienhuys & Thiemann [4].

THEOREM 2.1. Let S and T be countable everywhere dense subsets of R , let p be a continuous positive real function such that $\lim_{t \rightarrow \infty} t^{-n} p(t) = \infty$ for all $n \in N$ and let $f_0 \in EM$.

Then there exists a function $f \in EM$ such that

- i) f is strictly increasing on R and $f(S) = T$,
- ii) $|f(z) - f_0(z)| < p(|z|)$ for all $z \in C$.

LEMMA 2.3. Let $k \in Q$, $\beta \in R$ and $\alpha \in Q + ik$. Then there exists an entire function $f \in H_k$ such that $f(\alpha) = \beta$ and $f(Q + ik) = \{\beta\} \cup Q$.

PROOF. In Nienhuys and Thiemann's Theorem [4] take $S = Q$ and $T = \{\beta\} \cup Q$. Let $x_1, x_2, \dots$, be an enumeration of Q . Then as in the proof of this theorem there exists an entire function g such that $g(x_1) = \beta$ and $g(Q) = \{\beta\} \cup Q$. Let $x_1 = \alpha - ik$ and $h(z) = z - ik$. Then $f = goh$ is the desired function.

3. PROOF OF THE MAIN THEOREM.

It is easy to see that M is linear over the field of rational numbers and hence we have $-1 = M(-1) = M(i^2) = M(i)^2$ which implies $M(i) = \pm i$. We prove here only Part a) of the theorem and the proof of Part b) follows similarly.

Since $h = M(I)$ is a nonconstant analytic function on G , $h(G)$ is a nonempty open set in C and by Lemma 2.2, $h(G) \subset G$. Hence there exists $k \in Q$ such that $S = (R + ik) \cap h(G)$ has an interval parallel to real axis. Let $f \in H(G)$ and $h(z_0) \in (Q + ik) \cap G$. Then applying M on both sides and evaluating at z_0 in the following

$$f - f(h(z_0)) = (I - h(z_0)) \left(\frac{f - f(h(z_0))}{I - h(z_0)} \right)$$

we obtain

$$M(f - f(h(z_0)))(z_0) = 0$$

for all z_0 in G such that $h(z_0) \in Q + ik$. Thus we have

$$M(f)(z_0) = M(f(h(z_0)))(z_0), \text{ for all } f \in H(G) \text{ and for all } h(z_0) \in Q + ik. \quad (1)$$

Since a function f in H_k takes $Q + ik$ into rationals except for one point of $Q + ik$, we obtain $M(f(h(z_0))) = f(h(z_0))$ whenever $h(z_0)$ is in $(Q + ik) \cap G$. Since f is analytic we obtain

$$M(f) = foh, \text{ for all } f \in H_k. \quad (2)$$

For a given $\beta \in \mathbb{R}$ and each $h(z_0)$ in $Q + ik$, by Lemma 2.3 there exists an entire function in H_k such that $f(h(z_0)) = \beta$. Substituting this in (1) on the one hand we obtain

$$M(f)(z_0) = M(\beta)(z_0)$$

and evaluating (2) at z_0 on the other we obtain

$$M(f)(z_0) = foh(z_0) = f(h(z_0)) = \beta$$

Thus we obtain from the above two relations that

$$M(\beta)(z_0) = \beta \text{ for all } z_0 \in h^{-1}(Q + ik) \cap G.$$

Since $M(\beta)$ is analytic we have $M(\beta) = \beta$. Thus we have $M(k) = k$ for all $k \in \mathbb{C}$. This implies M is linear.

4. ACKNOWLEDGEMENTS.

The author wishes to acknowledge Professor Lee A. Rubel and Professor Peter J. de Paepe for their helpful suggestions.

REFERENCES

1. HENRIKSEN, M., On the Ideal Structure of the Ring of Entire Functions, Pacific J. Math., 2 (1952), 179-184.
2. BERS, L., Rings of Analytic Functions. Bull Amer. Math. Soc., 54 (1948), 311-315.
3. RUDIN, W., Some Theorems on Bounded Analytic Functions, Trans. Amer. Math. Soc., 78 (1955), 333-342.
4. NIENHUYIS, J.W and THIEMANN, J.G.F., On the Existence of Entire Functions Mapping Countable Dense Subsets on Each Other, Proc. Kon. Ned. Akad. Van Wetensch., A 79 (1976), 331-334.