Internat. J. Math. & Math. Sci.
VOL. 13 NO. 1 (1990) 129-134

ON SEMI-HOMEOMORPHISMS

J.P. LEE

Department of Mathematics
State University of New York, College at Old Westbury
Old Westbury, New York 11568

(Received May 5, 1989 and in revised form July 7, 1989)

ABSTRACT. In the first part of our work we show a condition for a semi-homeomorphism in the sense of Crossley and Hildebrand (s.h.C.H) to be a semi-homeomorphism in the sense of Biswas (s.h.B). Certain relevant examples are provided. Next, we define strong semi-homeomorphisms via "nice" restrictions of semi-homeomorphisms ("global condition") and we show that the new class of functions actually coincides with semi-homeomorphisms. Then, in the third part we introduce local semi-homeomorphisms (l.s.h.C.H.) via a corresponding "local condition" for restrictions. A few results pertaining to the preservation of some topological properties under this new class of functions are examined.

KEY WORDS AND PHRASES. semi-homeomorphisms
1980 AMS Subject Classification Code. 54C10

1. s.h.C.H. VERSUS s.h.B.

We shall start with the following definitions.

A subset $S \subset X$ is said to be *semi-open* if there is an open set $U \subset X$ such that $U \subset S \subset \bar{U}$.

A function $f: X \to Y$ is said to be a *semi-homeomorphism in the sense of Crossley and Hildebrand* (or simply, s.h.C.H.) [1] if:

1. f is bijective
2. f is irresolute (i.e. inverse images of semi-open sets are semi-open)
3. f is pre-semi-open (i.e. images of semi-open sets are semi-open)

Further, a function $f: X \to Y$ is said to be a *semi-homeomorphism in the sense of Biswas* (or simply s.h.B. ([2]), if

1. f is bijective
2. f is continuous
3. f is semi-open

Clearly every homeomorphism is both s.h.B and s.h.C.H. .

T. Neubrunn [3] has shown that there are s.h.C.H. that are not s.h.B. . Answering his question Z. Piotrowski [4] has shown an example of a s.h.B. which is not s.h.C.H. Further he also obtained certain conditions for a s.h.C.H. to be a s.h.B.

In this paragraph we shall prove the following.

Proposition 1. Assume Y has a clopen base. If $f: X \to Y$ is one-to-one, semi-open and somewhat continuous then f is irresolute.

PROOF: Let $A \subset Y$ be semi-open. Let $x \in f^{-1}(A)$ i.e. $f(x) = y \in A$. We shall show that $x \in \overline{\operatorname{Int} f^{-1}(A)}$.

For any open set U containing x, the set f(U) is semi-open and contains y. Further $f(U) \cap A \neq \emptyset$. Since f(U) is open, Y having a clopen base and A is open - all semi-open and open sets coincide, under the assumption upon Y, there is a nonempty open set G such that

$$G = f(U) \cap A. \tag{1.1}$$

Clearly,

$$f^{-1}(G) \subset f^{-1}(f(U) \cap A) \subset f^{-1}(f(U)) = U \tag{1.2}$$

f being one-to-one.

Now, somewhat continuity of f implies that there is an open set $V \subset f^{-1}(G)$, $V \neq \emptyset$. Therefore $V \subset U$, $V \subset f^{-1}(A)$. And since U is an arbitrary neighborhood of x, we have $x \in \overline{\operatorname{Int} f^{-1}(A)}$. Thus $f^{-1}(A)$ is semi-open. □

REMARK: The author is indebted to the referee for pointing out that Proposition 1 generalizes Theorem 2.2 of [5].

The assumption upon Y to have a *clopen base* is essential. In fact:

EXAMPLE 2. There is a semi-open, semi-continuous (hence somewhat continuous!) bijection $f: [0,1] \longrightarrow [0,1]$ which is *not* irresolute. Take $f(x) = x$, if $x \in \left[0,\frac{1}{3}\right)$, $f(x) = x + \frac{1}{3}$, if $x \in \left[\frac{1}{3}, \frac{2}{3}\right]$. $f(x) = -x + \frac{4}{3}$, if $x \in \left(\frac{2}{3},1\right]$. Observe that $f^{-1}\left[\frac{1}{3}, \frac{2}{3}\right]$ is not semi-open.

In fact, there is even a *continuous*, semi-open injective function between two topological spaces which is not irresolute. We shall provide here such an example, originally designed for a different purpose.

EXAMPLE 3. ([4], Example 19, p. 8) Let $X = Y = \{a, b, c, d\}$. Let $\mathcal{O}_1$ and $\mathcal{O}_2$ denote the topologies for X and Y, respectively, such that $\mathcal{O}_1 = \{\emptyset, X, \{a\}, \{b\}, \{a,b\}, \{b,c,d\}\}$ and $\mathcal{O}_2 = \{\emptyset, Y, \{a\}, \{b\}, \{a,b\}\}$. Let $f:(X,\mathcal{O}_1) \longrightarrow (Y,\mathcal{O}_2)$ be the identity function. It is easy to see that f is continuous and semi-open but not irresolute, since $\{a,c\}$ is semi-open in Y while it is not semi-open in X.

REMARK 4. Example 2 above is *the best possible* in the class of semi-continuous bijections $f: [0,1] \longrightarrow [0,1]$ (or more generally, $f: X \to Y$, X-compact, Hausdorff and Y

being Hausdorff,) in the sense that if f is to be additionally continuous, then, being continuous bijection from a compact, Hausdorff space onto a Hausdorff space, it is a homeomorphism, (see [6], Thm 2.1, p. 226). Now, every homeomorphism (actually openess and continuity suffices) implies irresoluteness of f - we leave the proof of this fact to the reader, also see [1].

Since it is well-known that continuity and somewhat openess imply pre-semi-openess, see also [1] we have the following Corollary from Proposition 1.

COROLLARY 5. Assume Y has a clopen base. If $f: X \to Y$ is s.h.B, then f is s.h.C.H. .

2. STRONG SEMI-HOMEOMORPHISMS ARE PRECISELY SEMI-HOMEOMORPHISMS.

In this paragraph a "semi-homeomorphism" stands for s.h.C.H. .

The following, seemingly stronger conditions (*) and (**) which define - what we call - a *strong semi-homeomorphism* are actually equivalent (!) to the semi-homeomorphicity of f, see the following.

THEOREM 6. A function $f: X \to Y$ is a semi-homeomorphism if and only if:

(*) f is bijective and

(**) $\forall\, U \subset X$, U-open, $f|U$ is a semi-homeomorphism

PROOF: In fact, it is easy to see that if f satisfies (*) and (**), then f is a semi-homeomorphism - take $U = X$ in (**). Conversely, let $X = \cup \{U_\alpha : \alpha \in A\}$, where each U_α is open and suppose that each restriction $f|U_\alpha$ is both pre-semi-open and irresolute. We shall show that f is also such.

Let $(f|U_\alpha) : U_\alpha \to Y$ denote the restriction of f to U_α. We shall show that f is irresolute. Really, given a semi-open set $K \subset Y$ we have:

$$f^{-1}(K) = \cup \{f^{-1}(K) \cap U_\alpha : \alpha \in A\} = \cup \{(f|U_\alpha)^{-1}(K) : \alpha \in A\}. \qquad (2.1)$$

The latter set is semi-open as the sum of semi-open sets.

Similarly, we shall prove that f is pre-semi-open. Let $L \subset X$ be semi-open, in X. Then $L = \cup \{L \cap U_\alpha : \alpha \in A\}$. Then:

$$\begin{aligned} f(L) &= f(\cup \{L \cap U_\alpha : \alpha \in A\}) = \\ &= \cup f\{L \cap U_\alpha : \alpha \in A\} = \\ &= \cup \{(f|U_\alpha)(L) : \alpha \in A\}. \end{aligned} \qquad (2.2)$$

And again, the latter set is semi-open, in Y. □

The following example shows that the assumption "for *every* open" in (**) is real. As one can see, the restrictions $f|U_\alpha$, $\alpha \in A$ are even *homeomorphisms* (!) for every $U_\alpha \neq X$.

EXAMPLE 7. (See Example 3 of §1.) There is a function $f: X \to Y$ such that

1. f is bijective and
2. $\forall\, U_\alpha \subsetneq X$, U_α-open, $\alpha \in A$, $f|U_\alpha$ is a homeomorphism

(hence, a semi-homeomorphism) whereas $f: X \to Y$ is not a semi-homeomorphism.

Really, $f|\{a\}$, $f|\{b\}$ and $f|\{a,b\}$ are homeomorphisms. Now, consider $f|\{b,c,d\}$. We have $X = Y = \{b,c,d\}$ and $\mathcal{O}_1 \cap X = \{\emptyset, \{b,c,d\}, \{b\}\}$, whereas $\mathcal{O}_2 \cap Y = \{\emptyset, \{b,c,d\}, \{b\}\}$. And, here again, $f|\{b,c,d\}$ is a homeomorphism.

3. LOCAL SEMI-HOMEOMORPHISMS.

Local homeomorphisms, being a very natural generalization of homeomorphisms, occupy an important place in topology, especially in the theory of 1-dimensional continua (curves) as well as some parts of algebraic topology, see also [7] for an extensive treatment of this topic.

Let us define our new class of functions. We say that a function $f: X \to Y$ is a *local semi-homeomorphism* in the sense of Crossley and Hildebrand if:

1. f is bijective and
2. $\forall x \in X \; \exists$ U-open, $x \in U \subset X$ such that $f|U$ is a semi-homeomorphism in the sense of Crossley and Hildebrand.

Well, it is easy to see that every semi-homeomorphism is a local semi-homeomorphism; take $U = X$. Since every homeomorphism is a semi-homeomorphism, see [1] we have the following diagram:

$$\text{homeomorphism} \Rightarrow \overset{\text{strong}}{\text{semi-homeomorphism}} \Leftrightarrow \text{semi-homeomorphism} \Rightarrow \text{local semi-homeomorphism}$$

We shall now provide an example of a local semi-homeomorphism which is not a semi-homeomorphism, showing that the arrow to the right is, in general, not reversable.

EXAMPLE 8. Consider Example 3, see §1. Take $\{a\}$, $\{b\}$, $\{b,c,d\}$, $\{b,c,d\}$, respectively for open neighborhoods of a, b, c and d, respectively. Using arguments similar to ones applied in Example 7 we prove that f is a local semi-homeomorphism; it has been shown in [4], p. 508 that f is not a semi-homeomorphism.

LEMMA 9. If for every $x \in X$ there is an open set $U \subset X$, $x \in U$ such that $f|U$ is a semi-homeomorphism in the sense of Crossley and Hildebrand, then f is somewhat continuous (inverse images of every nonempty open set if nonempty it has the nonempty interior) and f is somewhat open (image of every open nonempty set has the nonempty interior).

PROOF: Let D be a dense set in X. We shall show that $f(D)$ is dense in $f(X)$. This, in turn, shows that f is somewhat continuous.

In fact, suppose $y \in f(X)\backslash f(D)$ and assume further that there is an open neighborhood V containing y, such that:

$$(*) \qquad V \cap f(D) = \emptyset. \qquad (3.1)$$

Since f is "onto", there is $x \in X$, such that $f(x) = y$. There is an open set $U \ni x$ such that $f|U$ is a semi-homeomorphism, f being a local semi-homeomorphism. Clearly

$D \cap U$ is dense in U; further $f(D \cap U)$ is dense in $f(U)$, f being semi-homeomorphism on U. Now, $f(U)$ is a semi-open set containing $f(x) = y$. By an elementary property of semi-open sets, $f(D \cap U)$ is dense in $V \cap \operatorname{Int} f(U)$, and hence, also in $V \cap f(U)$. So, $V \cap f(D) \neq \emptyset$, contradicting (*).

Now, for somewhat openess part, consider a dense set D contained in $f(X)$. We shall show that $f^{-1}(D)$ is dense (in X). Suppose $f^{-1}(D)$ is not dense. So, there is a point $x \in X$ and an open neighborhood $U \ni x$ such that

$$(**) \qquad U \cap f^{-1}(D) = \emptyset . \qquad (3.2)$$

Without loss of generality we may assume that U is the open neighborhood of x from the definition of local homeomorphism (or, simply, take the intersection of the two sets, in question). Then $f(U)$ is a semi-open set, free of points of D. For otherwise the set:

$$f^{-}(f(U) \cap D) = f^{-1}(f(U)) \cap f^{-1}(D) = U \cap f^{-1}(D) \neq \emptyset, \qquad (3.3)$$

contradicting (**), which finishes the proof.

COROLLARY 10. Baireness is a local semi-topological property.

PROOF: See [8], Corollary 2, p. 410 and Lemma 9, above.

COROLLARY 11. Separability is a local semi-topological property.

PROOF: Every local semi-homeomorphism is somewhat continuous, and this implies (see [9]) that dense subsets are preserved, which in turn proves our claim.

We will close this work with the following natural

Question 12. What are the topological conditions for X and/or Y so that every local semi-homeomorphism $f: X \to Y$ is a semi-homeomorphism?

ACKNOWLEDGEMENT. This research was supported by the State University of New York Graduate and Research Initiative Grant.

REFERENCES

1. CROSSLEY, S.G. and HILDEBRAND, S. K., Semi-topological properties, Fund. Math. 74(1972), 233-254.

2. BISWAS, N., On some mappings in topological spaces, Ph.D. Thesis, University of Calcutta, 1971.

3. NEUBRUNN, T., On semi-homeomorphisms and related mappings, Acta Fac. Rerum Natur. Comenian. Mat. 33(1977), 133-137.

4. PIOTROWSKI, Z., On semi-homeomorphisms, Boll. Un. Mat. Ital., 16-A(1979), 501-509.

5. NOIRI, T., A note on semi-homeomorphisms, Bull. Calcutta Math. Soc. 76(1984), 1-3.

6. DUGUNDJI, J., Topology, Allyn and Bacon, Boston, 1966.

7. JUNGCK, G., Local homeomorphisms, Dissertationes Math. 209 (1983), Warsaw.

8. DOBOŠ, J. J., A note on the invariance of Baire spaces under mappings, Časopis pro pěst. matem. 108(1983), 409-411.

9. FROLIK, Z., Remarks concerning the invariance of Baire spaces under mappings, Czechoslovak Math. J. 11(86), (1961), 381-385.