INVERSES OF MEASURES ON A CLASS OF DISCRETE GROUPS

C. KARANIKAS

Department of Mathematics
Faculty of Science
Aristotle University of Thessaloniki
54006 Thessaloniki, Greece

(Received February 12, 1990)

ABSTRACT. We examine a class of groups G , having a certain growth condition. We given an estimate for the norm of the inverse of an element in $l_1(G)$ in terms of the spectral radius and the cardinality of the support.

KEY WORDS AND PHRASES. Discrete groups, spectral radius and measure algebra.

1980 AMS SUBJECT CLASSIFICATION CODES. 43A05, 43A10, 22D05.

1. INTRODUCTION.

Throughout this work G is a discrete group and $M(G)$ is the usual measure algebra on G : we write δ for the unit of $M(G)$, $\mu * \nu$ for the (convolution) product of two measures $\mu, \nu \in M(G)$.

The purpose of this paper is to investigate the following problems: Let $\mu \in M(G)$ have finite support and the (convolution) inverse of $(\delta - \mu)$ exist in $M(G)$. Is it possible to estimate the norm $\|(\delta - \mu)^{-1}\|$ of it?

One can easily realize that this problem becomes more interesting in the "limit" case, where the support of μ is infinite. One also can realize the connection of this with the general problem of the invertibility in $M(G)$; namely the characterization of the class of Hermitian groups G (see [1]) or the equality of different norms and spectrums in $M(G)$ (see [2], [3] and [4]).

Our work here can be separated into two parts. In the first part we consider the class of groups A and we give an estimate of $\|(\delta - \mu)^{-1}\|$ and of $\|\exp \mu\|$ in terms of the spectral radius $r(\mu)$ and of the cardinality of the support of μ .

In the second part we examine the relation of the class A , with the class of nilpotent groups and [FC] groups (groups having finite conjugacy class) (see [1]). We show that nilpotent groups are A -groups and that the class A is closed under finite extension. We should note that by Gromov's well known result, any finitely generated group G with polynomial growth is a finite extension of a nilpotent group, and so it is an A -group. It is remarkable to note that all known Hermitian groups, as they are

referred to in [1], are A-groups. We shall complete this introduction with some definitions and notations.

We say that G is an A-group if there is a map $\kappa: J \rightarrow \mathbb{N}$, where J is the set of all finite subsets of G , and a $\lambda \in \mathbb{N}$ such that for any $F \in J$

$$\#(F^n) \leq \binom{\kappa(F)+n-1}{n} \lambda, \quad n \in \mathbb{N}$$

where $F^n = F F \dots F$ (n -times) and $\#F$ is the cardinality of F .

An n -word in the elements of F is any (reduced) word of length n . We denote by $(F^n)'$ a collection of n -words in the elements of F which as a subset of G consists of all distinct elements of F^n . Finally we shall denote by μ^n the convolution product $\mu * \mu \dots * \mu$ (n -times); the spectral radius $r(\mu)$ of μ is the limit $\lim_{n \rightarrow \infty} \|\mu^n\|^{1/n}$.

2. NORMS OF CERTAIN INVERTIBLE MEASURES.

We are going to show the following:

THEOREM 2.1. Let G be an A-group and let μ be an element in $M(G)$, with finite support F and $r(\mu) < 1$. Then $\delta - \mu$ is invertible such that

$$\|(\delta - \mu)^{-1}\| \leq (1 - r(\mu))^{-\kappa(F)\lambda} \quad (2.1)$$

where $\kappa(F)$ and λ are constants determined from the A-group structure. Furthermore

$$\text{if } \exp \mu = \sum_{n=0}^{\infty} \frac{\mu^n}{n!} \text{ we have} \quad \|\exp \mu\| \leq \exp(\kappa(F)\lambda r(\mu)) \quad (2.2)$$

PROOF. First we observe that for any $n \in \mathbb{N}$

$$\begin{aligned} \|\mu^n\| &= \sum_{x \in G} |\mu^n(x)| \\ &\leq \#(F^n)^{1/2} \|\mu^n\|_2 \\ &\leq \#(F^n)^{1/2} \sup \| \mu^n * g \|_2, (g \in l_2(G), \|g\|_2 = 1) \\ &\leq \#(F^n)^{1/2} \rho(\mu)^n \end{aligned}$$

where $\|\cdot\|_2$ is the norm in $l_2(G)$ and $\rho(\mu)$ is the norm of the left regular representation, i.e. the norm of the operator $\mu: l_2(G) \rightarrow l_2(G): f \mapsto \mu * f$. Since $\rho(\mu) \leq r(\mu)$ we have

$$\|\mu^n\| \leq \#(F^n)^{1/2} r(\mu)^n. \quad (2.3)$$

Now by (2.3)

$$\begin{aligned}
 \left| \left| \delta + \mu + \mu^2 + \dots \right| \right| &\leq 1 + \#(F)^{1/2} r(\mu) + \\
 &+ \#(F^2)^{1/2} r(\mu)^2 + \dots \\
 &< \sum_{n=0}^{\infty} \binom{\kappa(F)+n-1}{n}^{\lambda} r(\mu)^n
 \end{aligned}$$

Now let $K = \kappa(F)$ we see that $\binom{K+n-1}{n}^{\lambda} < \binom{K^{\lambda}+n-1}{n}$. In fact,

$$\begin{aligned}
 \binom{K+n-1}{n}^{\lambda} &= \prod_{j=1}^n \left(\frac{K-1}{j} + 1 \right)^{\lambda} \\
 &= \prod_{j=1}^n \left(1 + \sum_{m=1}^{\lambda} \binom{\lambda}{m} \left(\frac{K-1}{j} \right)^m \right) \\
 &< \prod_{j=1}^n \left(\frac{1}{j} [(K-1)+1]^{\lambda} + \frac{j-1}{j} \right) \\
 &< \prod_{j=1}^n \left(\frac{K^{\lambda}-1}{j} + 1 \right) \\
 &< \binom{K^{\lambda}+n-1}{n}
 \end{aligned}$$

Thus

$$\left| \left| \delta + \mu + \mu^2 + \dots \right| \right| < \sum_{n=0}^{\infty} \binom{\kappa(F)^{\lambda}+n-1}{n} r(\mu)^n$$

and since $r(\mu) < 1$, by the binomial formula, $(\delta - \mu)^{-1}$ exists and we obtain (2.1).

To see (2.2) we observe that since $\kappa > 1$ and $j \geq 1$, $\left(\frac{\kappa-1}{j} + 1 \right) \leq \kappa$ and

$$\begin{aligned}
 \binom{\kappa(F)+n-1}{n}^{\lambda} &< \prod_{j=1}^n \left(\frac{\kappa(F)-1}{j} + 1 \right)^{\lambda} \\
 &< \kappa(F)^{n\lambda}
 \end{aligned} \tag{2.4}$$

Thus by (2.3) and (2.4)

$$\begin{aligned}
 \left| \left| \exp \mu \right| \right| &< \sum_{n=0}^{\infty} \frac{\#(F^n)}{n!} r(\mu)^n \\
 &< \sum_{n=0}^{\infty} \frac{1}{n!} \binom{\kappa(F)+n-1}{n}^{\lambda} r(\mu)^n \\
 &< \sum_{n=0}^{\infty} \frac{1}{n!} (K(F)^{\lambda} r(\mu))^n \\
 &= \exp(K(F)^{\lambda} r(\mu)).
 \end{aligned}$$

3. THE CLASS OF GROUPS A.

In this section, we show that [FC] groups and nilpotent groups are A-groups. We also show that the class A is closed under finite extension.

First we examine the growth of [FC] groups.

PROPOSITION 3.1. Let G be an $[FC]$ group and F be a finite subset of G . Then

$$\#(F^r) \leq \binom{k+r-1}{r} \quad (r \in \mathbb{N})$$

where $k = \# \left(\bigcup_{f \in F} [f] \right)$, $[f]$ is the conjugacy class of f .

PROOF. We show that if $[F] = \bigcup_{f \in F} [f]$

$$\#(F^2) \leq \#([F]^2) \leq \binom{k+1}{2}$$

Given $f, g \in F$, $f \neq g$, there is an element g_1 (say) in $[g] \subset [F]$ such that $fgf^{-1} = g_1$, and so $fg = g_1f$.

Hence any 2-word in the elements of $[F]$ consisting of two different letters is equal to another 2-word in the elements of $[F]$.

Thus $[F]^2$ has no more than k elements f^2 and $k(k-1)/2$ elements fg where $f, g \in F$, $f \neq g$. It is clear that

$$\#(F^2) \leq k + \frac{k(k-1)}{2} = \binom{k+1}{2}.$$

We suppose that the Theorem is true for any $r < n$, we show that it is also true for $r = n$.

We denote by $\phi(g)$ the number of all appearances of a $g \in [F]$ in the words of $([F]^n)'$.

If all elements of $[F]$ had the same chance to appear in $([F]^n)'$ then for each $g \in [F]$, $\phi(g) = \frac{n}{k} ([F]^n)'$.

Thus we may consider a $g \in [F]$ such that

$$\#([F]^n) \leq \frac{k}{n} \phi(g) \quad (3.1)$$

Since for any $f \in [F]$ $f \neq g$, there is some f_1 (say) such that $fg = gf_1$. Hence without loss of generality we may assume that in any word of $([F]^n)'$, either there is no g or all g 's keep the left place of the word. Now from each word of $([F]^n)'$, where g appears, cancel one g . The resulting $(n-1)$ -words form a subset of distinct elements of $[F]^{n-1}$; we denote this set by $g^{-1}([F]^n)'$. Hence from our hypothesis it is clear that

$$\phi(g) \leq \binom{k+n-2}{n-1} + \phi_1(g) \quad (3.2)$$

where $\phi_1(g)$ is the number of all appearances of g in $g^{-1}([F]^n)'$. Suppose that

$$\phi_1(g) \leq \frac{n-1}{k} \binom{k+n-2}{n-1} \quad (3.3)$$

then by (3.1) and (3.2) we obtain

$$\#([F]^n) \leq \frac{k}{n} \left(1 + \frac{n-1}{k} \right) \binom{k+n-2}{n-1} \leq \binom{k+n-1}{n}$$

and so in this case there is nothing to show.

If the inequality (3.3) does not occur, from (3.1) and (3.2) we have

$$([F]^n) \leq \frac{k}{n} \left(\frac{k}{n-1} + 1 \right) \phi_1(g) \leq \frac{k(k+n-1)}{n(n-1)} \phi_1(g)$$

In a similar way we define $\phi_i(g)$ ($1 \leq i \leq n-1$) i.e. the number of appearances of g in the collection $g^{-1}([F]^n)$.

As in (3.2)

$$\phi_{i-1}(g) \leq \left(\frac{k+n-i-1}{n-i} \right) + \phi_i(g) \quad (i=2,3,\dots,n-1)$$

and if for some $i \leq n-1$

$$\phi_i(g) \leq \frac{n-i}{k} \left(\frac{k+n-i-1}{n-i} \right) \quad (3.4)$$

it is nothing to show. If the inequality (3.4) does not occur for any $i \leq n-1$ we observe that

$$\phi_{n-1}(g) \leq \frac{1}{k} \binom{k}{1}$$

In this case we write

$$\#([F]^n) \leq \frac{k(k+n-1) \dots (k+1)}{n \cdot (n-1) \dots 1} = \frac{(k+n-1)!}{(k-1)!n!}$$

and this completes the proof.

COROLLARY 3.1. If G is abelian and F is a finite subset of G then,

$$(F^r) \leq \binom{\#F+r-1}{r}, \quad r \in \mathbb{N}$$

PROOF. Clear

LEMMA 3.1. Let G be a discrete group with a normal subgroup K such that G/K is abelian, and let π be the canonical map $\pi: G \rightarrow G/K$. If $F \subset G$ is such that $\#F = \#\pi(F)$, then the number of all r -words in the elements of F ($r \in \mathbb{N}$), in a given class of G modulo K can not be greater than

$$\binom{\#F+r-1}{r}.$$

PROOF. Let $\tilde{x} \in G/K$ fixed and $J_r = \pi^{-1}(\tilde{x}) \cap F^r$

Note that J_r and F^r in this proof mean collections of r -words, which as elements of G may not be distinct.

We shall denote by $\#J_r$ the cardinality of J_r , and we shall show that

$$\#([F]^n) \leq \binom{\#F+1}{2} \quad (3.5)$$

Let $x, y \in F$ and $xy \in J_2$, then $yx \in J_2$; in fact since G/K is abelian $\pi(xy) = \pi(x)\pi(y) = \pi(yx)$. Now suppose that there is a $z \in F$ such that x, z (or z, x) is in J_2 , i.e. $\pi(x, y) = \pi(x, z)$ and so $\pi(y) = \pi(z)$ and $\#\pi(F) < \#F$ -contradiction. Hence it is clear that $\#J_2 < \#F$ and (3.5) follows.

We suppose that Lemma 3.1 is true for any $r > n-1$ and we show this for $r=n$. For

for some $\tilde{g} \in G/K$, let

$$J_n = \pi^{-1}(\tilde{g}) \cap F^n.$$

If all elements of F had the same chance to appear in J_n , then the number of all appearances $\phi(x)$, $x \in F$, of x in the words of J_n should be

$$\phi(x) = \frac{n}{\#F} \#(J_n)$$

We consider a $x \in F$ such that

$$\#J_n < \frac{\#F}{n} \phi(x) \quad (3.6)$$

From the set of all words in J_n where x has at least one entry we cancel one x . We denote by ${}_x J_n$ the collection of all the resulting $(n-1)$ -words. We show that ${}_x J_n$ is in a class of G modulo K .

Let w_1, w_2, w'_1, w'_2 , be words in the elements of F such that $w_1 \times w_2, w'_1 \times w'_2 \in J_n$. Since $\pi(w_1 \times w_2) = \pi(w'_1 \times w'_2)$, we have $\pi(w_1 w_2) = \pi(w'_1 w'_2)$ and ${}_x J_n$ is as we claimed.

Now, the set ${}_x J_n$ by our inductive hypothesis has cardinality no greater than $\binom{\#F+n-2}{n-1}$ and so

$$\phi(x) < \binom{\#F+n-2}{n-1} + \phi_1(x)$$

where $\phi_1(x)$ is the number of appearances of x in ${}_x J_n$.

As in Proposition (3.1), (3.7) in the case where

$$\phi_1(x) < \frac{n-1}{\#F} \binom{\#F+n-2}{n-1}$$

it is nothing to show. If the inequality above is not true by (3.6) and (3.7) we obtain

$$\#J_n < \frac{\#F}{n} \frac{\#F+n-1}{n-1} \phi_1(x)$$

We complete the proof in the same arguments as in Proposition 3.1.

PROPOSITION 3.2. Any nilpotent group G is an A -group with $\kappa(F) = \#F$ and

$\lambda = 2q-1$, where q is the index of G and F is a finite set.

PROOF. Let $G = A_0 \supset A_1 \supset \dots \supset A_{q-1} \supset A_q = \{e\}$ be the normal series of a nilpotent group G of index q . We note that A_{i-1}/A_i is the center of G/A_i ($1 < i < q-1$) and we denote by π_i the canonical map $G \rightarrow G/A_i$.

It is obvious that for any $F \subset G$ and $r \in \mathbb{N}$

$$\#(F^r) < \#(\pi_1(F^r)) \max \{ \# \pi^{-1}(\tilde{g}) \cap F^r : \tilde{g} \in G/A_1 \}. \quad (3.8)$$

We denote by $|F^r|_1$ the RHS of (3.8).

We shall show that there are q positive integers $m_1, m_2, \dots, m_q$ such that

$$\#(F) = m_1 + m_2 + \dots + m_q \text{ and}$$

$$|F^r|_1 \leq \binom{m_1+r-1}{r} \binom{m_2+r-1}{r} \dots \binom{m_{q-1}+r-1}{r} \binom{m_q+r-1}{r} \quad (3.9)$$

If $q=i$, i.e. G is abelian. Corollary (3.1) implies (3.9). We suppose that (3.9) is true for each nilpotent group of index $q = p - 1$.

Let G be nilpotent of index p . Since G/A_1 is abelian by Lemma 3.1 if $\#F < \# \pi_1(F)$.

$$\text{We have, } |F^r|_1 \leq \binom{\#F+r-1}{r}.$$

Let $\#(\pi_1(F)) = m_1 < \#(F)$, then F can be written as

$$F = \{x_i \alpha_j : 1 \leq i \leq m_1, 1 \leq j \leq \#(F)-m_1\}$$

where $\pi_1(x_i) \neq \pi_1(x_j)$ $i \neq j$, $1 \leq j \leq m_1$, and all α_j 's are in A_1 . By (3.8) we obtain,

$$|F^r|_1 \leq \binom{m_1+r-1}{r} \#J_1 \quad (3.10)$$

where $J_1 = {}^{-1}_1(g) F^r$, for some $g \in G/A_1$. Any element of J_1 can be written as

$$x_{i_1} \alpha_{j_1} x_{i_2} \alpha_{j_2} \dots x_{i_r} \alpha_{j_r} \quad (3.11)$$

where $(x_{i_1} x_{i_2} \dots x_{i_r}) \in \tilde{g}$, each i_t ($1 \leq t \leq r$) is one of $1, 2, \dots, m_1$ and each j_t is one of $1, 2, \dots, \#F-m_1$.

By Lemma (3.1) the cardinality of all $x_{i_1} \dots x_{i_r}$ in $\tilde{g}$ is $< \binom{m_1+r-1}{r}$ and so

$$\#J'_1 \leq \binom{m_1+r-1}{r} \#J'_1$$

where

$$J'_1 = x_{i_1} F_1 x_{i_2} F_1 \dots x_{i_r} F_1$$

$x_{i_1} x_{i_2} \dots x_{i_r}$ is fixed suitably choosen from (3.11) and belongs to $\tilde{g}$;

$$F_1 = \{\alpha_j : 1 \leq j \leq \#F-m_1\}$$

Note that if $q = 2$, then A_1 is the center of G , all α_j 's commute with x_i 's and by Corollary (3.1), $|F^r|_1 \leq \binom{\#F-m_1+r-1}{r}$; thus by (3.10) and (3.12), (3.9) follows.

As in (3.8), for some $\tilde{g} \in G/A_2$ we have

$$\#J'_1 \leq \# \pi_2(J'_1) \# \{\pi_2^{-1}(\tilde{g}) \cap J'_1\} \quad (3.13)$$

Since A_1/A_2 is the center of G/A_2 and $F_1 \subset A_1$, by (3.10), (3.12) and (3.13) we obtain

$$|F^r|_1 \leq \binom{m_1+r-1}{r} \cdot |F^r|_1 \quad (3.14)$$

where $|F_1^r|_1$ is defined as $|F^r|_1$.

Since A_1 is a nilpotent of index $p-1$ we apply our inductive hypothesis in (3.14) and for $q = p$ we obtain (3.9).

Now if we replace $m_1, \dots, m_q$ in (3.9) by $\#F$ we see that G is an A group with constants $k = 1$ and $\lambda = 2q-1$.

PROPOSITION 3.3. The class of A -groups is closed under extensions by finite groups.

PROOF. We may write $G/A = \{d_1A, d_2A, \dots, d_sA\}$ where $d_1, d_2, \dots, d_s$ are s representatives of all the different classes of G/A ; without loss of generality let $d_s = e$ the unit of G . We may also write

$$d_i d_j = \alpha(i, j) d(i, j) \quad (1 \leq i, j \leq s)$$

where each $d(i, j)$ is one of $d_1, \dots, d_s$ and each $\alpha(i, j)$ is in A .

Let $F = \{d_{i_1} x_1, \dots, d_{i_m} x_m\}$, $\#F = m$, each d_{i_t} is one of $d_1, \dots, d_s$ and $x_t \in A$ ($1 \leq t \leq m$).

Let $\langle x_{i_1} \rangle = \{d_j^{-1} x_{i_1} d_j : j = 1, 2, \dots, s\}$ ($1 \leq i \leq m$) and

$$d_{j_1} x_{i_1} d_{j_2} x_{i_2} \dots d_{j_r} x_{i_r} \quad (3.15)$$

be a typical r -word in the elements of F . Each word in (3.15) is in the set

$\langle x_{i_1} \rangle \alpha(j_1, j_2) d(j_1, j_2) x_{i_2} \dots d_{j_r} x_{i_r}$ or in

$\langle x_{i_1} \rangle \alpha(j_1, j_2) \langle x_{i_2} \rangle \alpha(\cdot, j_3) \dots d_{j_r} x_{i_r}$ or finally in

$$\langle x_{i_1} \rangle \langle x_{i_2} \rangle \dots \langle x_{i_r} \rangle a.d. \quad (3.16)$$

where $a \in A$ and $d \in \{d_1, d_2, \dots, d_s\}$.

It is clear that the cardinality of the r -words in the elements of F , as in (3.15) is less than the cardinality of the words in (3.16) in the elements of $\{\langle x_{i_1} \rangle, \dots, \langle x_{i_m} \rangle\}$, which is a subset of the A -group, A . Thus G inherits the growth of A .

REFERENCES

1. PALMER, T. Classes of Non-Abelian, Non-Compact, Locally Compact Groups, Rocky Moun. J. Math. 8(4) (1978), 683-741.
2. FOUNTAIN, J.B., RUMSAY, R.W. and WILLIAMSON, J.H. "Functions of Measures on Compact Groups, Proc. Roy. Irish Acad. Sect. A., 76 (1976), 235-251.
3. KARANIKAS, C. and WILLIAMSON, J.H. Norms and Spectral for Certain Subalgebras of $M(G)$, Math. Proc. Camb. Phil. Soc. 95 (1984), 109-122.
4. KARANIKAS, C. Wiener Pairs of Measure Algebras, Pac. Jour. Math. 139(1), (1989), 79-86.