INTEGRAL REPRESENTATIONS OF GENERALIZED LAURICELLA HYPERGEOMETRIC FUNCTIONS

VU KIM TUAN
Institute of Mathematics
P.O. Box 631, B5 H5
Hanoi, Vietnam

AND

R.G. BUSCHMAN
Department of Mathematics
Box 3036, University Station
Laramie, WY 82071, USA

(Received August 10, 1990 and in revised form March 24, 1992)

ABSTRACT. The generalized hypergeometric function was introduced by Srivastava and Daoust. In the present paper a new integral representation is derived. Similarly new integral representations of Lauricella and Appell function are obtained.

KEY WORDS. Lauricella, Appell, integral representations, Mellin transformation.

1991 AMS SUBJECT CLASSIFICATION CODES. 33A35, 44A30

1. INTRODUCTION.

The generalized Lauricella function of several complex variables was introduced by Srivastava and Daoust [7,8]. The only integral representation which seems to be known for this function is in terms of a Mellin-Barnes integral [6,8]. From a fundamental result about Mellin transforms in n -dimensions, we obtain a new representation for the generalized Lauricella function under suitable restrictions on the parameters. In a similar manner new integral representations are obtained for the Lauricella functions $F_A^{(n)}$, $F_C^{(n)}$ and $F_D^{(n)}$, and consequently for the Appell functions F_1 , F_2 , and F_4 . From these derivations it is clear that the method does not provide representations for $F_B^{(n)}$ and F_3 .

2. THE FUNDAMENTAL THEOREM.

Although the following theorem is quite simple, nevertheless it has basic importance.

THEOREM. Let $f(x)x^{s-1} \in L(0, \infty)$ and

$$M_1[f(x)](s) = \int_0^\infty f(x)x^{s-1}dx = f^*(s). \quad (1)$$

If $\text{Res}_j > 0$, $j = 1, 2, \dots, n$ and $\text{Re}\left(\sum_{j=1}^n s_j\right) = \text{Res}$, then

$$\begin{aligned} M_n[f(\max\{x_1, \dots, x_n\})](s_1, \dots, s_n) &= \int_0^\infty \dots \int_0^\infty f(\max\{x_1, \dots, x_n\}) \prod_{i=1}^n x_i^{s_i-1} dx_1 \dots dx_n = \\ &= \frac{s_1 + \dots + s_n}{s_1 \dots s_n} f^*(s_1 + \dots + s_n). \end{aligned} \quad (2)$$

Here M_n represents the n -dimensional Mellin integral transformation [2].

PROOF. In fact we have

$$\begin{aligned} \int_0^\infty \dots \int_0^\infty f(\max\{x_1, \dots, x_n\}) \prod_{i=1}^n x_i^{s_i-1} dx_1 \dots dx_n &= \\ = \sum_{i=1}^n \int_0^\infty f(x_i) \int_0^{x_i} \dots \int_0^{x_i} \prod_{j=1}^n x_j^{s_j-1} dx_1 \dots dx_n &= \end{aligned}$$

$$= \sum_{i=1}^n \frac{s_i}{s_1 s_2 \cdots s_n} \int_0^\infty f(x_i) x_i^{s_1 + \cdots + s_n - 1} dx_i = \frac{s_1 + \cdots + s_n}{s_1 \cdots s_n} f^*(s_1 + \cdots + s_n).$$

Thus the formula (2) is proved.

The only related results which we have found in the literature are those for the two-dimensional Laplace transformation in the tables of Voelker and Doetsch [9; p. 165, (30), (32)] and in the work of Černov [4; p. 145]. In an analogous manner to our theorem, those results easily can be derived and extended to higher dimensions.

3. THE INTEGRAL REPRESENTATION OF THE GENERALIZED LAURICELLA FUNCTION.

The generalized hypergeometric function of Srivastava and Daoust [6] is defined by

$$\begin{aligned} & F_{q, q_1, \dots, q_n}^{p, p_1, \dots, p_n} \left(\begin{matrix} a_p; c_{p_1}^1; \dots; c_{p_n}^n \\ b_q; d_{q_1}^1; \dots; d_{q_n}^n \end{matrix}; x_1, \dots, x_n \right) = \\ &= \sum_{m_1, \dots, m_n=0}^{\infty} \frac{\prod_{j=1}^p (a_j)_{m_1 + \dots + m_n}}{\prod_{j=1}^q (b_j)_{m_1 + \dots + m_n}} \prod_{i=1}^n \left\{ \frac{\prod_{j=1}^{p_i} (c_j^i)_{m_i}}{\prod_{j=1}^{q_i} (d_j^i)_{m_i}} \frac{x_i^{m_i}}{m_i!} \right\} \end{aligned} \quad (3)$$

where for absolute convergence it is sufficient that

$$1 + q + q_k - p - p_k \geq 0; \quad k = 1, 2, \dots, n.$$

The function in (3) is a special case of the H -functions of several variables which were defined in [3, 6]. The fundamental theorem leads to the following result which involves the G -function of Meijer [8]. Let

$$f(x) = G_{q, p}^p \left(x \mid \begin{matrix} b_1, \dots, b_q \\ a_1, \dots, a_p \end{matrix} \right) = \frac{1}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{\prod_{j=1}^p \Gamma(a_j + s)}{\prod_{j=1}^q \Gamma(b_j + s)} x^{-s} ds$$

where $\operatorname{Re} a_j > 0$, $j = 1, 2, \dots, p$; and either $p = q$ and $\operatorname{Re} \sum_{j=1}^p a_j < \operatorname{Re} \sum_{j=1}^q b_j$ or $q < p$. It is known [5] that

$$f^*(s) = \frac{\prod_{j=1}^p \Gamma(a_j + s)}{\prod_{j=1}^q \Gamma(b_j + s)}, \quad \operatorname{Re} s > 0.$$

Therefore (2) becomes

$$\begin{aligned} & \int_0^\infty \cdots \int_0^\infty G_{q, p}^p \left(\max\{x_1, \dots, x_n\} \mid \begin{matrix} b_1, \dots, b_q \\ a_1, \dots, a_p \end{matrix} \right) \prod_{i=1}^n x_i^{s_i-1} dx_1 \cdots dx_n = \\ &= \frac{\prod_{j=1}^p \Gamma(a_j + s_1 + \cdots + s_n)}{\prod_{j=1}^q \Gamma(b_j + s_1 + \cdots + s_n)} \frac{(s_1 + \cdots + s_n)}{s_1 \cdots s_n}, \quad \operatorname{Re} s_1, \dots, \operatorname{Re} s_n > 0. \end{aligned} \quad (4)$$

The representation for the generalized Lauricella function can now be obtained as follows. Let $p_i \leq q_i$, $i = 1, 2, \dots, n$, and either $p = q$ and $\operatorname{Re} \sum_{j=1}^p a_j < \operatorname{Re} \sum_{j=1}^q b_j$ or $p > q$. Then

$$\begin{aligned} & F_{q, q_1, \dots, q_n}^{p, p_1, \dots, p_n} \left(\begin{matrix} a_p; c_{p_1}^1; \dots; c_{p_n}^n \\ b_q; d_{q_1}^1; \dots; d_{q_n}^n \end{matrix}; x_1, \dots, x_n \right) \\ &= \frac{1}{n} \sum_{m_1, \dots, m_n=0}^{\infty} \frac{\prod_{j=1}^p (a_j)_{m_1 + \dots + m_n} (n)_{m_1 + \dots + m_n} (n + m_1 + \cdots + m_n)}{\prod_{j=1}^q (b_j)_{m_1 + \dots + m_n} (n + 1)_{m_1 + \dots + m_n} (m_1 + 1) \cdots (m_n + 1)} \\ & \quad \cdot \prod_{i=1}^n \left\{ \frac{\prod_{j=1}^{p_i} (c_j^i)_{m_i} (2)_{m_i} x_i^{m_i}}{\prod_{j=1}^{q_i} (d_j^i)_{m_i} (1)_{m_i} m_i!} \right\} \\ &= \frac{1}{n} \frac{\Gamma(b_1) \cdots \Gamma(b_q)}{\Gamma(a_1) \cdots \Gamma(a_p)} \int_0^\infty \cdots \int_0^\infty G_{q+1, p+1}^{p+1, 0} \left(\max\{t_1, \dots, t_n\} \mid \begin{matrix} b_1 - n, \dots, b_q - n, 1 \\ a_1 - n, \dots, a_p - n, 0 \end{matrix} \right) \\ & \quad \cdot \prod_{i=1}^n \left\{ \sum_{m_i=0}^{\infty} \frac{\prod_{j=1}^{p_i} (c_j^i)_{m_i} (2)_{m_i} (x_i t_i)^{m_i}}{\prod_{j=1}^{q_i} (d_j^i)_{m_i} (1)_{m_i} m_i!} \right\} dt_1 \cdots dt_n. \end{aligned}$$

Thus we have proved

$$\begin{aligned} & F_{q, q_1, \dots, q_n}^{p, p_1, \dots, p_n} \left(\begin{matrix} a_p; c_1^1, \dots, c_n^n \\ b_q; d_1^1, \dots, d_n^n \end{matrix}; x_1, \dots, x_n \right) = \\ &= \frac{1}{n} \frac{\Gamma(b_1) \dots \Gamma(b_q)}{\Gamma(a_1) \dots \Gamma(a_p)} \cdot n t_0^\infty \dots \int_0^\infty G_{q+1, p+1}^{p+1, 0} \left(\max \{t_1, \dots, t_n\} \right)_{\substack{b_1-n, \dots, b_q-n, 1 \\ a_1-p, \dots, a_p-n, 0}} \cdot \\ & \cdot \prod_{i=1}^n F_{q_i+1}^{p_i+1} \left(\begin{matrix} c_1^1, \dots, c_{p_i}^{p_i}, 2; \\ d_1^1, \dots, d_{q_i}^{q_i}, 1; \end{matrix} x_i t_i \right) dt_1 \dots dt_n. \end{aligned} \quad (5)$$

Formula (5) is valid when $\operatorname{Re} a_j > n$; $j = 1, 2, \dots, p$; $p_i \leq q_i$, $i = 1, 2, \dots, n$; $p \geq q$, $1 + q + q_i - p - p_i \geq 0$; $i = 1, 2, \dots, n$.
(The restriction $\operatorname{Re} \sum_{j=1}^q b_j + 1 > \operatorname{Re} \sum_{j=1}^p a_j$ is needed when $p = q$.)

4. INTEGRAL REPRESENTATIONS OF LAURICELLA HYPERGEOMETRIC FUNCTIONS.

Derivations similar to those in the previous section lead to representations for 3 of the Lauricella functions.
(Since in (5) $p \geq q$ we do not get $F_B^{(n)}$.) We introduce the operator

$$\mathcal{D}_i = \left(\frac{a-1}{n} + x_i \frac{d}{d(x_i)} \right).$$

(a) Using the formula (2) we get

$$\begin{aligned} & \int_0^\infty \dots \int_0^\infty \exp(-\max \{x_1, \dots, x_n\}) \prod_{i=1}^n x_i^{\frac{a-1}{n} + m_i - 1} dx_1 \dots dx_n = \\ &= \frac{\Gamma(a + m_1 + \dots + m_n)}{\left(\frac{a-1}{n} + m_1\right) \dots \left(\frac{a-1}{n} + m_n\right)}, \quad \operatorname{Re} a > 1. \end{aligned} \quad (6)$$

Consequently, for the Lauricella function $F_A^{(n)}$ we have

$$\begin{aligned} & F_A^{(n)}(a; b_1, \dots, b_n; c_1, \dots, c_n; x_1, \dots, x_n) = \\ &= \sum_{m_1, \dots, m_n=0}^\infty \frac{(a)_{m_1+\dots+m_n} (b_1)_{m_1} \dots (b_n)_{m_n}}{(c_1)_{m_1} \dots (c_n)_{m_n} m_1! \dots m_n!} x_1^{m_1} \dots x_n^{m_n} = \\ &= \sum_{m_1, \dots, m_n=0}^\infty \frac{1}{\Gamma(a)} \int_0^\infty \dots \int_0^\infty \prod_{i=1}^n t_i^{\frac{a-1}{n} + m_i - 1} \exp(-\max \{t_1, \dots, t_n\}) \cdot \\ & \cdot \prod_{i=1}^n \frac{\left(\frac{a-1}{n} + m_i\right) (b_i)_{m_i} x_i^{m_i}}{(c_i)_{m_i} m_i!} dt_1 \dots dt_n = \\ &= \frac{1}{\Gamma(a)} \int_0^\infty \dots \int_0^\infty (t_1 \dots t_n)^{\frac{a-1}{n} - 1} \exp(-\max \{t_1, \dots, t_n\}) \cdot \\ & \cdot \prod_{i=1}^n \left\{ \sum_{m_i=0}^\infty \left(\frac{a-1}{n} + m_i \right) \frac{(b_i)_{m_i} (t_i x_i)^{m_i}}{(c_i)_{m_i} m_i!} \right\} dt_1 \dots dt_n, \end{aligned}$$

Consequently

$$\begin{aligned} & F_A^{(n)}(a; b_1, \dots, b_n; c_1, \dots, c_n; x_1, \dots, x_n) = \\ &= \frac{1}{\Gamma(a)} \prod_{i=1}^n \mathcal{D}_i \int_0^\infty \dots \int_0^\infty \prod_{i=1}^n t_i^{\frac{a-1}{n} - 1} \exp(-\max \{t_1, \dots, t_n\}) \cdot \prod_{i=1}^n {}_1F_1(b_i, c_i; x_i t_i) dt_1 \dots dt_n, \\ & \operatorname{Re} a > 1. \end{aligned} \quad (7)$$

(b) From (2) we have

$$\begin{aligned} & \int_0^1 \dots \int_0^1 (1 - \max \{x_1, \dots, x_n\})^{b-a} \prod_{i=1}^n x_i^{\frac{a-1}{n} + m_i - 1} dx_1 \dots dx_n = \\ &= \frac{\Gamma(a + m_1 + \dots + m_n) \Gamma(b - a + 1)}{\Gamma(b + m_1 + \dots + m_n) \left(\frac{a-1}{n} + m_1\right) \dots \left(\frac{a-1}{n} + m_n\right)}, \quad 1 + \operatorname{Re} b > \operatorname{Re} a > 1. \end{aligned} \quad (8)$$

Therefore for the Lauricella function $F_D^{(n)}$ we get

$$\begin{aligned} F_D^{(n)}(a; b_1, \dots, b_n; c; x_1, \dots, x_n) &= \\ &= \sum_{m_1, \dots, m_n=0}^{\infty} \frac{(a)_{m_1+\dots+m_n} (b_1)_{m_1} (b_n)_{m_n} x_1^{m_1} \dots x_n^{m_n}}{(c)_{m_1+\dots+m_n} m_1! \dots m_n!} = \\ &= \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a+1)} \int_0^1 \dots \int_0^1 \prod_{i=1}^n t_i^{\frac{a-1}{n}-1} (1 - \max\{t_1, \dots, t_n\})^{c-a} \\ &\quad \prod_{i=1}^n \sum_{m_i=0}^{\infty} \left(\frac{a-1}{n} + m_i \right) \cdot \frac{(b_i)_{m_i} (x_i t_i)^{m_i}}{m_i!} dt_1 \dots dt_n . \end{aligned}$$

Hence

$$\begin{aligned} &F_D^{(n)}(a; b_1, \dots, b_n; c; x_1, \dots, x_n) \\ &= \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a+1)} \prod_{i=1}^n \mathcal{D}_i \int_0^1 \dots \int_0^1 \prod_{i=1}^n t_i^{\frac{a-1}{n}-1} (1 - \max\{t_1, \dots, t_n\})^{c-a} \prod_{i=1}^n (1 - x_i t_i)^{-b_i} dt_1 \dots dt_n, \end{aligned} \quad (9)$$

$$1 + \operatorname{Re} c > \operatorname{Re} a > 1 .$$

(c) Using the formula (2) again, we have

$$\begin{aligned} &\int_0^\infty \dots \int_0^\infty (\max\{x_1, \dots, x_n\})^{\frac{b-a+1}{2}} K_{b-a+1} \left(2(\max\{t_1 \dots t_n\})^{\frac{1}{2}} \right) \prod_{i=1}^n (x_i)^{\frac{a-1}{n}+m_i-1} dx_1 \dots dx_n = \\ &= \frac{1}{2} \frac{\Gamma(a+m_1+\dots+m_n) \Gamma(b+m_1+\dots+m_n)}{\left(\frac{a-1}{n} + m_1 \right) \dots \left(\frac{a-1}{n} + m_n \right)} , \quad \operatorname{Re} a > 1, \quad \operatorname{Re} b > -0 . \end{aligned} \quad (10)$$

Consequently, for the Lauricella hypergeometric function $F_C^{(n)}$, by a similar development we have the result

$$\begin{aligned} &F_C^{(n)}(a; b; c_1, \dots, c_n; x_1, \dots, x_n) = \\ &= \frac{2}{\Gamma(a)\Gamma(b)} \prod_{i=1}^n \mathcal{D}_i \int_0^\infty \dots \int_0^\infty \prod_{i=1}^n t_i^{\frac{a-1}{n}-1} (\max\{t_1, \dots, t_n\})^{\frac{b-a+1}{2}} K_{b-a+1} \left(2(\max\{t_1 \dots t_n\})^{\frac{1}{2}} \right) \\ &\quad \prod_{i=1}^n {}_0F_1(c_i; x_i t_i) dt_1 \dots dt_n, \\ &\quad \operatorname{Re} a > 1, \quad \operatorname{Re} b > -1 . \end{aligned} \quad (11)$$

For Appell functions these read

$$\begin{aligned} &F_1(a; b_1, b_2; c; x_1, x_2) = \frac{\Gamma(c)}{\Gamma(a)\Gamma(c-a+1)} \\ &\mathcal{D}_1 \mathcal{D}_2 \int_0^1 \int_0^1 (1 - \max\{t_1, t_2\})^{c-a} (t_1 t_2)^{(a-3)/2} \left\{ (1 - x_1 t_1)^{-b_1} \right\} \left\{ (1 - x_2 t_2)^{-b_2} \right\} dt_1 dt_2 , \\ &\quad 1 + \operatorname{Re} c > \operatorname{Re} a > 1 ; \end{aligned} \quad (12)$$

$$\begin{aligned} &F_2(a; b_1, b_2; c_1, c_2; x_1, x_2) = \\ &= \frac{1}{\Gamma(a)} \mathcal{D}_1 \mathcal{D}_2 \int_0^\infty \int_0^\infty (t_1 t_2)^{(a-3)/2} \exp(-\max\{t_1, t_2\}) \\ &\quad \{ {}_1F_1(b_1; c_1; x_1 t_1) \} \{ {}_1F_1(b_2; c_2; x_2 t_2) \} dt_1 dt_2 , \\ &\quad \operatorname{Re} a > 1 ; \end{aligned} \quad (13)$$

$$\begin{aligned}
 &F_4(a; b; c_1, c_2; x_1, x_2) = \\
 &= \frac{2\Gamma(c_1)\Gamma(c_2)}{\Gamma(a)\Gamma(b)} \mathcal{D}_1 \mathcal{D}_2 \int_0^\infty \int_0^\infty (\max\{t_1, t_2\})^{(b-a+1)/2} (t_1 t_2)^{(a-3)/2} K_{b-a+1} \left(2(\max\{t_1, t_2\})^{1/2} \right) \\
 &\quad \cdot \left\{ (x_1 t_1)^{(1-c_1)/2} I_{c_1-1} \left(2(x_1 t_1)^{1/2} \right) \right\} \left\{ (x_2 t_2)^{(1-c_2)/2} I_{c_2-1} \left(2(x_2 t_2)^{1/2} \right) \right\} dt_1 dt_2, \\
 &\quad \operatorname{Re} a > 1, \quad \operatorname{Re} b > 0.
 \end{aligned} \tag{14}$$

REFERENCES

1. P. APPELL AND J. KAMPÉ DE FÉRIET, *Fonctions hypergéométriques et hypersphériques, Polynômes d'Hermite*, Gauthier-Villars, Paris, 1926.
2. YU. A. BRYCHKOV, H.J. GLAESKE, A.P. PRUDNIKOV, AND VU KIM TUAN, *Multidimensional Integral Transforms*, Geest & Portig K.G., Leipzig and D. Reidel Publishing Company, Amsterdam, 1990.
3. R.G. BUSCHMAN, *The H-functions of N variables*, Ranchi Univ. Math. J. **10** (1979), 81-88.
4. V.M. ČERNOV, *Some questions of the operational calculus concerning the two-sided, two-dimensional Laplace-Carson transform*, Izv. Vysš. Učebn. Zaved. Mat. **2** (33) (1963), 140-151.
5. O.I. MARICHEV, *Handbook of Integral Transforms of Higher Transcendental Functions, Theory and Algorithmic Tables*, Ellis Horwood, Chichester, 1982.
6. O.I. MARICHEV AND VU KIM TUAN, *The problems of definitions and symbols of G- and H-functions of several variables*, Rev. Tecn. Fac. Ing. Univ. Zulia, Special Edition **6** (1983), 144-151.
7. H.M. SRIVASTAVA AND M.C. DAOUST, *Certain generalized Neumann expansions associated with the Kampé de Fériet function*, Nederl. Akad. Wetensch. Proc. A72 (1969), 449-457.
8. H.M. SRIVASTAVA, K.C. GUPTA, AND S.P. GOYAL, *The H-Functions of One and Two Variables with Applications*, South Asian Publishers Ltd., New Delhi, 1982.
9. D. VOELKER UND G. DOETSCH, *Die Zweidimensionale Laplace Transformation*, Verlag Birkhäuser, Basel, 1950.