MEASURE-THEORETIC CHARACTERIZATIONS OF HEREDITARILY-NORMAL SPACES

JOSEPH HERTZLINGER

Polytechnic University

(Received March 16, 1991 and in revised form March 30, 1992)

ABSTRACT. In this paper we characterize hereditarily-normal spaces in terms of the measure-theoretic properties of the lattice of closed sets. We then generalize from that lattice to other lattices. We apply the results to extremally-disconnected spaces.

KEY WORDS AND PHRASES. Hereditarily-normal topological spaces, two-valued measures, lattice of closed sets, extremally-disconnected topological space.

1991 AMS SUBJECT CLASSIFICATION CODES. 54D15, 54G05.

1. INTRODUCTION

We can devise two measure-theoretic characterizations of hereditarily-normal spaces. First, a space is hereditarily-normal if and only if every pre-measure is a measure. Second, a space is hereditarily-normal if and only if the set of measures greater than a given measure form a chain. We can then generalize these results to other lattices (the lattices of open sets, zero sets, cozero sets, etc.). When we apply the results to the lattice of open sets we can derive a few conclusions about extremally-disconnected spaces.

We begin by giving a few definitions (we note that our lattice methods and terminology are consistent with standard usage, e.g., Camacho [1], Eid [4], Szeto [8]):

DEFINITION 1. We can define a partial order on the set of two-valued nontrivial measures on a space X as follows: $\mu_1 \leq \mu_2$ if and only if $\mu_1(F) \leq \mu_2(F)$ for all closed sets F . We define $\geq$ in a similar manner. (We note that here we are referring to finitely-additive two-valued measures on the algebra generated by the lattice of closed sets.)

DEFINITION 2. A two-valued pre-measure Π on a space X is a set function from the collection of closed sets to $\{0, 1\}$ such that for any pair of closed sets F_1 and F_2 :

$$\Pi(\emptyset) = 0, \quad (1.1)$$

$$\text{if } F_1 \subseteq F_2 \text{ then } \Pi(F_1) \leq \Pi(F_2), \quad (1.2)$$

$$\text{if } \Pi(F_1) = 1 \text{ and } \Pi(F_2) = 1 \text{ then } \Pi(F_1 \cap F_2) = 1, \quad (1.3)$$

$$\text{if } F_1 \cup F_2 = X \text{ then either } \Pi(F_1) = 1 \text{ or } \Pi(F_2) = 1. \quad (1.4)$$

The last condition can be compared with the corresponding condition for measures:

$$\text{if } \Pi(F_1 \cup F_2) = 1 \text{ then either } \Pi(F_1) = 1 \text{ or } \Pi(F_2) = 1. \quad (1.5)$$

DEFINITION 3. A space X has the pre-measure property if and only if every pre-measure on X is a measure.

DEFINITION 4. A space X is hereditarily normal if and only if every subset Y of X is normal.

We can replace this with the equivalent but more useful definition:

DEFINITION 4A. A space is hereditarily normal if and only if for every subset Y of X and every pair of closed sets F_1 and F_2 the condition $F_1 \cap F_2 \cap Y = \emptyset$ implies that there exist closed sets F_3 and F_4 such that $F_3 \cap F_1 \cap Y = F_4 \cap F_2 \cap Y = \emptyset$ and $Y \subseteq F_3 \cup F_4$.

DEFINITION 5. A regular measure μ is a measure on the algebra generated by the lattice of closed sets such that $\mu(M)$ is the supremum of the measures of all closed sets inside M . A regular measure is also a maximal measure (i.e., there is no measure strictly greater than it). For a two-valued measure the sets of measure one form a closed ultrafilter.

DEFINITION 6. A space X has the measure-tree property if and only if for all two-valued measures μ, μ_1 , and μ_2 on the algebra generated by the lattice of closed sets the conditions $\mu \leq \mu_1$ and $\mu \leq \mu_2$ imply that $\mu_2 \leq \mu_1$ or $\mu_1 \leq \mu_2$. We note that the partial order $\leq$ is the same as that defined in Definition 1.

The term "measure-tree property" requires explaining. In a normal space the set of measures less than a given measure (see Theorem 1) form a partially ordered set in which measures branch outward from the regular measure. If the space does not have the measure-tree property those measures also branch inward. If the space has the measure-tree property the measures form a tree in which the branches do not reunite. This is equivalent to saying that all measures greater than a given measure form a chain.

We also note that this property closely resembles an axiom suggested by the noted mathematician Lewis Carroll [2] (*Sylvie and Bruno*, Chapter 18, p. 425): Things greater than the same are greater than one another.

2. MAIN RESULTS

The following (see Eid [4] and Frolik [5]) is a measure-theoretic characterization of normal spaces:

THEOREM 2.1. A space X is normal if and only if for every measure μ and every pair of regular measures μ_1 and μ_2 , if $\mu_1 \geq \mu$ and $\mu_2 \geq \mu$ then $\mu_1 = \mu_2$.

PROOF. Let us assume X is normal. Let us consider measure μ and regular measures μ_1 and μ_2 . Let $\mu_1 \geq \mu$ and $\mu_2 \geq \mu$. Let $\mu_1 \neq \mu_2$ and let us derive a contradiction. If $\mu_1 \neq \mu_2$ then there is a set M belonging to the algebra generated by the topology such that $\mu_1(M) = 1, \mu_2(M) = 0, \mu_1(M') = 0$, and $\mu_2(M') = 1$. Since μ_1 and μ_2 are regular measures there exist closed sets F_1 and F_2 such that $\mu_1(F_1) = \mu_2(F_2) = 1, \mu_1(F_2) = \mu_2(F_1) = 0, F_1 \subseteq M, F_2 \subseteq M'$, and $F_1 \cap F_2 = \emptyset$. We therefore have $\mu(F_1) \leq \mu_2(F_1) = 0$ and $\mu(F_2) \leq \mu_1(F_2) = 0$. Since X is normal there exist open sets G_1 and G_2 such that $G_1 \subseteq F_1, G_2 \subseteq F_2$, and $G_1 \cap G_2 = \emptyset$. Let their complements be $F_3 = G_1'$ and $F_4 = G_2'$. Since $\mu_1(G_1) \geq \mu_1(F_1) = 1$ and $\mu_2(G_2) \geq \mu_2(F_2) = 1$ we have $\mu_1(F_3) = \mu_2(F_4) = 0$. Therefore $\mu(F_1) = \mu(F_2) = 0$. Let $F_5 = F_1 \cup F_3$ and $F_6 = F_2 \cup F_4$. Since $\mu(F_5) = \mu(F_1 \cup F_3) = 0, \mu(F_5') = 1$. Similarly $\mu(F_6') = 1$. Since $F_3 \subseteq F_5$ and $F_4 \subseteq F_6$ we have $F_5' \subseteq F_3'$ and $F_6' \subseteq F_4'$. Since $F_3' \cap F_4' = \emptyset$ we have $F_5' \cap F_6' = \emptyset$. This means that $\mu(F_5' \cup F_6') = \mu(F_5') + \mu(F_6') = 2$, which is impossible. The necessary contradiction has been achieved.

For the other direction let X be not normal and let us construct a measure μ and regular measures μ_1 and μ_2 such that $\mu_1 \neq \mu_2$ but $\mu_1 \geq \mu$ and $\mu_2 \geq \mu$. Since X is not normal there are two disjoint closed sets F_1 and F_2 which cannot be separated by open sets. The collection of sets $\{G \text{ is open} \mid F_1 \subseteq G \text{ or } F_2 \subseteq G\}$ therefore has the finite intersection property. The open filter that it generates can be extended to an open ultrafilter which induces a measure μ such that $\mu(G) = 1$ for G in the collection. Let F_0 be an arbitrary closed set with measure one; $\mu(F_0') = 0$ which means F_0' is not in the collection. This means that $F_0' \not\subseteq F_1$ and $F_0' \not\subseteq F_2$ or — in other words — $F_1 \cap F_0' \neq \emptyset$ and $F_2 \cap F_0' \neq \emptyset$. This means that the collection of sets $\{F \text{ is closed} \mid \mu(F) = 1\} \cup \{F_1\}$ has the finite intersection property. The filter that it generates can be extended to a closed ultrafilter. The regular measure based on it will be called μ_1 ; $\mu \leq \mu_1$ and $\mu_1(F_1) = 1$. Using similar reasoning we can see there is a regular measure μ_2 such that $\mu \leq \mu_2$ and $\mu_2(F_2) = 1$. Since $F_1 \cap F_2 = \emptyset, \mu_1 \neq \mu_2$ which means the required measures have been constructed. ■

THEOREM 2.2. If a space X has the pre-measure property then it has the measure-tree property.

PROOF. Let X have the pre-measure property and let μ, μ_1 , and μ_2 be measures on X such that:

$$\mu \leq \mu_1 \text{ and } \mu \leq \mu_2. \quad (2.1)$$

Let $\Pi = \min(\mu_1, \mu_2)$. It is easy to see that $\mu \leq \Pi$.

The function Π is also a pre-measure. Conditions (1.1)-(1.3) in Definition 2 are obviously fulfilled. To prove condition (1.4), we let $F_1 \cup F_2 = X$. Because $\mu(X) = 1$ we also have $\mu(F_1 \cup F_2) = 1$. The function μ is also a measure and therefore $\mu(F_1) = 1$ or $\mu(F_2) = 1$. Since $\mu \leq \Pi$, $\Pi(F_1) = 1$ or $\Pi(F_2) = 1$. Therefore condition (1.4) is fulfilled. The function Π is thus a pre-measure. Since X has the pre-measure property, Π is a measure. This is possible only if either

$$\mu_2 \leq \mu_1 \text{ or } \mu_1 \leq \mu_2. \quad (2.2)$$

(If this weren't the case then there would be closed sets F_1 and F_2 such that $\mu_1(F_1) = \mu_2(F_2) = 1$ and $\mu_1(F_2) = \mu_2(F_1) = 0$. This means that $\mu_1(F_1 \cup F_2) = \mu_2(F_1 \cup F_2) = 1$. Since $\Pi = \min(\mu_1, \mu_2)$, we have $\Pi(F_1 \cup F_2) = 1$ but $\Pi(F_1) = \Pi(F_2) = 0$. This is incompatible with Π being a measure.)

We have been able to derive (2.2) from (2.1). Therefore X has the measure-tree property. ■

THEOREM 2.3. If space X is hereditarily normal then it has the pre-measure property.

PROOF. Let Π be a pre-measure on X . In order to show that it is a measure we let F_1 and F_2 be closed sets such that $\Pi(F_1 \cup F_2) = 1$. Let $Y = X \setminus (F_1 \cap F_2)$. This means that $F_1 \cap F_2 \cap Y = \emptyset$. Since X is hereditarily normal this implies that there exist closed sets F_3 and F_4 such that $F_3 \cap F_1 \cap Y = F_4 \cap F_2 \cap Y = \emptyset$ and $Y \subseteq F_3 \cup F_4$. Let F_5 and F_6 be closed sets such that $F_5 = F_3 \cup (F_1 \cap F_2)$ and $F_6 = F_4 \cup (F_1 \cap F_2)$. $F_5 \cup F_6 = X$ and since Π is a pre-measure $\Pi(F_5) = 1$ or $\Pi(F_6) = 1$. Without loss of generality let $\Pi(F_5) = 1$. Let $F_7 = F_5 \cap (F_1 \cup F_2)$.

Let us now prove $F_2 = F_7$. We can do this point by point.

First, let point $x \in F_2$. Either $x \in F_1$ or $x \in F_1'$. If $x \in F_1$ then $x \in F_1 \cap F_2$ which means $x \in F_5$ and thus $x \in F_7$. If, on the other hand, $x \in F_1'$ then $x \in F_2 \setminus F_1$. Since F_4 is disjoint from $F_2 \setminus F_1$, $x \in F_4'$. Since $x \in Y$ and $F_3 \cup F_4 = Y$, $x \in F_3$. Therefore $x \in F_5$ and thus $x \in F_7$.

Second, let $x \in F_7$. $F_7 = (F_5 \cap F_2) \cup (F_5 \cap F_1)$. Either $x \in F_5 \cap F_2$ or $x \in F_5 \cap F_1$. If $x \in F_5 \cap F_2$ then $x \in F_2$. On the other hand if $x \in F_5 \cap F_1$ then $x \in F_5$. If $x \in F_5$ then either $x \in (F_1 \cap F_2)$ or $x \in F_1 \cap F_2$ or $x \in F_3$. If $x \in F_1 \cap F_2$ then $x \in F_2$. Now examine $x \in F_3$. If $x \in F_3$ then, since F_3 is disjoint from $F_1 \cap Y$, $x \in (F_1 \cap Y)'$. Since $F_3 \subseteq Y$, $x \in Y$. Therefore $x \in F_1'$. On the other hand, since $x \in F_7$, $x \in F_1 \cup F_2$. This means that $x \in F_2$.

We have thus shown that $F_2 = F_7$ and therefore $F_2 = F_5 \cap F_1 \cup F_2$. Since $\Pi(F_5) = 1$ and $\Pi(F_1 \cup F_2) = 1$ therefore $\Pi(F_2) = 1$.

Similarly if $\Pi(F_6) = 1$ then $\Pi(F_1) = 1$. In other words, either $\Pi(F_1) = 1$ or $\Pi(F_2) = 1$.

We have shown that if $\Pi(F_1 \cup F_2) = 1$ then either $\Pi(F_1) = 1$ or $\Pi(F_2) = 1$. Therefore Π is a measure. This applies to all pre-measures which means that X has the pre-measure property. ■

LEMMA 2.1. If μ is a measure on space X , Y is a subspace of X , and Z is a set in the algebra generated by the topology such that $Z \subseteq Y$ and $\mu(Z) = 1$ then the set function μ_Y defined as $\mu_Y(F \cap Y) = \mu(F)$ for all closed sets F is well-defined and μ_Y is a measure on X .

PROOF. Let F be a closed set in Y . Let $\mu_Y(F)$ be evaluated in two different ways:

$$\mu_Y(F) = \mu(F_1) \text{ where } F_1 \text{ is closed and } F_1 \cap Y = F, \quad (2.3)$$

$$\mu_Y(F) = \mu(F_2) \text{ where } F_2 \text{ is closed and } F_1 \cap Y = F. \quad (2.4)$$

Since $Z \subseteq Y$ and $F_1 \cap Y = F = F_2 \cap Y$ we obtain $F_1 \cap Z = F_2 \cap Z$. Since $\mu(Z) = 1$ we also obtain $\mu(F_1 \cap Z) = \mu(F_1)$ and $\mu(F_2 \cap Z) = \mu(F_2)$. This means that $\mu(F_1) = \mu(F_2)$ so that μ_Y is the same no matter how it is evaluated.

Most of the properties which determine if a set function is a measure are inherited by μ_Y . The remaining property is whether $\mu_Y(Y) = 1$. Since $Y = X \cap Y$, $\mu_Y(Y) = \mu(X) = 1$ which implies that μ_Y is a measure. ■

THEOREM 2.4. If a space X has the measure-tree property then it is hereditarily normal.

PROOF. It is easiest to show this using the contrapositive.

Let X be not hereditarily normal. There must exist a subset Y of X such that Y is not normal. We must have a measure μ on Y and a pair of regular measures μ_1 and μ_2 on Y such that $\mu \leq \mu_1$ and $\mu \leq \mu_2$ but $\mu_1 \neq \mu_2$. We must therefore have sets M_1 and M_2 in the algebra generated by the inherited topology on Y such that:

$$\mu_1(M_1) = \mu_2(M_2) = 1 \text{ and } \mu_2(M_1) = \mu_1(M_2) = 0. \quad (2.5)$$

Since these are regular measure we must have closed sets F_1 and F_2 in Y such that:

$$\mu_1(F_1) = \mu_2(F_2) = 1 \text{ and } \mu_2(F_1) = \mu_1(F_2) = 0. \quad (2.6)$$

Let μ_3, μ_4 , and μ_5 be two-valued functions of sets in the algebra generated by the topology on X such that for all M in the algebra:

$$\mu_3(M) = \mu(M \cap Y), \mu_4(M) = \mu_1(M \cap Y), \text{ and } \mu_5(M) = \mu_2(M \cap Y). \quad (2.7)$$

These are obviously measures. They also satisfy the conditions $\mu_3 \leq \mu_4$ and $\mu_3 \leq \mu_5$. If X had the measure-tree property then they would satisfy either $\mu_4 \leq \mu_5$ or $\mu_5 \leq \mu_4$. Since $F_1, F_2 \subseteq Y$ it follows that $F_1 = F_3 \cap Y$ and $F_2 = F_4 \cap Y$ where F_3 and F_4 are closed. We therefore have the following

$$\mu_4(F_3) = \mu_1(F_3 \cap Y) = \mu_1(F_1) = 1, \quad (2.8)$$

$$\mu_5(F_4) = \mu_2(F_4 \cap Y) = \mu_2(F_2) = 1, \quad (2.9)$$

$$\mu_4(F_4) = \mu_1(F_4 \cap Y) = \mu_1(F_2) = 0, \quad (2.10)$$

$$\mu_5(F_3) = \mu_2(F_3 \cap Y) = \mu_2(F_1) = 0. \quad (2.11)$$

The conclusion $\mu_4(F_3) \leq \mu_5(F_3)$ is not true for F_3 and $\mu_5(F_4) \leq \mu_4(F_4)$ is not true for F_4 . The space X therefore does not have the measure-tree property.

We have thus proved the contrapositive and thus the original theorem that if X has the measure-tree property then it is hereditarily normal. ■

THEOREM 2.5. (Summing up.) The following concepts are equivalent: a) hereditary normality; b) the measure-tree property; and c) the pre-measure property.

The above reasoning can also be applied to other lattices than topologies. For example, lattices of zero sets are hereditarily normal. (This is easy to see. Let Z_f and Z_g be zero sets of functions f and g , respectively. The functions remain continuous on the subspace. The function $(|f| - |g|)/(|f| + |g|)$ is also continuous because $(|f| + |g|)$ is nonzero if Z_f and Z_g are disjoint in the subspace. The cozero sets of the function's positive and negative portions produce the necessary cozero sets.) The theorems in this paper can be used to prove the well-known fact that pre-measures on such lattices are measures. Similarly, Boolean algebras are obviously hereditarily normal; they also have

the pre-measure property.

It is easy to see that the lattice of open sets is a normal lattice if and only if the space is extremally disconnected (i.e., in an extremally-disconnected space any two disjoint open sets can be separated by closed sets). It is also easy to see that if μ_1 and μ_2 are measures on a space then $\mu_1(G) \leq \mu_2(G)$ for all open sets G if and only if $\mu_1(F) \geq \mu_2(F)$ for all closed sets F . If we put these facts together we can easily derive the following theorems:

THEOREM 2.6. A space X is extremally disconnected if and only if for every measure μ and every pair of minimal measures (a measure is a minimal measure if and only if there is no nontrivial measure strictly less than it for every closed set) μ_1 and μ_2 , if $\mu_1 \leq \mu$ and $\mu_2 \leq \mu$ then $\mu_1 = \mu_2$.

THEOREM 2.7. A space X is hereditarily extremally disconnected if and only if for all two-valued measures μ, μ_1 , and μ_2 on X the conditions $\mu \geq \mu_1$ and $\mu \geq \mu_2$ imply that either $\mu_2 \geq \mu_1$ or $\mu_1 \geq \mu_2$.

These theorems can be used to prove:

THEOREM 2.8. A normal extremally-disconnected space is hereditarily normal if and only if it is hereditarily extremally disconnected.

PROOF. Let X be normal and hereditarily extremally disconnected and let us prove it is hereditarily normal. Let μ_1, μ_2 , and μ_3 be three measures such that $\mu_1 \leq \mu_2$ and $\mu_1 \leq \mu_3$. Let μ_4 be a regular measure greater than μ_2 , and let μ_5 be a regular measure greater than μ_3 . Both μ_4 and μ_5 are greater than μ_1 and therefore, since the space is normal, $\mu_4 = \mu_5$. This means that $\mu_2 \leq \mu_4$ and $\mu_3 \leq \mu_4$. The space is hereditarily extremally disconnected and therefore μ_2 and μ_3 are comparable (i.e., either $\mu_2 \leq \mu_3$ or $\mu_3 \leq \mu_2$). We have shown that any two measures greater than the same measure are comparable. The space is therefore hereditarily normal.

We can use similar reasoning to show that if a space is extremally disconnected and hereditarily normal then it is hereditarily extremally disconnected. ■

If we examine the lattice of zero sets in a Tychonoff space, we can see that since it is hereditarily normal, if the lattice of cozero sets is normal (i.e., if the space is an F -space – see Gillman and Jerison [6], Chapter 14) then it is hereditarily normal. We can also easily prove the well-known results that a z -ideal in an F -space belongs to a chain from a minimal ideal to a maximal ideal.

The obvious conclusion – the property of being an F -space is hereditary – is false. Dow [3] constructed an F -space with an open subspace which is not an F -space. This can be reconciled with the fact that the lattice of cozero sets in an F -space is hereditarily normal. The members of the normal descendant lattice on the subspace consist of those cozero sets inherited from the original space. However, not every cozero set in the subspace has been inherited from the original space. The cozero sets of a subspace can include additional sets if there are more continuous functions on the subspace than on the original space.

At this point we have two questions: 1) Are there any spaces which are normal and extremally disconnected but which are not hereditarily extremally disconnected? 2) Are there any spaces which are hereditarily normal and extremally disconnected but which are not discrete?

The answer to the first question is yes. The Stone–Čech compactification of the integers ($\beta\mathbb{N}$, Example 111 in Steen and Seebach [7]) is the Stone space of a complete algebra. It is therefore normal and extremally disconnected. The power set of the integers modulo the ideal of finite sets is incomplete so there its Stone space is not extremally disconnected even though it is a subspace of $\beta\mathbb{N}$. This means that $\beta\mathbb{N}$ is not hereditarily extremally disconnected even though it is normal and extremally disconnected.

The answer to the second question is also yes. If we consider the single ultrafilter (Example 114 in Steen and Seebach [7]) we can see that this space must be extremally disconnected because if there are two disjoint open sets, at least one of them does not belong to the particular ultrafilter and it must therefore be closed. The two sets can therefore be separated by clopen sets. The space must also be hereditarily normal. It is countable and thus hereditarily Lindelöf. A regular hereditarily-Lindelöf space is hereditarily normal.

REFERENCES

1. CAMACHO Jr., J. On maximal measures with respect to a lattice, Int. J. Math. Math. Sci. **14** (1991), 93.
2. CARROL, L. The Complete Lewis Carroll, Vintage Books, New York, 1976.
3. DOW, A. CH and open subspaces of F -spaces, Proc. Amer. Math. Soc., **89** (1983), 341.
4. EID G. On normal lattices and Wallman spaces, Int. J. Math. Math. Sci., **13** (1990), 31.
5. FROLIK, Z. Prime filters with the C.I.P., Comm. Math. Univ. Carolinae, **13** (1972), 553.
6. GILLMAN L. and JERISON M. Rings of Continuous Functions, Van Nostrand, New York, 1960.
7. STEEN L. and SEEBACH Jr, J. A. Counterexamples in Topology, Springer-Verlag, New York–Heidelberg–Berlin, 1978.
8. SZETO M. Measure repleteness and mapping preservations, J. Indian Math. Soc., **43** (1979), 35.