A WAVE EQUATION WITH DISCONTINUOUS TIME DELAY

JOSEPH WIENER

Department of Mathematics
The University of Texas - Pan American
Edinburg, Texas 78539
and

LOKENATH DEBNATH

Department of Mathematics
University of Central Florida
Orlando, Florida 32816

(Received October 21, 1991)

ABSTRACT. The influence of certain discontinuous delays on the behavior of the solutions of the wave equation is studied.

KEY WORDS AND PHRASES: Partial Differential Equation, Piecewise Constant Delay, Wave Equation, Boundary Value Problem, Fourier Method, Oscillation

1991 AMS SUBJECT CLASSIFICATION CODES. 35L05, 35L20, 35L50, 35L99.

1. INTRODUCTION.

Functional differential equations (FDE) with delay provide a mathematical model for a physical or biological system in which the rate of change of the system depends upon its past history. The theory of FDE with continuous argument is well developed and has numerous applications in natural and engineering sciences. This paper continues our earlier work [1-5] in an attempt to extend this theory to differential equations with discontinuous argument deviations. In these papers, ordinary differential equations with arguments having intervals of constancy have been studied. Such equations represent a hybrid of continuous and discrete dynamical systems and combine properties of both differential and difference equations. They include as particular cases loaded and impulse equations, hence their importance in control theory and in certain biomedical models. Continuity of a solution at a point joining any two consecutive intervals implies recursion relations for the values of the solution at such points. Therefore, differential equations with piecewise constant argument (EPCA) are intrinsically closer to difference rather than differential equations. In [6] boundary value problems for some linear EPCA in partial derivatives were considered and the behavior of their solutions studied. The results were also extended to equations with positive definite operators in Hilbert spaces. In [7] initial value problems were studied for EPCA in partial derivatives. A class of loaded equations that arise in solving certain inverse problems was explored within the general framework of differential equations with piecewise constant delay. The purpose of the present note is to investigate the influence of terms with piecewise constant time on the behavior of the solutions, especially their oscillatory properties, of the wave equation.

2. SEPARATION OF VARIABLES IN SYSTEMS OF PDE.

Consider the boundary value problem (BVP) consisting of the equation

$$U_t(x, t) = AU_{xx}(x, t) + BU_{xx}(x, [t]), \quad (2.1)$$

the boundary conditions

$$U(0, t) = U(1, t) = O, \quad (2.2)$$

and the initial condition

$$U(x, 0) = U_0(x). \quad (2.3)$$

Here $U(x, t)$ and $U_0(x)$ are real $m \times m$ matrices, A and B are real constant $m \times m$ matrices, and $[.]$ designates the greatest integer function. According to [6], a matrix $U(x, t)$ is called a solution of BVP (2.1)-(2.3) if it satisfies the conditions: (i) $U(x, t)$ is continuous in $G = [0, 1] \times [0, \infty)$; (ii) U_t and U_{xx} exist and are continuous in G , with the possible exception of the points (x, n) , where one-sided derivatives exist ($n = 0, 1, 2, \dots$); (iii) $U(x, t)$ satisfies Eq. (2.1) in G , with the possible exception of the points (x, n) , and conditions (2.2), (2.3).

Looking for the solution of (2.1) - (2.3) in the form

$$U(x, t) = T(t)X(x) \quad (2.4)$$

gives

$$T'(t)X(x) = AT(t)X''(x) + BT([t])X''(x),$$

whence

$$(AT(t) + BT([t]))^{-1}T'(t) = X''(x)X^{-1}(x) = -P^2,$$

which generates the BVP

$$X''(x) + P^2X(x) = O, \quad (2.5)$$

$$X(0) = X(1) = O$$

and the equation with piecewise constant argument

$$T'(t) = -AT(t)P^2 - BT([t])P^2. \quad (2.6)$$

The general solution of Eq. (2.5) is

$$X(x) = \cos(xP)C_1 + \sin(xP)C_2,$$

where

$$\cos(xP) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n} P^{2n}}{(2n)!}, \quad \sin(xP) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1} P^{2n+1}}{(2n+1)!}$$

and C_1, C_2 are arbitrary constant matrices. From $X(0) = O$ we conclude that $C_1 = O$, and the condition $X(1) = O$ enables us to choose $\sin P = O$ (although this is not the necessary consequence of the equation $(\sin P)C_2 = O$). This can be written as $e^{iP} - e^{-iP} = O$ or $e^{2iP} = I$. Assuming that all eigenvalues $p_1, p_2, \dots, p_m$ of P are distinct and $S^{-1}PS = \mathcal{D} = \text{diag}(p_1, p_2, \dots, p_m)$, we have $e^{2iS\mathcal{D}S^{-1}} = I$, whence $Se^{2i\mathcal{D}}S^{-1} = I$ and $e^{2i\mathcal{D}} = I$. Therefore, $\mathcal{D} = \text{diag}(\pi j_1, \pi j_2, \dots, \pi j_m)$, where the j_k are integers, and $P = S\mathcal{D}S^{-1}$, $P^2 = S\mathcal{D}^2S^{-1} = S\text{diag}(\pi^2 j_1^2, \pi^2 j_2^2, \dots, \pi^2 j_m^2)S^{-1}$, $\sin(xP) = S \sin(x\mathcal{D})S^{-1} = S\text{diag}(\sin \pi j_1 x, \dots, \sin \pi j_m x)S^{-1}$. Furthermore, we can put

$$P_j = \text{diag}(\pi(m(j-1)+1), \dots, \pi mj), \quad (j = 1, 2, \dots) \quad (2.7)$$

in (2.5) and obtain the following result.

THEOREM 2.1. There exists an infinite sequence of matrix eigenfunctions for BVP (2.5)

$$X_j(x) = \sqrt{2} \text{diag}(\sin \pi(m(j-1)+1)x, \dots, \sin \pi m j x), \quad (j = 1, 2, \dots) \quad (2.8)$$

which is complete and orthonormal in the space $L^2[0, 1]$ of $m \times m$ matrices, that is

$$\int_0^1 X_j(x) X_k(x) dx = \begin{cases} 0, & j \neq k \\ I, & j = k \end{cases}$$

where I is the identity matrix.

REMARK. The matrices $SX_j(x)S^{-1}$ satisfy Theorem 2.1 for any nonsingular S .

THEOREM 2.2. Let $E(t)$ be the solution of the problem

$$T'(t) = -AT(t)P^2, \quad T(0) = I \quad (2.9)$$

and let

$$M(t) = E(t) + (E(t) - I)A^{-1}B. \quad (2.10)$$

If the matrix A is nonsingular, then Eq. (2.6) with the initial condition $T(0) = C_0$ has on $[0, \infty)$ a unique solution

$$T(t) = M(t - [t])M^{[t]}(1)C_0. \quad (2.11)$$

PROOF. On the interval $n \leq t < n+1$, where $n \geq 0$ is an integer, Eq. (2.6) turns into

$$T'(t) = -AT(t)P^2 - BC_nP^2, \quad C_n = T(n)$$

with the general solution

$$T(t) = E(t - n)C - A^{-1}BC_n.$$

At $t = n$ we have $C_n = C - A^{-1}BC_n$, whence $C = (I + A^{-1}B)C_n$ and $T(t) = (E(t - n) + (E(t - n) - I)A^{-1}B)C_n$, that is,

$$T(t) = M(t - n)C_n. \quad (2.12)$$

At $t = n+1$ we have $C_{n+1} = M(1)C_n$ and

$$C_n = M^n(1)C_0. \quad (2.13)$$

Hence,

$$T(t) = M(t - n)M^n(1)C_0,$$

which is equivalent to (2.11).

THEOREM 2.3. If $\|M(1)\| < 1$, then $\|T(t)\|$ exponentially tends to zero as $t \rightarrow +\infty$.

EXAMPLE. For the scalar parabolic equation

$$u_t(x, t) = a^2 u_{xx}(x, t) + bu_{xx}(x, [t])$$

we have $m = 1$ and $P_j = \pi_j$, according to (2.7). For Eq. (2.9), with $A = a^2$ and $P = P_j$, we have $E_j(t) = e^{-a^2 \pi_j^2 t}$

and $M_j(t) = e^{-a^2 \pi_j^2 t} - \frac{b}{a^2}(1 - e^{-a^2 \pi_j^2 t})$. Hence, the inequality $|M_j(1)| < 1$ is equivalent to

$$-1 < e^{-a^2 \pi_j^2} - \frac{b}{a^2}(1 - e^{-a^2 \pi_j^2}) < 1,$$

whence

$$-a^2 < b < a^2(1 + e^{-a^2 \pi_j^2})/(1 - e^{-a^2 \pi_j^2}).$$

Since the function $(1 + e^{-x})/(1 - e^{-x})$ is decreasing, all functions $T_j(t)$ exponentially tend to zero as $t \rightarrow \infty$ if and only if

$$-a^2 < b \leq a^2.$$

If $b < -a^2$, then all $T_j(t)$ monotonically tend to infinity as $t \rightarrow \infty$; and if $b > a^2(1 + e^{-a^2 \pi_j^2})/(1 - e^{-a^2 \pi_j^2})$, then all $T_j(t)$ are unbounded and oscillatory. For any $b > a^2$, there exists a positive integer j_0 such that the $T_{j_0}(t)$

are unbounded and oscillatory, for $j > j_0$. Indeed, letting $b = a^2 + \varepsilon$ and solving the inequality

$$a^2 + \varepsilon > a^2(1 + e^{-a^2 \pi^2 j^2})(1 - e^{-a^2 \pi^2 j^2})$$

gives

$$e^{-a^2 \pi^2 j^2} < \varepsilon / (2a^2 + \varepsilon),$$

which holds for any $\varepsilon > 0$ and sufficiently large j and implies $M_j(1) < -1$. If $b = -a^2$, then $M_j(t) = 1$, $T_j(t) = T_j(0)$ and $u(x, t) = u_0(x)$, for all t . Therefore, the condition $|b| \leq a^2$ is necessary and sufficient for the series

$$u(x, t) = \sum_{j=1}^{\infty} T_j(t) x_j(x) \quad (2.14)$$

to be a solution of the scalar BVP (2.1) - (2.3), with $A = a^2$ and $B = b$, if $u_0(x)$ is three times continuously differentiable. The coefficients $T_j(0)$ are given by

$$T_j(0) = \int_0^1 u_0(x) X_j(x) dx,$$

where $X_j(x) = \sqrt{2} \sin(\pi j x)$ and $u_0(x) \in C^3[0, 1]$ satisfies $u_0(0) = u_0(1) = 0$.

THEOREM 2.4. The solution $T = O$ of Eq. (2.6) is globally asymptotically stable as $t \rightarrow +\infty$ if and only if the eigenvalues λ_r of the matrix $M(1)$ satisfy the inequalities

$$|\lambda_r| < 1, \quad r = 1, \dots, m. \quad (2.15)$$

PROOF. There exists a nonsingular matrix Q such that $M(1) = QJQ^{-1}$, where J is a diagonal or Jordan matrix with the diagonal elements λ_r . Hence, from (2.13) it follows that

$$C_n = (QJQ^{-1})^n C_0 = QJ^n Q^{-1} C_0$$

or

$$C_n = \sum_{r=1}^k Q_r(n) \lambda_r^n, \quad k \leq m \quad (2.16)$$

where the entries of the matrices $Q_r(n)$ are polynomials of degree not exceeding $k - 1$. This implies $C_n \rightarrow O$ as $n \rightarrow \infty$ if and only if (2.15) holds, and the conclusion of the theorem follows from (2.12).

LEMMA. All entries of every solution of the equation

$$T'(t) = A_1 T(t) A_2, \quad (2.17)$$

with constant $m \times m$ matrices A_1 and A_2 are linear combinations of terms $t^k \exp(\alpha_i^{(1)} \alpha_j^{(2)} t)$, where $\alpha_i^{(1)}$ and $\alpha_j^{(2)}$ are eigenvalues of A_1 and A_2 and k is integer.

PROOF. Assume, for simplicity, that the eigenvalues of A_1 and A_2 are distinct, and let

$$S_1^{-1} A_1 S_1 = D_1 = \text{diag}(\alpha_1^{(1)}, \alpha_2^{(1)}, \dots, \alpha_m^{(1)}),$$

$$S_2^{-1} A_2 S_2 = D_2 = \text{diag}(\alpha_1^{(2)}, \alpha_2^{(2)}, \dots, \alpha_m^{(2)}).$$

Then the substitution $T = S_1 V$ changes (2.17) to

$$V'(t) = D_1 V(t) A_2 \quad (2.18)$$

and the substitution $V = WS_2^{-1}$ transforms (2.18) into

$$W'(t) = D_1 W(t) D_2.$$

The entries of $W(t)$ are $c_{ij} \exp(\alpha_i^{(1)} \alpha_j^{(2)} t)$, with arbitrary constants c_{ij} , and the completion of the proof follows

from the formula $T = S_1 W S_2^{-1}$. If some of the eigenvalues are multiple, polynomial factors will replace the constants c_{ij} .

THEOREM 2.5. If all eigenvalues of A have positive real parts, $U_0(x) \in C^3[0, 1]$, and $\|A^{-1}B\| < 1$, then BVP (2.1) - (2.3) has a solution (2.14). This series and all its term-by-term derivatives converge uniformly.

PROOF. The functions $T_j(t)$ satisfy Eq. (2.6), with P_j given by (2.7). Therefore, setting in the above lemma $A_1 = -A$, $A_2 = P_j^2$ and noting that $\operatorname{Re} \alpha_i^{(1)} < 0$, $\alpha_j^{(2)} > 0$, we conclude that $E_j(t) \rightarrow 0$ as $j \rightarrow \infty$ and all $t > 0$, where $E_j(t)$ is the solution of (2.9). Hence, (2.10) implies $M_j(t) \rightarrow -A^{-1}B$ as $j \rightarrow \infty$ and $\|M_j(t)\| < 1$, for sufficiently large j , by virtue of the condition $\|A^{-1}B\| < 1$. From the formula

$$T_j(t) = M_j(t - [t]) M_j^{[t]}(1) C_{0j}$$

we note that $\|T_j(t)\| \rightarrow 0$ exponentially as $t \rightarrow \infty$, for sufficiently large j . Therefore, for any $\varepsilon > 0$, there exist $t = t_0 > 0$ and $j = N$ such that $\|R_N(t)\| < \varepsilon$, for $t > t_0$, where $R_N(t)$ is the remainder of series (2.14) after the N th term. For $0 < t \leq t_0$ and large j , we have $\|T_j(t)\| < \|C_{0j}\|$, and in this case the uniform convergence of series (2.14), together with its respective derivatives in t and x , follows from the smoothness of the initial function $U_0(x)$ and the formula

$$T_j(0) = C_{0j} = \int_0^1 U_0(x) X_j(x) dx.$$

THEOREM 2.6. If all eigenvalues λ_r of the matrix $M(1)$ are negative, where $M(t)$ is defined by (2.10), then each entry of every matrix solution of Eq. (2.6) is oscillatory and, more precisely, has a zero in each interval $n \leq t \leq n + 1$, for sufficiently large n .

PROOF. Let $\tau^{(is)}(t)$ denote an entry of a solution $T(t)$ of Eq. (2.6) and let $c_n^{(is)} = \tau^{(is)}(n)$. Then from (2.16) we get

$$c_n^{(is)} = \sum_{r=1}^k q_r^{(is)}(n) \lambda_r^n, \quad c_{n+1}^{(is)} = \sum_{r=1}^k q_r^{(is)}(n+1) \lambda_r^{n+1},$$

where $q_r^{(is)}$ are the entries of the matrices Q_r . If $c_n^{(is)} = 0$, for infinitely many n , then $\tau^{(is)}(t)$ is oscillatory. Assuming $c_n^{(is)} \neq 0$, for all sufficiently large n , we conclude that

$$\lim_{n \rightarrow \infty} c_{n+1}^{(is)} / c_n^{(is)} = \lambda_\mu < 0$$

(since the coefficients $q_r^{(is)}(n)$ are polynomials of n), where λ_μ is one of the eigenvalues of $M(1)$. Hence, $c_{n+1}^{(is)} / c_n^{(is)} < 0$, starting with some n , which implies that $\tau^{(is)}(t)$ has a zero in each interval $n < t < n + 1$, for sufficiently large t .

THEOREM 2.7. If the matrix $M(1)$ has a negative eigenvalue, then there exists an initial matrix C_0 such that the corresponding solution $T(t)$ of Eq. (2.6) has an oscillatory entry with a zero in each interval $n < t < n + 1$, for sufficiently large n .

PROOF. Assume, for simplicity, that in the equation $M(1) = QJQ^{-1}$ the matrix J is diagonal, $J = \operatorname{diag}(\lambda_1, \lambda_2, \dots, \lambda_m)$, and that $\lambda_1 < 0$. In the equation $C_n = QJ^n Q^{-1} C_0$ put $C_0 = Q$; then $C_n = QJ^n$ and $c_n^{(1)} = (q_{11}\lambda_1^n, q_{21}\lambda_1^n, \dots, q_{m1}\lambda_1^n)$, where $c_n^{(1)}$ is the first column of C_n and $(q_{11}, q_{21}, \dots, q_{m1})$ is the first column of Q . Since Q is nonsingular, there exists an element $q_{11} \neq 0$. Hence, $\lim_{n \rightarrow \infty} c_{n+1}^{(1)} / c_n^{(1)} = \lambda_1 < 0$ as $n \rightarrow \infty$, which proves that the entry $\tau^{(1)}(t)$ of the solution $T(t)$ has a zero in each interval $(n, n+1)$, for sufficiently large n .

Following [8], we can prove that if the matrix $M(1)$ has no positive eigenvalues and no eigenvalues with equal moduli, then each entry of every solution of Eq. (2.6) is oscillatory. On the other hand, the availability of positive eigenvalues for $M(1)$ implies the existence of nonoscillatory entries for $T(t)$.

Separation of variables in the matrix equation with constant coefficients

$$U_j(x, t) = A_0 U(x, t) + A_2 U_{xx}(x, t) + B_0 U(x, [t]) + B_2 U_{xx}(x, [t])$$

leads to (2.5) and to the EPCA

$$T'(t) = A_0 T(t) - A_2 T(t) P^2 + B_0 T([t]) - B_2 T([t]) P^2,$$

which can be also investigated by the above method.

3. A SCALAR WAVE EPCA.

Separation of variables in the equation with constant coefficients

$$u_{tt}(x, t) = a^2 u_{xx}(x, t) - b u_{xx}(x, [t]) \quad (3.1)$$

and boundary conditions (2.2) yields $X_j(x) = \sqrt{2} \sin(\pi j x)$ and leads to the EPCA

$$T''_j(t) + a^2 \pi^2 j^2 T_j(t) = b \pi^2 j^2 T_j([t]). \quad (3.2)$$

For brevity, we omit the subindex j and make the substitution $T'(t) = V(t)$, which changes (3.2) to a vector EPCA

$$w'(t) = A w(t) + B w([t]), \quad (3.3)$$

where $w = \text{col}(T, V)$ and

$$A = \begin{pmatrix} 0 & 1 \\ -a^2 \pi^2 j^2 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 \\ b \pi^2 j^2 & 0 \end{pmatrix}.$$

Eq. (3.3) on the interval $n \leq t < n+1$ becomes

$$w'(t) = A w(t) + B c_n, \quad c_n = w(n)$$

with the solution

$$w(t) = M(t - n) c_n,$$

where

$$M(t) = e^{At} + (e^{At} - I) A^{-1} B. \quad (3.4)$$

Therefore, Eq. (3.3) with the initial condition $w(0) = c_0$ has on $[0, \infty)$ a unique solution given by the right-hand side of (2.11), where $M(t)$ is defined in (3.4).

THEOREM 3.1. For $b < 0$, the solution $w = 0$ of Eq. (3.3) is unstable.

PROOF. Computations show that

$$e^{At} = \cos(\omega t) I + \omega^{-1} \sin(\omega t) A$$

and

$$e^{At} - I = \begin{pmatrix} \cos \omega t - 1 & \omega^{-1} \sin \omega t \\ -\omega \sin \omega t & \cos \omega t - 1 \end{pmatrix},$$

where $\omega = a \pi j$. Also,

$$(e^{At} - I) A^{-1} B = \begin{pmatrix} b(1 - \cos \omega t)/a^2 & 0 \\ b \omega \sin \omega t/a^2 & 0 \end{pmatrix}.$$

Hence,

$$M(t) = \begin{pmatrix} \cos \omega t + b a^{-2} (1 - \cos \omega t) & \omega^{-1} \sin \omega t \\ (b a^{-2} - 1) \omega \sin \omega t & \cos \omega t \end{pmatrix}$$

and

$$\det M(1) = 1 - \frac{b}{a^2} + \frac{b}{a^2} \cos \omega.$$

The condition $b < 0$ implies $\det M(1) > 1$ and shows that at least one of the eigenvalues λ of $M(1)$ satisfies $|\lambda| > 1$. Therefore $\|w(t)\| \rightarrow \infty$ as $t \rightarrow +\infty$, for some initial vector $c_0 \neq 0$.

THEOREM 3.2. For $b > a^2$, the solution $w = 0$ of Eq. (3.3) is unstable.

PROOF. Calculations yield

$$\det(M(1) - \lambda J) = \lambda^2 - 2\left(\cos \omega + \frac{b}{a^2} \sin^2 \frac{\omega}{2}\right) \lambda + 1 - \frac{b}{a^2} + \frac{b}{a^2} \cos \omega$$

and the expressions $\lambda_1 = s + d$, $\lambda_2 = s - d$ for the eigenvalues λ_1, λ_2 of $M(1)$, where $s = \cos \omega + \frac{b}{a^2} \sin^2 \frac{\omega}{2}$, $d^2 = \left(\frac{b}{a^2} - 1\right) \sin^2 \omega + \frac{b^2}{a^4} \sin^4 \frac{\omega}{2}$. The condition $b > a^2$ shows that $d^2 > 0$ and $\lambda_1 > 1$. The latter inequality implies $\|w(t)\| \rightarrow \infty$ as $t \rightarrow +\infty$, for some initial vector $c_0 \neq 0$.

THEOREM 3.3. The solution $w = 0$ of Eq. (3.3) is asymptotically stable as $t \rightarrow +\infty$ if and only if

$$0 < b < a^2, \quad (3.5)$$

and $\omega \neq 2\pi n$, $n = 0, 1, 2, \dots$.

PROOF. The condition $d^2 < 0$, which means that the eigenvalues of $M(1)$ are complex, leads to

$$\cos^2 \frac{\omega}{2} > b^2 / (2a^2 - b)^2,$$

whence

$$b < a^2 \left(1 - \tan^2 \frac{\omega}{4}\right) \text{ or } b < a^2 \left(1 - \cot^2 \frac{\omega}{4}\right).$$

Since $|\lambda_1| = |\lambda_2|$ and $\det M(1) = \lambda_1 \lambda_2$, the inequality $|\lambda_1| < 1$ is equivalent to $\det M(1) < 1$, that is, to $b > 0$.

Therefore, in the case of complex eigenvalues, a criterion for asymptotic stability is $0 < b < \max\left(a^2\left(1 - \tan^2 \frac{\omega}{4}\right), a^2\left(1 - \cot^2 \frac{\omega}{4}\right)\right)$. The inequality $d^2 > 0$ in the case of distinct real eigenvalues leads to $b > \max\left(a^2\left(1 - \tan^2 \frac{\omega}{4}\right), a^2\left(1 - \cot^2 \frac{\omega}{4}\right)\right)$, and the inequalities $\lambda_1 < 1$, $\lambda_2 > -1$ yield $b < a^2$. Hence, in this case a criterion of asymptotic stability is

$$\max\left(a^2\left(1 - \tan^2 \frac{\omega}{4}\right), a^2\left(1 - \cot^2 \frac{\omega}{4}\right)\right) < b < a^2.$$

Finally, if $b = \max\left(a^2\left(1 - \tan^2 \frac{\omega}{4}\right), a^2\left(1 - \cot^2 \frac{\omega}{4}\right)\right)$, then $d = 0$ and $\lambda_1 = \lambda_2 = \cos \omega + ba^{-2} \sin^2 \omega / 2$ whence $\cos \omega < \lambda_1 < \cos^2 \omega / 2$ and $|\lambda_1| < 1$. According to (2.15), this implies asymptotic stability and completes the proof of criterion (3.5).

REMARK 1. If $b = a^2$, then $\lambda_1 = 1$, $\lambda_2 = \cos \omega$, and the solutions of (3.3) are bounded but not asymptotically stable.

REMARK 2. In Theorems 3.1 and 3.2 it was implicitly assumed that $\omega \neq 2\pi n$. If $\omega = 2\pi n$, then $\lambda_1 = \lambda_2 = 1$, which leads to the existence of unbounded solutions for (3.3).

COROLLARY 1. If the coefficient a is irrational, then (3.5) is a criterion of asymptotic stability of the solutions to (3.2) for all j .

PROOF. Recalling that $\omega = \omega_j = a\pi j$, we note that the equality $a\pi j = 2\pi n$ is impossible for any irrational a .

COROLLARY 2. For any rational a , there exist infinitely many integers j such that the corresponding solutions $w_j(t)$ of (3.3) are unbounded.

PROOF. The equation $\omega_j = 2\pi n$ implies $j = 2n/a$, and j will be an integer for infinitely many integers n .

THEOREM 3.4. Each component of every solution of Eq. (3.3) oscillates if and only if either of the following conditions holds true:

- (i) $b < \max\left(a^2\left(1 - \tan^2\frac{\omega}{4}\right), a^2\left(1 - \cot^2\frac{\omega}{4}\right)\right);$
- (ii) $\max\left(a^2\left(1 - \tan^2\frac{\omega}{4}\right), a^2\left(1 - \cot^2\frac{\omega}{4}\right)\right) < b < \frac{a^2}{2\sin^2\frac{\omega}{2}} \quad \text{and} \quad \cos\omega < -1/2.$

PROOF. Hypothesis (i) means that the eigenvalues of $M(1)$ are complex, and in this case all nontrivial solutions of (3.3) oscillate componentwise. In the case of real eigenvalues, we require $d^2 > 0$ and $\lambda_1 = s + d < 0$, which also implies $\lambda_2 = s - d < 0$, where s and d are defined in Theorem 3.2. The condition $d^2 > 0$ leads to the left-side inequality in (ii), and the condition $\lambda_1 < 0$ yields the right-side inequality in (ii). Finally, the restriction $\cos\omega < -1/2$ arises from the comparison of the inequalities $s < 0$ and $\lambda_1 < 0$.

REMARK 3. The restriction $\cos\omega < -1/2$ implies $a^2/2 \sin^2(\omega/2) < 2a^2/3$.

In conclusion, it is worth noting that the asymptotic properties of Eq. (3.2) depend on the algebraic nature of the coefficient a . For $b < 0$, all solutions of Eq. (3.2) are unstable and oscillatory; for $b > a^2$ all solutions of Eq. (3.2) are unstable and nonoscillatory. These two cases hold true for both rational and irrational values of a . For

$$0 < b < \max\left(a^2\left(1 - \tan^2\frac{\omega}{4}\right), a^2\left(1 - \cot^2\frac{\omega}{4}\right)\right),$$

all solutions of (3.2) are asymptotically stable and oscillatory, provided that $\omega \neq 2\pi n$. However, as indicated in Corollary 2, for any rational a , there exist infinitely many integers j such that $\omega_j = 2\pi n$, which leads to the existence of unbounded solutions for (3.2). Furthermore, since $\omega = \omega_j = a\pi j$, the inequality $\cos\omega < -1/2$ breaks down for infinitely many integers j . Therefore, under hypothesis (ii) of Theorem 3.4, there are infinitely many solutions of Eq. (3.2) which are asymptotically stable and oscillatory, as well as infinitely many solutions which are asymptotically stable and nonoscillatory ($\omega \neq 2\pi n$). Also, for $\omega \neq 2\pi n$ and $a^2/2 \sin^2(\omega/2) < b < a^2$, the solutions of (3.2) are asymptotically stable and nonoscillatory. Problems of this nature deserve further investigation.

ACKNOWLEDGEMENT. This research was partially supported by U.S. Army Grant DAAL03-89-G-0107, NASA Grant NAG 9-553, and by the University of Central Florida.

REFERENCES

1. WIENER, J. Differential equations with piecewise constant delays, in Trends in the Theory and Practice of Nonlinear Differential Equations, Lakshmikantham, V. (editor), Marcel Dekker, New York, 1983, 546-552.
2. COOKE, K. L. and WIENER, J. Retarded differential equations with piecewise constant delays, J. Math. Anal. Appl. **99**(1), (1984), 264-297.
3. SHAH, S. M. and WIENER, J. Advanced differential equations with piecewise constant argument deviations, Internat. J. Math. & Math. Sci. **6**(4), (1983), 671-703.
4. COOKE, K. L. and WIENER, J. Neutral differential equations with piecewise constant argument, Bollettino Unione Matematica Italiana **7** (1987), 321-346.
5. COOKE, K. L. and WIENER, J. An equation alternately of retarded and advanced type, Proc. Amer. Math. Soc. **99** (1987), 726-732.
6. WIENER, J. Boundary-value problems for partial differential equations with piecewise constant delay, Internat. J. Math. & Math. Sci. **14** (1991), 301-321.
7. WIENER, J. and DEBNATH, L. Partial differential equations with piecewise constant delay, Internat. J. Math. & Math. Sci. **14** (1991), 485-496.
8. WIENER, J. and COOKE, K. L. Oscillations in systems of differential equations with piecewise constant argument, J. Math. Anal. Appl. **137**(1), (1989), 221-239.