GENERIC SUBMANIFOLDS OF A LOCALLY CONFORMAL KAEHLER MANIFOLD-II

M. HASAN SHAHID

Department of Mathematics
Faculty of Natural Sciences
Jamia Millia Islamia
New Delhi-110 025, India

and

KOUEI SEKIGAWA

Department of Mathematics
Faculty of Science
Niigata University
Niigata, 950-21, Japan

(Received March 6, 1991 and in revised form August 8, 1991)

ABSTRACT. The purpose of this paper is to study generic submanifolds with parallel structures, generic product submanifolds and totally umbilical submanifolds of a locally conformal Kaehler manifold. Moreover, we give some examples of generic submanifolds of a locally conformal Kaehler manifold which are not CR -submanifolds.

KEY WORDS AND PHRASES. Locally conformal Kaehler manifold, generic submanifold, CR -submanifold.

1991 AMS SUBJECT CLASSIFICATION CODES. 53C35, 53C40.

1. INTRODUCTION.

Let $\bar{M}$ be an almost Hermitian manifold with almost Hermitian structure (J, g) . The manifold $\bar{M}$ is called a *local conformal Kaehler* (briefly, l.c.K.) manifold if for any $x \in \bar{M}$ there is an open neighborhood $\mathfrak{U}$ such that, for some differentiable function $\sigma: \mathfrak{U} \rightarrow \mathbb{R}$, $g' = e^{-\sigma} g|_{\mathfrak{U}}$ is a Kaehler metric on $\mathfrak{U}$. If $\mathfrak{U} = \bar{M}$ then the manifold is called a *globally conformal Kaehler* (briefly, g.c.K.) manifold. Let Ω be the Kaehler form of an almost Hermitian manifold $\bar{M}$ defined by $\Omega(U, V) = g(U, JV)$, for any vector fields U, V on $\bar{M}$. Then it is easy to see that $\bar{M}$ is a l.c.K. manifold if and only if there is a global 1-form ω (the Lee form of $\bar{M}$) such that

$$d\Omega = \omega \wedge \Omega, \quad d\omega = 0, \quad (1.1)$$

and $\bar{M}$ is a g.c.K. manifold if and only if ω is exact. For a l.c.K. manifold $\bar{M}$, the Lee vector field B is given by

$$g(B, U) = \omega(U) \quad (1.2)$$

for any vector field U on $\bar{M}$. We denote by $\bar{\nabla}$ the Levi-Civita connection of g . We define a torsion-free linear connection $\tilde{\nabla}$ on $\bar{M}$ by

$$\tilde{\nabla}_U V = \bar{\nabla}_U V - \frac{1}{2} \{ \omega(U)V + \omega(V)U - g(U, V)B \} \quad (1.3)$$

for any vector fields U, V on $\bar{M}$. The linear connection $\tilde{\nabla}$ is called the Weyl connection of $\bar{M}$. Then we may easily observe that the Weyl connection $\tilde{\nabla}$ satisfies the condition: $\tilde{\nabla} J = 0, \tilde{\nabla} g = 0$ on each neighborhood on which $(J, g' = e^{-\sigma} g|_{\mathcal{Q}})$ is a Kaehler structure.

In general, let $\bar{M}$ be a $2n$ -dimensional almost Hermitian manifold and M be an m -dimensional Riemann manifold isometrically immersed in $\bar{M}$. Let ∇ be the Levi-Civita connection on M induced by $\bar{\nabla}$. Then the Gauss and Weingarten formulas are given respectively by

$$\bar{\nabla}_U V = \nabla_U V + h(U, V), \quad (1.4)$$

$$\bar{\nabla}_U N = -A_N U + \nabla_U^\perp N \quad (1.5)$$

for any vector fields U, V tangent to M and N normal to M , where h is the second fundamental form of M in $\bar{M}$ and $\nabla^\perp$ is the normal connection on the normal bundle $T^\perp(M)$ with respect to the Levi-Civita connection $\bar{\nabla}$. Then we have $g(A_N U, V) = g(h(U, V), N)$, for any vector fields U, V tangent to M . For any vector field U tangent to M , we put

$$JU = PU + FU \quad (1.6)$$

where PU and FU are tangential and normal components of JU , respectively. Then P is an endomorphism of the tangent bundle $T(M)$ of M and F is a normal bundle valued 1-form on $T(M)$. For any vector field N normal to M , we put

$$JN = tN + fN, \quad (1.7)$$

where tN and fN are the tangential and normal components of JN , respectively. Then f is an endomorphism of the normal bundle $T^\perp(M)$ of M in $\bar{M}$ and t is a tangent bundle valued 1-form on $T^\perp(M)$.

DEFINITION: Let M be a submanifold of an almost Hermitian manifold $\bar{M}$. The holomorphic subspace D_x of $T_x M$ at $x \in M$ is defined by $D_x = T_x M \cap JT_x M$. D_x is the maximal complex subspace of $T_x \bar{M}$ which is contained in $T_x M$. If the dimension of D is constant along M , and furthermore, D defines a differentiable distribution on M , then M is called a generic submanifold of $\bar{M}$.

Let M be a generic submanifold of an almost Hermitian manifold $\bar{M}$. We call the distribution D the holomorphic distribution and the orthogonal complementary distribution $D^\perp$ the purely real distribution. They satisfy the following relations:

$$D_x \cap D_x^\perp = \{0\}, \quad D_x^\perp \cap JD_x^\perp = \{0\} \text{ for each } x \in M.$$

Let ν_x be the holomorphic normal space of M at x , i.e.,

$$\nu_x = T_x^\perp M \cap JT_x^\perp M.$$

Then $\nu_x (x \in M)$ defines a differentiable vector subbundle ν of $T^\perp(M)$ satisfying

$$T^\perp(M) = FD^\perp + \nu \text{ (direct sum)}, \quad t(T^\perp(M)) = D^\perp. \quad (1.8)$$

Furthermore, we have

$$D \perp D^\perp, PD = D \text{ and } D^\perp \supset PD^\perp. \quad (1.9)$$

We put $\dim D = 2p$ and $\dim D^\perp = q$. If $p, q \geq 1$, then the generic submanifold M is said to be proper. In the sequel, we shall consider only proper generic submanifolds. We put

$$(\nabla_U P)V = \nabla_U(PV) - P(\nabla_U V), \quad (1.10)$$

and

$$(\nabla_U F)V = \nabla_U^\perp(FV) - F\nabla_U V \quad (1.11)$$

for any vector fields U, V tangent to M . We say that P (resp. F) is parallel if $(\nabla_U P)V = 0$ (resp. $(\nabla_U F)V = 0$) for any vector fields U, V tangent to M . If a generic submanifold M of an almost Hermitian manifold $\bar{M}$ satisfies the condition $JD^\perp \subset T^\perp(M)$, then M is called a CR -submanifold of $\bar{M}$. Dragomir ([4]) studied CR -submanifolds of l.c.K. manifolds. The present paper is a continuation of the previous work [5].

2. PRELIMINARIES.

Let M be a generic submanifold of a l.c.K. manifold $\bar{M}$. For the Lee vector field B of $\bar{M}$, we put

$$B = B^T + B^\perp \text{ along } M, \quad (2.1)$$

where B^T (resp. $B^\perp$) is the tangential (resp. normal) component of B . Furthermore, we put

$$B^T = B^D + B^{D^\perp} \text{ along } M, \quad (2.2)$$

where B^D (resp. $B^{D^\perp}$) is the D -component (resp. $D^\perp$ -component) of $B^\perp$. Since $\tilde{\nabla} J = 0$ with respect to the Weyl connection $\tilde{\nabla}$, taking account of (1.3) ~ (1.7), (1.11), (1.12), (2.1) and (2.2), we have

$$\begin{aligned} (\nabla_X P)Y - \frac{1}{2}\omega(JY)X + \frac{1}{2}\omega(Y)JX - th(X, Y) \\ + \frac{1}{2}g(X, JY)B^T - \frac{1}{2}g(X, Y)PB^\perp - \frac{1}{2}g(X, Y)tB^\perp = 0, \end{aligned} \quad (2.3)$$

$$\begin{aligned} h(X, JY) - F\nabla_X Y + \frac{1}{2}g(X, JY)B^\perp \\ - \frac{1}{2}g(X, Y)FB^T - \frac{1}{2}g(X, Y)fB^\perp - fh(X, Y) = 0, \end{aligned} \quad (2.4)$$

$$(\nabla_X P)Z - A_{FZ}X - \frac{1}{2}\omega(JZ)X + \frac{1}{2}\omega(Z)JX - th(X, Z) = 0, \quad (2.5)$$

$$(\nabla_X F)Z + h(X, PZ) - fh(X, Z) = 0, \quad (2.6)$$

$$(\nabla_Z P)X - \frac{1}{2}\omega(JX)Z - \frac{1}{2}\omega(X)PZ - th(X, Z) = 0, \quad (2.7)$$

$$F\nabla_Z X - h(JX, Z) + fh(X, Z) = 0, \quad (2.8)$$

$$\begin{aligned} (\nabla_Z P)W - A_{FW}Z - \frac{1}{2}\omega(JW)Z + \frac{1}{2}\omega(W)PZ + \frac{1}{2}g(Z, JW)B^T \\ - \frac{1}{2}g(Z, W)PB^T - \frac{1}{2}g(Z, W)tB^\perp - th(Z, W) = 0, \end{aligned} \quad (2.9)$$

$$\begin{aligned} (\nabla_Z F)W + h(Z, PW) + \frac{1}{2}g(Z, JW)B^T + \frac{1}{2}\omega(W)FZ \\ - \frac{1}{2}g(Z, W)FB^T - \frac{1}{2}g(Z, W)fB^\perp - fh(Z, W) = 0, \end{aligned} \quad (2.10)$$

for any $X, Y \in D$ and $Z, W \in D^\perp$.

We recall the conditions for the distributions D and $D^\perp$ to be integrable.

PROPOSITION 2.1 ([5]). The distribution $D^\perp$ is integrable if and only if

$$g(h(X, JY) - h(JX, Y) + g(X, JY)B, FZ) = 0,$$

for any $X, Y \in D$ and $Z \in D^\perp$.

PROPOSITION 2.2 ([5]). The distribution $D^\perp$ is integrable if and only if

$$\nabla_Z(PW) - \nabla_W(PZ) + A_{FZ}W - A_{FW}Z + g(Z, JW)B \in D^\perp,$$

for any $Z, W \in D^\perp$.

Let M be a totally geodesic generic submanifold of a Kaehler manifold $\bar{M}$. Then it follows immediately that P and F are parallel, and furthermore D is integrable. So, it is worthwhile to study generic submanifolds with parallel structures and also totally umbilical generic submanifolds in a l.c.K. manifold.

3. GENERIC SUBMANIFOLDS WITH PARELLEL STRUCTURES.

In this section, we consider generic submanifolds with parallel P (resp. F) of a l.c.K. manifold.

THEOREM 3.1. Let M be a generic submanifold of a l.c.K. manifold $\bar{M}$. If P is parallel, then D is integrable and $B^D{}^\perp = 0$ along M . Moreover, if $\dim D \geq 4$, then $B^T = 0$ along M .

PROOF. By (1.11) and (2.3), we get

$$-\frac{1}{2}\omega(JY)X + \frac{1}{2}\omega(JX)Y + g(X, JY)B^T + \frac{1}{2}\omega(Y)JX - \frac{1}{2}\omega(X)JY = 0, \quad (3.1)$$

for $X, Y \in D$. Putting $Y = JX$ in (3.1), we get

$$\omega(X)X + \omega(JX)JX - g(X, X)B^\perp = 0, \quad (3.2)$$

for any vector field X on M . From (3.2), we get

$$(p-1)g(B^D, B^D) + pg(B^D{}^\perp, B^D{}^\perp) = 0. \quad (3.3)$$

First, we assume $p \geq 2$. Then, by (3.3), we have

$$B^D = 0, \quad B^D{}^\perp = 0 \text{ (and hence } B^\perp = 0). \quad (3.4)$$

Thus, by (2.3) and (3.4), we get

$$2th(X, Y) + g(X, Y)tB = 0, \quad (3.5)$$

for $X, Y \in D$. On one hand, by (1.11) and (2.4), we get

$$F\nabla_X(PY) + h(X, Y) + fh(X, JY) + \frac{1}{2}g(X, Y)B^\perp + \frac{1}{2}g(X, JY)fB^\perp = 0, \quad (3.6)$$

for $X, Y \in D$. By (1.11) and (3.6), we get

$$FP[X, Y] + f\{h(X, JY) - h(JX, Y)\} + g(X, JY)fB^\perp = 0, \quad (3.7)$$

for $X, Y \in D$. From (3.5), we get also

$$t\{h(X, JY) - h(JX, Y)\} + g(X, JY)tB^\perp = 0, \quad (3.9)$$

for $X, Y \in D$. Thus, by (3.7) and (3.8), we have

$$J\{h(X, JY) - h(JX, Y)\} + g(X, JY)JB^\perp = -FP[X, Y], \quad (3.9)$$

for $X, Y \in D$. By (3.9), we have

$$\begin{aligned} & g(h(X, JY) - h(JX, Y) + g(X, JY)B, JZ) \\ & = g(FP[X, Y], Z) = 0, \end{aligned} \quad (3.10)$$

for $X, Y \in D$ and $Z \in D^\perp$. Thus, from Proposition 3.1 and (3.10), it follows that D is integrable. Next, we assume that $p = 1$. Then, by (3.3), we have

$$B^D{}^\perp = 0. \quad (3.11)$$

By (2.3), we get

$$\begin{aligned} & \frac{1}{2}\omega(Y)X - \frac{1}{2}\omega(X)Y + \frac{1}{2}\omega(JY)JX - \frac{1}{2}\omega(JX)JY \\ & - t\{h(X, JY) - h(JX, Y)\} - g(X, JY)PB^T - g(X, JY)tB^T = 0, \end{aligned} \quad (3.12)$$

for $X, Y \in D$. On one hand, by (2.4) and (3.1), we get

$$FP[X, Y] - f\{h(X, JY) - h(JX, Y)\} + g(X, JY)fB^\perp = 0, \quad (3.13)$$

for $X, Y \in D$. By (3.12) and (3.13), we get

$$\begin{aligned} & J\{h(X, JY) - h(JX, Y)\} + g(X, JY)JB^T + g(X, JY)PB^T \\ & + \frac{1}{2}\omega(X)Y - \frac{1}{2}\omega(Y)X + \frac{1}{2}\omega(JX)JY - \frac{1}{2}\omega(JY)JX + FP[X, Y] = 0, \end{aligned} \quad (3.14)$$

for $X, Y \in D$. From (3.11), it follows that $PB^T = JB^T$.

Thus, (3.14) implies

$$\begin{aligned} & h(X, JY) - h(JX, Y) + g(X, JY)B \\ & = \frac{1}{2}\omega(X)JY - \frac{1}{2}\omega(Y)JX - \frac{1}{2}\omega(JX)Y + \frac{1}{2}\omega(JY)X + JFP[X, Y], \end{aligned} \quad (3.15)$$

for $X, Y \in D$. By (3.15), we have

$$g(h(X, JY) - h(JX, Y) + g(X, JY)B, FZ) = g(FP[X, Y], Z) = 0, \quad (3.16)$$

for $X, Y \in D$ and $Z \in D^\perp$. Thus, from (3.16) and Proposition 3.1, it follows that D is integrable.

THEOREM 3.2. Let M be a generic submanifold of a l.c.K. manifold $\bar{M}$ such that F is parallel. Then the distribution D is integrable and each leaf of D is totally geodesic in M .

PROOF. By (1.12), we have

$$0 = (\nabla_K F)Y = F\nabla_X Y, \text{ for } X, Y \in D. \quad (3.17)$$

By (3.17), we have $\nabla_X Y \in D$ for any $X, Y \in D$, from which the theorem follows immediately.

4. GENERIC PRODUCT SUBMANIFOLDS.

Let M be a generic submanifold of an almost Hermitian manifold $\bar{M}$. If M is locally expressed in the form $M = M_D \times M_{D^\perp}$, where M_D (resp. $M_{D^\perp}$) is a holomorphic submanifold (resp. a purely real submanifold) of $\bar{M}$, then M is called a generic product submanifold of $\bar{M}$. In this section, we consider generic product submanifold of a l.c.K. manifold $\bar{M}$.

THEOREM 4.1. Let M be a generic product submanifold of a l.c.K. manifold $\bar{M}$. If $B^D = 0$ along M , then we have

$$B^T = 0 \text{ along } M, \quad (4.1)$$

and

$$\nabla_X P = 0, \quad (\nabla_Z P)X = 0, \quad (4.2)$$

for $X \in D, Z \in D^\perp$.

PROOF. Since $(\nabla_X P)Z \in D^\perp$, for $X \in D, Z \in D^\perp$, by (2.5), we get

$$g(h(X, Y), FZ) + \frac{1}{2}\omega(JZ)g(X, Y) - \frac{1}{2}\omega(Z)g(JX, Y) = 0, \quad (4.3)$$

for $X, Y \in D$, $Z \in D^\perp$. By (4.3), we get immediately $B^{D^\perp} = 0$, and hence (4.1). Since $(\nabla_X P)Y \in D$, for $X, Y \in D$, by (2.3) and (4.1), we get

$$(\nabla_X P)Y = 0, \quad \text{for } X, Y \in D. \quad (4.4)$$

Since $(\nabla_Z P) \in D^\perp$, for $Z, W \in D^\perp$, by (2.9) and (4.1), we get

$$g(h(X, Z), FW) = 0, \text{ for } X \in D, Z \in D^\perp. \quad (4.5)$$

by (2.5), (4.1) and (4.5), we have

$$\begin{aligned} 0 &= g((\nabla_X P)Z, W) - g(h(X, W), FZ) - g(th(X, Z), W) \\ &= g((\nabla_X P)Z, W) - g(h(X, W), FZ) + g(h(X, Z), FW) \\ &= g((\nabla_X P)Z, W) \end{aligned} \quad (4.6)$$

for $X \in D$, $Z, W \in D^\perp$. By (4.3) and (4.6), we have the first equality of (4.2). Since $(\nabla_Z P)X \in D$, for $X \in D$, $Z \in D^\perp$, by (2.7), we have immediately the second equality of (4.2). *Q.E.D.*

COROLLARY 4.2. Let M be a CR -product submanifold of a l.c.K. manifold $\bar{M}$. If $B^D = 0$ along M , then P is parallel.

PROOF. Since $PW = 0$, and $\nabla_Z W, (\nabla_Z P)W \in D^\perp$, for $Z, W \in D^\perp$, we have immediately $(\nabla_Z P)W = 0$ for $Z, W \in D^\perp$. Thus, from this together with (4.2), the corollary follows. *Q.E.D.*

5. TOTALLY UMBILICAL GENERIC SUBMANIFOLDS.

A Riemannian submanifold M of a Riemannian manifold $\bar{M}$ is called a totally umbilical submanifold if

$$h(U, V) = g(U, V)H, \quad (5.1)$$

for any vector fields U, V tangent to M , where H is the mean curvature vector. In this section, we consider some totally umbilical generic submanifolds of a l.c.K. manifold.

THEOREM 5.1. Let M be a totally umbilical generic submanifold of a l.c.K. manifold $\bar{M}$ such that P is parallel. Then we have $B^{D^\perp} = 0$ and $2H + B^\perp = 0$ along M . In particular, if $\dim D \geq 4$, then $2H + B = 0$ along M .

PROOF. Since P is parallel, from Theorem 3.1 and (3.4), (3.11), it follows that D is integrable and

$$B^{D^\perp} = 0. \quad (5.2)$$

By (2.4), we have easily

$$2H + B^\perp = 0. \quad (5.3)$$

By (3.1), we get

$$\omega(X)^2 + \omega(JX)^2 = g(X, X)g(B^T, B^T), \quad \text{for } X \in D. \quad (5.4)$$

By (5.2) and (5.4), we have

$$(p-1)g(B^T, B^T) = 0. \quad (5.5)$$

By (5.5), if $p \geq 2$, we have $B^T = 0$. Therefore, the Theorem follows from (5.3). *Q.E.D.*

COROLLARY 5.2. Let M be a totally umbilical generic submanifold of a l.c.K. manifold $\bar{M}$ such that $B \in D$. If P is parallel, then M is totally geodesic and $B = 0$ along M .

THEOREM 5.3. Let M be a totally umbilical generic submanifold of a l.c.K. manifold $\bar{M}$ such

that $\dim FD^\perp < \dim D^\perp$ on a dense open subset in M . If P is parallel and $\dim D^\perp \geq 2$, then $2H + B = 0$ along M .

PROOF. By (1.5), (2.9), (5.1) and (5.2), we have

$$\begin{aligned} 0 &= -\frac{1}{2}\omega(JW)g(Z, B^T) + \frac{1}{2}\omega(JZ)g(W, B^T) \\ &\quad + \frac{1}{2}\omega(W)g(PZ, B^T) - \frac{1}{2}\omega(Z)g(PW, B^T) + g(Z, JW)g(B^T, B^T) \\ &= g(Z, JW)g(B^T, B^T), \end{aligned} \quad (5.6)$$

for $Z, W \in D^\perp$. From (5.2) and (5.6), taking account of Theorem 5.1, the theorem follows immediately. Q.E.D.

THEOREM 5.4. Let M be a totally umbilical generic submanifold of a l.c.K. manifold $\bar{M}$ such that $B \in D^\perp$. Then the purely real distribution $D^\perp$ is totally geodesic in M .

PROOF. For $X \in D$, $W \in D^\perp$ and $N \in T^\perp(M)$, by (1.3), (1.5) and (5.1), we have

$$\begin{aligned} 0 &= g((\tilde{\nabla}_W J)N, X) \\ &= g(\tilde{\nabla}_W(JN), X) - g(J\tilde{\nabla}_W N, X) \\ &= g(\tilde{\nabla}_W(JN), X) + g(\bar{\nabla}_W N, JX) \\ &= g(\bar{\nabla}_W(tN), X) + g(\bar{\nabla}_W(fN), X) \\ &= g(\nabla_W(tN), X), \end{aligned}$$

from which the theorem follows immediately. Q.E.D.

THEOREM 5.5. Let M be a totally umbilical generic submanifold of a l.c.K. manifold $\bar{M}$ such that F is parallel. Then we have $2H + B^\perp = 0$ along M .

PROOF. Since F is parallel, from Theorem 3.2, it follows that $D^\perp$ is integrable and each leaf of $D^\perp$ is totally geodesic in M . Thus, by (2.4) and (5.1), we have immediately $2H + B^\perp = 0$. Q.E.D.

6. EXAMPLES.

In this section, we give some examples of generic submanifolds of Hopf manifolds which are not CR-submanifolds. Let $\mathbf{R}^{2n+2}$ be a $(2n+2)$ -dimensional Euclidean space equipped with the canonical inner product $(\cdot, \cdot)$ and $\{e_1, \dots, e_{2n+1}, e_{2n+2}\}$ the canonical orthonormal basis of $\mathbf{R}^{2n+2}$. We denote by J_0 the complex structure on $\mathbf{R}^{2n+2}$ defined by

$$J_0 e_{2m-1} = e_{2m}, J_0 e_{2m} = -e_{2m-1}, \quad 1 \leq m \leq n+1. \quad (6.1)$$

Let $S^{2n+1} = \{x \in \mathbf{R}^{2n+2}; (x, x) = 1\}$ be a $(2n+1)$ -dimensional unit sphere with the canonical Sasakian structure (φ, ξ, η, h) induced from the Kaehler structure $(J_0, (\cdot, \cdot))$ on $\mathbf{R}^{2n+2}$. It is well known that the structure vector field ξ defines the Hopf fibration $\pi: S^{2n+1} \rightarrow CP^n$, where CP^n is a (complex) n -dimensional complex projective space equipped with the canonical Fubini-Study metric of constant holomorphic sectional curvature 4. Let $S^1 = \{e^{i\sqrt{-1}t}; t \in \mathbf{R}\}$ be a unit circle. We define an almost complex structure J on $M = S^{2n+1} \times S^1$ (resp. $\bar{M} = S^{2n+1} \times \mathbf{R}$) by

$$JT = \xi, J\xi = -T \text{ and } JU = \varphi U, \quad (6.2)$$

for any vector field U on $\bar{M}$ such that $\eta(U) = 0$, where $T = \frac{\partial}{\partial t}$ is the canonical unit vector field on S^1 (resp. $\mathbf{R}^1$). Then $(S^{2n+1} \times S^1, J)$ (resp. $(S^{2n+1} \times \mathbf{R}^1, J)$) is a l.c.K. manifold (resp. a g.c.K. manifold) together with the product metric $g = h + 1$ on $\bar{M} = S^{2n+1} \times S^1$ (resp. $\bar{M} = S^{2n+1} \times \mathbf{R}^1$). Then the Lee form ω of $\bar{M}$ is given by $\omega = 2dt$.

I. We denote by S_{pq} the Segre imbedding $S_{pq}: CP^p \times CP^q \rightarrow CP^{p+q+pq}$ ([2]). Let M_1 be any q -dimensional purely real submanifold of CP^q . Then $M = CP^p \times M_1$ is a generic product submanifold

of CP^{p+q+pq} in which CP^p is imbedded as a totally geodesic complex submanifold. We denote by the immersion $\iota: M_1 \rightarrow CP^q$. Let $M = \{S_{pq} \circ (1 \times \iota)^{-1}(S^{2(p+q+pq)+1})\}$ be pull-back of the Hopf bundle $\pi: S^{2(p+q+pq)+1}$ by the immersion $S_{pq} \circ (1 \times \iota): CP^p \times M_1 \rightarrow CP^{p+q+pq}$. Then we may easily observe that M is a generic submanifold of the Hopf manifold $\bar{M} = S^{2(p+q+pq)+1} \times S^1$. For example, let M_1 be the real submanifold of CP^q ($q > 1$) defined by

$M_1 = \{(x_0, \dots, x_{q-1}, x_q + \sqrt{-1} x_{q-1}) \in CP^q; (x_0, \dots, x_{q-1}, x_q) \text{ are homogeneous coordinates of a } q\text{-dimensional real projective space } \mathbf{RP}^q\}$. Then M_1 is a purely real submanifold of CP^q which is not totally real.

In the following II ~ IV, we assume that $\bar{M} = S^7 \times S^1$.

II. Let Π be the 5-dimensional linear subspace of $\mathbf{R}^8$ given by $\Pi = \text{span}_{\mathbf{R}}\{e_1, \dots, e_5\}$. We put

$S^4 = S^7 \cap \Pi$ and $M_2^4 = \{x = \sum_{i=1}^5 x_i e_i \in S^4; 0 < x^5 < 1\}$. For each point $x \in M_2^4$, let D'_x be the subspace of $T_x M_2^4$ defined by $D'_x = \{u \in T_x M_2^4; (u, J_0 x) = 0, (u, e_5) = 0\}$. We put $M = M_2^4 \times S^1 (\subset S^7 \times S^1)$. For each point $(x, e^{\sqrt{-1}t}) \in M$, let $D_{(x, e^{\sqrt{-1}t})}$ be the subspace of $T_{(x, e^{\sqrt{-1}t})} M$ defined by $D_{(x, e^{\sqrt{-1}t})} = \{(u, 0) \in T_{(x, e^{\sqrt{-1}t})} M; u \in D'_x\}$. Then we may easily observe that M is a totally geodesic generic submanifold of $\bar{M}$ with the holomorphic distribution D which is not a CR -submanifold of $\bar{M}$. We may easily check that the Lee form of $\bar{M}$ is tangent to M .

III. We put $M = M_2^4 \times \{1\} (\subset S^7 \times S^1)$. Then M is also a totally geodesic generic submanifold of $\bar{M}$ with holomorphic distribution D as in II (restricted to $M_2^4 \times \{1\}$) which is not CR -submanifold of $\bar{M}$. In this case, we may easily check that the Lee form of $\bar{M}$ is normal to M .

IV. We put $M_3^4 = \{x = \sum_{i=1}^5 x_i e_i + \frac{1}{\sqrt{2}} e_7 \in S^7; 0 < x_5 < \frac{1}{\sqrt{2}}\}$. For each point $x \in M_3^4$, let D''_x be the subspace of $T_x M_3^4$ defined by $D''_x = \{u \in T_x M_3^4; (u, J_0 x) = 0, (u, e_5) = 0\}$. We put $M = M_3^4 \times \{1\}$. For each point $(x, 1) \in M$, let $D_{(x, 1)}$ be the subspace of $T_{(x, 1)} M$ defined by $D_{(x, 1)} = \{(u, 0) \in T_{(x, 1)} M; u \in D''_x\}$. Then we may easily observe that M is a totally umbilical generic submanifold of $\bar{M}$ with holomorphic distribution D which is not a CR -submanifold of $\bar{M}$ and is not totally geodesic in $\bar{M}$.

ACKNOWLEDGEMENT. The authors would like to express their hearty thanks to Professor L. Vanhecke for his kind advices.

REFERENCES

1. BEJANCU, A., CR -submanifolds of a Kaehler manifold I, Proc. Amer. Math. Soc. **69** (1978), 135-142.
2. CHEN, B.Y., Geometry of Submanifolds, New York, 1973.
3. CHEN, B.Y., Differential geometry of real submanifolds in a Kaehler manifold, Monatsh. Math. **91** (1981), 257-274.
4. DRAGOMIR, S., Cauchy-Riemann submanifolds of locally conformal Kaehler manifold, Geometriae Dedicata **28** (1988), 181-197.
5. SHAHID, M.H. & HUSAIN, S.I., Generic submanifolds of a locally conformal Kaehler manifold, Soochow J. Math. **14** (1983), 111-117.
6. KASHIWADA, T., Some properties of locally conformal Kaehler manifolds, Hokkaido Math. J. **8** (1979), 191-198.
7. VAISMAN, I., On locally and globally conformal Kaehler manifolds, Trans. Amer. Math. Soc. **262** (1980), 533-542.