APPROXIMATING FIXED POINTS OF NONEXPANSIVE AND GENERALIZED NONEXPANSIVE MAPPINGS

M. MAITI and B. SAHA

Department of Mathematics
Indian Institute of Technology, Kharagpur, India

(Received May 20, 1991 and in revised form November 15, 1991)

ABSTRACT.

In this paper we consider a mapping S of the form

$$S = \alpha_0 I + \alpha_1 T + \alpha_2 T^2 + \dots + \alpha_k T^k,$$

where $\alpha_i \geq 0$, $\alpha_1 > 0$ with $\sum_{i=0}^k \alpha_i = 1$, and show that in a uniformly convex Banach space the Picard iterates of S converge to a fixed point of T when T is nonexpansive or generalized nonexpansive or even quasi-nonexpansive.

KEY WORDS AND PHRASES: Fixed points, nonexpansive mappings, uniformly convex Banach spaces.

1991 AMS SUBJECT CLASSIFICATION CODES. Primary 47H10; Secondary 54H25.

1. INTRODUCTION.

Let B be a Banach space and C a convex subset of B . A mapping $T: C \rightarrow C$ is said to be nonexpansive if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$. A mapping $T: C \rightarrow C$ is said to be quasi-nonexpansive if T has a fixed point p such that $\|Tx - p\| \leq \|x - p\|$ for all $x \in C$. The concept of quasi-nonexpansiveness is more general than that of nonexpansiveness. Indeed, a nonexpansive mapping with at least one fixed point is quasi-nonexpansive, but there exists quasi-nonexpansive mappings which are not nonexpansive. See, for example, Petryshyn and Williamson [7].

If T is nonexpansive, then the Picard iterates of T may not converge and, even if they do converge, they may not converge to a fixed point of T . However, to circumvent the difficulty, one may consider the mapping

$$T_\lambda = (1 - \lambda)I + \lambda T, \tag{1.1}$$

where I is the identity mapping and $0 < \lambda < 1$, and show that the Picard iterates of T_λ converge to a fixed point of T under certain restrictions, see [3,5,10]. Generalizing the idea Kirk [6] has introduced a mapping S given by

$$S = \alpha_0 I + \alpha_1 T + \alpha_2 T^2 + \dots + \alpha_k T^k \tag{1.2}$$

where $\alpha_i \geq 0$, $\alpha_1 > 0$ with $\sum_{i=0}^k \alpha_i = 1$, and has shown that the Picard iterates of S converge to a fixed point of T under conditions similar to those imposed in connection with the convergence of Picard iterates of T_λ .

Our purpose here is two-fold. First we show that the Picard iterates of S converge to a fixed point of T under conditions weaker than those imposed by Kirk [6]. Secondly, we establish that the Picard iterates of S converge to a fixed point of T even when T is generalized nonexpansive, i.e., when T satisfies

$$\|Tx - Ty\| \leq a\|x - y\| + b\{\|x - Tx\| + \|y - Ty\|\} + c\{\|y - Tx\| + \|x - Ty\|\} \quad (1.3)$$

for all $x, y \in C$, where $a, b, c \geq 0$ with $a + 2b + 2c \leq 1$. Then the analysis has been extended to a more general mapping resulting in generalization of some results obtained by Ray and Rhoades [9].

2. CONVERGENCE TO FIXED POINTS

It has been established by Kirk [6] that S and T have common fixed points if T is nonexpansive. Let the common fixed point set be denoted by F . Further, the set F is closed when T is nonexpansive or even when T is quasi-nonexpansive (see Dotson [2]). We now state the following conditions:

CONDITION-A. A mapping $T: C \rightarrow C$ with a nonempty fixed point set F is said to satisfy Condition-A if there is a nondecreasing function $f: [0, \infty) \rightarrow [0, \infty)$ with $f(0) = 0$ and $f(r) > 0$ for all $r \in (0, \infty)$ such that $\|x - Sx\| \geq f(d(x, F))$ for all $x \in C$, where $d(x, F) = \inf_{p \in F} \|x - p\|$.

CONDITION-B. A mapping $T: C \rightarrow C$ with a nonempty fixed point set F is said to satisfy Condition-B if there exists a number $\alpha > 0$ such that $\|x - Sx\| \geq \alpha d(x, F)$ for all $x \in C$.

It may be remarked that the mappings which satisfy Condition-B also satisfy Condition-A. However, Condition-B may be verified easily by giving examples. It may be further remarked that Conditions I and II of Senter and Dotson [11] are identical with Conditions A and B when $\alpha_2 = \alpha_3 = \dots = \alpha_k = 0$.

We now recall the following lemma due to Dotson [1]. This will be used later to establish our results.

LEMMA. If the sequences $\{s_n\}$ and $\{t_n\}$ are in the closed unit ball of a uniformly convex Banach space and $\{z_n\} = \{(1 - \alpha_n)s_n + \alpha_n t_n\}$ satisfies $\lim_{n \rightarrow \infty} \|z_n\| = 1$, where $0 < a \leq \alpha_n \leq b < 1$, then $\lim_{n \rightarrow \infty} \|s_n - t_n\| = 0$.

THEOREM 1. Let C be a nonempty, closed, convex and bounded subset of a uniformly convex Banach space B and $T: C \rightarrow C$ be a nonexpansive mapping. If T satisfies Condition-A, where F is the fixed point set of T in C , then for an arbitrary $x_0 \in C$, the Picard iterates $(S^n x_0)$ converge to a member of F .

PROOF. If $x_0 \in F$, then the result is trivial. We assume that $x_0 \in C - F$. Then, setting $x_n = S^n x_0$, we have for an arbitrary $p \in F$

$$\begin{aligned} \|x_{n+1} - p\| &= \|S^{n+1} x_0 - p\| = \|Sx_n - p\| \\ &= \|\alpha_0 x_n + \alpha_1 T x_n + \alpha_2 T^2 x_n + \dots + \alpha_k T^k x_n - p\| \\ &= \|\alpha_0(x_n - p) + \alpha_1(Tx_n - p) + \alpha_2(T^2 x_n - p) + \dots + \alpha_k(T^k x_n - p)\| \\ &\leq \alpha_0 \|x_n - p\| + \alpha_1 \|Tx_n - p\| + \alpha_2 \|T^2 x_n - p\| + \dots + \alpha_k \|T^k x_n - p\| \\ &\leq \|x_n - p\|. \end{aligned}$$

This implies that $d(x_{n+1}, F) \leq d(x_n, F)$ and hence that the sequence $\{d(x_n, F)\}$ is nonincreasing. Then $\lim_{n \rightarrow \infty} d(x_n, F)$ exists. In the sequel we shall show that this limit is zero.

Suppose that $\lim_{n \rightarrow \infty} d(x_n, F) = b > 0$. Then, for a $p \in F$, $\lim_{n \rightarrow \infty} \|x_n - p\| = b' \geq b > 0$. Choose a positive integer N such that $\|x_n - p\| \leq 2b'$ for $n \geq N$. Set $y_n^i = (T^i x_n - p) / \|x_n - p\|$ for all n and all $i = 0, 1, 2, \dots, k$ with $T^0 x_n = x_n$. Then $\|y_n^i\| \leq 1$. Further, set $z_n = \alpha_0 y_n^0 + (1 - \alpha_0) t_n$, where $t_n = \sum_{i=1}^k (\alpha_i y_n^i) / (1 - \alpha_0)$ with $\|t_n\| \leq 1$. Then $\{y_n^0\}$ and $\{T_n\}$ are in the closed unit ball. Now

$$\begin{aligned} \|z_n\| &= \left\| \alpha_0 \frac{x_n - p}{\|x_n - p\|} + \alpha_1 \frac{Tx_n - p}{\|x_n - p\|} + \alpha_2 \frac{T^2x_n - p}{\|x_n - p\|} + \dots + \alpha_k \frac{T^kx_n - p}{\|x_n - p\|} \right\| \\ &= \frac{\|\alpha_0x_n + \alpha_1Tx_n + \alpha_2T^2x_n + \dots + \alpha_kT^kx_n - p\|}{\|x_n - p\|} = \frac{\|x_{n+1} - p\|}{\|x_n - p\|}, \end{aligned} \quad (2.1)$$

implying $\|z_n\| \rightarrow 1$ as $n \rightarrow \infty$. But, for $n \geq N$, we have

$$\begin{aligned} \|y_n^0 - t_n\| &= \left\| \frac{x_n - p}{\|x_n - p\|} - \frac{1}{(1 - \alpha_0)} \sum_{i=1}^k \frac{\alpha_i(T^i x_n - p)}{\|x_n - p\|} \right\| \\ &= \left\| \frac{x_n - p}{\|x_n - p\|} - \frac{Sx_n - \alpha_0x_n - (1 - \alpha_0)p}{(1 - \alpha_0)\|x_n - p\|} \right\| \\ &= \frac{\|x_n - Sx_n\|}{(1 - \alpha_0)\|x_n - p\|} \\ &\geq \frac{f(d(x_n, F))}{(1 - \alpha_0)\|x_n - p\|} \geq \frac{f(b)}{2b'(1 - \alpha_0)} > 0, \end{aligned} \quad (2.2)$$

implying $\lim_{n \rightarrow \infty} \|y_n^0 - t_n\| \neq 0$, which contradicts the lemma. Hence $\lim_{n \rightarrow \infty} d(x_n, F) = 0$. This implies that $\{x_n\}$ converges to a member of F , since F is closed.

REMARK 1. It is obvious that the above theorem holds if Condition-B is satisfied instead of Condition-A.

REMARK 2. Condition-A is more general than the condition imposed by Kirk [6] in establishing the convergence of Picard iterates $\{S^n x_0\}$, see Senter and Dotson [11].

REMARK 3. In the above theorem the existence of a nonempty fixed point set F is not assumed and is ensured by the conditions assumed therein. However, if we assume that T has a nonempty fixed point set F , then T need not be assumed to be nonexpansive and it is enough for T to be quasi-nonexpansive. Further, C need not be bounded. Indeed, the following result holds.

THEOREM 2. Let C be a nonempty, closed and convex subset of a uniformly convex Banach space B and $T: C \rightarrow C$ be a quasi-nonexpansive mapping. If T satisfies Condition-A, where F is the fixed point set of T in C , then, for an arbitrary $x_0 \in C$, the Picard iterates $\{S^n x_0\}$ converge to a member of F .

The proof may be established exactly in the same way as in Theorem 1. It only remains to be shown here that S and T have common fixed points. A fixed point of T is obviously a fixed point of S . We now show that the converse is also so. Let p be a fixed point of S . Then from Condition-A it is obvious that $d(p, F) = 0$, implying $p \in \bar{F}$. Since T is quasi-nonexpansive, F is closed and hence $p \in F$, i.e., p is a fixed point of T . Hence the result.

Next, we show that the Picard iterates of S converge to a fixed point of T even when T is generalized nonexpansive. However, one need not assume Condition-A or Condition-B in this case. These conditions are automatically satisfied.

THEOREM 3. Let C be a nonempty, bounded, closed and convex subset of a uniformly convex Banach space B and $T: C \rightarrow C$ be a continuous mapping such that

$$\|Tx - Ty\| \leq a\|x - y\| + b\{\|x - Tx\| + \|y - Ty\|\} + c\{\|x - Ty\| + \|y - Tx\|\} \quad (2.3)$$

for all $x, y \in C$, where $a, c \geq 0$ and $b > 0$ with $a + 2b + 2c \leq 1$. Then for an arbitrary $x_0 \in C$, the Picard iterates $\{S^n x_0\}$ converges to the unique fixed point of T .

PROOF. By Theorem 2 of Goebel, Kirk and Shimi [4] the mapping T has a unique fixed point p , say. Setting $y = p$ in (2.3) we have

$$\begin{aligned}\|Tx - p\| &\leq (a + c)\|x - p\| + b\|x - Tx\| + c\|Tx - p\| \\ &\leq (a + b + c)\|x - p\| + (b + c)\|Tx - p\|,\end{aligned}$$

implying

$$\|Tx - p\| \leq \frac{a + b + c}{1 - b - c}\|x - p\| \leq \|x - p\|, \quad (2.4)$$

since $a + 2b + 2c \leq 1$. Thus T is quasi-nonexpansive. Further, it is easy to verify that

$$\|Sx - p\| \leq \|Tx - p\| \leq \|x - p\|, \quad (2.5)$$

implying S is also quasi-nonexpansive.

It is obvious that p is also a fixed point of S . We now show that S cannot have a fixed point other than p . If possible, let $q (\neq p)$ be a fixed point of S . Then

$$\begin{aligned}\|q - Tq\| &= \|Sq - Tq\| \\ &= \|\alpha_0 q + \alpha_1 Tq + \alpha_2 T^2 q + \dots + \alpha_k T^k q - Tq\| \\ &= \|\alpha_0(q - Tq) + \alpha_2(T^2 q - Tq) + \dots + \alpha_k(T^k q - Tq)\| \\ &\leq \alpha_0\|q - Tq\| + \alpha_2\|T^2 q - Tq\| + \dots + \alpha_k\|T^k q - Tq\| \\ &\leq \alpha_0\{\|q - p\| + \|Tq - p\|\} + \alpha_2\{T^2 q - p\| + \|Tq - p\|\} \\ &\quad + \dots + \alpha_k\{\|T^k q - p\| + \|Tq - p\|\} \\ &\leq 2(\alpha_0 + \alpha_2 + \dots + \alpha_k)\|q - p\| = 2(1 - \alpha_1)\|q - p\|.\end{aligned} \quad (2.6)$$

Since T is generalized nonexpansive, we have

$$\begin{aligned}\|Tq - p\| &= \|Tq - Tp\| \\ &\leq a\|q - p\| + b\|Tq - q\| + c\{\|Tq - p\| + \|q - p\|\} \\ &\leq (a + 2c)\|q - p\| + b\|Tq - q\|.\end{aligned} \quad (2.7)$$

Substituting from (2.6) into (2.7) and noting that $a + 2c \leq 1 - 2b$ we obtain

$$\begin{aligned}\|Tq - p\| &\leq (1 - 2b)\|q - p\| + 2b(1 - \alpha_1)\|q - p\| \\ &= \|q - p\| - 2b\alpha_1\|q - p\| \\ &< \|q - p\| - 2b\alpha_1\|Tq - p\|,\end{aligned}$$

implying

$$\|Tq - p\| \leq \frac{1}{1 + 2b\alpha_1}\|q - p\| \quad (2.8)$$

Now from (2.8) we have

$$\|q - p\| = \|Sq - p\| \leq \|Tq - p\| \leq \frac{1}{1 + 2b\alpha_1}\|q - p\|, \quad (2.9)$$

which implies $q = p$, since $b, \alpha_1 > 0$. Thus S and T have a unique fixed point p .

Next, we show that T satisfies Condition-B. For $x \in C$ we have

$$\|Tx - p\| = \|Tx - Tp\| \leq a\|x - p\| + b\|x - Tx\| + c\{\|x - p\| + \|Tx - p\|\},$$

implying

$$\|Tx - p\| \leq \frac{a + c}{1 - c}\|x - p\| + \frac{b}{1 - c}\|x - Tx\|. \quad (2.10)$$

Now,

$$\begin{aligned}\|Tx - x\| &\leq \|Sx - Tx\| + \|Sx - x\| \\ &\leq \alpha_0\|x - Tx\| + \alpha_2\|T^2x - Tx\| + \dots + \alpha_k\|T^kx - Tx\| + \|Sx - x\| \\ &\leq 2(1 - \alpha_1)\|x - p\| + \|Sx - x\|.\end{aligned}\quad (2.11)$$

Also we observe that

$$\|Sx - p\| \leq \alpha_0\|x - p\| + (\alpha_1 + \alpha_2 + \dots + \alpha_k)\|Tx - p\| \quad (2.12)$$

and that

$$\|Sx - x\| \geq \|x - p\| - \|Sx - p\|. \quad (2.13)$$

Now, substituting from (2.12) into (2.13) we derive

$$\begin{aligned}\|Sx - x\| &\geq \|x - p\| - \alpha_0\|x - p\| - (\alpha_1 + \alpha_2 + \dots + \alpha_k)\|Tx - p\| \\ &= (1 - \alpha_0)\{\|x - p\| - \|Tx - p\|\},\end{aligned}$$

whence, using (2.10), we obtain

$$\begin{aligned}\|Sx - x\| &\geq (1 - \alpha_0)\left\{\|x - p\| - \frac{a+c}{1-c}\|x - p\| - \frac{b}{1-c}\|x - Tx\|\right\} \\ &= (1 - \alpha_0)\left\{\frac{1-a-2c}{1-c}\|x - p\| - \frac{b}{1-c}\|x - Tx\|\right\} \\ &\geq (1 - \alpha_0)\left\{\frac{2b}{1-c}\|x - p\| - \frac{b}{1-c}\|x - Tx\|\right\} \\ &= \frac{b(1 - \alpha_0)}{1 - c}\{2\|x - p\| - \|x - Tx\|\}.\end{aligned}\quad (2.14)$$

Now, substituting from (2.11) into (2.14) we get

$$\begin{aligned}\|Sx - x\| &\geq \frac{b(1 - \alpha_0)}{1 - c}\{2\|x - p\| - 2(1 - \alpha_1)\|x - p\| - \|Sx - x\|\} \\ &= \frac{b(1 - \alpha_0)}{1 - c}\{2\alpha_1\|x - p\| - \|Sx - x\|\},\end{aligned}$$

implying

$$\|Sx - x\| \geq \alpha\|x - p\|, \quad (2.15)$$

where

$$\alpha = \frac{2b\alpha_1(1 - \alpha_0)}{1 - c + b(1 - \alpha_0)} > 0,$$

since $b, \alpha_1 > 0$. Thus T satisfies Condition-B. Hence, by Theorem 2, the result follows.

REMARK 4. It may be noted that the stipulation $\alpha_1 > 0$ in S is necessary to rule out the possibility that fixed point of S is a point at which T may be periodic.

REMARK 5. If we do not restrict $b > 0$ in Theorem 3, then the fixed pint set of T is not a singleton, and Condition-A is to be imposed to ensure the convergence of $\{S^n x_0\}$.

The present analysis can be extended to a more general mapping T which satisfies

$$\|Tx - Ty\| \leq \max \{\|x - y\|, [\|x - Tx\| + \|y - Ty\|]/2, [\|y - Tx\| + \|x - Ty\|]/2\} \quad (2.16)$$

for all $x, y \in C$. This mapping includes nonexpansive and generalized nonexpansive mappings (see Rhoades [8]). It is easy to verify that T is quasi-nonexpansive. It has been proved by Ray and Rhoades [9] that S and T have the same fixed point set. Further, they have established the following theorem in this connection.

THEOREM 4. ([9, Theorem 2]). Let C be a nonempty, closed convex and bounded subset of a uniformly convex Banach space B and T a self-mapping of C which satisfies (2.16). If I - S maps bounded closed subsets of C into closed sets of B , then, for each $x_0 \in C$, the sequence $\{S^n x_0\}$ converges to a fixed point of T in C .

However, the fact that I - S maps bounded closed subsets of C into closed sets implies Condition-A (see Senter and Dotson [11]). Thus Condition-A is more general and incorporating this condition we may obtain the following as generalizations of Theorems 2 and 3 of Ray and Rhoades [9].

THEOREM 5. Let C be a nonempty, closed and convex subset of a uniformly convex Banach space B and T a self-mapping of C which satisfies (2.16). If T satisfies Condition-A, where F is the nonempty fixed point set of T in C , then, for an arbitrary $x_0 \in C$, the Picard iterates $\{S^n x_0\}$ converge to a member of F .

We may note that C need not be bounded in Theorem 5. Because we have assumed the existence of nonempty fixed point set of T and Condition-A (see [11]). But the boundedness of C cannot be omitted from the statement of Theorem 4.

ACKNOWLEDGEMENT. The authors are indebted to Professor B. E. Rhoades for his valuable suggestions for the improvement of the paper.

REFERENCES

- [1] DOTSON, JR., W. G. On the Mann iterative process, Trans. Amer. Math. Soc. **149** (1970), 65-73.
- [2] DOTSON, JR., W. G. Fixed points of quasi-nonexpansive mappings, J. Austral. Math. Soc. **13** (1972), 167-170.
- [3] EDELSTEIN, M. A remark on a theorem of M. A. Krasnoselskii, Amer. Math. Monthly **73** (1966), 509-510.
- [4] GOEBEL, K., KIRK, W. A. and SHIMI, T. N. A fixed point theorem in uniformly convex Banach spaces, Boll. Un. Mat. Ital. **7** (1973), 67-75.
- [5] KRASNOSELSKII, M. A. Two remarks about the method of successive approximations, Uspehi Mat. Nauk **10** (1955), 123-127.
- [6] KIRK, W. A. On successive approximations for nonexpansive mappings in Banach spaces, Glasgow Math. Jour. **12** (1971), 6-9.
- [7] PETRYSHYN, W. V. and WILLIAMSON, JR., T. E. Strong and weak convergence of the sequence of successive approximations for quasi-nonexpansive mappings, J. Math. Anal. and Appl. **43** (1973), 459-497.
- [8] RHOADES, B. E. A comparison of various definitions of contractive mappings, Trans. Amer. Math. Soc. **226** (1977), 257-290.
- [9] RAY, B. K. and RHOADES, B. E. A class of fixed point theorems, Math. Semi. Notes **7** (1979), 477-489.
- [10] SCHAEFER, H. Ober Die Methode seksessiver Approximation, Jber. Deutsch. Math. Verein. **59** (1957), 131-140.
- [11] SENTER, H. F. and DOTSON, JR., W. G. Approximating fixed points of nonexpansive mappings, Proc. Amer. Math. Soc. **44** (1974), 375-380.