ON CERTAIN CLASSES OF p -VALENT ANALYTIC FUNCTIONS

NAK EUN CHO

Department of Applied Mathematics
National Fisheries University of Pusan
Pusan 608-737
KOREA

(Received November 5, 1991)

ABSTRACT. The objective of the present paper is to introduce a certain general class $P(p, \alpha, \beta)$ ($p \in N = \{1, 2, 3, \dots\}$, $0 \leq \alpha < p$ and $\beta \geq 0$) of p -valent analytic functions in the open unit disk U and we prove that if $f \in P(p, \alpha, \beta)$ then $J_{p,c}(f)$, defined by

$$J_{p,c}(f) = \frac{c+p}{z^c} \int_0^z t^{c-1} f(t) dt \quad (c \in N),$$

belongs to $P(p, \alpha, \beta)$. We also investigate inclusion properties of the class $P(p, \alpha, \beta)$. Furthermore, we examine some properties for a class $T_p(\alpha, \beta)$ of analytic functions with negative coefficients.

KEY WORDS AND PHRASES. p -valent analytic function, Hadamard product, integral operator, multiplier transformation, p -valently convex of order δ .

1991 AMS SUBJECT CLASSIFICATION CODE. Primary 30C45.

1. INTRODUCTION.

Let A_p denote the class of functions of the form

$$f(z) = z^p + \sum_{n=1}^{\infty} a_{n+p} z^{n+p} \quad (p \in N = \{1, 2, 3, \dots\}) \quad (1.1)$$

which are analytic in the unit disk $U = \{z: |z| < 1\}$. We also denote by S_p the subclass of A_p consisting of functions which are p -valent in U .

A function $f \in A_p$ is said to be in the class $P(p, \alpha)$ ($0 \leq \alpha < p$) if and only if it satisfies the inequality

$$Re \left\{ \frac{f(z)}{z^{p-1}} \right\} > \alpha \quad (0 \leq \alpha < p, z \in U). \quad (1.2)$$

The classes $P(1, 0)$ and $P(p, 0)$ were investigated by MacGregor [7] and Umezawa [11], respectively. In fact, the class $P(p, \alpha)$ is a subclass of the class S_p [11].

Let f and g be in the class A_p , with $f(z)$ given by (1.1), and $g(z)$ defined by

$$g(z) = z^p + \sum_{n=1}^{\infty} b_{n+p} z^{n+p}. \quad (1.3)$$

The convolution or Hadamard product of f and g is defined by

$$(f * g)(z) = z^p + \sum_{n=1}^{\infty} a_{n+p} b_{n+p} z^{n+p}. \quad (1.4)$$

For a function $f \in A_p$ given by (1.1), Reddy and Padmanabhan [10] defined the integral operator $J_{p,c}$ ($p, c \in N$) by

$$\begin{aligned} J_{p,c}(f) &= \frac{c+p}{z^c} \int_0^z t^{c-1} f(t) dt \\ &= z^p + \sum_{n=1}^{\infty} \frac{c+p}{c+p+n} a_{n+p} z^{n+p}. \end{aligned} \quad (1.5)$$

The operator $J_{1,c}$ was introduced by Bernardi [2]. In particular, the operator $J_{1,1}$ were studied by Libera [5] and Livingston [6].

Clearly, (1.5) yields

$$f \in A_p \Rightarrow J_{p,c} \in A_p \quad (1.6)$$

Thus, by applying the operator $J_{p,c}$ successively, we can obtain

$$J_{p,c}^n(f) = \begin{cases} J_{p,c}(J_{p,c}^{n-1}(f)) & (n \in N), \\ f(z) & (n = 0). \end{cases} \quad (1.7)$$

We now recall the following definition of a multiplier transformation (or fractional integral and fractional derivative).

DEFINITION 1([3]). Let the function

$$\phi(z) = \sum_{n=0}^{\infty} c_{n+p} z^{n+p} \quad (1.8)$$

be analytic in U and let λ be a real number. Then the multiplier transformation $I^\lambda \phi$ is defined by

$$I^\lambda \phi(z) = \sum_{n=0}^{\infty} (n+p+1)^{-\lambda} c_{n+p} z^{n+p} \quad (z \in U). \quad (1.9)$$

The function $I^\lambda \phi$ is clearly analytic in U . It may be regarded as a fractional integral (for $\lambda > 0$) or fractional derivative (for $\lambda < 0$) of ϕ . Furthermore, in terms of the Gamma function, we have

$$I^\lambda \phi(z) = \frac{1}{\Gamma(\lambda)} \int_0^1 (\log \frac{1}{t})^{\lambda-1} \phi(zt) dt \quad (\lambda > 0). \quad (1.10)$$

DEFINITION 2. The fractional derivative $D^\lambda \phi$ of order $\lambda \geq 0$, for an analytic function ϕ given by (1.8), is defined by

$$D^\lambda \phi(z) = I^{-\lambda} \phi(z) = \sum_{n=0}^{\infty} (n+p+1)^\lambda c_{n+p} z^{n+p} \quad (\lambda \geq 0, z \in U). \quad (1.11)$$

Making use of Definition 2, we now introduce an interesting generalization of the class $P(p, \alpha)$ of functions in A_p which satisfy the inequality (1.2).

DEFINITION 3. A function $f \in A_p$ is said to be in the class $P(p, \alpha, \beta)$ if and only if

$$(p+1)^{-\beta} D^\beta f \in P(p, \alpha) \quad (0 \leq \alpha < p, \beta \geq 0)$$

Observe that $P(p, \alpha, 0) = P(p, \alpha)$. Furthermore, since $f \in A_p$, it follows from (1.1) and (1.9) that

$$(p+1)^{-\beta} D^\beta f(z) = z^p + \sum_{n=1}^{\infty} \left(\frac{n+p+1}{p+1} \right)^\beta a_{n+p} z^{n+p}, \quad (1.12)$$

which shows that $(p+1)^{-\beta} D^\beta f \in A_p$ if $f \in A_p$. In particular, the class $P(1, \alpha, \beta)$ was introduced by Kim, Lee, and Srivastava [4].

2. SOME INCLUSION PROPERTIES.

In our present investigation of the general class $P(p, \alpha, \beta)$ ($0 \leq \alpha < p, \beta \geq 0$), we need the following lemma.

LEMMA 2.1([1]). Let $M(z)$ and $N(z)$ be analytic in U , $N(z)$ map U onto a many sheeted starlike region of order γ ($0 \leq \gamma < p$) and

$$M(0) = N(0) = 0, \quad \frac{M'(0)}{N'(0)} = p, \quad \operatorname{Re} \left(\frac{M'(z)}{N'(z)} \right) > \gamma.$$

Then we have

$$\operatorname{Re} \left(\frac{M(z)}{N(z)} \right) > \gamma \quad (0 \leq \gamma < p, p \geq 1).$$

By using Lemma 2.1, we can prove

THEOREM 2.1. Let the function $f(z)$ be in the class $P(p, \alpha, \beta)$. Then $J_{p,c}(f)$ defined by (1.5) is also in the class $P(p, \alpha, \beta)$.

PROOF. A simple calculation shows that

$$\frac{d}{dz} D^\beta (J_{p,c}(f)) = \frac{c+p}{z^{c+p}} \int_0^z t^c \left\{ \frac{d}{dt} D^\beta f(t) \right\} dt \quad (2.1)$$

where the operators $J_{p,c}$ ($c \in N$) and D^λ ($\lambda \geq 0$) are defined by (1.5) and (1.11), respectively. In view of (2.1), we get

$$M(z) = \frac{c+p}{(p+1)^\beta} \int_0^z t^c \left\{ \frac{d}{dt} D^\beta f(t) \right\} dt \text{ and } N(z) = z^{p+c}, \quad (2.2)$$

so that

$$\operatorname{Re} \left\{ \frac{M'(z)}{N'(z)} \right\} = \operatorname{Re} \left\{ \frac{(p+1)^{-\beta} \frac{d}{dz} D^\beta f(z)}{z^{p-1}} \right\}. \quad (2.3)$$

Since, by hypothesis, $f \in P(p, \alpha, \beta)$, the second member of (2.3) is greater than α , and hence

$$\operatorname{Re} \left\{ \frac{M'(z)}{N'(z)} \right\} > \alpha \quad (0 \leq \alpha < p). \quad (2.4)$$

Thus, by Lemma 2.1, we have

$$\operatorname{Re} \left\{ \frac{M(z)}{N(z)} \right\} = \operatorname{Re} \left\{ \frac{(p+1)^{-\beta} \frac{d}{dz} D^\beta (J_{p,c}(f))}{z^{p-1}} \right\} > \alpha \quad (0 \leq \alpha < p, \beta \geq 0), \quad (2.5)$$

which completes the proof of Theorem 2.1.

Let $f \in A_p$ be given by (1.1). Suppose also that

$$\begin{aligned}
 F_m(f) &= J_{p, c_1} \left(\dots \left(J_{p, c_m}(f) \right) \right) \\
 &= z^p + \sum_{n=1}^{\infty} \frac{(c_1+p) \dots (c_m+p)}{(c_1+p+n) \dots (c_m+p+n)} a_{n+p} z^{n+p} \quad (c_j \in N (j=1, 2, \dots, m), m \in N).
 \end{aligned} \tag{2.6}$$

Then, by Theorem 2.1, we have

COROLLARY 2.1. Let the function $f(z)$ be in the class $P(p, \alpha, \beta)$. Then the function $F_m(f)$ defined by (2.6) is also in the class $P(p, \alpha, \beta)$.

The next inclusion property of the class $P(p, \alpha, \beta)$, contained in Theorem 2.2 below, would involve the operator $J_{p,1}^\lambda (\lambda > 0)$ defined by

$$J_{p,1}^\lambda(f) = (1+p)^\lambda I^\lambda f(z) \quad (\lambda > 0, f \in A_p). \tag{2.7}$$

For $\lambda = m \in N$, we have

$$\begin{aligned}
 J_{p,1}^m(f) &= (1+p)^m I^m f(z) \\
 &= \frac{(1+p)^m}{(m-1)!} \int_0^1 (\log \frac{1}{t})^{m-1} f(t) dt.
 \end{aligned} \tag{2.8}$$

Clearly, we have

$$f \in A_p \Rightarrow J_{p,1}^\lambda(f) \in A_p \quad (\lambda > 0). \tag{2.9}$$

THEOREM 2.2. Let the function $f(z)$ be in the class $P(p, \alpha, \beta)$. Then the function $J_{p,1}^\lambda (\lambda > 0)$ defined by (2.7) is also in the class $P(p, \alpha, \beta)$.

PROOF. Making use of (1.9) and (1.11), the definition (2.7) yields

$$(p+1)^{-\beta} D^\beta (J_{p,1}^\lambda(f)) = J_{p,1}^\lambda((p+1)^{-\beta} D^\beta f) \quad (\beta \geq 0, \lambda > 0, f \in A_p) \tag{2.10}$$

Therefore, setting

$$g(z) = (p+1)^{-\beta} D^\beta f \text{ and } G(z) = J_{p,1}^\lambda(g), \tag{2.11}$$

we must show that

$$Re \left\{ \frac{G'(z)}{z^{p-1}} \right\} > \alpha \quad (0 \leq \alpha < p) \tag{2.12}$$

whenever $f \in P(p, \alpha, \beta)$.

From the integral representation in (1.10), we obtain

$$G'(z) = \frac{(p+1)^\lambda}{\Gamma(z)} \int_0^1 (\log \frac{1}{t})^{\lambda-1} t g'(zt) dt \quad (\lambda > 0), \tag{2.13}$$

so that

$$Re \left\{ \frac{G'(z)}{z^{p-1}} \right\} = \frac{(p+1)^\lambda}{\Gamma(\lambda)} \int_0^1 (\log \frac{1}{t})^{\lambda-1} t^p Re \left\{ \frac{g'(zt)}{(zt)^{p-1}} \right\} dt \quad (\lambda > 0), \tag{2.14}$$

Since $f \in P(p, \alpha, \beta)$, we have

$$Re \left\{ \frac{g'(zt)}{(zt)^{p-1}} \right\} > \alpha \quad (0 \leq \alpha < p, 0 \leq t \leq 1), \tag{2.15}$$

and hence (2.14) yields

$$\operatorname{Re}\left\{\frac{G'(z)}{z^{p-1}}\right\} = \frac{(p+1)^\lambda}{\Gamma(\lambda)} \alpha \int_0^1 (\log \frac{1}{t})^{\lambda-1} t^p dt = \alpha \quad (0 \leq \alpha < p, \lambda > 0), \quad (2.16)$$

which completes the proof of Theorem 2.2.

COROLLARY 2.2. If $0 \leq \alpha < p$ and $0 \leq \beta < \gamma$, then $P(p, \alpha, \gamma) \subset P(p, \alpha, \beta)$.

PROOF. Setting $\lambda = \gamma - \beta > 0$ in Theorem 2.2, we observe that

$$\begin{aligned} f \in P(p, \alpha, \gamma) &\Rightarrow J_{p,1}^{\gamma-\beta}(f) \in P(p, \alpha, \gamma) \\ &\Leftrightarrow (p+1)^{-\gamma} D^\gamma (J_{p,1}^{\gamma-\beta}(f)) \in P(p, \alpha) \\ &\Leftrightarrow (p+1)^{-\beta} D^\beta f \in P(p, \alpha) \\ &\Leftrightarrow f \in P(p, \alpha, \beta), \end{aligned}$$

and the proof of Corollary 2.2 is completed.

Next we define a function $h \in A_p$ by

$$h(z) = z^p + \sum_{n=1}^{\infty} \left(\frac{n+p+1}{p+1} \right) z^{n+p} \quad (z \in U). \quad (2.18)$$

Then, in terms of the Hadamard product defined by (1.4), we have

$$(h * f)(z) = \frac{1}{p+1} \{f(z) + z f'(z)\} \quad (2.19)$$

which, when compared with (1.11) with $m = 1$, yields

$$(h * f)(z) = \frac{1}{p+1} D^1 f. \quad (2.20)$$

We now need the following lemma for another inclusion property of the class $P(p, \alpha, \beta)$.

LEMMA 2.2([8]). Let $\phi(u, v)$ be a complex valued function such that

$$\phi: D \rightarrow C, \quad D \subset C \times C (C \text{ is the complex plane}),$$

and let $u = u_1 + iu_2$, $v = v_1 + iv_2$. Suppose that the function $\phi(u, v)$ satisfies

- (i) $\phi(u, v)$ is continuous in D ,
- (ii) $(1, 0) \in D$ and $\operatorname{Re}\{\phi(1, 0)\} > 0$,
- (iii) for all $(iu_2, v_1) \in D$ such that $v_1 \leq -\frac{1+u_2^2}{2}$, $\operatorname{Re}\{\phi(iu_2, v_1)\} \leq 0$.

Let $p(z) = 1 + p_1 z + p_2 z^2 + \dots$ be analytic in the unit disk U such that $(p(z), zp'(z)) \in D$ for all $z \in U$. If

$$\operatorname{Re}\{(p(z), zp'(z))\} > 0 \quad (z \in U),$$

then $\operatorname{Re}\{p(z)\} > 0 (z \in U)$.

THEOREM 2.3. If $0 \leq \alpha < p$ and $\beta \geq 0$, then

$$P(p, \alpha, \beta + 1) \subset P(p, \mu, \beta) \quad \left(\mu = \frac{2\alpha(p+1) + p}{2(p+1) + 1} \right) \quad (2.21)$$

PROOF. Let the function

$$F(z) = \frac{1}{p+1} \{f(z) + z f'(z)\} \quad (f \in A_p). \quad (2.22)$$

First, we shall show that

$$Re\left\{\frac{f'(z)}{z^{p-1}}\right\} > \frac{2\alpha(p+1)+p}{2(p+1)+1} \quad (0 \leq \alpha < p, z \in U), \quad (2.23)$$

whenever

$$Re\left\{\frac{F'(z)}{z^{p-1}}\right\} > \alpha \quad (0 \leq \alpha < p, z \in U). \quad (2.24)$$

By the differentiation of $F(z)$, we obtain

$$F'(z) = \frac{1}{p+1}\{2f'(z) + zf''(z)\}. \quad (2.25)$$

We define the function $p(z)$ by

$$\frac{f'(z)}{pz^{p-1}} = \gamma + (1-\gamma)p(z) \quad (2.26)$$

with $\gamma = \frac{2\alpha(p+1)+p}{2p(p+1)+p}$ ($0 \leq \gamma < 1$). Then $p(z) = 1 + p_1z + p_2z^2 + \dots$ is analytic in U . By using (2.25) and (2.26), we obtain

$$\frac{F'(z)}{z^{p-1}} = \frac{1}{p+1} \{(p^2+p)(\gamma + (1-\gamma)p(z)) + p(1-\gamma)zp'(z)\}. \quad (2.27)$$

Hence, in view of $Re\left\{\frac{F'(z)}{z^{p-1}}\right\} > \alpha$ ($0 \leq \alpha < p$), we have

$$Re\{\phi(p(z), zp'(z))\} > 0, \quad (2.28)$$

where $\phi(u, v)$ is defined by

$$\phi(u, v) = \frac{1}{p+1}\{(p^2+p)(\gamma + (1-\gamma)u) + p(1-\gamma)v\} - \alpha \quad (2.29)$$

with $u = u_1 + iu_2$ and $v = v_1 + iv_2$. Then we see that

- (i) $\phi(u, v)$ is continuous in $D = C \times C$,
- (ii) $(1, 0) \in D$ and $Re\{\phi(1, 0)\} = p - \alpha > 0$,
- (iii) for all $(iu_2, v_1) \in D$ such that $v_1 \leq \frac{-(1+u_2^2)}{2}$,

$$\begin{aligned} Re\{\phi(iu_2, v_1)\} &= \frac{1}{p+1}\{(p^2+p)\gamma + p(1-\gamma)v_1\} - \alpha \\ &\leq \frac{1}{p+1}\left\{(p^2+p)\gamma - \frac{p(1-\gamma)(1+u_2^2)}{2}\right\} - \alpha \leq 0 \end{aligned}$$

for $\gamma = \frac{2\alpha(p+1)+p}{2p(p+1)+p}$. Consequently, $\phi(u, v)$ satisfies the conditions in Lemma 2.2. Therefore, we have

$$Re\left\{\frac{f'(z)}{z^{p-1}}\right\} > p\gamma = \frac{2\alpha(p+1)+p}{2(p+1)+1}. \quad (2.30)$$

Next, in view of (2.20) and above arguments, we have

$$\begin{aligned} f \in P(p, \alpha, \beta+1) &\Leftrightarrow (p+1)^{-\beta-1}D^{\beta+1}f \in P(p, \alpha) \\ &\Rightarrow h * \{(p+1)^{-\beta}D^{\beta}f\} \in P(p, \alpha) \\ &\Rightarrow (p+1)^{-\beta}D^{\beta}f \in P(p, \mu) \quad \left(\mu = \frac{2\alpha(p+1)+p}{2(p+1)+1}\right) \\ &\Leftrightarrow f \in P(p, \mu, \beta), \end{aligned} \quad (2.31)$$

which evidently proves Theorem 2.3.

REMARK. Since $0 \leq \alpha < p$, we have

$$\mu = \frac{2\alpha(p+1)+p}{2(p+1)+1} > \alpha,$$

and hence $P(p, \mu, \beta) \subset P(p, \alpha, \beta)$.

3. THE CONVERSE PROBLEM.

Let T_p denote the class of functions of the form

$$f(z) = z^p - \sum_{n=1}^{\infty} a_{n+p} z^{n+p} \quad (p \in N = \{1, 2, 3, \dots\}, a_{n+p} \geq 0)$$

which are analytic in U and let $T_p(\alpha, \beta) = T_p \cap P(p, \alpha, \beta)$.

In this section, we investigate the converse problem of integrals defined by (1.5) for the class $T_p(\alpha, \beta)$.

LEMMA 3.1. Let $f \in T_p$. Then $f \in T_p(\alpha, \beta)$ if and only if

$$\sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+1}{p+1} \right)^{\beta} a_{n+p} \leq p - \alpha. \quad (3.1)$$

PROOF. Suppose that

$$\sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+1}{p+1} \right)^{\beta} a_{n+p} \leq p - \alpha.$$

It is sufficient to show that the values for $\frac{(p+1)^{-\beta}(D^{\beta}f)'}{z^{p-1}}$ lie in a circle centered at p whose radius is $p - \alpha$. Indeed, we have

$$\begin{aligned} \left| \frac{(p+1)^{-\beta}(D^{\beta}f)'}{z^{p-1}} - p \right| &= \left| - \sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+1}{p+1} \right)^{\beta} a_{n+p} z^n \right| \\ &\leq \sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+1}{p+1} \right)^{\beta} a_{n+p} |z|^n \\ &< \sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+1}{p+1} \right)^{\beta} a_{n+p} \leq p - \alpha. \end{aligned} \quad (3.2)$$

Conversely, assume that

$$\operatorname{Re} \left\{ \frac{(p+1)^{-\beta}(D^{\beta}f)'}{z^{p-1}} \right\} > \alpha \quad (0 \leq \alpha < p), \quad (3.3)$$

which is equivalent to

$$\operatorname{Re} \left\{ \sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+1}{p+1} \right)^{\beta} a_{n+p} z^n \right\} < p - \alpha. \quad (3.4)$$

Choose values of z on the real axis so that

$$\sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+1}{p+1} \right)^{\beta} a_{n+p} z^n$$

is real. Letting $z \rightarrow 1$ along the real axis, we obtain

$$\sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+1}{p+1} \right)^{\beta} a_{n+p} \leq p - \alpha.$$

The proof is completed.

THEOREM 3.1. Let $F \in T_p(\alpha, \beta)$ and $f(z) = \left[\frac{z^{1-c}}{p+c} \right] [z^c F(z)]'$ ($c \in N$). Then the function $f(z)$ belongs to the class $T_p(\delta, \beta)$ ($0 \leq \delta < p$) for $|z| < r$, where

$$r = \inf_{n \geq 1} \left[\frac{(p-\delta)(p+c)}{(p-\alpha)(n+p+c)} \right]^{\frac{1}{n}}. \quad (3.5)$$

The result is sharp.

PROOF. Let $F(z) = z^p - \sum_{n=1}^{\infty} a_{n+p} z^{n+p}$. Then it follows from (1.5) that

$$\begin{aligned} f(z) &= \left[\frac{z^{1-c}}{p+c} \right] \frac{d}{dz} [z^c F(z)] \\ &= z^p - \sum_{n=1}^{\infty} \left(\frac{n+p+c}{p+c} \right) a_{n+p} z^{n+p}. \end{aligned} \quad (3.6)$$

To prove the result, it suffices to show that

$$\left| \frac{(p+1)^{-\beta} (D^\beta f)'}{z^{p-1}} - p \right| \leq p - \delta \quad (3.7)$$

for $|z| \leq r$. Now

$$\begin{aligned} \left| \frac{(p+1)^{-\beta} (D^\beta f)'}{z^{p-1}} - p \right| &= \left| - \sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+1}{p+1} \right)^\beta \left(\frac{n+p+c}{p+c} \right) a_{n+p} z^n \right| \\ &\leq \sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+1}{p+1} \right)^\beta \left(\frac{n+p+c}{p+c} \right) a_{n+p} |z|^n. \end{aligned} \quad (3.8)$$

Thus we have

$$\left| \frac{(p+1)^{-\beta} (D^\beta f)'}{z^{p-1}} - p \right| \leq p - \delta \quad (3.9)$$

if

$$\sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+1}{p+1} \right)^\beta \left(\frac{n+p+c}{p+c} \right) a_{n+p} |z|^n \leq p - \delta. \quad (3.10)$$

But Lemma 3.1 confirms that

$$\sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+1}{p+1} \right)^\beta a_{n+p} \leq p - \alpha. \quad (3.11)$$

Therefore (3.10) will be satisfied if

$$\left(\frac{n+p}{p-\delta} \right) \left(\frac{n+p+c}{p+c} \right) |z|^n \leq \left(\frac{n+p}{p-\alpha} \right) \quad (3.12)$$

for each $n \in N$, or if

$$|z| \leq \left[\left(\frac{p-\delta}{p-\alpha} \right) \left(\frac{p+c}{n+p+c} \right) \right]^{\frac{1}{n}}. \quad (3.13)$$

The required result follows now from (3.13). Sharpness follows if we take

$$F(z) = z^p - \left(\frac{p-\alpha}{n+p} \right) \left(\frac{p+1}{n+p+1} \right)^\beta z^{n+p} \quad (3.14)$$

for each $n \in N$.

THEOREM 3.2. Let $F \in T_p(\alpha, \beta)$ and $f(z) = \left[\frac{z^{1-c}}{p+c} \right] [z^c F(z)]'$ ($c \in N$). Then the function $f(z)$ p -valently convex of order δ ($0 \leq \delta < p$) in the disk

$$|z| < r^* = \inf_{n \geq 1} \left[\frac{p(p-\delta)}{(n+p+\delta)(p-\alpha)} \left(\frac{n+p+c}{p+c} \right) \left(\frac{n+p+1}{p+1} \right)^\beta \right]^{\frac{1}{n}}. \quad (3.15)$$

The result is sharp.

PROOF. To prove the theorem, it is sufficient to show that

$$\left| \left(1 + \frac{zf''(z)}{f'(z)} \right) - p \right| \leq p - \delta \quad (3.16)$$

for $|z| \leq r^*$. In view of (3.6), we have

$$\begin{aligned} \left| \left(1 + \frac{zf''(z)}{f'(z)} \right) - p \right| &= \left| \frac{-\sum_{n=1}^{\infty} n(n+p) \left(\frac{n+p+c}{p+c} \right) a_{n+p} z^{n+p-1}}{(p - \sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+c}{p+c} \right) a_{n+p} z^n) z^{p-1}} \right| \\ &\leq \frac{\sum_{n=1}^{\infty} n(n+p) \left(\frac{n+p+c}{p+c} \right) a_{n+p} |z|^n}{p - \sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+c}{p+c} \right) a_{n+p} |z|^n}. \end{aligned} \quad (3.17)$$

Thus

$$\left| \left(1 + \frac{zf''(z)}{f'(z)} \right) - p \right| \leq p - \delta \quad (3.18)$$

if

$$\frac{\sum_{n=1}^{\infty} n(n+p) \left(\frac{n+p+c}{p+c} \right) a_{n+p} |z|^n}{p - \sum_{n=1}^{\infty} (n+p) \left(\frac{n+p+c}{p+c} \right) a_{n+p} |z|^n} \leq p - \delta, \quad (3.19)$$

or

$$\sum_{n=1}^{\infty} \frac{(n+p)(n+p+\delta)}{p(p-\delta)} \left(\frac{n+p+c}{p+c} \right) a_{n+p} |z|^n \leq 1. \quad (3.20)$$

But from Lemma 3.1, we obtain

$$\sum_{n=1}^{\infty} \left(\frac{n+p}{p-\alpha} \right) \left(\frac{n+p+1}{p+1} \right)^\beta a_{n+p} \leq 1. \quad (3.21)$$

Hence $f(z)$ is p -valently convex of order δ ($0 \leq \delta < p$) if

$$\frac{(n+p)(n+p+\delta)}{p(p-\delta)} \left(\frac{n+p+c}{p+c} \right) |z|^n \leq \left(\frac{n+p}{p-\alpha} \right) \left(\frac{n+p+1}{p+1} \right)^\beta, \quad (3.22)$$

or

$$|z| \leq \left[\frac{p(p-\delta)}{(n+p+\delta)(p-\alpha)} \left(\frac{p+c}{n+p+c} \right) \left(\frac{n+p+1}{p+1} \right)^\beta \right]^{\frac{1}{n}} \quad (3.23)$$

for each $n \in \mathbb{N}$. This completes the proof of the theorem. The result is sharp for the function given by (3.14).

REFERENCES

1. AOUF, M.K., OBRADOVIC, M. & OWA, S., On certain class of p -valent functions with reference to the Bernardi integral operator, *Math. Japonica* 35 (1990), 839-848.
2. BERNARD, S.D., Convex and starlike univalent functions, *Trans. Amer. Math. Soc.* 135 (1969), 429-446.

3. FLETT, T.M., The dual of an inequality of Hardy and Littlewood and some related inequalities, J. Math. Anal. Appl. **38** (1972), 746-765.
4. KIM, Y.C., LEE, S.H., & SRIVASTAVA, H.M., Some properties of convolution operators in the class $P_\alpha(\beta)$, submitted.
5. LIBERA, R.J., Some classes of regular univalent functions, Proc. Amer. Math. Soc. **16** (1965), 755-758.
6. LIVINGSTON, A.E., On the radius of univalence of certain analytic functions, Proc. Amer. Math. Soc. **17** (1966), 352-357.
7. MACGREGOR, T.H., Functions whose derivatives has a positive real part, Trans. Amer. Math. Soc. **104** (1962), 532-537.
8. MILLER, S.S., Differential inequalities and Caratheodory functions, Bull. Amer. Math. Soc. **81** (1975), 79-81.
9. OWA, S. & NUNOKAWA, M., Properties of certain analytic functions, Math. Japonica **33** (1988), 577-582.
10. REDDY, G.L. & PADMANABHAN, K.S., On analytic functions with reference to the Bernardi integral operator, Bull. Austral. Math. Soc. **25** (1982), 387-396.
11. UMEZAWA, T., Multivalently close-to-convex functions, Proc. Amer. Math. Soc. **8** (1957), 869-874.