CHARACTERISTIC POLYNOMIALS OF SOME WEIGHTED GRAPH BUNDLES AND ITS APPLICATION TO LINKS

MOO YOUNG SOHN

Department of Mathematics
Changwon University
Changwon 641-773, Korea

and

JAEUN LEE

Department of Mathematics
Kyungpook National University
Taegu 702-701, Korea

(Received February 20, 1992)

ABSTRACT. In this paper, we introduce weighted graph bundles and study their characteristic polynomial. In particular, we show that the characteristic polynomial of a weighted K_2 ($\bar{K}_2$)-bundles over a weighted graph Γ_ω can be expressed as a product of characteristic polynomials two weighted graphs whose underlying graphs are Γ . As an application, we compute the signature of a link whose corresponding weighted graph is a double covering of that of a given link.

KEY WORDS AND PHRASES. Graphs, weighted graphs, graph bundles, characteristic polynomials, links, signature.

1991 AMS SUBJECT CLASSIFICATION CODES. 05C10, 05C50, 57M25.

1. INTRODUCTION.

Let Γ be a simple graph with vertex set $V(\Gamma)$ and edge set $E(\Gamma)$. Let $\mathbf{R}$ be the field of real numbers. A *weighted graph* is a pair $\Gamma_\omega = (\Gamma, \omega)$, where Γ is a graph and $\omega: V(\Gamma) \cup E(\Gamma) \rightarrow \mathbf{R}$ is a function. We call Γ the *underlying graph* of Γ_ω and ω the *weight function* of Γ_ω . In particular, if $\omega(E(\Gamma)) \subset \{1, -1\}$ and $\omega(V(\Gamma)) = \{0\}$, then we call Γ_ω a *signed graph*.

Let $V(\Gamma) = \{u_1, \dots, u_n\}$. The *adjacency matrix* of Γ_ω is an $n \times n$ matrix $A(\Gamma_\omega) = (a_{ij})$ defined as follows:

$$a_{ij} = \begin{cases} \omega(e) & \text{if } e = u_i u_j \in E(\Gamma) \text{ and } i \neq j, \\ \omega(u_i) & \text{if } i = j, \\ 0 & \text{otherwise,} \end{cases}$$

for $1 \leq i, j \leq n$.

The characteristic polynomial $P(\Gamma_\omega; \lambda) = |\lambda I - A(\Gamma_\omega)|$ of the adjacency matrix $A(\Gamma_\omega)$ is called the *characteristic polynomial* of the weighted graph Γ_ω . A root of $P(\Gamma_\omega; \lambda)$ is called an *eigenvalue* of Γ_ω .

Note that if the weight function $\mathbf{l}$ of Γ is defined by $\mathbf{l}(e) = -1$ for $e \in E(\Gamma)$ and $\mathbf{l}(u) = \deg(u)$ for $u \in V(\Gamma)$, where $\deg(u)$ denotes the degree of u , that is, the number of edges incident to u , then the weighted adjacency matrix $A(\Gamma_{\mathbf{l}})$ is called the *Laplacian matrix* of Γ . We call $\mathbf{l}$ the *Laplacian function* of Γ . The number of spanning trees of a connected graph Γ is the

value of any cofactor of $A(\Gamma_{\mathbf{L}})$ [*Matrix tree theorem*] and is equal to the value $\frac{1}{n} \prod_{\lambda \neq 0} \lambda$, where λ runs through all non-zero eigenvalues of $A(\Gamma_{\mathbf{L}})$. Moreover, the eigenvalues of $A(\Gamma_{\mathbf{L}})$ may be used to calculate the radius of gyration of a Gaussian molecule. For more applications of the eigenvalues of $A(\Gamma_{\mathbf{L}})$, the reader is suggested to refer [5].

2. WEIGHTED GRAPH BUNDLES.

First, we introduce a weighted graph bundle. Every edge of a graph Γ gives rise to a pair of oppositely directed edges. We denote the set of directed edges of Γ by $D(\Gamma)$. By ϵ^{-1} we mean the reverse edge to an edge $\epsilon \in D(\Gamma)$. For any finite group G , a G -voltage assignment of Γ is a function $\phi: D(\Gamma) \rightarrow G$ such that $\phi(\epsilon^{-1}) = \phi(\epsilon)^{-1}$ for all $\epsilon \in D(\Gamma)$. We denote the set of all G -voltage assignments of Γ by $C^1(\Gamma; G)$. Let Λ be another graph and let $\phi \in C^1(\Gamma; \text{Aut}(\Lambda))$, where $\text{Aut}(\Lambda)$ is the group of all graph automorphisms of Λ . Now, we construct a graph $\Gamma \times^\phi \Lambda$ as follows: $V(\Gamma \times^\phi \Lambda) = V(\Gamma) \times V(\Lambda)$. Two vertices (u_1, v_1) and (u_2, v_2) are adjacent in $\Gamma \times^\phi \Lambda$ if either $u_1 u_2 \in D(\Gamma)$ and $v_2 = \phi(u_1 u_2) v_1$ or $u_1 = u_2$ and $v_1 v_2 \in E(\Lambda)$. We call $\Gamma \times^\phi \Lambda$ the Λ -bundle over Γ associated with ϕ and the natural map $p^\phi: \Gamma \times^\phi \Lambda \rightarrow \Gamma$ the bundle projection. We also call Γ and Λ the base and the fibre of $\Gamma \times^\phi \Lambda$, respectively. Note that the map p^ϕ maps vertices to vertices but an image of an edge can be either an edge or a vertex. If Λ is the complement $\overline{K_n}$ of the complete graph K_n of n vertices, then every Λ -bundle over Γ is an n -fold covering graph of Γ .

Let ω and μ be two weighted graphs and let $\phi \in C^1(\Gamma; \text{Aut}(\Lambda))$. We define the product of μ and ω with respect to ϕ , $\omega \times^\phi \mu$, as follows:

- (1) For each vertex (u, v) of $V(\Gamma \times^\phi \Lambda)$, $(\omega \times^\phi \mu)(u, v) = \omega(u) + \mu(v)$.
- (2) For each edge $e = (u_1, v_1)(u_2, v_2)$ of $E(\Gamma \times^\phi \Lambda)$,

$$(\omega \times^\phi \mu)(e) = \begin{cases} \omega(u_1 u_2) & \text{if } u_1 u_2 \in D(\Gamma) \text{ and } v_2 = \phi(u_1 u_2) v_1 \\ \mu(v_1 v_2) & \text{if } u_1 = u_2 \text{ and } v_1 v_2 \in E(\Gamma). \end{cases}$$

We call the weighted graph $(\Gamma \times^\phi \Lambda)_{\omega \times^\phi \mu}$ the Λ_μ -bundle over Γ_ω associated with ϕ . Briefly, we call it a weighted graph bundle.

FIGURE 1. The graphs $C_4 \times^\phi K_2$ and $(C_4 \times^\phi K_2)_{\omega \times^\phi \mu}$.

3. CHARACTERISTIC POLYNOMIALS.

In this section, we give a computation for the characteristic polynomial of a weighted graph bundle $\Gamma \times^\phi \Lambda$, where Λ is either complete graph K_2 of two vertices or its complement $\overline{K_2}$, and study their related topics. Note that $\text{Aut}(K_2) = \text{Aut}(\overline{K_2}) = \mathbb{Z}_2$.

For a given graph Γ with weight function ω and for a $\phi \in C^1(\Gamma; \mathbb{Z}_2)$, we define a new weight function ω^ϕ on Γ as follows:

(1) For $e \in E(\Gamma)$,

$$\omega^\phi(e) = \begin{cases} \omega(e) & \text{if } \phi(e) = 1 \\ -\omega(e) & \text{if } \phi(e) = -1 \end{cases}$$

(2) For $v \in V(\Gamma)$, $\omega^\phi(v) = w(v)$.

A subgraph of Γ is called an *elementary configuration* if its components are either complete graph K_1 or K_2 or a cycle C_m ($m \geq 3$). We denote by E_k the set of all elementary configurations of Γ having k vertices. In [3], the characteristic polynomial of a weighted graph Γ_ω is given as follows:

$$P(\Gamma_\omega; \lambda) = \sum_{k=0}^n a_k(\Gamma_\omega) \lambda^{n-k},$$

where

$$a_k(\Gamma_\omega) = \sum_{S \in E_k} (-1)^{\kappa(S)} 2^{|C(S)|} \prod_{u \in I_v(S)} \omega(u) \prod_{e \in I_E(S)} \omega(e)^2 \prod_{e \in C(S)} \omega(e).$$

In the above equation, symbols have the following meaning: $\kappa(S)$ is the number of components of S , $C(S)$ the set of all cycles, C_m ($m \geq 3$), in S , and $I_v(S)$ ($I_E(S)$) is the set of all isolated vertices (edges) in S . Moreover, the product over empty index set is defined to be 1.

For a fixed voltage assignment $\phi \in C^1(\Gamma; \mathbb{Z}_2)$, we denote by $E_{\phi-1}$ the set of edges of Γ such that $\phi(e) = -1$, i.e., $E_{\phi-1} = \{e \in E(\Gamma) : \phi(e) = -1\}$. Let $\Gamma(E_{\phi-1})$ be the edge subgraph of Γ induced by $E_{\phi-1}$ having weight zero in vertices. If Γ_ω is a weighted graph, then the weight function of its subgraph S is the restriction of ω on S .

THEOREM 1. Let $\overline{K}_2$ be a constant weighted graph, say $\mu(v) = c$ for $v \in \overline{K}_2$. Then, for each $\phi \in C^1(\Gamma; \mathbb{Z}_2)$, we have

$$P((\Gamma \times {}^\phi \overline{K}_2)_{\omega \times \phi_c}; \lambda) = P(\Gamma_\omega; \lambda - c) P(\Gamma_{\omega^\phi}; \lambda - c).$$

PROOF. Let $A(\Gamma_\omega)$ be the adjacency matrix of Γ_ω and let $A(\Gamma_{\omega^\phi})$ the adjacency matrix of Γ_{ω^ϕ} . Then we have

$$\begin{aligned} A(\Gamma_\omega) &= A(\Gamma \setminus (E_{\phi-1}))_\omega + A(\Gamma(E_{\phi-1})_\omega), \\ A(\Gamma_{\omega^\phi}) &= A(\Gamma \setminus (E_{\phi-1}))_{\omega^\phi} - A(\Gamma(E_{\phi-1})_\omega). \end{aligned}$$

Let $V(\Gamma \times {}^\phi \overline{K}_2) = \{(u_1, 1), \dots, (u_n, 1), (u_1, -1), \dots, (u_n, -1)\}$. It is not difficult to show that

$$\begin{aligned} A((\Gamma \times {}^\phi \overline{K}_2)_{\omega \times \phi_c}) &= \left(A(\Gamma_\omega) - A(\Gamma(E_{\phi-1})_\omega) + \begin{bmatrix} c & & 0 \\ & c & \\ 0 & & \ddots \\ & & & c \end{bmatrix} \right) \otimes \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \\ &\quad + (A(\Gamma(E_{\phi-1})_\omega)) \otimes \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \end{aligned}$$

Let M be a regular matrix of order 2 satisfying

$$M^{-1} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} M = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Put

$$X = A(\Gamma_\omega) - A(\Gamma(E_{\phi-1})_\omega) + \begin{bmatrix} c & & 0 \\ & c & \\ 0 & & \ddots \\ & & & c \end{bmatrix}$$

$$Y = A(\Gamma(E_{\phi-1})_\omega).$$

Then

$$\begin{aligned}
 & (I \otimes M^{-1}) A \left((\Gamma \times {}^\phi \overline{K_2})_{\omega \times \phi_c} \right) (I \otimes M) \\
 &= \begin{bmatrix} X+Y & 0 \\ 0 & X-Y \end{bmatrix} \\
 &= \begin{bmatrix} A(\Gamma_\omega) + \begin{bmatrix} c & & 0 \\ & c & \\ 0 & & \ddots \\ & & & c \end{bmatrix} & 0 \\ 0 & A(\Gamma_{\omega\phi}) + \begin{bmatrix} c & & 0 \\ & c & \\ 0 & & \ddots \\ & & & c \end{bmatrix} \end{bmatrix}.
 \end{aligned}$$

Since $|(I \otimes M^{-1})(I \otimes M)| = 1$ and

$$\left| \left[\lambda I - A \left((\Gamma \times {}^\phi \overline{K_2})_{\omega \times \phi_c} \right) \right] \right| = \left| \left[\lambda I - (I \otimes M^{-1}) A \left((\Gamma \times {}^\phi \overline{K_2})_{\omega \times \phi_c} \right) (I \otimes M) \right] \right|,$$

we have our theorem. $\square$

THEOREM 2. Let $K_{2,\mu} = (K_2, \mu)$ be a weighted graph having constant weight on vertices. Then, for each $\phi \in C(\Gamma; Z_2)$, we have

$$P\left((\Gamma \times {}^\phi K_2)_{\omega \times \phi_\mu}; \lambda\right) = P(\Gamma_\omega; \lambda - c_v - c_e) P(\Gamma_{\omega\phi}; \lambda - c_v + c_e),$$

where $c_v = \mu(v_1) = \mu(v_2)$ for the vertices v_1, v_2 and $c_e = \mu(e)$ for the edge e in K_2 .

PROOF. Clearly, we have

$$\begin{aligned}
 A\left((\Gamma \times {}^\phi K_2)_{\omega \times \phi_\mu}\right) &= \left(A(\Gamma_\omega) - A(\Gamma(E_{\phi-1})_\omega) + \begin{bmatrix} c_v & & 0 \\ & c_v & \\ 0 & & \ddots \\ & & & c_v \end{bmatrix} \right) \otimes \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \\
 &+ \left(A(\Gamma(E_{\phi-1})_\omega) + \begin{bmatrix} c_e & & 0 \\ & c_e & \\ 0 & & \ddots \\ & & & c_e \end{bmatrix} \right) \otimes \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}
 \end{aligned}$$

where $c_v = \mu(v_1) = \mu(v_2)$ and $c_e = \mu(e)$ for the edge e in K_2 . Let M be a regular matrix of order 2 satisfying

$$M^{-1} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} M = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

Then

$$\begin{aligned}
 & (I \otimes M^{-1}) A \left((\Gamma \times {}^\phi K_2)_{\omega \times \phi_\mu} \right) (I \otimes M) \\
 &= \begin{bmatrix} X+Y + \begin{bmatrix} c_e & & 0 \\ & c_e & \\ 0 & & \ddots \\ & & & c_e \end{bmatrix} & 0 \\ 0 & X-Y + \begin{bmatrix} c_e & & 0 \\ & c_e & \\ 0 & & \ddots \\ & & & c_e \end{bmatrix} \end{bmatrix}.
 \end{aligned}$$

$$= \begin{bmatrix} A(\Gamma_\omega) + Z_1 & 0 \\ 0 & A(\Gamma_{\omega\phi}) + Z_2 \end{bmatrix}$$

where X and Y are the same matrices as in the proof of Theorem 1 and for $i = 1, 2$,

$$Z_i = \begin{bmatrix} c_v + (-1)^{i-1} c_e & & 0 \\ & c_v + (-1)^{i-1} c_e & \\ 0 & & c_v + (-1)^{i-1} c_e \end{bmatrix}$$

Using method similar to the proof of Theorem 1, we have our theorem. $\square$

Note that for any $\phi \in C^1(\Gamma; \text{Aut}(\Lambda))$, the Laplacian function of $\Gamma \times^\phi \Lambda$ is the product of Laplacian functions of Γ and Λ with respect to ϕ . Clearly, the Laplacian function of the $\overline{K}_2$ is the zero function; and the Laplacian function of the K_2 has value 1 and -1 for each of its vertices and its edge, respectively. We shall denote the Laplacian function of a graph by $\mathcal{L}$ if it makes no confusion. Then Theorem 1 and Theorem 2 give the following corollary.

COROLLARY 1. For any $\phi \in C^1(\Gamma; Z_2)$,

$$(1) \quad P(\Gamma \times^\phi \overline{K}_2; \mathcal{L}; \lambda) = P(\Gamma; \mathcal{L}; \lambda) P(\Gamma_{\mathcal{L}\phi}; \lambda).$$

$$(2) \quad P(\Gamma \times^\phi K_2; \mathcal{L}; \lambda) = P(\Gamma; \mathcal{L}; \lambda) P(\Gamma_{\mathcal{L}\phi}; \lambda - 2). \quad \square$$

Now, we consider another invariant of weighted graphs called the *signature*. Since $A(\Gamma_\omega)$ is symmetric, $A(\Gamma_\omega)$ can be diagonalized through congruence over $\mathbb{R}$. Let d_+ denote the number of positive diagonal entries, and d_- the number of negative diagonal entries. The *signature* of a weighted graph (Γ_ω) is defined by $\sigma(A(\Gamma_\omega)) = d_+ - d_-$ and is denoted by $\sigma(\Gamma_\omega)$. It is an invariant for weighted 2-isomorphic graphs (see [7]).

From now on, we will consider the weight function on $\overline{K}_2$ as zero function and the weight function μ on K_2 as the map defined by $\mu(v) = 0$ for each $v \in V(K_2)$ and $\mu(e) = c_e$ for the edge e of K_2 . Then we can compute the signature of a double covering of Γ .

COROLLARY 2. $\sigma((\Gamma \times^\phi \overline{K}_2)_{\omega \times \phi_0}) = \sigma(\Gamma_\omega) + \sigma(\Gamma_{\omega\phi})$ for $\phi \in C^1(\Gamma; Z_2)$. $\square$

For convenience, we adapt the following notations. For a real number c , a weighted graph Γ_η and an eigenvalue λ of Γ_η ,

$$P(c)_\eta^- = \{\lambda < 0: \lambda + c > 0\},$$

$$P(c)_\eta^+ = \{\lambda > 0: \lambda + c > 0\},$$

$$Z(c)_\eta = \{\lambda \neq 0: \lambda + c = 0\},$$

$$N(c)_\eta^- = \{\lambda < 0: \lambda + c < 0\},$$

$$N(c)_\eta^+ = \{\lambda > 0: \lambda + c < 0\}.$$

We also denote the multiplicity of λ by $m_\eta(\lambda)$.

By using the above notations and Theorem 2, we get the signature of a K_2 -bundle over Γ .

COROLLARY 3. For $\phi \in C^1(\Gamma; Z_2)$,

(1) if $c_e \geq 0$, then

$$\begin{aligned} \sigma(\Gamma \times^\phi K_2)_{\omega \times \phi_\mu} &= \sigma(\Gamma_\omega) + \sigma(\Gamma_{\omega\phi}) \\ &\quad + (2 \sum_{\lambda \in P(c_e)_\omega^-} m_\omega(\lambda) + m_\omega(0) + \sum_{\lambda \in Z(c_e)_\omega} m_\omega(\lambda)) \\ &\quad - (2 \sum_{\lambda \in N(-c_e)_{\omega\phi}^+} m_{\omega\phi}(\lambda) + m_{\omega\phi}(0) + \sum_{\lambda \in Z(-c_e)_{\omega\phi}} m_{\omega\phi}(\lambda)). \end{aligned}$$

(2) if $c_e < 0$, then

$$\begin{aligned} \sigma(\Gamma \times {}^\phi K_2)_{\omega \times {}^\phi \mu} &= \sigma(\Gamma_\omega) + \sigma(\Gamma_{\omega^\phi}) \\ &\quad - \left(2 \sum_{\lambda \in N(c_e)_\omega^+} m_\omega(\lambda) + m_\omega(0) + \sum_{\lambda \in Z(c_e)_\omega} m_\omega(\lambda) \right) \\ &\quad + \left(2 \sum_{\lambda \in P(-c_e)_{\omega^\phi}^-} m_{\omega^\phi}(\lambda) + m_{\omega^\phi}(0) + \sum_{\lambda \in Z(-c_e)_{\omega^\phi}} m_{\omega^\phi}(\lambda) \right). \end{aligned}$$

□

REMARK. Though the results in this section stated only for a simple graph, it remains true for any graph.

4. APPLICATIONS TO LINKS.

In a signed graph Γ_ω , an edge e of Γ is said to be *positive* if $\omega(e) = 1$ and *negative* otherwise. For a signed graph Γ_ω , we define a new weight function $\tilde{\omega}$ of Γ by $\tilde{\omega}(e) = \omega(e)$ for any edge $e \in \Gamma_\omega$ and $\tilde{\omega}(u_i) = \sum_{j=1, i \neq j}^n a_{ij}$, where a_{ij} is the number of positive edges minus the number of negative edges which have two end vertices u_i and u_j . Given a knot or link L in $\mathbf{R}^3$, we project it into $\mathbf{R}^2$ so that each crossing point has proper double crossing. The image of L is called a *link (or knot) diagram* of L , and we do not distinguish between a diagram and the image of L .

We may assume without loss of generality that a link diagram $\tilde{L}$ of L intersects itself transversely and has only finitely many crossings. The link diagram $\tilde{L}$ divides $\mathbf{R}^2$ into finitely many domains, which will be classified as shaded or unshaded. No two shaded or unshaded domains have an edge in common. We now construct a signed planar graph Γ_ω from $\tilde{L}$ as follows: take a point v_i from each unshaded domain D_i . These points form the set of vertices $V(\Gamma_\omega)$ of Γ_ω . If the boundaries of D_i and D_j intersect k -times, say, crossing at $c_{\ell_1}, c_{\ell_2}, \dots, c_{\ell_k}$, then we form multiple edges $e_{\ell_1}, e_{\ell_2}, \dots, e_{\ell_k}$ on $\mathbf{R}^2$ with common end vertices v_i and v_j , where each edge e_{ℓ_m} passes through a crossing c_{ℓ_m} , for $m = 1, 2, \dots, k$. To define the weight of an edge, first, we define the index $\epsilon(c)$ to each crossing c of the link diagram as in Figure 2. To each edge of Γ passes through exactly one crossing, say c , of $\tilde{L}$, the weight $\omega(e)$ will be defined as $\omega(e) = \epsilon(c)$. (See Figure 3).

FIGURE 2. The index $\epsilon(c)$.

FIGURE 3. The correspondence between $\tilde{L}$ and $\Gamma_\omega(\tilde{L})$.

The resulting signed planar is called the *graph of a link with respect to $\tilde{L}$* and is denoted by $\Gamma_\omega(\tilde{L})$. The signed planar graph $\Gamma_\omega(\tilde{L})$ depends not only on $\tilde{L}$ but also on shading. Conversely, given a signed planar graph Γ_ϑ , one can construct uniquely the link diagram $L(\tilde{L}_\vartheta)$ of a link so that $\Gamma_\omega(L(\tilde{L}_\vartheta)) = \Gamma_\vartheta$.

FIGURE 4. The index $\omega(c)$.

Suppose that we are given an oriented link L . The orientation of L induces the orientation of a diagram $\tilde{L}$. We then define the second index $\omega(c)$, called the *twist or writhe* at each crossing c as show in Figure 4. We now need the third index $\eta_\rho(c)$ at crossing c . Let $\tilde{L}$ be an oriented diagram and ρ shading on $\tilde{L}$. Let $\eta_\rho(c) = \omega(c)\delta_{\epsilon(c)\omega(c)}$, where δ denotes Kronecker's delta. We define $\eta_\rho(\tilde{L}) = \sum \eta_\rho(c)$, where the summation runs over all crossing in $\tilde{L}$. The index $\eta_\rho(\tilde{L})$ depends not only on the shading ρ but also on the orientation of $\tilde{L}$. The following Lemma can be found in ([7], [4]).

LEMMA 1. The signature $\sigma(L)$ of a link L is $\sigma(L) = \sigma(\Gamma(\tilde{L})) - \eta_\rho(\tilde{L})$. $\square$

Let $\tilde{L}_1$ and $\tilde{L}_2$ be link diagrams of L_1 and L_2 , respectively. The link L_2 is called a *double covering* of the link L_1 if $\Gamma_\omega(\tilde{L}_2)$ is a double covering of $\Gamma_\omega(\tilde{L}_1)$ as weighted graphs and it can be extended to a branched covering on $\mathbb{R}^2$. Let ϕ be a voltage assignment in $C^1(\Gamma_\omega(\tilde{L}); \mathbb{Z}_2)$ such that $\phi(e) = -1$ for some edge e and $\phi(e) = 1$ otherwise, then $\Gamma_\omega(\tilde{L}) \times {}^\phi K_2$ is a planar double covering of $\Gamma_\omega(\tilde{L})$ of which the corresponding link is a double covering of L .

Therefore, one can construct the double covering link diagram $\tilde{L}(\Gamma_\omega(\tilde{L}) \times {}^\phi K_2)$ of $\tilde{L}$. Moreover, we can give an orientation on $\tilde{L}(\Gamma_\omega(\tilde{L}) \times {}^\phi K_2)$ so that the covering map from $\tilde{L}$ to $\tilde{L}(\Gamma_\omega(\tilde{L}) \times {}^\phi K_2)$ preserves the orientation. We have $\eta_\rho(\tilde{L}(\Gamma_\omega(\tilde{L}) \times {}^\phi K_2)) = 2\eta_\rho(\tilde{L})$ (see Figure 5).

FIGURE 5. Covering graph and covering link.

Therefore, by using Lemma 1 and Corollary 2, we get the following theorem.

THEOREM 3. For any oriented link diagram $\tilde{L}$,

$$\sigma(\tilde{L}(\Gamma_\omega(\tilde{L}) \rtimes^\phi \overline{K_2})) = \sigma(\Gamma_\omega(\tilde{L})) + \sigma(\Gamma_\omega(\tilde{L})) - 2\eta_\rho(\tilde{L})$$

for each $\phi \in C^1(\Gamma; \mathbb{Z}_2)$ such that $\phi(e) = -1$ for some edge $e \in \Gamma_\omega(\tilde{L})$ and $\phi(e) = 1$ otherwise.

ACKNOWLEDGEMENT. The first author was supported by KOSEF and the second author was supported by TGRC-KOSEF.

REFERENCES

1. BIGGS, N., *Algebraic Graph Theory*, Cambridge University Press, 1974.
2. CHAE, Y.; KWAK, J.H. and LEE, J., Characteristic polynomials of some graph bundles, preprint.
3. CVETKOVIĆ, D.M.; DOOB, M. and SACHS, H., *Spectra of Graphs*, Academic Press, New York, 1979.
4. GORDON, C.M. and LITTERLAND, R.A., On the signature of a link, *Invent. Math.* **47** (1978), 53-69.
5. GRONE, R.; MERRIS, R. and SUNDER, V.S., The Laplacian spectrum of a graph, *SIAM J. Matrix Anal. Appl.* **11** (1990), 218-238.
6. KWAK, J.H. and LEE, J., Isomorphism classes of graph bundles, *Canad. J. Math.* **42** (1990), 747-761.
7. MURASUGI, K., On invariants of graphs with applications to knot theory, *Trans. Amer. Math. Soc.* **314** (1989), 1-49.
8. MURASUGI, K., On the signature of graphs, *C.R. Math. Rep. Acad. Sci. Canada*, **10** (1989), 107-111.
9. MURASUGI, K., On the certain numerical invariant of link type, *Trans. Amer. Math. Soc.* **117** (1965), 387-422.
10. SCHWENK, A.J., Computing the characteristic polynomial of a graph, *Lecture Notes in Mathematics* **406**, Springer-Verlag (1974), 153-172.