GENERALIZED COMMON FIXED POINT THEOREMS FOR A SEQUENCE OF FUZZY MAPPINGS

B.S. LEE

Department of Mathematics
Kyungsoong University
Pusan 608-736, Korea

G.M. LEE, S.J. CHO

Department of Natural Sciences
Pusan National University of Technology
Pusan 608-739, Korea

and

D.S. KIM

Department of Applied Mathematics
National Fisheries University of Pusan
Pusan 608-737, Korea

(Received November 13, 1992 and in revised form May 6, 1993)

ABSTRACT. We obtain generalized common fixed point theorems for a sequence of fuzzy mappings, which is a generalization of the result of Lee and Cho [6].

KEY WORDS AND PHRASES. Fuzzy set, fuzzy mapping, upper semi-continuous, common fixed point.

1991 AMS SUBJECT CLASSIFICATION CODES. 54H25, 47H10.

1. INTRODUCTION. Heilpern [3] first introduced the concept of fuzzy mappings and proved a fixed point theorem for fuzzy contraction mappings, which is a fuzzy analogue of the fixed point theorems for multi-valued mappings ([2], [4], [9]) and the well-known Banach fixed point theorem. Bose and Sahani [1], in their first theorem, extended Heilpern's result for a pair of generalized fuzzy contraction mappings. They also, in their second theorem, proved a fixed point theorem for non-expansive fuzzy mappings on a compact star-shaped subset of a Banach space. Lee and Cho [5] proved a fixed point theorem for a contractive-type fuzzy mapping which is an extension of the result of Heilpern [3]. Also, they [6] obtained common fixed point theorems for a sequence of fuzzy mappings which are generalizations of their result in [5]. Lee et al. [7] obtained a common fixed point theorem for a sequence of fuzzy mappings satisfying certain conditions, which is a generalization of the second theorem of Bose and Sahani. They also showed common fixed point theorems for a pair of fuzzy mappings in [8], which is an extension of the first theorem of Bose and Sahani [1].

In this paper, we prove generalized common fixed point theorems for a sequence of fuzzy mappings satisfying certain conditions which are generalizations of the result of Lee and Cho [6].

2. PRELIMINARIES.

Let (X, d) be a linear metric linear space. A fuzzy set A in X is a function from X into $[0, 1]$. If $x \in X$, the function value $A(x)$ is called the *grade of membership* of x in A . The α -level set of A , denote by A_α , is defined by

$$A_\alpha = \{x: A(x) \geq \alpha\} \text{ if } \alpha \in (0, 1], \quad A_0 = \overline{\{x: A(x) > 0\}},$$

where $\bar{B}$ denotes the closure of the nonfuzzy set of B .

Let $W(X)$ be the collection of all the fuzzy sets A in X such that A_α is compact and convex for each $\alpha \in [0, 1]$, and $\sup_{x \in X} A(x) = 1$. For $A, B \in W(X)$, $A \subset B$ means $A(x) \leq B(x)$ for each $x \in X$.

DEFINITION 2.1. Let $A, B \in W(X)$ and $\alpha \in [0, 1]$. Then we define

$$P_\alpha(A, B) = \inf_{x \in A_\alpha, y \in B_\alpha} d(x, y), \quad P(A, B) = \sup_\alpha P_\alpha(A, B)$$

and

$$D(A, B) = \sup_\alpha d_H(A_\alpha, B_\alpha),$$

where d_H is the Hausdorff metric induced by the metric d . We note that P_α is a nondecreasing function of α and D is a metric on $W(X)$.

DEFINITION 2.2. Let X be an arbitrary set and Y be any linear metric space. F is called a *fuzzy mapping* if and only if F is a mapping from the set X into $W(Y)$.

In the following section, we will use the following lemmas.

LEMMA 2.1 [5]. Let (X, d) be a complete linear metric space, F a fuzzy mapping from X into $W(X)$ and $x_0 \in X$, then there exists $x_1 \in X$ such that $\{x_1\} \subset F(x_0)$.

LEMMA 2.2 [8]. Let $A, B \in W(X)$. Then for each $\{x\} \subset A$ there exists $\{y\} \subset B$ such that $D(\{x\}, \{y\}) \leq D(A, B)$.

We can easily prove the following lemma.

LEMMA 2.3. Let $x \in X$ and $B \in W(X)$. If $\{y\} \subset B$, then $P(\{x\}, B) \leq d(x, y)$.

3. COMMON FIXED POINTS THEOREMS FOR A SEQUENCE OF FUZZY MAPPINGS.

THEOREM 3.1. Let g be a non-expansive mapping from a complete linear metric space (X, d) into itself. If $(F_i)_{i=1}^\infty$ is a sequence of fuzzy mappings from X into $W(X)$ satisfying the following condition: For each pair of fuzzy mappings, F_i, F_j , and for any $x \in X$, $\{u_x\} \subset F_i(x)$, there exists $\{v_y\} \subset F_j(y)$ for all $y \in X$ such that

$$\begin{aligned} D(\{u_x\}, \{v_y\}) &\leq a_1 d(g(x), g(u_x)) + a_2 d(g(y), g(v_y)) \\ &\quad + a_3 d(g(y), g(u_x)) + a_4 d(g(x), g(v_y)) + a_5 d(g(x), g(y)), \end{aligned}$$

where a_1, a_2, a_3, a_4, a_5 are nonnegative real numbers, $a_1 + a_2 + a_3 + a_4 + a_5 < 1$ and $a_3 \geq a_4$. Then there exists $p \in X$ such that $\{p\} \subset \bigcap_{i=1}^\infty F_i(p)$.

PROOF. Let $x_0 \in X$. Then we can choose $x_1 \in X$ such that $\{x_1\} \subset F_1(x_0)$ by Lemma 2.1. By our assumptions, there exists $x_2 \in X$ such that $\{x_2\} \subset F_2(x_1)$ and

$$\begin{aligned} D(\{x_1\}, \{x_2\}) &\leq a_1 d(g(x_0), g(x_1)) + a_2 d(g(x_1), g(x_2)) + a_3 d(g(x_1), g(x_1)) \\ &\quad + a_4 d(g(x_0), g(x_2)) + a_5 d(g(x_0), g(x_1)) \\ &\leq a_1 d(x_0, x_1) + a_2 d(x_1, x_2) + a_3 d(x_1, x_1) + a_4 d(x_0, x_2) + a_5 d(x_0, x_1). \end{aligned}$$

Again we can find $x_3 \in X$ such that $\{x_3\} \subset F_3(x_2)$ and

$$D(\{x_2\}, \{x_3\}) \leq a_1 d(x_1, x_2) + a_2 d(x_2, x_3) + a_3 d(x_2, x_2) + a_4 d(x_1, x_3) + a_5 d(x_1, x_2).$$

Inductively, we obtain a sequence (x_n) in X such that $\{x_{n+1}\} \subset F_{n+1}(x_n)$ and

$$\begin{aligned} D(\{x_n\}, \{x_{n+1}\}) &\leq a_1 d(x_{n-1}, x_n) + a_2 d(x_n, x_{n+1}) + a_3 d(x_n, x_n) \\ &\quad + a_4 d(x_{n-1}, x_{n+1}) + a_5 d(x_{n-1}, x_n). \end{aligned} \tag{3.1}$$

Since $D(\{x_n\}, \{x_{n+1}\}) = d(x_n, x_{n+1})$, by (3.1)

$$\begin{aligned} d(x_n, x_{n+1}) &\leq a_1 d(x_{n-1}, x_n) + a_2 d(x_n, x_{n+1}) \\ &\quad + a_4 d(x_{n-1}, x_n) + a_4 d(x_n, x_{n+1}) + a_5 d(x_{n-1}, x_n). \end{aligned}$$

Hence

$$d(x_n, x_{n+1}) \leq [(a_1 + a_4 + a_5)/(1 - a_2 - a_4)]d(x_{n-1}, x_n).$$

Let $r = (a_1 + a_4 + a_5)/(1 - a_2 - a_4)$. Since $a_3 \geq a_4$, $0 < r < 1$. Moreover, we have $d(x_n, x_{n+1}) \leq r^n d(x_0, x_1)$. We can easily show that $(x_n)_{n=1}^\infty$ is a Cauchy sequence in X . Since X is complete, there exists $p \in X$ such that $\lim_{n \rightarrow \infty} x_n = p$. Let F_m be an arbitrary member of $(F_i)_{i=1}^\infty$. Since $\{x_n\} \subset F_n(x_{n-1})$ for all n , there exists $v_n \in X$ such that $\{v_n\} \subset F_m(p)$ for all n and

$$D(\{x_n\}, \{v_n\}) \leq a_1 d(x_{n-1}, x_n) + a_2 d(p, v_n) + a_3 d(p, x_n) + a_4 d(x_{n-1}, v_n) + a_5 d(x_{n-1}, p). \quad (3.2)$$

From (3.2), we have

$$\begin{aligned} d(x_n, v_n) &\leq a_1 d(x_{n-1}, x_n) + a_2 d(p, x_n) + a_2 d(x_n, v_n) + a_3 d(p, x_n) \\ &\quad + a_4 d(x_{n-1}, x_n) + a_4 d(x_n, v_n) + a_5 d(x_{n-1}, p) \end{aligned}$$

Thus we have

$$(1 - a_2 - a_4)d(x_n, v_n) \leq a_1 d(x_{n-1}, x_n) + a_2 d(p, x_n) + a_3 d(p, x_n) + a_4 d(x_{n-1}, x_n) + a_5 d(x_{n-1}, p).$$

Since $x_n \rightarrow p$ as $n \rightarrow \infty$, $(1 - a_2 - a_4)d(x_n, v_n) \rightarrow 0$ as $n \rightarrow \infty$. Hence $d(x_n, v_n) \rightarrow 0$ as $n \rightarrow \infty$. Since $d(p, v_n) \leq d(p, x_n) + d(x_n, v_n)$, $v_n \rightarrow p$ as $n \rightarrow \infty$. Since $F_m(p) \in W(X)$, $F_m(p)$ is upper semi-continuous and thus

$$\lim_{n \rightarrow \infty} \sup [F_m(p)](v_n) \leq [F_m(p)](p).$$

Since $\{v_n\} \subset F_m(p)$ for all n , $[F_m(p)](p) = 1$. Hence $\{p\} \subset F_m(p)$. Since F_m is arbitrary, $\{p\} \subset \bigcap_{i=1}^\infty F_i(p)$.

Putting $g(x) = x$, we get the following corollary from Theorem 3.1.

COROLLARY 3.1. Let (X, d) be a complete linear metric space. If $(F_i)_{i=1}^\infty$ is a sequence of fuzzy mappings from X into $W(X)$ satisfying the following condition (*): For each pair of fuzzy mapping F_i, F_j and for any $x \in X$, $\{u_x\} \subset F_i(x)$, there exists $\{v_y\} \subset F_j(y)$ for all $y \in X$ such that

$$D(\{u_x\}, \{v_y\}) \leq a_1 d(x, u_x) + a_2 d(y, v_y) + a_3 d(y, u_x) + a_4 d(x, v_y) + a_5 d(x, y),$$

where a_1, a_2, a_3, a_4, a_5 are nonnegative real numbers, $a_1 + a_2 + a_3 + a_4 + a_5 < 1$ and $a_3 \geq a_4$. Then there exists $p \in X$ such that $\{p\} \subset \bigcap_{i=1}^\infty F_i(p)$.

By Lemmas 2.2 and 2.3, we can obtain the following corollary from Corollary 3.1.

COROLLARY 3.2. Let (X, d) be a complete linear metric space and let $(F_i)_{i=1}^\infty$ be a sequence of fuzzy mappings from X into $W(X)$ satisfying the following condition (**): For each pair of fuzzy mappings F_i, F_j ,

$$D(F_i(x), F_j(y)) \leq a_1 P(x, F_i(x)) + a_2 P(y, F_j(y)) + a_3 P(y, F_i(x)) + a_4 P(x, F_j(y)) + a_5 d(x, y),$$

for all x, y in X , where a_1, a_2, a_3, a_4, a_5 are nonnegative real numbers, $a_1 + a_2 + a_3 + a_4 + a_5 < 1$ and $a_3 \geq a_4$. Then there exists $p \in X$ such that $\{p\} \subset \bigcap_{i=1}^\infty F_i(p)$.

The following example shows that the condition (*) in Corollary 3.1 does not imply the condition (**) in Corollary 3.2.

EXAMPLE 3.1. Let $(F_i)_{i=1}^\infty$ be a sequence of fuzzy mappings from $[0, \infty)$ into $W([0, \infty))$, where $F_i(x): [0, \infty) \rightarrow [0, 1]$ is defined as follows

$$\begin{aligned} \text{if } x = 0, \quad [F_i(x)](z) &= \begin{cases} 1, & z = 0 \\ 0, & z \neq 0, \end{cases} \\ \text{otherwise, } [F_i(x)](z) &= \begin{cases} 1, & 0 \leq z \leq x/2 \\ 1/2, & x/2 < z \leq ix \\ 0, & z > ix. \end{cases} \end{aligned}$$

Then the sequence $(F_i)_{i=1}^\infty$ satisfies the condition $(*)$ when $a_1 = a_2 = a_3 = a_4 = 0$, but does not satisfy the condition $(**)$.

Putting $a_1 = a_2 = a_3 = a_4 = 0$, we get the following corollary from Theorem 3.1.

COROLLARY 3.3 [6]. Let g be a non-expansive mapping from a complete linear metric space (X, d) into itself and $(F_i)_{i=1}^\infty$ a sequence of fuzzy mappings from X into $W(X)$ satisfying the following condition: There exists a constant k with $0 < k < 1$ such that for each pair of fuzzy mappings F_i, F_j and for any $x \in X, \{u_x\} \subset F_i(x)$, there exists $\{v_y\} \subset F_j(y)$ for all $y \in X$ such that

$$D(\{u_x\}, \{v_y\}) \leq kd(g(x), g(y)).$$

Then there exists $p \in X$ such that $\{p\} \subset \bigcap_{i=1}^\infty F_i(p)$.

By Lemma 2.2, we get the following corollary from Corollary 3.3.

COROLLARY 3.4 [6]. Let g be a non-expansive mapping from a complete linear metric space (X, d) into itself and $(F_i)_{i=1}^\infty$ a sequence of fuzzy mappings from X into $W(X)$ satisfying the following condition: There exists a constant k with $0 < k < 1$ such that for each pair of fuzzy mappings F_i, F_j ,

$$D(F_i(x), F_j(y)) \leq kd(g(x), g(y)) \quad \text{for all } x, y \text{ in } X,$$

Then there exists $p \in X$ such that $\{p\} \subset \bigcap_{n=1}^\infty F_n(p)$.

Putting $g(x) = x$, we get the following corollary from Corollary 3.4.

COROLLARY 3.5 [6]. Let (X, d) be a complete linear metric space and $(F_i)_{i=1}^\infty$ be a sequence of fuzzy mappings from X into $W(X)$ satisfying the following condition. There exists a constant k with $0 < k < 1$ such that for each pair of fuzzy mappings F_i, F_j ,

$$D(F_i(x), F_j(y)) \leq kd(x, y) \quad \text{for all } x, y \text{ in } X$$

Then there exists $p \in X$ such that $\{p\} \subset \bigcap_{n=1}^\infty F_n(p)$.

REFERENCES

1. BOSE, R.K. & SAHANI, D., Fuzzy mappings and fixed point theorems, *Fuzzy Sets and Systems* **21** (1987), 53-58.
2. CHEN, M.P. & SHIN, M.H., Fixed point theorems for point-to-point and point-to-set maps, *J. Math. Anal. Appl.* **71** (1979), 512-524.
3. HEILPERN, S., Fuzzy mappings and fixed point theorem, *J. Math. Anal. Appl.* **83** (1981), 566-569.
4. HUSAIN, T. & LATIF, A., Fixed points of multivalued nonexpansive maps, *Internat. J. Math. & Math. Sci.* **14** (1991), 421-430.
5. LEE, B.S. & CHO, S.J., A fixed point theorem for contractive-type fuzzy mappings, *Fuzzy Sets and Systems* (submitted).
6. LEE, B.S. & CHO, S.J., Common fixed point theorems for sequences of fuzzy mappings, *Internat. J. Math. & Math. Sci.* (to appear).
7. LEE, B.S., CHO, S.J., LEE, G.M. & KIM, D.S., On the common fixed point theorems for a sequence of fuzzy mappings, *Fuzzy Sets and Systems* (submitted).
8. LEE, B.S., LEE, G.M., CHO, S.J. & KIM, D.S., A common fixed point theorems for a pair of fuzzy mappings, *Information Sciences* (submitted).
9. NADLER, JR., S.B., Multi-valued contraction mappings, *Pacific. J. Math.* **30** (1969), 475-488.