FIXED POINT THEOREMS FOR NON-SELF MAPS I

TROY L. HICKS and LINDA MARIE SALIGA

Department of Mathematics and Statistics
University of Missouri-Rolla
Rolla, MO 65401

(Received July 29, 1992)

ABSTRACT. Suppose $f:C \rightarrow X$ where C is a closed subset of X . Necessary and sufficient conditions are given for f to have a fixed point. All results hold when X is complete metric space. Several results hold in a much more general setting.

KEY WORDS AND PHRASES. Commuting, compatible, d -complete topological spaces, fixed points, non-self maps, pairs of mappings.

1991 AMS SUBJECT CLASSIFICATION CODE. 47H10, 54H25.

1. INTRODUCTION.

Fixed point theorems for non-self maps are unusual. We surely require that $C \cap f(C)$ is non-empty. $f(x) = x + 1$ for X in $[0, 1]$ is a linear isometry from the compact space $[0, 1]$ into the compact space $[0, 2]$ but f is fixed point free. The mapping $f(x) = x + \frac{1}{x}$ for x in $[1, \infty)$ is a continuous mapping from $[1, \infty)$ into $[0, \infty)$. It is fixed point free but $|f(x) - f(y)| < |x - y|$ for $x \neq y$.

THEOREM (Brouwer [1]). If E is a non-empty convex compact subset of E^n and $f:E \rightarrow E$ is continuous, then $f(x) = x$ for some x in E .

2. RESULTS.

THEOREM 1. Let C be a closed subset of a complete metric space X and suppose f maps C onto X or f maps C into X with $C \subset f(C)$. If for some $k > 1$, $d(f(x), f(y)) \geq k d(x, y)$ for every x, y in C , then f has a unique fixed point in C .

PROOF. Clearly, f is one-to-one. Let $g = f^{-1}$ restricted to C . Now g maps C into C . For x, y in C , $d(x, y) = d(f(gx), f(gy)) \geq k d(g(x), g(y))$ or $d(g(x), g(y)) \leq \frac{1}{k} d(x, y)$ and $0 < \frac{1}{k} < 1$. g has a unique fixed point from Banach's fixed point theorem. But $f(x_0) = f(g(x_0)) = x_0$. If $x_1 = f(x_1)$, then $g(x_1) = g(f(x_1)) = x_1$ and $c_1 = x_0$.

The above result suggests that one should consider non-self maps that satisfy $C \subset f(C)$. It is well known that a continuous function from an arc onto a containing arc must have a fixed point. $[0, 1]$ or any homeomorphic image is called an arc. Thus Brouwer's theorem extends to the case $C \subset f(C)$ for $n = 1$. In [7], Sam Nadler showed that for $n \geq 2$ Brouwer's theorem does not extend. For $n \geq 2$, let A and B be closed balls in E^n with $A \subset B$ and $A \neq B$. He showed that there exists f and g such that:

- (a) $f:A \rightarrow B$ where f is continuous, onto, $f(\partial A) = B$, and f is fixed point free,
- (b) $g:A \rightarrow B$ where g is continuous, onto, $g^{-1}(\partial B) = \partial A$, and g if fixed point free.

THEOREM 2. Let C be a closed bounded, and convex subset of a uniformly convex Banach space and suppose f maps C onto X or f maps C into X with $C \subset f(C)$. If for every x, y in C $\|f(x) - f(y)\| \geq \|x - y\|$, then f has a fixed point in C .

PROOF. Clearly, f is one-to-one. Let $g = f^{-1}$ restricted to C and observe that $\|g(x) - g(y)\| \leq \|x - y\|$ where $g: C \rightarrow C$. From Kirk's theorem [6], g has a fixed point x_0 in C . Clearly, $f(x_0) = x_0$.

The following is an example of a mapping f that takes a closed, bounded, and convex subset C of a Banach space X into X where $C \subset f(C)$, $\|f(x) - f(y)\| = \|x - y\|$ for all $x, y \in C$, and f has no fixed points.

EXAMPLE 1. Let X be the space of sequences which converge to zero with $\|x\| = \sup_n |x_n|$ for x in X . Let $C = \{x \in X: \|x\| = 1 \text{ and } x_0 = 1\}$. C is closed, bounded, and convex. Define $f: C \rightarrow X$ by $f(x) = y$ where $y_n = x_{n+1}$, $n = 0, 1, 2, \dots$. $\|f(x) - f(y)\| = \|x - y\|$ and f is linear. To see that $C \subset f(C)$ consider the following. For $z \in C$, define r to be the sequence where $r_0 = 1$ and $r_n = z_{n-1}$, $n = 1, 2, 3, \dots$. Then $r \in C$, and $f(r) = z$ so $C \subset f(C)$. If $s = \{1, 0, 0, \dots\}$, $s \in C$ but $f(s) = \{0, 0, 0, \dots\} \notin C$. Hence $C \neq f(C)$. If $f(x) = x$ for some x in C , then $x_n = x_{n+1}$ for $n = 0, 1, 2, \dots$. Since $x_0 = 1$, $x_n = 1$ for all n and $x \notin C$. Therefore, f does not have a fixed point in C .

The following example shows that Banach's fixed point theorem does not generalize to non-self maps.

EXAMPLE 2. Let $X = C(\mathbf{R}, \mathbf{R})$ with $\|f\| = \sup_{t \in \mathbf{R}} |f(t)|$ for $f \in X$. Let $C = \{f \in X: f(t) = 0 \text{ for all } t \leq 0 \text{ and } \lim_{t \rightarrow \infty} f(t) \geq 1\}$. C is a closed and convex subset of X . Define $T: C \rightarrow X$ by $(Tf)(t) = \frac{1}{2} f(t+1)$. To see that $C \subset T(C)$ consider the following. For f in C set $g(t) = 2f(t-1)$. $g(t) = 0$ for $t \leq 0$ since $t-1 < 0$ and $f(t) = 0$ for all $t \leq 0$. $\lim_{t \rightarrow \infty} g(t) = \lim_{t \rightarrow \infty} 2f(t-1) \geq 2$. Thus $g \in C$ and $(Tg)(t) = f(t)$. Hence $C \subset T(C)$. Let $f(t)$ be defined as 0 if $t \leq 0$, t if $0 < t < 1$, and 1 if $t \geq 1$. Then $f \in C$. Now $(Tf)(t)$ is 0 if $t \leq -1$, $\frac{1}{2}(t+1)$ if $-1 < t < 0$, and $\frac{1}{2}$ if $t \geq 0$. Therefore, $Tf \notin C$ and $C \neq T(C)$. For $f, g \in C$, $\|Tf - Tg\| = \frac{1}{2} \|f - g\|$. If $Tf = f$ for some $f \in C$, then $f(t) = \frac{1}{2} f(t+1)$ and it follows that $f(n) = 0$ for all integers n . Hence $\lim_{t \rightarrow \infty} f(t) \not\geq 1$ and $f \notin C$. Therefore T does not have a fixed point in C . Note that T is linear, one-to-one, and $T(C)$ is closed.

We now turn to finding necessary and sufficient conditions for a non-self map to have a fixed point. Then it becomes clear that $C \subset f(C)$ is a natural assumption.

Let (X, t) be a topological space and $d: X \times X \rightarrow [0, \infty)$ such that $d(x, y) = 0$ if and only if $x = y$. X is said to be d -complete if $\sum_{n=1}^{\infty} d(x_n, x_{n+1}) < \infty$ implies that the sequence $\{x_n\}$ is convergent in (X, t) . These spaces include complete (quasi) metric spaces and d -complete (symmetric) semi-metric spaces. In [2] and [3] several basic metric space fixed point theorems were extended to this setting. $f: X \rightarrow X$ is w -continuous at x if $x_n \rightarrow x$ as $n \rightarrow \infty$ implies $f(x_n) \rightarrow f(x)$ as $n \rightarrow \infty$.

The following definition was given by G. Jungck in [5].

DEFINITION 1. Two maps f and g are compatible if, for any sequence $\{x_n\}$ such that $\lim_n f(x_n) = \lim_n g(x_n) = t$ it follows that $\lim_n d(f(gx_n), g(fx_n)) = 0$. Commuting maps are compatible but the converse is false.

DEFINITION 2. Given a map f , a map g is compatible with f , if for any sequence $\{x_n\}$ such that $\lim_n f(x_n) = \lim_n g(x_n) = t$ it follows that $\lim_n f(g(x_n)) = g(t)$.

REMARK 1. If f and g are w -continuous and (X, d) is a metric space, then, using definition

2, f is compatible with g is equivalent to g is compatible with f . In this case, we say that f and g are compatible.

PROOF. Assume f and g are w -continuous and that $\lim_n f(x_n) = \lim_n g(x_n) = t$ implies $\lim_n f(g(x_n)) = g(t)$. If we are in a metric space,

$$d(f(gx_n), g(fx_n)) \leq d(f(gx_n), g(t)) + d(g(t), g(fx_n))$$

and

$$d(g(fx_n), f(t)) \leq d(g(fx_n), f(gx_n)) + d(f(gx_n), f(t)).$$

It follows that f is compatible with g implies that g is compatible with f . Interchanging g and f above gives the converse.

It also follows from the above argument that if f and g are w -continuous and (X, d) is a metric space, then the two definitions of compatibility are equivalent.

REMARK 2. If (X, t) is a d -complete topological space, g is w -continuous, and f and g commute, then g is compatible with f using definition 2. We use definition 2 for d -complete topological spaces.

Theorem 3 and its corollaries are generalizations of theorems due to Hicks and Rhoades [4] which are generalizations of theorems due to Jungck [5].

THEOREM 3. Let (X, t) be a Hausdorff d -complete topological space and suppose $f: C \rightarrow X$ where f is w -continuous and C is a closed subset of X . Then f has a fixed point in C if and only if there exists $\alpha \in (0, 1)$ and a w -continuous function $g: C \rightarrow C$ such that $g(C) \subset f(C)$, g is compatible with f on $f^{-1}(C)$ and

$$(1) \quad d(g(x), g(y)) \leq \alpha d(f(x), f(y)) \text{ for all } x, y \in C.$$

Indeed, if (1) holds, f and g have a unique common fixed point.

PROOF. If $f(a) = a$ for some $a \in C$, set $g(x) = a$ for every $x \in C$. If $x \in f^{-1}(C)$, $f(x) \in C$ and $g(f(x)) = a = f(a) = f(g(x))$. If $x \in C$, $g(x) = a = f(a)$ gives $g(C) \subset f(C)$. Also, $0 = d(a, a) = d(g(x), g(y)) \leq \alpha d(f(x), f(y))$ for all $x, y \in C$.

Suppose there exists $\alpha \in (0, 1)$ and a w -continuous function $g: C \rightarrow C$ such that $g(C) \subset f(C)$, g is compatible with f on $f^{-1}(C)$ and $d(g(x), g(y)) \leq \alpha d(f(x), f(y))$ for all $x, y \in C$. Let $x_0 \in C$. $g(x_0) = f(x_1)$ for some $x_1 \in C$ since $g(C) \subset f(C)$. Construct a sequence $\{x_n\}$ with $\{x_n\} \subset C$ and $f(x_n) = g(x_{n-1})$ for $n = 1, 2, 3, \dots$. Since

$$d(f(x_n), f(x_{n+1})) = d(g(x_{n-1}), g(x_n)) \leq \alpha d(f(x_{n-1}), f(x_n)),$$

it follows that $d(f(x_n), f(x_{n+1})) \leq \alpha^{n-1} d(f(x_1), f(x_2))$. Hence $\sum_{n=1}^{\infty} d(f(x_n), f(x_{n+1})) < \infty$. The space is d -complete so there exists $p \in X$ with $\lim_n f(x_n) = p$. $f(x_n) = g(x_{n-1}) \in C$ gives $p \in cl(C) = C$. Now f is w -continuous gives $f(g(x_{n-1})) \rightarrow f(p)$ as $n \rightarrow \infty$. Since g is compatible with f on $f^{-1}(C)$ and $p \in f^{-1}(C)$ we get $\lim_n f(g(x_n)) = g(p)$. The space is Hausdorff so $f(p) = g(p)$ and $p \in f^{-1}(C)$. Consider the sequence $y_n = p$ for all n . Then $f(y_n) \rightarrow f(p)$ as $n \rightarrow \infty$, $g(y_n) \rightarrow g(p)$ as $n \rightarrow \infty$, and compatibility give $f(g(p)) = f(g(y_n)) \rightarrow g(f(p))$ as $n \rightarrow \infty$. Thus, $f(g(p)) = g(f(p))$. Therefore, $f(f(p)) = f(g(p)) = g(f(p)) = g(g(p))$ together with $d(g(p), g(gp)) \leq \alpha d(f(p), f(gp)) = \alpha d(g(p), g(gp))$ implies $g(p) = g(g(p))$. Hence $g(p) = g(g(p)) = f(g(p))$ and $g(p)$ is a common fixed point of f and g .

If $x = f(x) = g(x)$, then $d(x, g(p)) = d(g(x), g(gp)) \leq \alpha d(f(x), f(gp)) = \alpha d(x, g(p))$ gives $x = g(p)$.

COROLLARY 1. Let (X, t) be a Hausdorff d -complete topological space and C be a closed

subset of X . Suppose $f: C \rightarrow X$ and $g: C \rightarrow C$, where f and g are w -continuous, commute on $f^{-1}(C)$, and $g(C) \subset f(C)$. If there exists $\alpha \in (0, 1)$ and a positive integer k such that $d(g^k(x), g^k(y)) \leq \alpha d(f(x), f(y))$ for all $x, y \in C$, then f and g have a unique common fixed point.

PROOF. Clearly, g^k commutes with f on $f^{-1}(C)$ and $g^k(C) \subset g(C) \subset f(C)$. Applying the theorem to g^k and f gives a unique $p \in C$ such that $p = g^k(p) = f(p)$. Since f and g commute on $f^{-1}(C)$ and $p \in f^{-1}(C)$, $g(p) = g(f(p)) = f(g(p)) = g^k(g(p))$ or $g(p)$ is a common fixed point of f and g^k . Uniqueness of the common fixed point of f and g^k gives $g(p) = p = f(p)$. If $q = g(q) = f(q)$ then $g^k(q) = f(q)$ and $q = p$.

COROLLARY 2. Let n be a positive integer and let $\alpha > 1$. Suppose C is a closed subset of a Hausdorff d -complete topological space and $f: C \rightarrow X$ with $C \subset f(C)$. If $d(f^n(x), f^n(y)) \geq \alpha d(x, y)$ for all x, y in $(f^{n-1})^{-1}(C)$, then f has a fixed point in C .

PROOF. For $n = 1$, this follows from corollary 1 by letting $g = I$. f^n is one-to-one. $C \subset f(C)$ implies $C \subset f^n(C)$. Let h be the restriction of $(f^n)^{-1}$ to C . $h: C \rightarrow C$ and $d(h(x), h(y)) \leq \frac{1}{\alpha} d(x, y)$ for all $x, y \in C$. From corollary 1 with $k = 1$, $h = g^k = g$ and $f = I$, there exists a unique x_0 such that $h(x_0) = x_0$. Hence $f(x_0) = f^{n+1}(x_0) = f^n(f(x_0))$ or $h(f(x_0)) = (f^n)^{-1}(f(x_0)) = f(x_0)$. Uniqueness of the fixed point for h gives $x_0 = f(x_0)$. If $f(y) = y$ then $f^n(y) = y$ and $y = h(y)$. Again, uniqueness of the fixed point for h gives $x_0 = y$.

REFERENCES

1. BROUWER, L.E.J., Über abbildung von mannigfaltigkeiten, *Math. Ann.* **71** (1910), 97-115.
2. HICKS, T.L., Fixed point theorems for d -complete topological spaces I, *Internat. J. Math. and Math. Sci.* **15** (1992), 435-440.
3. HICKS, T.L. and RHOADES, B.E., Fixed point theorems for d -complete topological spaces II, *Math. Japonica* **37** (1992), 847-853.
4. HICKS, T.L. and RHOADES, B.E., Fixed points for pairs of mappings in d -complete topological spaces, *Internat. J. Math. and Math. Sci.* **16** (1993), 259-266.
5. JUNGCK, G., Compatible mappings and common fixed points, *Internat. J. Math. and Math. Sci.* **9** (1986), 771-779.
6. KIRK, W.A., A fixed point theorem for mappings which do not increase distance, *Amer. Math. Monthly* **72** (1965), 1004-1006.
7. NADLER, JR., S.B., Examples of fixed point free maps from cells onto larger cells and spheres, *Rocky Mnt. J. Math.* **11** (1981), 319-325.