THE T -STATISTICALLY CONVERGENT SEQUENCES ARE NOT AN FK -SPACE

JEANNETTE KLINE

DEPARTMENT OF MATHEMATICS
OHIO UNIVERSITY-CHILlicothe
CHILlicothe, OH 45601

(Received September 26, 1994 and in revised form November 30, 1994)

Abstract In this note we show that under certain restrictions on a nonnegative regular summability matrix T , the space of T -statistically convergent sequences cannot be endowed with a locally convex FK topology.

KEYWORDS AND PHRASES. Statistical convergence, FK -space
1990 AMS SUBJECT CLASSIFICATION CODE. 40H05

1. INTRODUCTION

Statistical convergence, introduced by Fast [5], has most recently been studied by Fridy and Orhan [7] [8], and Kolk [9], among others [3] [4] [6] [11]. In [3], it is shown that the space of statistically convergent sequences cannot be endowed with a locally convex FK topology. In this note, we establish that under certain restrictions on a nonnegative regular summability matrix T , the space of T -statistically convergent sequences cannot be endowed with a locally convex FK topology.

An infinite matrix $T = (t_{nk})$ is *nonnegative* if $t_{nk} \geq 0$ for all n and k , and *regular* if, for a convergent sequence x with limit l , $\lim_n \sum_{k=1}^{\infty} t_{nk} x_k = l$. Throughout this note T denotes a nonnegative regular matrix. We say that the rows of T *spread* if $\lim_n \max_k t_{nk} = 0$. We let ω denote the space of all real valued sequences, φ denote the finitely nonzero elements of ω and $\mathbb{N}$ denote the positive integers. For $\epsilon > 0$ and a scalar l , we let $A_{\epsilon,l} = \{k : |x_k - l| < \epsilon\}$. A sequence $x \in \omega$ is *T -statistically convergent to l* provided that for all $\epsilon > 0$,

$$\lim_n \sum_{k=1}^{\infty} t_{nk} \chi_{A_{\epsilon,l}}(k) = 1, \quad (1.1)$$

(where χ_A is the characteristic function of A). The space of T -statistically convergent sequences is denoted by S_T . Note that for $T = C_1$, the Cesàro matrix, this definition concurs with the definition of statistical convergence [6].

2. THE MAIN RESULT

A common theme in summability is the quest for "soft" methods to apply to classical type problems. An example of this is the " FK program," in which a summability space is given an FK topology ([13], pg. 54). An FK space X is a subspace of ω with a complete locally convex Fréchet topology such that the inclusion map from X into ω is continuous [13]. Our result shows that we cannot apply the FK -program to T -statistical convergence and improves Theorem 3.3 of [3].

THEOREM 1. *For a nonnegative regular summability matrix T whose rows spread, the space of T -statistically convergent sequences cannot be endowed with a locally convex FK topology.*

The proof of this theorem depends upon the following result of Bennett and Kalton [2].

THEOREM 2. *Let S be a dense subspace of ω . Then the following are equivalent:*

- 1) S is barrelled.
- 2) If E is a locally convex FK space that contains S , then $E = \omega$.

Note that the above statement is a restricted version of the result in [2], and an exposition can be found in ([12], pg. 253).

The proof of the main result follows that of Theorem 3.3 in [3].

PROOF OF THEOREM 1:

We show that S_T is a dense barrelled subspace of ω . Recall that S_T is barrelled if and only if every $\sigma(\varphi, S_T)$ -bounded subset of φ is $\sigma(\varphi, \omega)$ -bounded ([12], pg.248). Thus, to show S_T is barrelled it suffices to show that if E is not $\sigma(\varphi, \omega)$ -bounded, then E is not $\sigma(\varphi, S_T)$ -bounded.

We may assume that E is $\sigma(\varphi, \varphi)$ -bounded, since otherwise E is not $\sigma(\varphi, S_T)$ -bounded and we are done. Thus, there exists a sequence of integers $\langle B_n \rangle$ such that for x an element of E with $spt(x) = \{k \in \mathbb{N} : x_k \neq 0\} \subseteq \{1, 2, \dots, n\}$, we have $\sup_{1 \leq i \leq n} |x_i|$ is less than or equal to B_n . Since E is not $\sigma(\varphi, \omega)$ -bounded, we can choose s in ω such that $\sup_{x \in E} |\sum_{i=1}^{\infty} x_i s_i|$ is infinite. Note that for all x in E , we have $|\sum_{i=1}^n x_i s_i| \leq n B_n \sup_{i \leq n} |s_i|$.

Select $x^1 \in E$ such that $|\sum_{i=1}^{\infty} x_i^1 s_i| > B_1 |s_1|$, and select $j_1 > 1$ such that $x_{j_1}^1 \neq 0$ (such a j_1 exists since E is not $\sigma(\varphi, \omega)$ -bounded and since $|x_1| \leq B_1$). Assume that $\{x^1, x^2, \dots, x^n\}$ and $j_1 < j_2 < \dots < j_n$ have been chosen so that $x_{j_n}^n \neq 0$ and $j_n > \max\{k \in \mathbb{N} : k \in spt(x^{n-1})\}$. Set $t = \max\{j_n, \max\{k \in \mathbb{N} : k \in spt(x^n)\}\}$, and select x^{n+1} such that

$$|\sum_{i=1}^{\infty} x_i^{n+1} s_i| > t B_t \sup_{k \leq t} |s_k|. \quad (2.1)$$

Now select j_{n+1} such that $t < j_{n+1}$ and $x_{j_{n+1}}^{n+1} \neq 0$, and proceed inductively.

By [10], since the rows of T spread there is a subsequence $\langle j_{p_m} \rangle$ of $\langle j_n \rangle$ such that $\lim_n \sum_{k=1}^{\infty} t_{nk} \chi_{\{j_{p_m}, m \in \mathbb{N}\}}(k) = 0$. Since $x_{j_{p_m}}^{p_m} \neq 0$ for all m , it is possible to construct a sequence $\alpha = (\alpha_k)$ such that $\sum_{k=1}^{\infty} \alpha_k x_{j_{p_k}}^{p_k} \rightarrow \infty$ as $m \rightarrow \infty$. Then we set

$$z = (z_r) = \begin{cases} \alpha_k & \text{if } r = j_{p_k}, \text{ for } k = 1, 2, \dots \\ 0 & \text{else.} \end{cases} \quad (2.2)$$

Now, z is T -statistically convergent to 0 (because the non-zero entries of z occur on the subsequence $\langle j_{p_m} \rangle$ and by the definition of T -statistical convergence). Note also that $\sum_{k=1}^{\infty} x_k^{p_m} z_k = \sum_{k=1}^{\infty} x_{j_{p_k}}^{p_k} \alpha_k$. Since the right hand side of this equation tends to infinity as m does, it follows that E is not $\sigma(\varphi, S_T)$ -bounded. Since $\varphi \subseteq S_T$, S_T is a dense barrelled subspace of ω . Now, by the result of Bennett and Kalton, we have that S_T cannot be endowed with a locally convex FK topology.

The following examples illustrate the necessity of the hypothesis that the rows of T spread. Consider the identity matrix $I = (i_{nk})$, where $i_{nn} = 1$ and $i_{nk} = 0$ for all $k \neq n$. The space

of I -statistically convergent sequences is the space of convergent sequences c , a well-known FK space. For a less trivial example, consider the matrix $T = (t_{nk})$ where $t_{11} = 1$, $t_{1k} = 0$ for $k \geq 2$, and for $n \geq 2$,

$$t_{nk} = \begin{cases} \frac{1}{2} & \text{if } k = n \text{ or } k = n - 1, \\ 0 & \text{else.} \end{cases} \quad (2.3)$$

Note that the rows of T spread, and that this method is regular. In fact, T is stronger than convergence since the sequence $x = \langle (-1)^n \rangle$ has $\lim_n \sum_{k=1}^{\infty} t_{nk} x_k = 0$. As in the case of the identity matrix, the T -statistically convergent sequences are again the FK space c of convergent sequences.

A consequence of this result is that we cannot employ the FK program when studying T -statistical convergence for a matrix T whose rows spread. Instead, the Stone-Ćech compactification of the integers has been used [4][1] as an avenue for "soft" methods for treating T -statistical convergence of bounded sequences. This result also includes Corollary 4.4 of [9], where it is shown that under certain restrictions, a matrix B maps the space of T -statistically convergent sequences into a sequence space Y if and only if B has at most finitely many non-zero columns which belong to Y .

ACKNOWLEDGEMENT. The author wishes to thank Jeff Connor for critically reading an earlier version of this manuscript and making many helpful suggestions.

REFERENCES

- 1 ATTALLA, R., *On the multiplicative behaviour of regular matrices*, Proc. Amer. Math. Soc. **26** 3, 437-446
- 2 BENNETT, G. and KALTON, N. J., *Inclusion theorems for K spaces*, Canadian J. Mathematics **25**, 511-524
- 3 CONNOR, J. S., *The statistical and strong p -Cesàro convergence of sequences*, Analysis **8**, 47-63
- 4 CONNOR, J. S., *R -type summability methods, Cauchy criteria, P -sets, and statistical convergence*, Proc. Amer. Math. Soc. **115** 2, 319-327
- 5 FAST, H., *Sur la convergence statistique*, Colloq. Math. **2**, 241-244
- 6 FRIDY, J., *On statistical convergence*, Analysis **5**, 301-313.
- 7 FRIDY, J. and ORHAN, C., *Lacunary statistical convergence*, Pacific Jour. Math **160**, 1187-1192.
- 8 FRIDY, J. and ORHAN, C., *Lacunary statistical summability*, J. Math. Analysis and Applications **173**, 497-504.
- 9 KOLK, E., *Matrix summability of statistically convergent sequences*, Analysis **13**, 77-83.
- 10 LORENTZ, G. G., *A contribution to the theory of divergent sequences*, Acta Mathematica **80**, 167-190.
- 11 MILLER, H., *A measure theoretical subsequence characterization of statistical convergence*, preprint.
- 12 WILANSKY, A., "Modern Methods in Topological Vector Spaces," McGraw-Hill Inc..
- 13 WILANSKY, A., "Summability through Functional Analysis," North-Holland.