MINIMAL CR-SUBMANIFOLDS OF A SIX-DIMENSIONAL SPHERE

M. HASAN SHAHID

and

S. I. HUSAIN

Department of Mathematics
Faculty of Natural Sciences
Jamia Millia Islamia, New Delhi - 110025
INDIA

Department of Mathematics
Aligarh Muslim University
Aligarh, 202002, INDIA

(Received February 18, 1993)

ABSTRACT. We establish several formulas for a 3-dimensional CR-submanifold of a six-dimensional sphere and state some results obtained by making use of them.

KEY WORDS AND PHRASES. CR-submanifold, D-minimal and $D^\perp$ -minimal CR-submanifold.
1991 AMS SUBJECT CLASSIFICATION CODE. 53C40.

1. INTRODUCTION. Among all submanifolds of a Kaehler manifold there are three typical classes: the complex submanifolds, the totally real submanifolds and the CR-submanifolds. The notion of a CR-submanifold of a Kaehler manifold was introduced by Bejancu [1] and it includes the other two classes as special cases. A Riemannian submanifold M of an almost Hermitian manifold $\bar{M}$ is called a CR-submanifold if there exists a pair of orthogonal complementary distribution D and $D^\perp$ on M satisfying $JD = D$ and $JD^\perp \subset \nu$, where ν is the normal bundle of M . If M is a real hypersurface of a Kaehler manifold, then M is obviously a CR-submanifold.

It is known that every Kaehler manifold is nearly Kaehler but the converse is not true in general. The most typical example of nearly Kaehler manifolds is a six-dimensional sphere S^6 . It is because of this nearly Kaehler, non-Kaehler, structure that S^6 has attracted attention.

The object of the present paper is to establish several formulas for a 3-dimensional CR-submanifold of a six-dimensional sphere and state some result obtained by making use of them.

2. PRELIMINARIES.

Let $\bar{M}$ be an almost complex manifold with almost complex structure J , and Hermitian metric g . $\bar{M}$ is called a nearly Kaehler manifold if

$$(\bar{\nabla}_X J)(Y) + (\bar{\nabla}_Y J)(X) = 0 \quad (2.1)$$

for $X, Y \in (\bar{M})$, where $\bar{\nabla}$ is Riemannian connection on $\bar{M}$.

In [5], K. Takamatsu and T. Sato proved the following theorem:

THEOREM. Let $\bar{M} = (\bar{M}, J, g)$ be a non-Kaehler, nearly Kaehler manifold of constant holomorphic sectional curvature. Then $\bar{M}$ is a six-dimensional space of positive constant sectional curvature.

If a nearly Kaehler manifold $\bar{M}$ is constant holomorphic sectional curvature c , then by the above result, the curvature tensor $\bar{R}$ of $\bar{M}$ is given by

$$R(X, Y, Z, W) = c(g(Y, Z)g(X, W) - g(X, Z)g(Y, W)). \quad (2.2)$$

Let M be an m -dimensional CR-submanifolds of a six-dimensional sphere $\bar{M}$ and let us denote by the same g the Riemannian metric tensor field induced on M from that of $\bar{M}$. Let P and Q be the projection operators corresponding to D and $D^\perp$ respectively.

For a vector field X tangent to M , we put

$$JX = PX + QX \quad (2.3)$$

where PX (resp. QX) denote the tangent (resp. normal) component of JX .

We now denote by $\bar{\nabla}$ (resp. ∇) the Riemannian connection in $\bar{M}$ (resp. M) with respect to the Riemannian metric g . The linear connection induced by $\bar{\nabla}$ on the normal bundle $T^\perp M$ is denoted by $\nabla^\perp$. Thus the Gauss and Weingarten formulas are given by

$$\bar{\nabla}_X Y = \nabla_X Y + h(X, Y), \quad \bar{\nabla}_X N = -A_N X + \nabla_X^\perp N \quad (2.4)$$

for all $X, Y \in TM$ and $N \in T^\perp M$, where h is the second fundamental form of M and A_N is the fundamental tensor with respect to the normal section N . These tensor fields are related by

$$g(h(X, Y), N) = g(A_N X, Y). \quad (2.5)$$

The equation of Gauss is given by

$$\begin{aligned} \bar{R}(X, Y, Z, W) &= R(X, Y, Z, W) - g(h(X, W), h(Y, Z)) \\ &\quad + g(h(X, Z), h(Y, W)). \end{aligned} \quad (2.6)$$

DEFINITION. A CR-submanifold M is called D -minimal (resp. D -minimal) if $\sum_{i=1}^{2p} h(E_i, E_i) = 0$ (resp. $\sum_{i=1}^q h(F_i, F_i) = 0$) where $\{E_1, E_2, \dots, E_{2p}\}$ (resp. $\{F_1, F_2, \dots, F_q\}$) is a local field of frames of D (resp. $D^\perp$).

DEFINITION. A CR-submanifold M is called D -totally geodesic (resp. $D^\perp$ -totally geodesic) if $h(X, Y) = 0$ for each $X, Y \in D$ (resp. $X, Y \in D^\perp$). M is called a mixed totally geodesic if $h(X, Z) = 0$ for each $X \in D, Z \in D^\perp$.

3. THREE-DIMENSIONAL CR-SUBMANIFOLDS OF S^6 .

Let M be a 3-dimensional CR-submanifold of S^6 . It is known that S^6 is nearly Kaehler manifold of constant type 1. Suppose $\dim D = 2, \dim D^\perp = 1$, and $\{E_1, JE_1\}$ be a local frame in D and $\{F\}$ that of $D^\perp$.

The mean curvature vector H is defined by

$$H = \frac{1}{3} \left\{ \sum_{i=1}^2 h(E_i, E_i) + h(F, F) \right\}. \quad (3.1)$$

If $H = 0$, then M is said to be minimal. Now we define

$$H_D = \frac{1}{2} \sum_{i=1}^2 h(E_i, E_i), H_{D^\perp} = h(F, F). \quad (3.2)$$

If $H_D = 0$, then M is said to be D -minimal and if $H_{D^\perp} = 0$, then M is said to be $D^\perp$ -minimal.

Let U, V be any vector field tangent to CR-submanifold M . The Ricci tensor and the scalar curvature are respectively given by

$$S(U, V) = \sum_{i=1}^2 g(R(E_i, U)V, E_i) + g(R(F, U)V, F). \quad (3.3)$$

$$\rho = \sum_{i=1}^2 S(E_i, E_i) + S(F, F). \quad (3.4)$$

Also

$$S_D(U, V) = g(R(E_i, U)V, E_i), S_{D^\perp}(U, V) = g(R(F, U)V, F). \quad (3.5)$$

$$\rho_{DD} = \sum_{i=1}^2 S_D(E_i, E_i), \rho_{DD^\perp} = S_D(F, F). \quad (3.6)$$

$$\rho_{D^\perp D} = \sum_{i=1}^2 S_{D^\perp}(E_i, E_i), \quad \rho_{D^\perp D^\perp} = S_{D^\perp}(F, F). \quad (3.7)$$

Now using (2.2) and (2.6), we have for $X, Y \in TM$

$$S_D(X, Y) = 2g(X, Y) - g(PX, PY) + 2g(H_D, h(X, Y)) \quad (3.8)$$

$$- \sum_{i=1}^2 g(h(E_i, X), h(E_i, Y)),$$

$$S_{D^\perp}(X, Y) = g(X, Y) - g(QX, QY) + g(H_{D^\perp}, h(X, Y)) \quad (3.9)$$

$$- g(h(F, X), h(F, Y)),$$

$$\rho_{DD} = 2 + 4g(H_D, H_D) - \sum_{i,j=1}^2 \|h(E_i, E_j)\|^2, \quad (3.10)$$

$$\rho_{DD^\perp} = 2 + 2g(H_{D^\perp}, H_D) - \sum_{i=1}^2 \|h(E_i, F)\|^2, \quad (3.11)$$

$$\rho_{D^\perp D^\perp} = g(H_{D^\perp}, H_{D^\perp}) - \|h(F, F)\|^2. \quad (3.12)$$

It is easy to see that

$$\rho_{DD^\perp} = \rho_{D^\perp D}$$

Now we prove

THEOREM 1. Let M be a D -minimal CR-submanifold of a 6-dimensional sphere S^6 . Then the following hold:

$$(a) \quad S_D(X, X) - 2\|X\|^2 + \|PX\|^2 \leq 0, \text{ for } X \in TM$$

$$(b) \quad \rho_{DD} \leq 2$$

$$(b') \quad \rho_{DD^\perp} \leq 2.$$

The equality in (a) for $X \in D$, and in (b) holds if and only if M is D -totally geodesic.

The equality in (a) for $X \in D^\perp$ and in (b') holds if and only if M is mixed totally geodesic.

PROOF. Since M is D -minimal, from (3.8), we have

$$S_D(X, X) - 2\|X\|^2 + \|PX\|^2 = - \sum_{i=1}^2 g(h(E_i, X), h(E_i, X)).$$

This proves (a) and (b), (b') follow from (3.10) and (3.11). Similarly, we have

THEOREM 2. Let M be a $D^\perp$ -minimal CR-submanifold of a 6-dimensional sphere S^6 .

Then the following hold:

$$(a) \quad S_{D^\perp}(X, X) - \|X\|^2 + \|QX\|^2 \leq 0, \text{ for } X \in TM$$

$$(b) \quad \rho_{D^\perp D^\perp} \leq 2,$$

$$(b') \quad \rho_{D^\perp D^\perp} \leq 2.$$

The equality for $X \in D^\perp$ in (a) and (b') holds if and only if M is $D^\perp$ -totally geodesic.

The equality for $X \in D$ in (a) and in (b) holds if and only if M is mixed totally geodesic.

PROOF. Since M is $D^\perp$ -minimal, so from (3.9), we have

$$S_{D^\perp}(X, X) - \|X\|^2 + \|QX\|^2 = -g(h(F, X), h(F, X)),$$

which proves (a) and (b), (b') follows from (3.11) and (3.12).

REMARKS. The example given by Sekigawa [6] is an example of D -totally geodesic and $D^\perp$ -totally geodesic (and hence minimal) proper CR-submanifold of a 6-dimensional sphere and this illustrates the Theorem 1 in the sense that $S_f^2 \times S^1$, where f is a function on S^2 , is a D -minimal CR-submanifold of S^6 in which it is easily verified that $\rho_{DD} = 2$. The equality arises because it is also D -totally geodesic in S^6 .

ACKNOWLEDGEMENT. The author wishes to express their hearty thanks to Professor K. Sekigawa who kindly checked the original manuscripts.

REFERENCES

1. BEJANCU, A., CR-Submanifolds of Kaehler manifold 1, *Proc. Amer. Math. Soc.* **69** (1987), 135-142.
2. CHEN, B.Y., *Geometry of Submanifolds*, Marcel Dekkar, New York, 1973.
3. GRAY, A., Classification des varietes approximativement Kaehleriennes de courbure sectionnelle constant, *C.R. Acad. Sc. Paris* **279** (1979), 797-800.
4. DESHMUKH, S.; SHAHID, M.H. & ALI, S., CR-Submanifolds of a nearly Kaehler manifold, *Tamkang J. Maths.* **17** (1986), 17-27.
5. TAKAMATSU, K. & SATO, T., A K-space of constant holomorphic sectional curvature, *Kadai Math. Sem. Rep.* **27** (1976), 116-127.
6. SEKIGAWA, K., Some CR-submanifolds in a 6-dimensional sphere, *Tensor N.S.* **41** (1984), 13-19.
7. SWAKI, K.; WATANABE, Y. & SATO, T., Notes on a K-space of constant holomorphic sectional curvature, *Kadai Math. Sem. Rep.* **26** (1975), 438-445.