COEFFICIENT SUBRINGS OF CERTAIN LOCAL RINGS WITH PRIME-POWER CHARACTERISTIC

TAKAO SUMIYAMA

Department of Mathematics
Aichi Institute of Technology
Yakusa-chô, Toyota
470-03 Japan

(Received October 13, 1993 and in revised form September 15, 1994)

ABSTRACT. If R is a local ring whose radical $J(R)$ is nilpotent and $R/J(R)$ is a commutative field which is algebraic over $GF(p)$, then R has at least one subring S such that $S = \bigcup_{i=1}^{\infty} S_i$, where each S_i is isomorphic to a Galois ring and $S/J(S)$ is naturally isomorphic to $R/J(R)$. Such subrings of R are mutually isomorphic, but not necessarily conjugate in R .

KEY WORDS AND PHRASES: Coefficient ring, Galois ring, local ring, Szele matrix.

1991 AMS SUBJECT CLASSIFICATION CODES: 16L30, 16P10, 16D70

0. INTRODUCTION

Let p be a fixed prime. For any positive integers n and r , there exists up to isomorphism a unique r -dimensional separable extension $GR(p^n, r)$ of $\mathbb{Z}/p^n\mathbb{Z}$, which is called the Galois ring of characteristic p^n and rank r (see [9, p. 293, Theorem XV.2]). This ring was first noticed by Krull [8], and was later rediscovered by Janusz [6] and Raghavendran [12].

By Wedderburn-Malcev theorem (see, for instance, [4, p. 491]), if R is a finite dimensional algebra over a field K such that $\bar{R} = R/J(R)$ is a separable algebra over K , then R contains a semisimple subalgebra S such that $R = S \oplus J(R)$ (direct sum as vector spaces). Such subalgebras of R are conjugate each other.

Concerning the case R is not an algebra over a field, Raghavendran [12, Theorem 8], Clark [3] and Wilson [17, Lemma 2.1] have proved the following: If R is a finite local ring with characteristic p^n whose residue field is $GF(p')$, then R contains a subring S such that S is isomorphic to $GR(p^n, r)$ (hence $R = S + J(R)$). Such a subring S of R is called a coefficient ring of R . Coefficient rings of R are conjugate each other. We can embed R to a ring of Szele matrices over S (see §1).

If R is a finite ring of characteristic p^n , then R contains a subring T (unique up to isomorphism) such that (1) $R = T \oplus N$ (as abelian groups), where N is an additive subgroup of $J(R)$, (2) T is a direct sum of matrix rings over Galois rings, (3) $J(T) = T \cap J(R) = pT$, and (4) $R = T + J(R)$.

The purpose of this paper is to extend these results to certain rings which are not necessarily finite.

1.

In what follows, when S is a set, $|S|$ will denote the cardinal number of S . When A is a ring, for any subset S of A , $\langle S \rangle$ denotes the subring of A generated by S . A ring A is called locally finite if any finite subset of A generates a finite subring. When A is a ring with 1, for B to be called a subring of A , B must contain 1. Let $J(A)$ denote the Jacobson radical of A , $Aut(A)$ the automorphism group of A , and $(A)_{n \times n}$ the ring of $n \times n$ matrices having entries in A . If $A \ni 1$, A^* denotes the group of units of A . For $a \in A^*$, $o(a)$ denotes the multiplicative order of a .

The Galois ring $GR(p^n, r)$ is characterized as a ring isomorphic to $(\mathbb{Z}/p^n\mathbb{Z})[X]/(f(X)(\mathbb{Z}/p^n\mathbb{Z})[X])$, where $f(X) \in (\mathbb{Z}/p^n\mathbb{Z})[X]$ is a monic polynomial of degree r , and is irreducible modulo $p\mathbb{Z}/p^n\mathbb{Z}$ (see [7, Chapter XVI]). By [12, Proposition 1], any subring of $GR(p^n, r)$ is isomorphic to $GR(p^n, s)$, where

s is a divisor of r . Conversely, if s is a divisor of r , then there is a unique subring of $GR(p^n, r)$ which is isomorphic to $GR(p^n, s)$.

The following lemma is easily deduced from [16, Theorem 3 (I)] and its proof.

LEMMA 1.1. Let R be a finite local ring of characteristic p^n whose residue field is $GF(p')$. If $a \in R^*$ satisfies $o(a) = p' - 1$, then the subring $\langle a \rangle$ of R is isomorphic to $GR(p^n, r)$.

A ring R will be called an IG-ring if there exists a sequence $\{R_i\}_{i=1}^\infty$ of subrings of R such that $R_i \subset R_{i+1}$, $R_i \cong GR(p^n, r_i)$ ($i \geq 1$) and $R = \bigcup_{i=1}^\infty R_i$, where $\{r_i\}_{i=1}^\infty$ is a sequence of positive integers such that $r_i \mid r_{i+1}$ ($i \geq 1$). If R is an IG-ring described above, then R_i is the only subring of R which is isomorphic to $GR(p^n, r_i)$. So we can write $R = \bigcup_{i=1}^\infty GR(p^n, r_i)$.

Let p be a prime, n a positive integer and $1 = r_1 \leq r_2 \leq \dots$ an infinite sequence of positive integers such that $r_i \mid r_{i+1}$. By the fact we observed above, there exists a natural embedding $\iota_i^{i+1} : GR(p^n, r_i) \rightarrow GR(p^n, r_{i+1})$ for each $i \geq 1$. Let us put $\iota_i^i = id_{GR(p^n, r_i)}$ and $\iota_i^j = \iota_{j-1}^j \circ \iota_{j-2}^{j-1} \circ \dots \circ \iota_i^{i+1}$ for $1 \leq i \leq j$. Then we see that $\{GR(p^n, r_i), \iota_i^j\}$ is an inductive system. The ring $R = \lim_{\leftarrow} GR(p^n, r_i)$ is an IG-ring. Conversely, any IG-ring can be constructed in this way. An IG-ring $R = \bigcup_{i=1}^\infty GR(p^n, r_i)$ is a Galois ring if $|R|$ is finite. When A is a ring with 1, a subring S of A is called an IG-subring of A if S is an IG-ring.

PROPOSITION 1.2. Let $R = \bigcup_{i=1}^\infty GR(p^n, r_i)$ be an IG-ring. Then:

- (I) R is a commutative local ring with radical $J(R) = pR$. The residue field R/pR is $\bigcup_{i=1}^\infty GF(p')$.
- (II) If e is a positive integer such that $1 \leq e \leq n$, then $R/p^e R$ is naturally isomorphic to the IG-ring $\bigcup_{i=1}^\infty GR(p^e, r_i)$.
- (III) R is a proper homomorphic image of a discrete valuation ring whose radical is generated by p .
- (IV) Any ideal of R is of the form $p^e R$ ($0 \leq e \leq n$).
- (V) R is self-injective.
- (VI) $Aut(R) \cong \lim_{\leftarrow} Aut(GR(p^n, r_i)) \cong \lim_{\leftarrow} Aut(GF(p')) \cong Aut\left(\bigcup_{i=1}^\infty GF(p')\right)$.

PROOF. (I) and (II). For each $i \geq 1$,

$$0 \rightarrow p^e GR(p^n, r_i) \rightarrow GR(p^n, r_i) \rightarrow GR(p^e, r_i) \rightarrow 0$$

is an exact sequence of $GR(p^n, r_i)$ -modules. So we get the result by [2, Chapitre 2, §6, n° 6, Proposition 8].

(III) Let us put $K = \bigcup_{i=1}^\infty GF(p')$. Let $W_n(K)$ be the ring of Witt vectors over K of length n (see [15, Chapter II, §6] or [5, Kapitel II, §10.4]. By (I) and [5, Kapitel II, §10.4], both R and $W_n(K)$ are elementary complete local rings (in [14], elementare vollständige lokale Ringe) of characteristic p^n whose residue fields are K . Since an elementary complete local ring is uniquely determined by its characteristic and residue field (see [14, Anhang 2]), we see that R is isomorphic to $W_n(K)$. Let $W(K)$ be the ring of Witt vectors over K of infinite length. By [7, Chapter V, §7], $W(K)$ is a discrete

valuation ring whose radical is generated by p . Since $W(K)$ is the projective limit of $\{W_n(K)\}_{n=1}^\infty$ (see [15, Chapter II, §6]), $W_n(K)$ is a homomorphic image of $W(K)$.

(IV) If R is a discrete valuation ring with radical pR , then any ideal of R is of the form $p^j R$ ($j \geq 0$), so the result is clear from (III).

(V) Clear from (III), since any proper homomorphic image of a principal ideal domain is self-injective.

(VI) Immediate by [9, p. 294, Corollary XV.3].

Let $\{r_l\}_{l=1}^\infty$ be an infinite sequence of positive integers such that $r_1 = 1$ and $r_l \mid r_{l+1}$ ($l \geq 1$), and $S = \bigcup_{l=1}^\infty GR(p^n, r_l)$ be an IG-ring of characteristic p^n . Let $n = n_1 \geq n_2 \geq \dots \geq n_t$ be a nonincreasing sequence of positive integers. Let us put $S_j = \bigcup_{l=1}^\infty GR(p^{n_j}, r_l)$ for $1 \leq j \leq t$. Let $\phi_j : S \rightarrow S_j$ be the natural homomorphism followed by the isomorphism $S/p^{n_j}S \cong S_j$ of Proposition 1.2 (II). Let us put $U(S; n_1, n_2, \dots, n_t) = \{(\alpha_{ij}) \in (S)_{t \times t} \mid \alpha_{ij} \in p^{n_j - n_i} S \text{ if } i > j\}$. It is easy to see that $U(S; n_1, n_2, \dots, n_t)$ forms a subring of $(S)_{t \times t}$. Let $M(S; n_1, n_2, \dots, n_t)$ denote the set of all $t \times t$ matrices (a_{ij}) , where $a_{ij} \in S_j$, and $a_{ij} \in p^{n_j - n_i} S_j$ for $i > j$. Let Φ be the mapping of $U(S; n_1, n_2, \dots, n_t)$ onto $M(S; n_1, n_2, \dots, n_t)$ defined by $(\alpha_{ij}) \mapsto (a_{ij})$ where $a_{ij} = \phi_j(\alpha_{ij})$. It is easy to check that addition and multiplication in $M(S; n_1, n_2, \dots, n_t)$ can be defined by stipulating that Φ preserves addition and multiplication. Following [17], we call $M(S; n_1, n_2, \dots, n_t)$ a ring of Szele matrices over S .

LEMMA 1.3. (cf. [17, Lemma 2.1]) Let R be a ring with 1 which contains an IG-subring S of characteristic p^n . If R is finitely generated as a left S -module, then there exists a nonincreasing sequence $n = n_1 \geq n_2 \geq \dots \geq n_t$ of positive integers such that R is isomorphic to a subring of $M(S; n_1, n_2, \dots, n_t)$.

PROOF. By Proposition 1.2 (V), there exists a submodule N of R such that $R = S \oplus N$ as left S -modules. By Proposition 1.2 (III), there are a discrete valuation ring W and a homomorphism ϕ of W onto S . By defining

$$ay = \phi(a)y \quad (a \in W, y \in N),$$

N is a finitely generated W -module. Since a finitely generated module over a principal ideal domain is a direct sum of cyclic modules, there exist $y_1, y_2, \dots, y_s \in N$ such that $N = \bigoplus_{i=1}^s Wy_i$. Let $t = s + 1$, $x_1 = 1$ and $x_i = y_{i-1}$ ($2 \leq i \leq t$). Then we get $R = \bigoplus_{i=1}^t Sx_i$. Let $Sx_i \cong S/p^{n_i}S$ as S -modules ($n_1 = n$). Without loss of generality, we may assume $n_1 \geq n_2 \geq \dots \geq n_t$. For each $a \in R$, we can write

$$x_i a = \sum_{j=1}^t \alpha_{ij} x_j \quad (\alpha_{ij} \in S).$$

Since

$$0 = p^{n_i} x_i a = \sum_{j=1}^n p^{n_i} \alpha_{ij} x_j,$$

by Proposition 1.2 (IV), $\alpha_{ij} \in p^{n_j - n_i} S$ if $i > j$. As α_{ij} is uniquely determined modulo $p^{n_j} S$ by a , we can define $\psi : R \rightarrow M(S; n_1, n_2, \dots, n_t)$ by $a \mapsto (\psi_j(\alpha_{ij}))$. It is easy to see that ψ is an injective ring homomorphism.

2.

Let G be a group, and N a normal subgroup of G . Let $\rho : G \rightarrow H = G/N$ be the natural homomorphism. A monomorphism $\lambda : H \rightarrow G$ will be called a right inverse of ρ if $\rho \circ \lambda = id_H$. If λ is a right inverse of ρ , then G is a semidirect product of N and $\lambda(H)$.

The following lemma is a variation of Schur-Zassenhaus theorem [13, Chapter 9, 9.1.2].

LEMMA 2.1. Let G be a group, and N a normal subgroup of G . Let $\rho : G \rightarrow H = G/N$ be the natural homomorphism. Assume that N is locally finite, and there exists a sequence $\{H_i\}_{i=1}^{\infty}$ of finite subgroups of H such that $H_i \subset H_{i+1}$ ($i \geq 1$), $\bigcup_{i=1}^{\infty} H_i = H$ and, for any $a \in N$ and any $i \geq 1$, $o(a)$ and $|H_i|$ are coprime. Then:

- (I) There exists a right inverse $\lambda : H \rightarrow G$ of ρ .
- (II) If, for some $m \geq 1$, there exists a monomorphism $\mu' : H_m \rightarrow G$ such that $\rho \circ \mu' = id_{H_m}$, then there exists a right inverse $\mu : H \rightarrow G$ of ρ such that $\mu|_{H_m} = \mu'$.
- (III) There exists a unique right inverse of ρ if and only if G is a nilpotent group.
- (IV) If $\mu' : H_m \rightarrow G$ and $\mu'' : H_m \rightarrow G$ are monomorphisms such that $\rho \circ \mu' = \rho \circ \mu'' = id_{H_m}$, then $\mu'(H_m)$ and $\mu''(H_m)$ are conjugate in G .

PROOF. (I) For each $x \in H_1$, we can choose an element g_x of G such that $\rho(g_x) = x$. As G is locally finite (see [13, Chapter 14, 14.3.1]), the subgroup G_1 of G generated by $\{g_x\}_{x \in H_1}$ is finite, and $\rho|_{G_1}$ is a homomorphism of G_1 onto H_1 . Let us put $N_1 = \text{Ker}(\rho|_{G_1})$. Since $|N_1|$ and $|H_1|$ are coprime, by Schur-Zassenhaus theorem [13, Chapter 9, 9.1.2], there exists a right inverse $\lambda_1 : H_1 \rightarrow G_1$ of $\rho|_{G_1}$. Next, let $\{g'_y\}_{y \in H_2}$ be a set of elements of G such that $\rho(g'_y) = y$ for any $y \in H_2$, and $\{g_x\}_{x \in H_1} \subset \{g'_y\}_{y \in H_2}$. Let G_2 be the finite subgroup of G generated by $\{g'_y\}_{y \in H_2}$. Then $\rho|_{G_2}$ is a homomorphism of G_2 onto H_2 . By [13, Chapter 9, 9.1.3], there exists a complement subgroup L of $N_2 = \text{Ker}(\rho|_{G_2})$ in G_2 such that $L \supset \lambda_1(H_1)$. The mapping $\lambda_2 : H_2 \rightarrow G_2$ defined by $H_2 = G_2/N_2 \ni bN_2 \mapsto b$ ($b \in L$) is a right inverse of $\rho|_{G_2}$. For any $a \in H_1$, $\lambda_2(a)^{-1}\lambda_1(a) \in N_2 \cap L = \{1\}$, hence we see $\lambda_2|_{H_1} = \lambda_1$. Continuing this process inductively, we get a sequence $G_1 \subset G_2 \subset \dots$ of finite subgroups of G and a sequence $\{\lambda_i\}_{i=1}^{\infty}$ of right inverses $\lambda_i : H_i \rightarrow G_i$ of $\rho|_{G_i}$ such that $\lambda_j|_{H_i} = \lambda_i$ for any $1 \leq i \leq j$. Then $\lambda = \lim_{i \rightarrow \infty} \lambda_i : H = \bigcup_{i=1}^{\infty} H_i \rightarrow G$ is a right inverse of ρ .

(II) can also be proved in the same way by starting from $\mu' : H_m \rightarrow \mu'(H_m)$.

(III) Assume that $\lambda : H \rightarrow G$ is the unique right inverse of ρ . Then G is a semidirect product of N and $\lambda(H)$. We shall show that this is the direct product. Suppose that there exist $c \in N$ and $z \in H$ such that $c\lambda(z) \neq \lambda(z)c$. Let us define $\mu : H \rightarrow G$ by $\mu(b) = z^{-1}\lambda(b)z$. Then μ is a right inverse of ρ different from λ , which contradicts our hypothesis. So G is the direct product of N and $\lambda(H)$. Hence G is nilpotent.

Conversely, let us suppose that G is nilpotent, and λ and μ are right inverses of ρ . For each $i \geq 1$, let G_i be the subgroup of G generated by $\lambda(H_i) \cup \mu(H_i)$. Then $\rho|_{G_i}$ is a homomorphism of G_i onto H_i . Both $\lambda(H_i)$ and $\mu(H_i)$ are complement subgroups for $N_i = \text{Ker}(\rho|_{G_i})$ in G_i . Since G_i is a finite nilpotent group, for each prime divisor q of $|G_i|$, G_i contains a unique q -Sylow subgroup. Each G_i is the direct product of such Sylow subgroups. As $|H_i|$ and $|N_i|$ are coprime, we have $\lambda(H_i) = \mu(H_i)$. So $\lambda|_{H_i} = \mu|_{H_i}$. Since this holds for each $i \geq 1$, we see $\lambda = \mu$.

(IV) Let L be the finite subgroup of G generated by $\mu'(H_m) \cup \mu''(H_m)$. Then $\rho|_L$ is a homomorphism of L onto H_m . Since $|\text{Ker}(\rho|_L)| = |N \cap L|$ and $|H_m|$ are coprime, by Schur-Zassenhaus theorem, $\mu'(H_m)$ and $\mu''(H_m)$ are conjugate in L .

Let G, N, H and $\rho : G \rightarrow H$ be as in Lemma 2.1. We say that G has property (GC) with respect to N if, for any two right inverses μ and ν of ρ , $\mu(H)$ and $\nu(H)$ are conjugate in G . If H is finite, then by Lemma 2.1 (IV), G has the property (GC) with respect to N .

Let R be a ring with 1. Let S be a subring of R , and $I = J(R) \cap S$. The homomorphism of S/I to $R/J(R)$ defined by $a + I \mapsto a + J(R)$ ($a \in S$) is injective. We shall say that S/I is naturally isomorphic to $R/J(R)$ if this homomorphism is onto. If S is a local subring of a local ring R and if $J(S)$ is nilpotent, then $J(S) = J(R) \cap S$.

Now we shall state main theorems of this section, which generalize the result of R. Raghavendran [9, p. 373, Theorem XIX.4].

THEOREM 2.2. Let R be a local ring with radical M . Assume that M is nilpotent, and $K = R/M$ is a commutative field of characteristic p (p a prime) which is algebraic over $GF(p)$. Then there exists an IG-subring S of R such that S/pS is naturally isomorphic to K .

PROOF. Since K is algebraic over $GF(p)$, $|K|$ is either finite or countably infinite. So there exists a sequence $\{K_i\}_{i=1}^{\infty}$ of finite subfields of K such that $K_i \subset K_{i+1}$ ($i \geq 1$) and $\bigcup_{i=1}^{\infty} K_i = K$. Let $K_i = GR(p^{r_i})$. The natural homomorphism $\pi: R \rightarrow K$ induces a group homomorphism $\pi^* = \pi|_{R^*}$ of R^* onto K^* . Each $(1 + M^i)/(1 + M^{i+1})$ is isomorphic to the additive group M^i/M^{i+1} . As $pM^i \subset M^{i+1}$, the order of each element of $1 + M = \text{Ker } \pi^*$ is a power of p . Furthermore, $K^* = \bigcup_{i=1}^{\infty} K_i^*$, where $|K_i^*| = p^{r_i} - 1$ is coprime to p . So, by Lemma 2.1 (I), there exists a right inverse $\lambda: K^* \rightarrow R^*$ of π^* . For each $i \geq 1$, let α_i be a generator of K_i^* . By Lemma 1.1, the subring $S_i = \langle \lambda(\alpha_i) \rangle$ of R is isomorphic to $GR(p^{r_i}, r_i)$, where p^{r_i} is the characteristic of R . Consequently, $S = \langle \lambda(K^*) \rangle = \bigcup_{i=1}^{\infty} S_i$ is an IG-subring of R , and S/pS is naturally isomorphic to K .

Such a subring S of R stated in Theorem 2.2 will be called a coefficient subring of R . When R is a commutative local ring satisfying the assumption of Theorem 2.2, S coincides with the subring described in [11, p. 106, Theorem 31.1].

Let R, M, S and $K = \bigcup_{i=1}^{\infty} GF(p^{r_i})$ be as in Theorem 2.2, where $\{r_i\}_{i=1}^{\infty}$ is a sequence of positive integers such that $r_i | r_{i+1}$ ($i \geq 1$). Let p^n be the characteristic of R . Let S' be another coefficient subring of R . From what was stated in §1, $S' \cong \bigcup_{i=1}^{\infty} GR(p^{r_i}, r_i)$, which is isomorphic to S . By Proposition 1.2 (V), there exists a left S' -submodule N of R such that $R = S' \oplus N$ as left S' -modules.

If $\lambda: K^* \rightarrow R^*$ is a right inverse of π^* , then by the proof of Theorem 2.3, $S = \langle \lambda(K^*) \rangle$ is a coefficient subring of R .

We shall show that, if λ and μ are different right inverses of π^* , then $\langle \lambda(K^*) \rangle \neq \langle \mu(K^*) \rangle$. Let us suppose $\langle \lambda(K^*) \rangle = \langle \mu(K^*) \rangle$ and denote it by S . Let $\{K_i\}_{i=1}^{\infty}$ be a sequence of finite subfields of K such that $K_i \cong GF(p^{r_i})$, $K_i \subset K_{i+1}$ ($i \geq 1$) and $\bigcup_{i=1}^{\infty} K_i = K$. As $\lambda \neq \mu$, there exist a number $j \geq 1$ and an element α of K_j such that $\lambda(\alpha) \neq \mu(\alpha)$. By Lemma 1.1, both $T = \langle \lambda(K_j^*) \rangle$ and $T' = \langle \mu(K_j^*) \rangle$ are isomorphic to $GR(p^{r_j}, r_j)$. As $S = \bigcup_{i=1}^{\infty} \langle \lambda(K_i^*) \rangle$, there exists a number $l \geq 1$ such that $T \cup T' \subset \langle \lambda(K_l^*) \rangle$. Since $\langle \lambda(K_l^*) \rangle$ is a Galois ring, $T \cong T'$ implies $T = T'$. The restriction $\pi|_{T^*}$ is a homomorphism of T^* onto K_j^* . Both $\lambda|_{K_j^*}$ and $\mu|_{K_j^*}$ are right inverses of $\pi|_{T^*}$, so T^* is the direct product of $\lambda(K_j^*)$ and $\text{Ker}(\pi|_{T^*}) = 1 + pT$, and is also the direct product of $\mu(K_j^*)$ and $1 + pT$. As $|K_j^*|$ and $|1 + pT|$ are coprime, we have $\lambda(K_j^*) = \mu(K_j^*)$. So there exists some $\beta \in K_j^*$ such that $\lambda(\alpha) = \mu(\beta)$. Then $\alpha = \pi^* \circ \lambda(\alpha) = \pi^* \circ \mu(\beta) = \beta$, which means $\lambda(\alpha) = \mu(\alpha)$. This contradicts our choice of α .

By making use of Lemma 2.1 (I), we can easily see that, if S is a coefficient subring of R , there exists a right inverse $\lambda: K^* \rightarrow S^*$ of π^* such that $S = \langle \lambda(K^*) \rangle$.

Summarizing the above, we obtain the following theorem.

THEOREM 2.3. Let R be a local ring with radical M . Assume that M is nilpotent, and $K = R/M$ is a commutative field of characteristic p (p a prime) which is algebraic over $GR(p)$. Let $\pi^*: R^* \rightarrow K^*$

be the group homomorphism induced by the natural ring homomorphism $\pi : R \rightarrow K$. Then:

(I) If S' is a coefficient subring of R , then there exists a S' -submodule N of R such that $R = S' \oplus N$ as left S' -modules.

(II) All coefficient subrings of R are isomorphic.

(III) If $\lambda : K^* \rightarrow R^*$ is a right inverse of π^* , then $S = \langle \lambda(K^*) \rangle$ is a coefficient subring of R . Conversely, if S is a coefficient subring of R , then there exists uniquely a right inverse $\lambda : K^* \rightarrow R^*$ of π^* such that $S = \langle \lambda(K^*) \rangle$.

(IV) All coefficient subrings of R are conjugate in R if and only if R^* has property (GC) with respect to $1 + M$.

With the same notation as in Theorem 2.2, M/M^2 is regarded as a left K -space by the operation

$$\bar{a} \bar{x} = \overline{ax} \quad (\bar{a} \in K = R/M, \bar{x} \in M/M^2).$$

THEOREM 2.4. Let R be a local ring with radical M . Assume that M is nilpotent, and $K = R/M$ is a commutative field of characteristic p (p a prime) which is algebraic over $GF(p)$. Let S be a coefficient subring of R . Then R is finitely generated as a left S -module if and only if M/M^2 is a finite dimensional left K -space. In this case, there exists a finitely generated left S -submodule N of M such that $R = S \oplus N$ as left S -modules, and there exists a nonincreasing sequence $n_1 \geq n_2 \geq \dots \geq n_t$ of positive integers (p^{n_i} is the characteristic of R) such that R is isomorphic to a subring of $M(S; n_1, n_2, \dots, n_t)$.

PROOF. Assume that R is finitely generated as left S -module. Then R is a Noetherian left S -module, since S is a Noetherian ring by Proposition 1.2 (IV). As M is a left S -submodule of R , M is a finitely generated left S -module. This implies that M/M^2 is a finite dimensional left K -space.

Conversely, let us assume that M/M^2 is a finite dimensional left K -space. Let ω be the nilpotency index of M . Let $x_1, x_2, \dots, x_d$ be elements of M whose images modulo M^2 form a K -basis of M/M^2 . As S/pS is naturally isomorphic to K , any element of y of M is written as

$$y = \sum_{i=1}^d a_i x_i + y' \quad (a_i \in S, y' \in M^2).$$

Let

$$z = \sum_{j=1}^d b_j x_j + z' \quad (b_j \in S, z' \in M^2)$$

be another element of M . Then

$$yz = \sum_{i,j=1}^d a_i x_i b_j x_j + w'' \quad (w'' \in M^3).$$

Each $x_i b_j$ is written as

$$x_i b_j = \sum_{k=1}^d c_{kij} x_k + w_{ij}' \quad (c_{kij} \in S, w_{ij}' \in M^2).$$

So we see that any element v' of M^2 can be written as

$$v' = \sum_{i,j=1}^d a_{ij} x_i x_j + v'' \quad (a_{ij} \in S, v'' \in M^3).$$

Continuing in this way, we see that any element of M is written as an S -coefficient linear combination of distinct products of $\omega - 1$ or fewer x_i 's. So M is a finitely generated left S -module. Also $K = R/M$ is a finitely generated left S -module, hence R is a finitely generated left S -module.

Now suppose that R is finitely generated as left S -module. By Theorem 2.3 (I), there exists a finitely generated left S -submodule N' of R such that $R = S \oplus N'$ as left S -modules. By Proposition 1.2 (III), there exist a discrete valuation ring V and a homomorphism ξ of V onto S . Defining $ay = \xi(a)y$ ($a \in V, y \in N'$), we can regard N' as a left V -module. Then there exist $x_1, x_2, \dots, x_t \in N'$

such that $N' = \bigoplus_{i=1}^t Vx_i = \bigoplus_{i=1}^t Sx_i$. By putting $x_0 = 1$, we get $R = \bigoplus_{i=0}^t Sx_i$. Let $c_1, c_2, \dots, c_t$ be elements of S such that $\bar{c}_i = \bar{x}_i$ under the natural homomorphism $\pi: R \rightarrow K$. Let us put $y_0 = 1$ and $y_i = x_i - c_i$ for $1 \leq i \leq t$. Then $y_i \in M$ ($1 \leq i \leq t$) and $R = \bigoplus_{i=0}^t Sx_i = \bigoplus_{i=0}^t Sy_i$. So $N = \bigoplus_{i=1}^t Sy_i$ has the desired property. The last statement is immediate from Lemma 1.3.

3.

Let R be a local ring described in Theorem 2.2. Then R may have more than one coefficient subring. Concerning this subject, first we can state the following.

THEOREM 3.1. Let T be an IG-ring of characteristic p^n different from $GR(p^n, 1)$. Then, for any infinite cardinal number χ , there exists a local ring R such that

- (1) $M = J(R)$ is nilpotent,
- (2) $K = R/M$ is a commutative field of characteristic p (p a prime) which is algebraic over $GF(p)$,
- (3) coefficient subrings of R are isomorphic to T ,
- (4) all coefficient subrings of R are conjugate in R , and
- (5) χ is the number of all coefficient subrings of R .

PROOF. Let $T = \bigcup_{i=1}^{\infty} GR(p^n, r_i)$, where $\{r_i\}_{i=1}^{\infty}$ is a sequence of positive integers such that $r_i \mid r_{i+1}$ ($i \geq 1$). Let $K = T/pT$ and $\pi': T \rightarrow K$ be the natural homomorphism. As K is a proper extension of $GF(p)$, there exists an automorphism $\bar{\sigma}$ of K different from id_K . Let σ be the automorphism of T which induces $\bar{\sigma}$ modulo pT (see Proposition 1.2 (VI)). Let A be a set of cardinality χ , and $V = \bigoplus_{\alpha \in A} T$ be a free T -module. The abelian group $T \oplus V$ together with the multiplication

$$(a, x)(a', x') = (aa', ax' + \sigma(a')x)$$

forms a ring, which we denote by R . Let $\pi: R \rightarrow K$ be the homomorphism defined by $(a, x) \mapsto \pi'(a)$, and $M = \text{Ker } \pi$. As $R/M \cong K$ and $M^{n+1} = 0$, R is a local ring with radical M whose residue field is K . By Theorem 2.3 (III), there exists a one-to-one correspondence between the set of all coefficient subrings of R and the set Y of all right inverses of $\pi^* = \pi|_{R^*}: R^* \rightarrow K^*$.

By the embedding $T \ni a \rightarrow (a, 0) \in R$, T is regarded as a coefficient subring of R . So, by Theorem 2.3 (III), there exists a right inverse $\lambda: K^* \rightarrow R^*$ of π^* such that $\langle \lambda(K^*) \rangle = T$. Since $K = \bigcup_{i=1}^{\infty} GF(p^{r_i})$, there exists a number $j \geq 1$ such that $\bar{\sigma}$ is not the identity on $GF(p^{r_j})$. Let γ be a generator of $GF(p^{r_j})^*$, and $c = \lambda(\gamma)$. It is easy to see that, for any $z \in V$, $R^* \ni h = (c, z)$ is of multiplicative order $p^{r_j} - 1$. So, for each $z \in V$, we can define a group homomorphism

$\mu_z': GF(p^{r_j})^* \rightarrow R^*$ by $\gamma' \mapsto (c, z)^{\gamma'}$. By Lemma 2.1 (II), we can extend μ_z' to $\mu_z \in Y$. If $V \ni z_1, z_2$ and $z_1 \neq z_2$, then $\mu_{z_1} \neq \mu_{z_2}$. So $|Y| \geq |V| = \chi$.

Let S be a coefficient subring of R . We shall show that S is conjugate to T . By Theorem 2.3 (III), there exists a right inverse $\lambda': K^* \rightarrow R^*$ of π^* such that $S = \langle \lambda'(K^*) \rangle$. Let $\lambda'(\gamma) = (c', z)$, where $c' \in T$ and $z \in V$. Let U be the finite subgroup of R^* generated by $\lambda(\gamma)$ and $\lambda'(\gamma)$. As the restriction $\pi|_U$ is a homomorphism of U onto $GF(p^{r_j})^*$, by Schur-Zassenhaus theorem, there exists $(b, w) \in R^*$ ($b \in T, w \in V$) and an integer i such that $\lambda'(\gamma) = (b, w)^{-1} \lambda(\gamma) (b, w)$. Then, $(c', z) = (b, w)^{-1} (c', 0) (b, w)$, which implies $c' = c$. As $\pi'(c') = \pi(\lambda'(c')) = \gamma = \pi(\lambda(\gamma)) = \pi'(c)$, so $c' = c$ and $\lambda'(\gamma) = (c, z)$. Let $x = \{c - \sigma(c)\}^{-1}z$. Suppose that $\alpha \in K$ satisfies $\alpha^m = \gamma$ for some integer m . Let $\lambda(\alpha) = a$. Then, by the same reason as above, we can write $\lambda'(\alpha) = (a, y)$ for some $y \in V$.

As

$$(c, z) = \lambda'(\gamma) = \lambda'(\alpha^m) = (a, y)^m = (a^m, \{a^m - (\sigma(a))^m\} \{a - \sigma(a)\}^{-1} y),$$

we get $c = a^m$ and $z = \{c - \sigma(c)\} \{a - \sigma(a)\}^{-1} y$. So $(1, x)\lambda'(\alpha) = (a, y + \sigma(a)x) = (a, ax) = \lambda(\alpha)(1, x)$.

As K^* is the union of cyclic subgroups generated by such α which contain $GF(p')^*$ (generated by γ), this proves $S = \langle \lambda'(K^*) \rangle = (1, x)^{-1}T(1, x)$. So $|Y|$, the number of all coefficient subrings of R , does not exceed χ . As we have seen $|Y| \geq \chi$, we get $|Y| = \chi$.

Next we shall consider the uniqueness of coefficient subrings.

A finite local ring T is called of type (1) if T is generated by two units a and b such that

- (1) $ab \neq ba$,
- (2) $a - b \in J(T)$, and
- (3) $o(a) = o(b) = |T/J(T)| - 1$.

If T is a finite local ring of type (1), then T^* is not a nilpotent group. For, let us suppose that T is a finite local ring of type (1). Let a and b be generators of T satisfying (1)-(3). Let A and B be cyclic subgroups of T^* generated by a and b respectively. Let $K = T/J(T) = GF(p')$. Then $|A| = |B| = p' - 1$ is coprime to $|J(T)|$. If T^* is nilpotent, then $A = B$, as both A and B are complement subgroups of $1 + J(T)$ in T^* . This contradicts (1), so we see that T^* is not nilpotent.

THEOREM 3.2. Let R be a local ring with radical M . Assume that M is nilpotent, and $K = R/M$ is a commutative field of characteristic p (p a prime) which is algebraic over $GF(p)$. Then the following are equivalent.

- (i) R has a unique coefficient subring.
- (ii) R^* is a nilpotent group.
- (iii) R^* is isomorphic to the direct product of K^* and $1 + M$.
- (iv) R^* has no finite local subring of type (1).

PROOF. (i) $\Leftrightarrow$ (ii). Clear from Lemma 2.1 (III) and Theorem 2.3 (III).

(i) $\Rightarrow$ (iii). Let $\pi^* = \pi|_{R^*} : R^* \rightarrow K^*$ be the group homomorphism induced by the natural homomorphism $\pi : R \rightarrow K$. Since R has a unique coefficient subring, by Theorem 2.3 (III), there exists a unique right inverse λ of π^* . Then R^* is a semidirect product of $1 + M$ and K^* . Let z be any fixed element of $1 + M$. The mapping $\mu : K^* \rightarrow R^*$ defined by $K^* \ni \alpha \mapsto z^{-1}\lambda(\alpha)z$ is a right inverse of π^* , so $\mu = \lambda$ by our hypothesis. This implies that each element of $\lambda(K^*)$ commutes with each element of $1 + M$. Hence R^* is the direct product of $1 + M$ and $\lambda(K^*)$.

(iii) $\Rightarrow$ (iv). Let us suppose that R contains a finite local subring U of type (1). By the proof of [10, Lemma 1], $1 + M$ is a nilpotent group. If R^* is isomorphic to the direct product of K^* and $1 + M$, then R^* is nilpotent. So U^* is nilpotent, which is a contradiction.

(iv) $\Rightarrow$ (i). Assume that R has at least two different coefficient subrings. Then there exist at least two different right inverses λ and μ of π^* . Let $\{K_i\}_{i=1}^\infty$ be a sequence of finite subfields of K such that $K_i \subset K_{i+1}$ and $\bigcup_{i=1}^\infty K_i = K$. There exists a number j such that $\lambda|_{K_j} \neq \mu|_{K_j}$. Let γ be a generator of K_j^* . Then the subring $\langle \lambda(\gamma), \mu(\gamma) \rangle$ of R is a finite local ring of type (1).

4.

From [9, p. 373, Theorem XIX.4 (b)] and the proof of Theorem 3.1, one may expect that, in Theorem 2.3, any two coefficient subrings of R are always conjugate. However, from the following example, we see that this is incorrect.

Let $K = \bigcup_{i=1}^{\infty} GF(p^{r_i})$, where $\{r_i\}_{i=1}^{\infty}$ is a strictly increasing sequence of positive integers such that $r_i \mid r_{i+1}$ ($i \geq 1$). Let $\{\sigma_i\}_{i=1}^{\infty}$ be automorphisms of K such that σ_i is not the identity on $GF(p^{r_i})$ ($i \geq 1$) and, for $j < i$, σ_i is the identity on $GF(p^{r_j})$. Let $V = \bigoplus_{i=1}^{\infty} Kx_i$ be a left K -vector space with basis $\{x_i\}_{i=1}^{\infty}$. We can regard V as a (K, K) -bimodule by defining

$$(\sum_i c_i x_i) a = \sum_i c_i \sigma_i(a) x_i, \quad (\sum_i c_i x_i \in V, a \in K).$$

The abelian group $R = K \oplus V$ together with the multiplication

$$(a, y)(b, z) = (ab, az + yb) \quad (a, b \in K, y, z \in V)$$

forms a local ring with radical $M = (0, V)$, which satisfies the assumption of Theorem 2.3. The homomorphism $\pi: R \rightarrow K$ defined by $(a, x) \mapsto a$ gives the isomorphism $R/M \cong K$. The subring $S = \{(a, 0) \mid a \in K\}$ of R is a coefficient subring of R .

For each $i \geq 1$, let γ_i be a generator of $GF(p^{r_i})^*$. Then we can write $\gamma_i = \gamma_{i+1}^{m_i}$ for a suitable integer m_i . We shall define elements $\{u_i\}_{i=1}^{\infty}$ of R^* inductively as follows: Let $u_1 = (\gamma_1, x_1)$. For $u_n = (\gamma_n, \sum_{j=1}^n r_j x_j)$ ($r_j \in K$), let

$$a_j = \{\gamma_n - \sigma_j(\gamma_n)\}^{-1} \{\gamma_{n+1} - \sigma_j(\gamma_{n+1})\} r_j \quad (1 \leq j \leq n)$$

and

$$u_{n+1} = (\gamma_{n+1}, \sum_{j=1}^n a_j x_j + x_{n+1}).$$

Then it is easy to check that $\phi(u_i) = p^{r_i} - 1$ and $u_i = u_{i+1}^{m_i}$. Let $f_i: GF(p^{r_i})^* \rightarrow R^*$ be defined by $\gamma_i \mapsto u_i$ ($i \in \mathbb{Z}$). Since $f_i|_{GF(p^{r_j})} = f_j$ for $j \leq i$, there exists $f = \lim_{\rightarrow} f_i: K^* \rightarrow R^*$. As f is a right inverse of $\pi^* = \pi|_{R^*}: R^* \rightarrow K^*$, so $S_1 = \langle f(K^*) \rangle$ is a coefficient subring of R .

We shall show that S_1 and S are not conjugate in R . Let us suppose that there exists an element $v = (s, \sum_i d_i x_i) \in R^*$ ($s \in K^*, d_i \in K$) such that $S_1 = v^{-1} S v$. Then, for each $i \geq 1$, there exists some $b_i \in K^*$ such that $f(\gamma_i) = v^{-1}(b_i, 0)v$. Then,

$$\begin{aligned} u_i &= (\gamma_i, \sum_{j=1}^{i-1} r'_j x_j + x_i) \quad (r'_j \in K) \\ &= v^{-1}(b_i, 0)v \\ &= (s^{-1}, -s^{-1}(\sum_i d_i x_i)s^{-1})(b_i, 0)(s, \sum_i d_i x_i) \\ &= (b_i, \sum_{j=1}^i (s^{-1} b_i d_j - s^{-1} d_j \sigma_j(b_i)) x_j), \end{aligned}$$

which yields

$$1 = s^{-1} \{b_i - \sigma_i(b_i)\} d_i.$$

So, for any $i \geq 1$, we see $d_i \neq 0$. This contradicts that $\sum_i d_i x_i$ is an element of the direct sum $V = \bigoplus_{i=1}^{\infty} Kx_i$.

In conclusion, we shall state a theorem which is a generalization of [3, Theorem].

THEOREM 4.1. (cf [9, p. 376, Theorem XIX.5] and [4, p. 491, Theorem 72.19]) Let R be a ring with 1. Assume that $J(R)$ is nilpotent. Let

$$R/J(R) = (K_1)_{n_1 \times n_1} \oplus (K_2)_{n_2 \times n_2} \oplus \dots \oplus (K_d)_{n_d \times n_d},$$

where each K_i ($1 \leq i \leq d$) is a commutative field of characteristic p (p a prime) which is algebraic over $GF(p)$. Then there exists a subring T of R which satisfies the following.

- (i) $R = T \oplus N$ (as abelian groups), where N is an additive subgroup of R .
- (ii) T is isomorphic to a finite direct sum of matrix rings over IG-rings.
- (iii) $J(T) = T \cap J(R) = pT$.
- (iv) T/pT is naturally isomorphic to $R/J(R)$.

Moreover,, if T' is another subring of R satisfying (ii)-(iv), then T' is isomorphic to T .

PROOF. Let $\bar{R} = R/J(R) = \bar{R}\bar{e}_1 \oplus \bar{R}\bar{e}_2 \oplus \dots \oplus \bar{R}\bar{e}_d$, where each $\bar{R}\bar{e}_i (1 \leq i \leq d)$ is a simple component of $\bar{R}$ and $\bar{e}_i$ is a central idempotent of $\bar{R}$. Let $\bar{R}\bar{e}_i = (K_i)_{n_i \times n_i}$, where K_i is a commutative field which is algebraic over $GF(p)$. Let $\pi: R \rightarrow \bar{R}$ be the natural homomorphism. There are mutually orthogonal idempotents $e_1, e_2, \dots, e_d$ of R such that $e_1 + e_2 + \dots + e_d = 1$ and $\pi(e_i) = \bar{e}_i (1 \leq i \leq d)$. Then,

$$R = e_1 R e_1 \oplus e_2 R e_2 \oplus \dots \oplus e_d R e_d \oplus (\oplus_{i \neq j} e_i R e_j)$$

as abelian groups. Since each $e_i R e_i$ is semiperfect and $e_i R e_i / J(e_i R e_i) \cong \bar{R}\bar{e}_i = (K_i)_{n_i \times n_i}$, there exist a local ring S_i and an isomorphism ϕ_i of $e_i R e_i$ onto $(S_i)_{n_i \times n_i}$ (see, for instance, [1, p. 160, Theorem 21]). Let

$$\begin{aligned} \phi &= \phi_1 + \phi_2 + \dots + \phi_d : e_1 R e_1 \oplus e_2 R e_2 \oplus \dots \oplus e_d R e_d \rightarrow \\ A &= (S_1)_{n_1 \times n_1} \oplus (S_2)_{n_2 \times n_2} \oplus \dots \oplus (S_d)_{n_d \times n_d} \end{aligned}$$

be the isomorphism. Since $S_i/J(S_i) \cong K_i$, by Theorem 2.2 and Theorem 2.3 (I), there exist an IG-subring T_i and a left T_i -submodule N_i of S_i such that $S_i = T_i \oplus N_i$ (as abelian groups), and T_i/pT_i is naturally isomorphic to $S_i/J(S_i)$. Then

$$B = (T_1)_{n_1 \times n_1} \oplus (T_2)_{n_2 \times n_2} \oplus \dots \oplus (T_d)_{n_d \times n_d}$$

is a subring of A . Let $T = \phi^{-1}(B)$. As $J(e_i R e_i) \cap \phi^{-1}((T_i)_{n_i \times n_i}) = J(\phi^{-1}((T_i)_{n_i \times n_i}))$, we see $J(t) = T \cap J(R) = pT$ and that T/pT is naturally isomorphic to

$$(e_1 R e_1 \oplus e_2 R e_2 \oplus \dots \oplus e_d R e_d) / J(e_1 R e_1 \oplus e_2 R e_2 \oplus \dots \oplus e_d R e_d) = R/J(R).$$

Let us put

$$N = \phi^{-1}\left\{ (N_1)_{n_1 \times n_1} \oplus (N_2)_{n_2 \times n_2} \oplus \dots \oplus (N_d)_{n_d \times n_d} \right\} \oplus \{ \oplus_{i \neq j} e_i R e_j \}.$$

Then we see $R = T \oplus N$.

Now, let us suppose that T' is a subring of R satisfying (ii)-(iv). Let e and f be primitive idempotents of T' . We claim that $Re \cong Rf$ (as left R -modules) if and only if $T'e \cong T'f$ (as left T' -modules). Let $\pi(e) = \bar{e}$ and $\pi(f) = \bar{f}$. Assume that $Re \cong R'f$. Then $\bar{R}\bar{e} \cong \bar{R}\bar{f}$ as left $\bar{R}$ -modules. Both $\bar{R}\bar{e}$ and $\bar{R}\bar{f}$ are minimal left ideals of $\bar{R}$, so they are contained in the same simple component of $\bar{R}$, which implies that $J(R)$ does not include eRf . Conversely, if $J(R)$ does not include eRf , then $\bar{R}\bar{e} \cong \bar{R}\bar{f}$, which means $Re \cong Rf$ (see, for instance, [1, p. 158, Theorem 16]). Thus we see that $Re \cong Rf$ (as left R -modules) if and only if $J(R)$ does not include eRf . Similarly, $T'e \cong T'f$ (as left T' -modules) if and only if $J(T') = pT'$ does not include $eT'f$. Since T'/pT' is naturally isomorphic to $R/J(R)$, $J(R)$ include eRf if and only if pT' includes $eT'f$. So we see that $Re \cong Rf$ (as left R -modules) if and only if $T'e \cong T'f$ (as left T' -modules).

By making use of matrix units, 1 of R is written in T as

$$1 = (e_{11} + e_{12} + \dots + e_{1n_1}) + (e_{21} + e_{22} + \dots + e_{2n_2}) + \dots + (e_{d1} + e_{d2} + \dots + e_{dn_d}),$$

where e_{ki} are mutually orthogonal primitive idempotents of T , and $Te_{ki} \cong Te_{lj}$ (as left T -modules) if and

only if $k = l$. Similarly,

$$1 = (f_{11} + f_{12} + \dots + f_{1m_1}) + (f_{21} + f_{22} + \dots + f_{2m_2}) + \dots + (f_{d1} + f_{d2} + \dots + f_{dm_d}),$$

where f_{ki} are mutually orthogonal primitive idempotents of T' , and $T'f_{ki} \cong T'f_{li}$ (as left T' -modules) if and only if $k = l$.

As $e_{ki}Te_{ki}/pe_{ki}Te_{ki} \cong e_{ki}Re_{ki}/e_{ki}J(R)e_{ki}$, we see that e_{ki} and f_{li} are primitive idempotents of R . Then $R = \oplus Re_{ki} = \oplus Rf_{li}$ are indecomposable decompositions.

By what was stated above, Krull-Schmidt theorem tells us that there exists a permutation σ of $\{1, 2, \dots, d\}$ such that $n_i = m_{\sigma(i)}$ and $Re_{ik} \cong Rf_{\sigma(i)l}$ as left R -modules ($1 \leq i \leq d, 1 \leq k, l \leq n_i$). By renumbering, we may assume $n_i = m_i$ and $Re_{ik} \cong Rf_{il}$ ($1 \leq i \leq d, 1 \leq k, l \leq n_i$). Now,

$$T \cong (e_{11}Te_{11})_{n_1 \times n_1} \oplus (e_{21}Te_{21})_{n_2 \times n_2} \oplus \dots \oplus (e_{d1}Te_{d1})_{n_d \times n_d}$$

and

$$T' \cong (f_{11}T'f_{11})_{n_1 \times n_1} \oplus (f_{21}T'f_{21})_{n_2 \times n_2} \oplus \dots \oplus (f_{d1}T'f_{d1})_{n_d \times n_d},$$

where $e_{i1}Te_{i1}$ and $f_{i1}T'f_{i1}$ are IG-rings. Hence, to complete the proof it will suffice to show $e_{i1}Te_{i1} \cong f_{i1}T'f_{i1}$.

As $e_{i1}Te_{i1}$ is an IG-ring which is naturally isomorphic to $e_{i1}Re_{i1}/e_{i1}J(R)e_{i1}$, so $e_{i1}Te_{i1}$ is a coefficient subring of $e_{i1}Re_{i1}$. Similarly, $f_{i1}T'f_{i1}$ is a coefficient subring of $f_{i1}Rf_{i1}$. As $e_{i1}Re_{i1} \cong \text{End}({}_R Re_{i1}) \cong \text{End}({}_R Rf_{i1}) \cong f_{i1}Rf_{i1}$, we see $e_{i1}Te_{i1} \cong f_{i1}T'f_{i1}$ by Theorem 2.3 (II).

Note. It is unknown when the subring T of Theorem 4.1 is unique up to inner automorphism of R (see [3, Problem]).

ACKNOWLEDGEMENT. The author would like to express his indebtedness and gratitude to Prof. Y. Hirano and Dr. H. Komatsu for their helpful suggestions and valuable comments.

REFERENCES

- [1] BEHRENS, E. A., *Ring Theory*, Academic Press (1972).
- [2] BOURBAKI, N., *Éléments de Mathématique*, Algèbre, Hermann (1962).
- [3] CLARK, W. E., A coefficient ring for finite noncommutative rings, *Proc. Amer. Math. Soc.* **33** (1972), 25-28.
- [4] CURTIS, C. W. & REINDER, I., *Representation Theory of Finite Groups and Associative Algebras*, Pure & Appl. Math. Ser. 11, Wiley-Interscience (1962).
- [5] HASSE, H., *Zahlentheorie*, 2, Auflage, Akademie-Verlag (Berlin, 1963).
- [6] JANUSZ, G. T., Separable algebras over commutative rings, *Trans. A. M. S.* **122** (1966), 461-479.
- [7] JACOBSON, N., *Lectures in Abstract Algebra*, vol. III, Van Nostrand (1964).
- [8] KRULL, W., Algebraische Theorie der Ringe, I, II and III, *Math. Annalen* **88** (1923), 80-122, **91** (1924), 1-46 and **92** (1924), 183-213.
- [9] MCDONALD, B. R., *Finite Rings with Identity*, Pure & Appl. Math. Ser. 28, Marcel Dekker (1974).
- [10] MOTOSE, K. & TOMINAGA, H., Group rings with nilpotent unit groups, *Math. J. Okayama Univ.* **14** (1969), 43-46.
- [11] NAGATA, M., *Local Rings*, Wiley-Interscience (1962).
- [12] RAGHAVENDRAN, R., Finite associative rings, *Compositio Math.* **21** (1969), 195-229.
- [13] ROBINSON, D. J. S., *A Course in the Theory of Groups*, Springer-Verlag (1982).
- [14] ROQUETTE, P., Abspaltung des Radikals in vollständigen lokalen Ringen, *Abh. Math. Sem. Univ. Hamburg* **23** (1959), 75-113.
- [15] SERRE, J. P., *Local Fields*, Springer-Verlag (1979).

- [16] SUMIYAMA, T., On double homothetisms of rings and local rings with finite residue fields, *Math. J. Okayama Univ.* **33** (1991), 13-20.
- [17] WILSON, R. S., Representations of finite rings, *Pacific J. Math.* **53** (1974), 643-649.